

C57Bl/6N mice have an attenuated lung inflammatory response to dsRNA compared to C57Bl/6J and BALB/c mice

Sofia Malm Tillgren, Juan José Nieto-Fontarigo, Samuel Cerps, Sangeetha Ramu, Mandy Menzel, Irma Mahmutovic Persson, Anja Meissner, Hamid Akbarshahi, Lena Uller

Angaben zur Veröffentlichung / Publication details:

Malm Tillgren, Sofia, Juan José Nieto-Fontarigo, Samuel Cerps, Sangeetha Ramu, Mandy Menzel, Irma Mahmutovic Persson, Anja Meissner, Hamid Akbarshahi, and Lena Uller. 2023. "C57Bl/6N mice have an attenuated lung inflammatory response to dsRNA compared to C57Bl/6J and BALB/c mice." *Journal of Inflammation* 20 (1): 6.
<https://doi.org/10.1186/s12950-023-00331-4>.

RESEARCH

Open Access



C57Bl/6N mice have an attenuated lung inflammatory response to dsRNA compared to C57Bl/6J and BALB/c mice

Sofia Malm Tillgren¹, Juan José Nieto-Fontarigo¹, Samuel Cerps¹, Sangeetha Ramu¹, Mandy Menzel¹, Irma Mahmutovic Persson¹, Anja Meissner^{2,3,4}, Hamid Akbarshahi⁵ and Lena Uller^{1*}

Abstract

Background Lower respiratory infections caused by ssRNA viruses are a major health burden globally. Translational mouse models are a valuable tool for medical research, including research on respiratory viral infections. In in vivo mouse models, synthetic dsRNA can be used as a surrogate for ssRNA virus replication. However, studies investigating how genetic background of mice impacts the murine lung inflammatory response to dsRNA is lacking. Hence, we have compared lung immunological responses of BALB/c, C57Bl/6N and C57Bl/6J mice to synthetic dsRNA.

Methods dsRNA was administered intranasally to BALB/c, C57Bl/6N and C57Bl/6J mice once/day for three consecutive days. Lactate dehydrogenase (LDH) activity, inflammatory cells, and total protein concentration were analyzed in bronchoalveolar lavage fluid (BALF). Pattern recognition receptors levels (TLR3, MDA5 and RIG-I) were measured in lung homogenates using RT-qPCR and western blot. Gene expression of IFN- β , TNF- α , IL-1 β and CXCL1 was assessed in lung homogenates by RT-qPCR. ELISA was used to analyze protein concentrations of CXCL1 and IL-1 β in BALF and lung homogenates.

Results BALB/c and C57Bl/6J mice showed infiltration of neutrophils to the lung, and an increase in total protein concentration and LDH activity in response to dsRNA administration. Only modest increases in these parameters were observed for C57Bl/6N mice. Similarly, dsRNA administration evoked an upregulation of MDA5 and RIG-I gene and protein expression in BALB/c and C57Bl/6J, but not C57Bl/6N, mice. Further, dsRNA provoked an increase in gene expression of TNF- α in BALB/c and C57Bl/6J mice, IL-1 β only in C57Bl/6N mice and CXCL1 exclusively in BALB/c mice. BALF levels of CXCL1 and IL-1 β were increased in BALB/c and C57Bl/6J mice in response to dsRNA, whereas the response of C57Bl/6N was blunt. Overall, inter-strain comparisons of the lung reactivity to dsRNA revealed that BALB/c, followed by C57Bl/6J, had the most pronounced respiratory inflammatory responses, while the responses of C57Bl/6N mice were attenuated.

Conclusions We report clear differences of the lung innate inflammatory response to dsRNA between BALB/c, C57Bl/6J and C57Bl/6N mice. Of particular note, the highlighted differences in the inflammatory response of C57Bl/6J and C57Bl/6N substrains underscore the value of strain selection in mouse models of respiratory viral infections.

Keywords BALB/c, C57Bl/6J, C57Bl/6N, dsRNA, Poly(I:C), Inflammation, Lung, Mouse models, Respiratory viral infection

*Correspondence:

Lena Uller

Lena.uller@med.lu.se

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Human viral respiratory infections and lower respiratory tract infections (LRTI) are a leading cause of mortality and morbidity globally [1]. LRTIs are most commonly caused by single stranded-RNA (ssRNA) viruses including respiratory syncytial virus (RSV), parainfluenza, rhinovirus (RV) and coronavirus [2]. Respiratory infections are also the most predominant triggers of asthma and chronic obstructive pulmonary disease (COPD) exacerbations, accounting for the majority of acute hospitalizations among these patients [3, 4].

As ssRNA viruses replicate inside host cells, double stranded RNA (dsRNA) is formed as an intermediate molecule [5]. dsRNA will then be recognized by pattern recognition receptors (PRRs) including endosomal toll-like receptor 3 (TLR3) and cytosolic receptors melanoma differentiation-associated protein 5 (MDA5) and retinoic acid-inducible gene I (RIG-I), which will initiate and orchestrate innate anti-viral and inflammatory responses [6]. Poly(I:C) is a synthetic dsRNA molecule widely used to mimic the innate inflammatory responses to ssRNA virus. Poly(I:C) has previously been shown to induce inflammation and impair lung function in mice, and serves as a suitable surrogate for ssRNA viral infection [7, 8]. Mouse models using poly(I:C) is therefore valuable when studying early innate responses orchestrated by TLRs.

Mouse models of respiratory viral infections have contributed extensively to knowledge on the pathogenesis and identification of host factors that are crucial for protection against severe viral-induced disease [9–11]. The inbred BALB/c mouse strain is well used to study viral infections, and have contributed to knowledge on the inflammatory response to different RV serotypes, as well as SARS-CoV-2 pathogenesis and vaccine development [9, 12]. Similarly, the C57Bl/6 mouse strain is also frequently used and have been used to investigate inflammatory responses to RSV, as well as mechanisms of RV-induced asthma exacerbations [10, 13]. Previous studies have highlighted that BALB/c and C57Bl/6 strains have separate susceptibility and inflammatory responses to some respiratory viruses. Yasui et al. have demonstrated different cytokine expression patterns of BALB/c and C57Bl/6J mice towards SARS-CoV-2 infection, and Watkiss et al. have reported different susceptibilities to pneumonia virus of mice (PVM) [14, 15].

The C57Bl/6 strain was the first strain to have its genome sequenced and is one of the most frequently used strains in medical research. Although the substrains of C57Bl/6 are genetically similar, differences have been identified, including 34 SNPs and 2 indels that separate the coding sequences of the C57Bl/6N and C57Bl/6J strains [16]. Among these, the C57Bl/6J strain have an

in-frame five-exon deletion of the nicotinamide nucleotide transhydrogenase (nnt), and loss of this protein has been attributed to an increase in inflammation [17]. As a consequence, comparing results from different substrains of C57Bl/6 can be misleading. Unfortunately, inadequate reporting of substrain identity is common. Hence, discrepancies in published results from similar experiments can be found, a possible consequence of the use of different substrains [18, 19].

Reports comparing the immune responses to respiratory viral infections of C57Bl/6N and C57Bl/6J in a single study are lacking. Comparisons of the inflammatory signature of the brain in response to intraperitoneal administration of dsRNA have been done by Warden et al., where they report distinct differences in the immune signature of these two strains [20]. Moreover, Eisfeld et al. have compared the susceptibility and pathology of influenza infection in C57Bl/6J and C57Bl/6N mice [21]. The C57Bl/6N is however a substrain of the C57Bl/6N, and hence, these strains also differ genetically. Furthermore, in some studies of BALB/c and C57Bl/6 comparisons, proper substrain specification is missing, as in the study by Watkiss et al. referenced above [15]. The aim of this study was therefore to compare the pulmonary innate inflammatory responses of C57Bl/6J, C57Bl/6N and BALB/c, three of the most frequently used mouse strains, in one single study.

We chose to use the viral mimic poly(I:C) to enable a wider applicability and to feature early innate responses. By doing this, we aimed to generate a study that can be used as an aid in strain selection for mouse studies involving respiratory ssRNA virus. By highlighting the differences and similarities in the immunological response, we believe that our study can improve consistency of experimental data, and improve translational discovery research.

Methods and materials

Animals

8–10-week-old female C57Bl/6N (Janvier Labs, Le-Genest-Saint-Isle, France), C57Bl/6J (Charles River Laboratories, Sulzfeld, Germany) and BALB/c (Taconic Biosciences, Ejby, Denmark) mice were used in this study. Mice were acclimatized for 1 week prior experiments. Mice were housed and experiments were performed according to the European Parliament and Council Directive 2010/63/EU, the Swedish Animal Welfare Act (Djurskyddslag 1988:534), the Swedish Animal Welfare Ordinance (Djurskydds-förordning 1988:539) and Institutional Animal Care and Use Committee (IACUC) guidelines, and was approved by the Malmö/Lund Animal Experimental Ethics Committee at the Lund District Court in Sweden (approval number M 6436/2017). The

mice were housed in groups of 4–5, with a 12-h light and dark cycle and had free access to food and water at the animal facility at Lund University.

Hundred micrograms (100 µg) polyinosine-polycytidylic acid (poly(I:C); InVivogen, San Diego, USA) dissolved in 25 µL saline, or 25 µL saline (as vehicle control) were administered intranasally (i.n.) for three consecutive days as previously described [22, 23]. Twenty-four hours after last administration mice were euthanized by intraperitoneal injection of 100 µL of 60 mg/ml pentobarbital (Apoteket AB, Stockholm, Sweden) (Fig. 1).

In total, 30 mice were divided into 6 different groups: C57Bl/6N:saline ($n=5$), C57Bl/6N:dsRNA ($n=5$), C57Bl/6J:saline ($n=5$), C57Bl/6J:dsRNA ($n=5$), BALB/c:saline ($n=5$) and BALB/c:dsRNA ($n=5$).

Bronchoalveolar lavage fluid and lung dissection

Bronchoalveolar lavage fluid (BALF) was collected by rinsing lungs with PBS. Following BALF collection, right lung lobes were dissected and snap frozen in liquid nitrogen followed by storage in -80°C . Firstly, a mechanical homogenization of the lung was performed using an OmniPrep Rotor Stator Generator (Omni International, Waterbury, USA). For ELISA, samples were homogenized in PBS with a protease inhibitor cocktail (1 tablet/20 mL PBS) (Roche Diagnostics GmbH, Mannheim, Germany). Samples for western blot were homogenized in lysis buffer (1% TritonX-100, 10 mM Tris-HCl, 50 mM NaCl, 5 mM EDTA, 30 mM sodium pyrophosphate, 50 mM NaF, 0.1 mM Na_3VO_4) with 1% protease and phosphatase inhibitors (Sigma-Aldrich, Stockholm, Sweden). RT-qPCR samples were homogenized in lysis buffer (Macherey–Nagel, Düren, Germany) with 1% β -mercaptoethanol (Sigma-Aldrich, Stockholm, Sweden).

Differential cell count in BALF

BALF was centrifuged at $230 \times g$ 10 min at 4°C , supernatants were collected and cell pellets were resuspended in PBS. Cell counts were determined using a NucleoCounter (Chemometec, Allerød, Denmark) and 50,000 cells were centrifuged (6 min at $450 \times g$) onto microscope slides using cytopsin funnels and a Shandon Cytospin 3 (Marshall Scientific, Hampton, USA). Cytospin slides were stained with Kwik-Diff stain (EpreDia, Kalamazoo, USA) and differential cell count of 400 cells was determined under a light microscope (Olympus CKX53, LRI Olympus, Lund, Sweden).

RNA extraction and gene expression analysis

RNA from homogenized lungs was extracted using the Nucleospin® RNA II kit (Macherey–Nagel, Düren, Germany), and 1 µg total RNA was reversely transcribed using the High-capacity cDNA reverse transcription kit (Applied Biosystems by Thermo Fisher Scientific Baltics UAB, Vilnius, Lithuania). Gene expression analysis was performed by RT-qPCR on an AriaMx qPCR system (Stratagen, La Jolla, USA) with standard cycling parameters. Primers were obtained from Qiagen (Stockholm, Sweden) and PrimerDesign (Southampton, United Kingdom), a full list can be found in Table 1. ΔCt for each gene was calculated by normalization to the house-keeping gene β -actin and $\Delta\Delta\text{Ct}$ was calculated by normalization to the mean ΔCt of each control group (saline).

Total protein and Enzyme-Linked Immunosorbent Assays (ELISA) analysis

Total protein concentration was determined in BALF supernatants and lung homogenates using Pierce BCA assay (Thermo Scientific, Waltham, USA). Concentrations of CXCL1 and IL-1 β were measured using ELISA




Fig. 1 Study design for intranasal (i.n.) dsRNA or saline administration to mice. dsRNA (100 µg) or saline were i.n. administered to C57Bl/6J, C57Bl/6N or BALB/c mice once per day for three consecutive days. At day 3, mice were euthanized following BALF collection and lung tissue dissection for further analysis

Table 1 Primer sequence list

Gene	Vendor	Sequence
TNF- α	PrimerDesign	AGCCAGGAGGGAGAACAGA (forward) CAGTGAGTGAAAGGGACAGAAC (reverse)
RIG-I	PrimerDesign	CGATATTTTGAAAGACTTGGGTACA (forward) ATGGCTCCGTTGTTGAGATTG (reverse)
TLR3	PrimerDesign	AAGTTATTCGCCCTCCTCTTGA (forward) AGATTCTGGATGCTTGTTTGA (reverse)
IFN- β	PrimerDesign	ATGGAAAGATCAACCTCACCTAC (forward) GGATGGCAAAGGCAGTGTA (reverse)
IFN- λ	PrimerDesign	ACCAGGCTCCAGTGGAA (forward) and TTTTTGAAGGCCTGCAGCTCT (reverse)
IL-1 β	Qiagen	Mm_Il1b_2_SG QuantiTect Primer Assay (QT01048355)
MDA5	Qiagen	Mm_lfh1_1_SG QuantiTect Primer Assay (QT00156338)
β -actin	Qiagen	Mm_Actb_1_SG QuantiTect Primer Assay (QT00095242)
CXCL1	Qiagen	Mm_Cxcl1_1_SG QuantiTect Primer Assay (QT00115647)
Viperin	Qiagen	Mm_Rsad2_1_SG QuantiTect Primer Assay (QT00109431)

Table 1. List indicating primer sequence (where available) and accession number for primers used in this work

with kits from R&D systems (Abingon, UK), according to manufacturer's instructions.

Lactate dehydrogenase activity assay

A cytotoxicity Detection kit (Roche, Stockholm, Sweden) was used to measure lactate dehydrogenase (LDH) activity in BALF, according to manufacturer's instructions.

Western blot analysis

Lung homogenates were analyzed for protein expression of PRRs by western blot. Equal amounts of protein were boiled at 95° (5 min) in laemmli buffer (Sigma Aldrich, Stockholm, Sweden), and loaded onto Mini-protean TGX stain free, 4–20% gels (Bio-rad, Stockholm, Sweden) and separated by gel electrophoresis using a Mini protean tetra cell electrophoresis chamber (Bio-Rad). Proteins were transferred to Trans-Blot turbo (Bio-Rad) membranes using the Trans-Blot turbo system (Bio-Rad). Membranes were cut at 50 kDa according to the Spectra Broad Range protein ladder (Thermo Scientific, Waltham, USA) and blocked in EveryBlot blocking buffer (Bio-Rad) for 10 min. Membranes were then incubated at 4 °C over-night with primary anti-mouse rabbit antibodies from Cell Signaling (Danvers, USA): GAPDH (#D16H11), MDA5 (#D74E4), RIG-I (#D14G6) and TLR3 (#D10F10), all at 1:1000 dilution in TBST + 5% BSA (20 mM TRIS-base, 137 mM NaCl, 0.1% Tween 20). Membranes were washed 3 \times 15 min in TBST and incubated 1 h at room temperature with an HRP-conjugated secondary anti-rabbit goat antibody (1:5000 in TBST + 5% BSA)(#09/2029)(Cell Signaling), followed by a second wash of 3 \times 15 min in TBST and 5 min incubation with Clarity Max Western ECL Substrate (Bio-Rad).

Optical density was quantified by a LI-COR odyssey Fc imager (LI-COR, Lincoln, USA) and the Image Studio Lite software (LI-COR), and optical ratios of target protein and GAPDH were calculated for each sample. For each membrane, TLR3 was first detected. Membranes were then stripped with Restore PLUS western blot stripping buffer (Thermo Scientific), blocked with EveryBlot blocking buffer and incubated with primary antibody against MDA5. This procedure was repeated for RIG-I. All samples were then normalized to the mean target/GAPDH ratio for each saline group.

Statistical analysis

All statistical analysis was performed using GraphPad Prism version 9 (GraphPad software, San Diego, USA). Data are presented as mean \pm standard deviation (SD). Multiple comparisons of all groups (strain and treatment) were made using a 2way ANOVA with Tukey's multiple comparisons test. *P*-values of <0.05 were considered statistically significant.

Results

Intranasal administration of dsRNA increases lung inflammation in BALB/c and C57Bl/6J, but not C57Bl/6N mice

To investigate the inflammatory response to dsRNA of BALB/c, C57Bl/6J and C57Bl/6N mice, 100 μ g dsRNA (poly(I:C)) or saline were administered i.n. once per day for three consecutive days (Fig. 1). To assess the degree of inflammation in the lung, cell concentration and composition was determined by an automated cell counter and subsequent differential cell count of BALF cells, respectively. There was a trend towards an increase in the

concentration of cells in response to dsRNA in BALB/c ($p=0.0632$), but not in C57Bl/6J or C57Bl/6N mice (Fig. 2a). A key feature of the lung inflammatory response to viral infections is neutrophil infiltration [24]. Here, differential cell count revealed a significant increase in the percentage of neutrophils in response to dsRNA in BALB/c and C57Bl/6J mice, while only a modest (non-significant) increase in neutrophil percentage was seen in C57Bl/6N (Fig. 2b-c), suggesting a reduced capacity of C57Bl/6N to respond to viral stimuli. The same pattern

was found when determining the total number of neutrophils in the BALF from each strain (Additional file 5b). No changes in lymphocyte percentages or total numbers in any strain (Fig. 2b-c, Additional file 5a) were found.

Total protein concentration and LDH activity in BALF are increased in response to dsRNA in BALB/c and C57Bl/6J mice, but not C57Bl/6N mice

Increases in total protein concentration is a feature of lung inflammation in mice [23]. dsRNA administration




Fig. 2 Effect of i.n. dsRNA administration on BALF cellular infiltration in mice. dsRNA (100 μg) or saline were i.n. administered to C57Bl/6J, C57Bl/6N or BALB/c mice once per day for three consecutive days, and cell concentration (cells/ml) (**A**) and cellular composition (%) (**B**) were determined in BALF. Representative pictures of BALF composition are shown in (**C**) (100X magnification). Orange arrows depict lymphocytes, purple arrows depict neutrophils and green arrows depict macrophages. Scale bars indicate 100 μm. Data are presented as mean ± SD. Differences between groups were tested using a 2way ANOVA with Tukey's multiple comparisons test. P -values of <0.05 are indicated in the graphs, each data point represents one mouse from one experiment ($n=4-5$ per group)

markedly increased total protein concentration about fourfold in both BALB/c and C57Bl/6J mice (Fig. 3a). However, only a modest increase (~twofold) was observed for C57Bl/6N mice, that did not reach statistical significance (Fig. 3a). Indeed, the increases in total protein concentration in response to dsRNA in BALB/c and C57Bl/6J mice were both significantly larger when compared to the dsRNA-induced increase in C57Bl/6N mice (Fig. 3a).

The activity of lactate dehydrogenase (LDH) in BALF is a marker of tissue damage which can be seen in inflammation, and is known to increase in viral-induced asthma exacerbations [25]. Here, dsRNA administration evoked increased LDH activity in BALB/c and C57Bl/6J, but not C57Bl/6N mice (Fig. 3b).

PRR expression is induced in response to dsRNA administration in BALB/c and C57Bl/6J mice, but not C57Bl/6N mice

As part of the innate immune system, PRRs in the lung sense virus components to orchestrate anti-viral and inflammatory responses [6]. Endosomal TLR3 and cytosolic receptors MDA5 and RIG-I sense viral dsRNA, the replication intermediate of ssRNA viruses [5]. Thus, we compared the induction of PRR expression in response to dsRNA in the three mouse strains. TLR3 mRNA was significantly upregulated in response to dsRNA in BALB/c mice (fourfold compared to saline control), but not in the other two strains (Fig. 4a). Furthermore, the induction of TLR3 gene expression in dsRNA-challenged mice was significantly greater in BALB/c compared to C57Bl/6J and C57Bl/6N (Fig. 4a). Similarly, MDA5 and RIG-I mRNA were significantly upregulated (6 and fourfold compared to saline control) in BALB/c mice in response

to dsRNA (Fig. 4b-c). Interestingly, dsRNA administration evoked similar MDA5 and RIG-I responses also in C57Bl/6J (Fig. 4b-c). No significant upregulations in response to dsRNA were observed in C57Bl/6N mice (Fig. 4b-c). Moreover, the dsRNA-induced increase of MDA5 gene expression in BALB/c mice was greater compared to that of C57Bl/6N (Fig. 4b).

On protein level, the TLR3 expression was surprisingly unaffected by dsRNA administration in all strains (Fig. 5a, d-f, Additional files 1 and 2). However, comparable to its mRNA levels, protein expression of MDA5 was upregulated in response to dsRNA in BALB/c and C57Bl/6J mice (20-fold compared to saline control) (Fig. 5b, d-e, Additional files 1 and 3). C57Bl/6N mice showed a small (fourfold compared to saline control), non-significant increase of MDA5 protein in response to dsRNA, and this increase was significantly lower compared to that of BALB/c and C57Bl/6J mice (Fig. 5b, f).

The RIG-I protein expression significantly increased in BALB/c (13-fold compared to saline control) and C57Bl/6J (sevenfold compared to saline control) but not in C57Bl/6N mice after dsRNA administration (Fig. 5c, d-f, Additional file 1 and 4). Here, the dsRNA response in BALB/c was significantly higher compared to both C57Bl/6J and C57Bl/6N mice, and the C57Bl/6J response was in turn significantly larger than the C57Bl/6N (Fig. 5c).

BALB/c and C57Bl/6J mice, but not C57Bl/6N, respond with pro-inflammatory cytokine expression after dsRNA administration

As a result of PRR sensing of dsRNA, production of anti-viral interferons and pro-inflammatory cytokines is initiated in infected cells [6]. Hence, as a measure of the




Fig. 3 Total protein concentration and LDH activity in BALF after i.n. dsRNA administration in mice. Total protein concentration (**A**) and LDH activity (**B**) in BALF were measured after 3 days of i.n. administration of 100 µg dsRNA or saline to BALB/c, C57Bl/6J and C57Bl/6N mice. Data are presented as mean ± SD. Differences between groups were tested using a 2way ANOVA with Tukey's multiple comparisons test. *P*-values of < 0.05 are indicated in the graphs, each data point represents one mouse from one experiment (*n* = 4–5 per group)




Fig. 4 The effect of i.n. dsRNA administration on gene expression of TLR3, MDA5 and RIG-I. dsRNA (100 μ g) or saline were i.n. administered to BALB/c, C57Bl/6J or C57Bl/6N mice once per day for three consecutive days, and gene expression of TLR3 (**A**), MDA5 (**B**), and RIG-I (**C**) was determined by RT-qPCR in lung homogenates. Data are presented as mean \pm SD, as relative to reference gene β -actin and fold change compared to the mean of saline within each strain. Differences between groups were tested using a 2way ANOVA with Tukey's multiple comparisons test. *P*-values of < 0.05 are indicated in the graphs, each data point represents one mouse from one experiment ($n = 5$ per group)

immunological response to dsRNA, we analyzed the gene expression of IFN- β , IFN- λ , the IFN-stimulated gene viperin and pro-inflammatory TNF- α , IL-1 β and CXCL1. Compared to saline controls, dsRNA administration did not evoke any IFN- β or IFN- λ responses, as evidenced by unchanged mRNA expression in all strains (Fig. 6a-b). In contrast, C57Bl/6J mice showed a significant induction of the IFN-stimulated gene viperin in response to dsRNA administration (73-fold compared to saline controls) (Fig. 6c). Furthermore, BALB/c mice responded with a mean 29-fold induction of viperin to dsRNA, however, this increase was not statistically significant ($p = 0.3241$). BALB/c and C57Bl/6J mice responded with increased mRNA levels of TNF- α to dsRNA administration (7 and eightfold compared to saline controls, respectively), where the C57Bl/6J response was also significantly larger compared to the C57Bl/6N response (Fig. 6d).

Surprisingly, C57Bl/6N was the only strain that significantly increased gene expression of IL-1 β in response to dsRNA (fivefold compared to saline control) (Fig. 6e). BALB/c mice also responded with an increased IL-1 β expression (fourfold compared to saline control), but this failed to reach significance ($p = 0.1017$) (Fig. 6e). Moreover, only BALB/c mice responded with significantly elevated levels of CXCL1 mRNA in response to dsRNA (sevenfold compared to saline controls) (Fig. 6f).

To confirm gene expression results, we analyzed the protein expression of CXCL1 and IL-1 β in both BALF and lung homogenates. In BALF, CXCL1 protein expression was significantly increased in response to dsRNA in BALB/c and C57Bl/6J strains (5 and threefold compared to saline control, respectively), with a significantly greater response in BALB/c mice compared to both C57Bl/6J and C57Bl/6N (Fig. 7a). In addition, the protein




Fig. 5 The effect of i.n. dsRNA administration on protein expression of TLR3, MDA5 and RIG-I. dsRNA (100 µg) or saline were i.n. administered to BALB/c, C57Bl/6J or C57Bl/6N mice once per day for three consecutive days and protein expression of PRRs in lung homogenates was analyzed by western blot. Quantification of immunoblots was performed for TLR3 (**A**), MDA5 (**B**) and RIG-I (**C**). Cropped immunoblots are presented for each of the strains; BALB/c (**D**), C57Bl/6J (**E**) and C57Bl/6N (**F**). Data are presented as mean ± SD, as relative to GAPDH and fold change compared to the mean of saline within each strain. Differences between groups were tested using a 2way ANOVA with Tukey's multiple comparisons test. *P*-values of <0.05 are indicated in the graphs, each data point represents one mouse from one experiment (*n* = 5 per group)

levels of CXCL1 after dsRNA administration were significantly higher in C57Bl/6J compared to C57Bl/6N mice (Fig. 7a). In lung homogenates, no significant differences in response to dsRNA administration were detected (Fig. 7b).

IL-1β expression in BALF was significantly increased after dsRNA administration in BALB/c (twofold compared to saline control), and increased (*p* = 0.0864) in C57Bl/6J mice (twofold compared to saline control)

(Fig. 7c). Contrary to its gene expression, dsRNA administration did not increase protein expression of IL-1β in C57Bl/6N mice (Fig. 7c). The dsRNA response of BALB/c was significantly greater compared to both C57Bl/6J and C57Bl/6N, and the C57Bl/6J response was greater than the C57Bl/6N response (Fig. 7c). No significant changes were found in lung homogenates within strains, but the dsRNA response in C57Bl/6J was significantly larger compared to the C57Bl/6N response (Fig. 7d).




Fig. 6 Gene expression of IFN- β , IFN- λ , viperin, TNF- α , IL-1 β and CXCL1 after i.n. dsRNA administration. dsRNA (100 μ g) or saline were i.n. administered to BALB/c, C57Bl/6J or C57Bl/6N mice once per day for three consecutive days and gene expression of IFN- β (**A**), IFN- λ (**B**), viperin (**C**), TNF- α (**D**), IL-1 β (**E**) and CXCL1 (**F**) was determined by RT-qPCR in lung homogenates. Data are presented as mean \pm SD, as relative to the reference gene β -actin and fold change compared to the mean of saline controls within each strain. Differences between groups were tested using a 2way ANOVA with Tukey's multiple comparisons test. *P*-values of <0.05 are indicated in the graphs, each data point represents one mouse from one experiment (*n* = 5 per group)




Fig. 7 Protein expression of IL-1 β and CXCL1 after i.n. dsRNA administration. dsRNA (100 μ g) or saline were i.n. administered to BALB/c, C57Bl/6J or C57Bl/6N mice once per day for three consecutive days, and protein expression of CXCL1 (**A**) and IL-1 β (**C**) in BALF and CXCL1 (**B**) and IL-1 β (**D**) in lung homogenates was determined by ELISA. Data are presented as pg/ml and mean \pm SD. Differences between groups were tested using a 2way ANOVA with Tukey's multiple comparisons test. *P*-values of < 0.05 are indicated in the graphs, each data point represents one mouse from one experiment (*n* = 5 per group)

Discussion

In this study, we compared the respiratory innate immune response to i.n. administration of dsRNA of BALB/c, C57Bl/6J and C57Bl/6N mice. We demonstrate that both BALB/c and C57Bl/6J mice display distinct upregulation of inflammatory markers in response to dsRNA, while the inflammatory response of C56Bl/6N mice is attenuated. In general, comparing the dsRNA response between strains, the BALB/c strain shows the most prominent inflammatory response. These data reveal distinct strain differences of the pulmonary immune response to a viral mimic and underscore the importance of strain selection when designing mouse studies of respiratory viral infections.

Immunological differences between C57Bl/6 and BALB/c mice have been previously reported. In a study by Watkiss et al., the immune responses toward the ssRNA pneumonia virus of mice (PVM) in BALB/c and C57Bl/6 (substrain not specified) mice were compared [15]. Watkiss et al. show that the BALB/c strain responds with higher levels of CXCL8/CXCL1 compared to the C57Bl/6 strain. In addition, they conclude that the immune response of BALB/c mice is both earlier and stronger compared to C57Bl/6 [15]. Our study, showing a larger immune response of BALB/c mice after 3 days of dsRNA compared to both C57Bl/6N and C57Bl/6 J, supports these observations. Furthermore, in a

study by Yasui et al., the immune response to SARS-CoV-2 infection was compared between BALB/c and C57Bl/6J mice. In analyzes of lung homogenates 3 days post infection, BALB/c mice had higher protein levels of IP-10, IL-1 β , TNF- α , M-CSF, IL-4, IL-12, IL-15 and IL-17 [14]. In contrast, in none of our lung homogenate analyzes (protein or mRNA), we were able to detect any differences in cytokine expression between the BALB/c and C57Bl/6J strains after dsRNA administration. The most marked differences we found were in BALF, where BALB/c had a higher concentration of IL-1 β and CXCL1 after dsRNA administration compared to C57Bl/6J. Nonetheless, our overall conclusion agrees with Yasui et al.: BALB/c mice exhibit a more pronounced inflammation compared to C57Bl/6J mice in response to viral stimuli. Finally, C57Bl/6 mice have been shown to have a much weaker immune reactivity of the lung towards allergens and development of asthma-like features compared to BALB/c mice [26]. This, together with a stronger response to viral stimulation, the main trigger of asthma exacerbations, makes BALB/c mice a suitable candidate for developing asthma exacerbation models.

C57Bl/6 mice are widely used in the field of molecular biology and biomedicine due to the extensive availability of commercial genetically modified mice, their easy breeding, robustness and availability of congenic strains. Despite several studies investigating the differences of viral-induced immune responses in BALB/c and C57Bl/6 mice, to our knowledge, this is the first study to compare the dsRNA-induced pulmonary immune response of two substrains of C57Bl/6: the C57Bl/6N and the C57Bl/6J. Our results show significant differences of general inflammation markers in response to dsRNA: the J-strain exhibits higher concentration of total protein, CXCL1 and IL-1 β protein levels in BALF, as well as larger increase in MDA5 and RIG-I protein expression and TNF- α gene expression in lung homogenates compared to the N-strain. A study by Warden et al. describes the inflammatory response of the brain of C57Bl/6J and C57Bl/6N mice at 3 and 24 h post intraperitoneal injection of dsRNA. Similar to our results, at 24 h, the response of C57Bl/6J was more pronounced than that of C57Bl/6N. However, at 3 h post dsRNA administration, inflammatory gene expression was induced to a similar extent in both strains. Thus, they conclude that the innate immune response of C57Bl/6N mice is shorter compared to C57Bl/6J [20]. Therefore, the discrepancies observed in our study could potentially be explained by the kinetics of the inflammatory response, and should be addressed in further studies.

In genetic analyses of the C57Bl/6J and N- mouse strains, several SNPs and structural variants have been identified [16, 27]. Indeed, Bezdek et al. have reported differences in the development of Aldara-induced psoriasiform dermatitis (AIPD) between C57Bl/6J and C57Bl/6N. Aldara is a TLR7 agonist, applied to the skin to induce skin inflammation. In agreement with our findings, Bezdek et al. show that the C57Bl/6J strain developed more severe skin inflammation than C57Bl/6N [28]. Of particular interest for these and our observations is an in-frame five-exon deletion of the nicotinamide nucleotide transhydrogenase (nnt) gene in C57Bl/6J mice, leading to a loss of the nnt protein [29]. Nnt over expression in murine macrophages suppresses reactive oxygen species (ROS) and cytokine release. In addition, transgenic C57Bl/6J mice expressing nnt produce less IL-1 β , IL-6 and CXCL1 and are more vulnerable to *S. pneumoniae* pulmonary infection compared to WT C57Bl/6J mice [17]. To our knowledge, there has been no investigations on the relationship between nnt and PRRs such as TLR7, TLR3, MDA5 and RIG-I, but in the light of our results, this is something that would be interesting to investigate in further studies.

PRRs are sensors of viral components, and orchestrate immune responses. In regards of dsRNA, TLR3 is responsible for the initial recognition and signaling, while cytosolic MDA5 and RIG-I are important for later responses. In a study employing RV infection of human bronchial epithelial cells, TLR3 was constitutively expressed while MDA5 and RIG-I were induced later during infection in response to TLR3 and IFN signaling [30]. In accordance with these observations, although TLR3 mRNA was significantly induced in BALB/c mice, we found no induction of TLR3 protein expression upon dsRNA administration in any of the investigated mouse strains. In contrast, MDA5 and RIG-I mRNA and protein levels were upregulated in response to dsRNA in BALB/c and C57Bl/6J mice. In terms of protein analysis, BALB/c and C57Bl/6J had a significantly larger increase of MDA5 and RIG-I in response to dsRNA compared to the C57Bl/6N strain. Thus, our findings suggest that TLR3 downstream signaling might be restricted in C57Bl/6N mice. Further studies are warranted to identify the underlying cause and the role of TLR3 in the attenuated response of C57Bl/6N mice.

We failed to detect any significant induction of IFN- β and IFN- λ by dsRNA in any strain. However, the IFN-stimulated gene viperin was significantly induced by dsRNA administration in C57Bl/6J mice. This could be explained by the rapid nature of the IFN response. In fact, in the study by Warden et al., the dsRNA-induced IFN- β expression in the

brain was highest at 3 h post dsRNA challenge in C57Bl/6J mice [20]. Thus, the increase in the IFN-stimulated genes viperin, but also MDA5 and RIG-I, are probably secondary effects of IFN expression at earlier time-points.

Both a strength and limitation of the present study is the use of dsRNA as a viral mimic. The strength lies in the reproducibility and uniformity of dsRNA use, as the “pathogen burden” can be better controlled compared to real viral infection. This eliminates confounding factors of viral susceptibility and variation in pathogen load when merely analyzing the immune response of mice. The limitation covers the fact that different viruses evoke different immune responses. For instance, infection of human bronchial epithelial cells with HRV-C has been shown to induce a lower cytokine release of CCL5, MCP-1 and IL-8 compared to infection with RSV [31]. Hence, dsRNA is a suitable viral mimic for some, but perhaps not all ssRNA viruses. As a consequence, it is of importance to be aware of this limitation when selecting mouse strains for future research.

Conclusions

Our study highlights remarkable discrepancies in the pulmonary innate immune responses towards dsRNA of BALB/c, C57Bl/6J and C57Bl/6N mice. Altogether, our results indicate that C57Bl/6N have an attenuated inflammatory response to dsRNA and that BALB/c present the strongest pro-inflammatory response to dsRNA. C57Bl/6J mice present similar features as BALB/c mice, although slightly less pronounced. Especially for C57Bl/6N and C57Bl/6J, one must be aware of the genetic background when comparing studies, as false results might be a consequence of different genetic backgrounds. Similarly, results might be hard to validate if different substrains are used. Overall, we suggest that the present results contribute to important progress in research on respiratory viral infections.

Abbreviations

AIPD	Aldara-induced psoriasiform dermatitis
BALF	Bronchoalveolar lavage fluid
CXCL	C-X-C motif chemokine ligand
dsRNA	Double stranded RNA
IL	Interleukin
IFN	Interferon
LDH	Lactate dehydrogenase
LRTI	Lower respiratory tract infections
MDA5	Melanoma differentiation-associated protein 5
NNT	Nicotinamide nucleotide transhydrogenase
PRR	Pattern recognition receptor
RIG-I	Retinoic acid-inducible gene I
ROS	Reactive oxygen species
RSV	Respiratory syncytial virus
RV	Rhinovirus
ssRNA	Single stranded RNA
TLR	Toll-like receptor
TNF	Tumor necrosis factor

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12950-023-00331-4>.

Additional file 1: Additional figure 1. Uncropped immunoblots of GAPDH blots. Chemiluminescence channel of C57Bl/6J blot (A), BALB/c blot (C) and C57Bl/6N blot (E). Chemiluminescence and 700 nm channel merge of C57Bl/6J blot (B), BALB/c blot (D) and C57Bl/6N (F), to visualize ladder. Ladder size is indicated.

Additional file 2: Additional figure 2. Uncropped immunoblots of TLR3 blots. Chemiluminescence channel of C57Bl/6J blot (A), BALB/c blot (C) and C57Bl/6N blot (E). Chemiluminescence and 700 nm channel merge of C57Bl/6J blot (B), BALB/c blot (D) and C57Bl/6N (F), to visualize ladder. Ladder size is indicated.

Additional file 3: Additional figure 3. Uncropped immunoblots of MDA5 blots. Chemiluminescence channel of C57Bl/6J blot (A), BALB/c blot (C) and C57Bl/6N blot (E). Chemiluminescence and 700 nm channel merge of C57Bl/6J blot (B), BALB/c blot (D) and C57Bl/6N (F), to visualize ladder. Ladder size is indicated.

Additional file 4: Additional figure 4. Uncropped immunoblots of RIG-I blots. Chemiluminescence channel of C57Bl/6J blot (A), BALB/c blot (C) and C57Bl/6N blot (E). Chemiluminescence and 700 nm channel merge of C57Bl/6J blot (B), BALB/c blot (D) and C57Bl/6N (F), to visualize ladder. Ladder size is indicated.

Additional file 5: Additional figure 5. Effect of i.n. dsRNA administration on BALF cellular infiltration in mice. dsRNA (100 µg) or saline were i.n. administered to C57Bl/6J, C57Bl/6N or BALB/c mice once per day for three consecutive days, and the total number of lymphocytes (A), neutrophils (B) and macrophages (C) were determined in BALF using cytopsin, differential cell counting and cell concentration measurements. Data are presented as mean ± SD. Differences between groups were tested using a 2way ANOVA with Tukey's multiple comparisons test. *P*-values of <0.05 are indicated in the graphs, each data point represents one mouse from one experiment (*n* = 4-5).

Acknowledgements

Not applicable.

Authors' contributions

ST and JJNF contributed with conception, design, acquisition, analysis, interpretation of data and have drafted and revised the work. SC, SR and AM contributed with conception, design, acquisition, analysis, interpretation of data and have revised the work. IMP, MM, HA and LU contributed with design, interpretation of data and have revised the work. All authors have approved the submitted version and have agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work, even ones in which the author was not personally involved, are appropriately investigated, resolved, and the resolution documented in the literature.

Funding

Open access funding provided by Lund University. This research was funded by Vetenskapsrådet (Swedish medical research council) grant numbers: 2020-00922_VR and 2017-00806_VR as well as Swedish Heart and lung foundation grant number: 20180207_HLF.

Availability of data and materials

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations

Ethics approval and consent to participate

This study was performed in accordance with the European Parliament and Council Directive 2010/63/EU, the Swedish Animal Welfare Act (Djurskyddslag 1988:534), the Swedish Animal Welfare Ordinance (Djurskyddsförordning

1988:539) and Institutional Animal Care and Use Committee (IACUC) guidelines, and approved the Malmö/Lund Animal Experimental Ethics Committee at the Lund District Court in Sweden (approval number M 6436/2017).

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Experimental Medical Science, Unit of Respiratory immunopharmacology, Lund University, Lund, Sweden. ²Wallenberg Center for Molecular Medicine, Lund University, Lund, Sweden. ³Department of Physiology, Institute of Theoretical Medicine, Medical Faculty, University of Augsburg, Augsburg, Germany. ⁴Department of Experimental Medical Science, Lund University, Lund, Sweden. ⁵Department of Clinical Sciences, Division of Respiratory Medicine and Allergology, Lund University, Lund, Sweden.

Received: 15 September 2022 Accepted: 26 January 2023

Published online: 21 February 2023

References

- Collaborators GBDLRI. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory infections in 195 countries, 1990–2016 a systematic analysis for the global burden of disease study 2016. *Lancet Infect Dis*. 2018;18(11):1191–210.
- Clark TW, Medina M-j, Batham S, Curran MD, Parmar S, Nicholson KG. Adults hospitalised with acute respiratory illness rarely have detectable bacteria in the absence of COPD or pneumonia; viral infection predominates in a large prospective UK sample. *J Infect*. 2014;69(5):507–15.
- Nicholson KG, Kent J, Ireland DC. Respiratory viruses and exacerbations of asthma in adults. *BMJ*. 1993;307(6910):982–6.
- Bouquet J, Tabor DE, Silver JS, Nair V, Tovchigrechko A, Griffin MP, et al. Microbial burden and viral exacerbations in a longitudinal multicenter COPD cohort. *Respir Res*. 2020;21(1):77.
- Rampersad S, Tennant P. Replication and Expression Strategies of Viruses. *Viruses*. 2018;55–82. <https://doi.org/10.1016/B978-0-12-811257-1.00003-6>. Epub 2018 Mar 30.
- Li D, Wu M. Pattern recognition receptors in health and diseases. *Signal Transduct Target Ther*. 2021;6(1):291.
- Stowell NC, Seideman J, Raymond HA, Smalley KA, Lamb RJ, Egenolf DD, et al. Long-term activation of TLR3 by Poly(I:C) induces inflammation and impairs lung function in mice. *Respir Res*. 2009;10(1):43.
- Mahmutovic-Persson I, Akbarshahi H, Bartlett NW, Glanville N, Johnston SL, Brandelius A, et al. Inhaled dsRNA and rhinovirus evoke neutrophilic exacerbation and lung expression of thymic stromal lymphopoietin in allergic mice with established experimental asthma. *Allergy*. 2014;69(3):348–58.
- Rajput C, Han M, Ishikawa T, Lei J, Goldsmith AM, Jazaeri S, Stroupe CC, Bentley JK, Hershenson MB. Rhinovirus C Infection Induces Type 2 Innate Lymphoid Cell Expansion and Eosinophilic Airway Inflammation. *Front Immunol*. 2021;12:649520.
- Bhoj VG, Sun Q, Bhoj EJ, Somers C, Chen X, Torres J-P, et al. MAVS and MyD88 are essential for innate immunity but not cytotoxic T lymphocyte response against respiratory syncytial virus. *Proc Natl Acad Sci U S A*. 2008;105(37):14046–51.
- Hu Z, Zhang Y, Wang Z, Wang J, Tong Q, Wang M, et al. Mouse-adapted H9N2 avian influenza virus causes systemic infection in mice. *Virology J*. 2019;16(1):135.
- Gu H, Chen Q, Yang G, He L, Fan H, Deng Y-Q, et al. Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine efficacy. *Sci*. 2020;369(6511):1603–7.
- Han M, Chung Y, Young Hong J, Rajput C, Lei J, Hinde JL, et al. Toll-like receptor 2-expressing macrophages are required and sufficient for rhinovirus-induced airway inflammation. *J Allergy Clin Immunol*. 2016;138(6):1619–30.
- Yasui F, Matsumoto Y, Yamamoto N, Sanada T, Honda T, Munakata T, et al. Infection with the SARS-CoV-2 B. 1. 351 variant is lethal in aged BALB/c mice. *Sci Rep*. 2022;12(1):4150.
- Watkiss ERT, Shrivastava P, Arsic N, Gomis S, van Drunen Littel-van den Hurk S. . Innate and adaptive immune response to pneumonia virus of mice in a resistant and a susceptible mouse strain. *Viruses*. 2013;5(1):295–320.
- Simon MM, Greenaway S, White JK, Fuchs H, Gailus-Durner V, Wells S, et al. A comparative phenotypic and genomic analysis of C57BL/6J and C57BL/6N mouse strains. *Genome Biol*. 2013;14(7):R82.
- Ripoll VM, Meadows NA, Bangert M, Lee AW, Kadioglu A, Cox RD. Nicotinamide nucleotide transhydrogenase (NNT) acts as a novel modulator of macrophage inflammatory responses. *Faseb J*. 2012;26(8):3550–62.
- Rochlitz S, Hoymann HG, Müller M, Braun A. No exacerbation but impaired anti-viral mechanisms in a rhinovirus-chronic allergic asthma mouse model. *Clin Sci (Lond)*. 2014;126(1):55–65.
- Han M, Bentley JK, Rajput C, Lei J, Ishikawa T, Jarman CR, et al. Inflammasome activation is required for human rhinovirus-induced airway inflammation in naive and allergen-sensitized mice. *Mucosal Immunol*. 2019;12(4):958–68.
- Warden AS, DaCosta A, Mason S, Blednov YA, Mayfield RD, Harris RA. Inbred Substrain Differences Influence Neuroimmune Response and Drinking Behavior. *Alcohol Clin Exp Res*. 2020;44(9):1760–8.
- Eisfeld AJ, Gasper DJ, Suresh M, Kawaoka Y. C57BL/6J and C57BL/6NJ Mice are Differentially Susceptible to Inflammation-Associated Disease Caused by Influenza A Virus. *Front Microbiol*. 2019;9:3307.
- Akbarshahi H, Menzel M, Ramu S, Mahmutovic Persson I, Björner L, Uller L. House dust mite impairs antiviral response in asthma exacerbation models through its effects on TLR3. *Allergy*. 2018;73(5):1053–63.
- Mahmutovic Persson I, Akbarshahi H, Menzel M, Brandelius A, Uller L. Increased expression of upstream TH2-cytokines in a mouse model of viral-induced asthma exacerbation. *J Transl Med*. 2016;14:52.
- Johansson C, Kirsebom FCM. Neutrophils in respiratory viral infections. *Mucosal Immunol*. 2021;14(4):815–27.
- Wark PAB, Johnston SL, Moric I, Simpson JL, Hensley MJ, Gibson PG. Neutrophil degranulation and cell lysis is associated with clinical severity in virus-induced asthma. *Eur Respir J*. 2002;19(1):68.
- Herz U, Renz H, Wiedermann U. Animal models of type I allergy using recombinant allergens. *Methods*. 2004;32(3):271–80.
- Zurita E, Chagoyen M, Cantero M, Alonso R, González-Neira A, López-Jiménez A, et al. Genetic polymorphisms among C57BL/6 mouse inbred strains. *Transgenic Res*. 2011;20(3):481–9.
- Bezdek S, Hdnah A, Sezin T, Mousavi S, Zillikens D, Ibrahim S, et al. The genetic difference between C57BL/6J and C57BL/6N mice significantly impacts Aldara™-induced psoriasiform dermatitis. *Exp Dermatol*. 2017;26(4):349–51.
- Freeman HC, Huggill A, Dear NT, Ashcroft FM, Cox RD. Deletion of nicotinamide nucleotide transhydrogenase: a new quantitative trait locus accounting for glucose intolerance in C57BL/6J mice. *Diabetes*. 2006;55(7):2153–6.
- Slater L, Bartlett NW, Haas JJ, Zhu J, Message SD, Walton RP, et al. Co-ordinated role of TLR3, RIG-I and MDA5 in the innate response to rhinovirus in bronchial epithelium. *PLoS Pathog*. 2010;6(11):e1001178-e.
- Yuan XH, Pang LH, Yang J, Jin Y. Comparison of immune response to human rhinovirus C and respiratory syncytial virus in highly differentiated human airway epithelial cells. *Virology J*. 2022;19(1):81.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.