

## Compact quotients of homogeneous negatively curved riemannian manifolds

Ernst Heintze

### Angaben zur Veröffentlichung / Publication details:

Heintze, Ernst. 1974. "Compact quotients of homogeneous negatively curved riemannian manifolds." *Mathematische Zeitschrift* 140 (1): 79–80. <https://doi.org/10.1007/bf01218648>.

### Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

**Deutsches Urheberrecht**

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



# Compact Quotients of Homogeneous Negatively Curved Riemannian Manifolds<sup>★</sup>

Ernst Heintze

## 1. Introduction

As it is well known (Borel [2]), a symmetric space of negative curvature admits compact quotients, also called compact Clifford-Klein forms. It is the purpose of this note to prove:

**Theorem.** *Let  $M$  be a connected homogeneous riemannian manifold of strictly negative curvature. If  $M$  admits a quotient of finite volume, then  $M$  is symmetric.*

This may be reformulated as follows:

**Theorem'.** *A locally homogeneous riemannian manifold of strictly negative curvature and finite volume is locally symmetric.*

The proof of the Theorem is a consequence of two results due to Eberlein-O'Neill [4] and Chen [3].

## 2. Preliminaries

Let  $H$  be a simply connected, complete  $n$ -dimensional riemannian manifold of curvature  $K \leq 0$  (Hadamard manifold). Let  $\bar{H}$  denote its naturally defined compactification (Klingenberg [6], Eberlein-O'Neill [4]), which is homeomorphic to the closed unit ball of  $\mathbb{R}^n$ . The group of isometries  $I(H)$  of  $H$  acts as a group of homeomorphisms on  $\bar{H}$ , extending its action on  $H$ . If  $G$  is a subgroup of  $I(H)$  denote its limit set by  $L(G)$ . This is defined by  $L(G) = \bar{G}(p) \cap H(\infty)$  where  $p$  is an arbitrary point in  $H$ ,  $H(\infty) = \bar{H} \setminus H$  are the points at infinity and  $\bar{G}(p)$  is the closure in  $\bar{H}$  of the orbit  $G(p)$ . The limit set is well defined, i.e. independent of  $p$ .

We will use the following two results:

**Theorem** (Eberlein-O'Neill). *Let  $M$  be a connected, simply connected complete riemannian manifold of curvature  $K \leq c < 0$  and  $\Gamma$  a group of isometries acting freely and properly discontinuously on  $M$ . Assume  $\Gamma$  has a fixed point  $x$  in  $M(\infty)$ . Then either:*

- (i)  $x$  is the only fixed point of  $\Gamma$  and  $L(\Gamma) = \{x\}$  or
- (ii)  $\Gamma$  has exactly two fixed points  $x$  and  $y$  and  $L(\Gamma) = \{x, y\}$ .

This is a combination of Proposition 8.9P and 8.9A in [4] observing that  $\Gamma$  can have at most two fixed points. Actually both propositions are stated there under a slightly weaker curvature assumption.

<sup>★</sup> This work was done under support of the "Deutsche Forschungsgemeinschaft" and the Sonderforschungsbereich "Theoretische Mathematik" at Bonn University.

**Theorem** (Chen [3]). *Let  $M$  be a connected, simply connected complete riemannian manifold with curvature  $K \leq c < 0$ . Suppose  $G$  is a connected subgroup of the group of isometries with  $L(G) = M(\infty)$ . If  $G$  does not have a common fixed point in  $M(\infty)$ , then  $G$  is semisimple.*

Chen's proof is based strongly on his Theorem 3.1 [3], whose proof seems to contain a gap at the end. But this may be filled using Lemma 4.7 of Eberlein-O'Neill [4].

### 3. Proof of the Theorem

Let  $M$  be a connected homogeneous riemannian manifold of strictly negative curvature. Let  $\bar{M} = M/\Gamma$  be a quotient of finite volume, where  $\Gamma$  is a group of isometries acting freely and properly discontinuously on  $M$ . We may assume  $\Gamma \subset I_0(M)$ , since  $I(M)$  has only finitely many components ( $M$  is homogeneous, thus  $I(M) = K \cdot I_0(M)$ , where  $K$  is a (compact) isotropy group). If  $I_0(M)$  is semisimple, then  $M$  is known to be symmetric. If  $I_0(M)$  is not semisimple, then it has a fixed point in the boundary by Chen's Theorem and the same holds for  $\Gamma$ . Thus  $L(\Gamma)$  contains at most two elements. On the other hand it is well known that  $L(\Gamma) = M(\infty)$ , if  $M/\Gamma$  has finite volume; a contradiction.

### 4. Remark

With results of the author [5] and Chen's Theorem it is possible to determine the group of isometries of a homogeneous manifold of strictly negative curvature  $M$ . If  $M$  is not symmetric, then  $I_0(M) = K \rtimes_\sigma G$ , i.e.  $I(M)$  is the semidirect product of a compact subgroup  $K$  and a normal solvable subgroup  $G$ , which is simply transitive on  $M$ . Furthermore  $G$  and hence  $I_0(M)$  are not unimodular. This implies again—by a result of Siegel [6]—that  $M$  has no quotients of finite volume. Recently Azencott and Wilson [1] announced that they have generalized this result to a class of homogeneous manifolds of non-positive curvature. Therefore we don't want to give the proof of the above statement.

### References

1. Azencott, R., Wilson, R.: Variétés homogènes à courbure négative. C.r. Acad. Sci., Paris **278**, 561–562 (1974)
2. Borel, A.: Compact Clifford-Klein forms of symmetric spaces. Topology **2**, 111–122 (1963)
3. Chen, S.S.: Complete homogeneous riemannian manifolds of negative curvature. To appear in Commentarii math. Helvet.
4. Eberlein, P., O'Neill, P.: Visibility manifolds. Pacific J. Math. **46**, 45–109 (1973)
5. Heintze, E.: On homogeneous manifolds of negative curvature. To appear in Math. Ann.
6. Klingenberg, W.: Geodätischer Fluß auf Mannigfaltigkeiten vom hyperbolischen Typ. Inventiones Math. **14**, 63–82 (1971)
7. Siegel, C.: Discontinuous groups. Ann. of Math. II. Ser. **44**, 674–689 (1943)

Ernst Heintze  
Mathematisches Institut der Universität Bonn  
D-5300 Bonn  
Wegelerstr. 10  
Federal Republic of Germany

(Received August 13, 1974)