

Stochastic resetting of active Brownian particles with Lorentz force

Iman Abdoli, Abhinav Sharma

Angaben zur Veröffentlichung / Publication details:

Abdoli, Iman, and Abhinav Sharma. 2021. "Stochastic resetting of active Brownian particles with Lorentz force." *Soft Matter* 17 (5): 1307–16.
<https://doi.org/10.1039/d0sm01773f>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



Stochastic resetting of active Brownian particles with Lorentz force

Iman Abdoli ^a and Abhinav Sharma *^{ab}

Equilibrium properties of a system of passive diffusing particles in an external magnetic field are unaffected by Lorentz force. In contrast, active Brownian particles exhibit steady-state phenomena that depend on both the strength and the polarity of the applied magnetic field. The intriguing effects of the Lorentz force, however, can only be observed when out-of-equilibrium density gradients are maintained in the system. To this end, we use the method of stochastic resetting on active Brownian particles in two dimensions by resetting them to the line $x = 0$ at a constant rate and periodicity in the y direction. Under stochastic resetting, an active system settles into a nontrivial stationary state which is characterized by an inhomogeneous density distribution, polarization and bulk fluxes perpendicular to the density gradients. We show that whereas for a uniform magnetic field the properties of the stationary state of the active system can be obtained from its passive counterpart, novel features emerge in the case of an inhomogeneous magnetic field which have no counterpart in passive systems. In particular, there exists an activity-dependent threshold rate such that for smaller resetting rates, the density distribution of active particles becomes non-monotonic. We also study the mean first-passage time to the x axis and find a surprising result: it takes an active particle more time to reach the target from any given point for the case when the magnetic field increases away from the axis. The theoretical predictions are validated using Brownian dynamics simulations.

1 Introduction

A fundamental feature of active Brownian particles (ABPs) is self-propulsion which requires a continual consumption of energy from the local environment.^{1–7} Since ABPs are internally driven, they do not require breaking the spatial symmetry to exist in an out-of-equilibrium state. An ABP is generally modelled as a particle which propels itself along a direction which randomizes *via* rotational diffusion. Given the simplicity of the model, it is not surprising that ABPs serve as a minimalistic model to study the effect of broken time-reversal symmetry and nonequilibrium steady states in general.^{8–14} The interest in ABPs is not purely theoretical as evident in the vast body of research in pharmaceutical and medical applications.^{15–22} Since an ABP adjusts its propulsion speed in response to the local fuel concentration,^{19,23} it is also used as a simple model to understand the emergence of chemotaxis in proto-forms of life.^{24–27}

Recently, the behaviour of diffusion systems subjected to an external magnetic field has attracted considerable interest.^{28–39} It has been shown that Lorentz force due to an external magnetic field induces additional Lorentz fluxes in diffusion systems

which are perpendicular to the typical diffusive fluxes.^{33,35} The Lorentz force generates dynamics which are different from those of a purely diffusive system. Interestingly, the unusual properties due to the Lorentz force persist in the small-mass limit in which the dynamics are overdamped. However, the equilibrium properties, as expected from the Bohr–van Leeuwen theorem,⁴⁰ are unaffected by the applied magnetic field due to no performance of work on the particle. Since the Lorentz force only influences the dynamics, there are essentially two conditions to observe its unusual effects: (i) the system is out of equilibrium and (ii) there are density gradients in the system. This has been recently demonstrated in a system of ABPs with a uniform activity subjected to an inhomogeneous magnetic field,³⁶ which satisfies the aforementioned conditions even in the stationary state. The nonequilibrium steady state in such a system is characterized by density inhomogeneity and bulk fluxes.

In order to ensure that there exists a nontrivial stationary state in a system of ABPs, one requires either a confining potential or periodic boundary conditions.³⁶ In recent years, stochastic resetting has emerged as a powerful framework which gives rise to nontrivial stationary states in diffusive systems characterized by a non-Gaussian probability distribution and steady-state fluxes.^{41–53} Stochastic resetting is unique in the sense that it renews the underlying process and therefore, in some sense, preserves the dynamics of the underlying process in the steady state. With the recent experimental demonstrations,^{54,55}

^a Leibniz-Institut für Polymerforschung Dresden, Institut Theorie der Polymere, 01069 Dresden, Germany. E-mail: sharma@ipfdd.de

^b Technische Universität Dresden, Institut für Theoretische Physik, 01069 Dresden, Germany

stochastic resetting is now no longer a pure theoretical pursuit but rather an alternative and practical method to drive and maintain a system out of equilibrium. We have recently shown that stochastically resetting a passive particle to the origin in the presence of Lorentz force gives rise to a novel stationary state which bears the unusual dynamical properties owing to the magnetic field.³⁷ While the stochastic resetting of passive particles has been thoroughly studied, much less is done about active particles.^{46,56,57} In a very recent work, the motion of an ABP under different resetting protocols has been studied⁵⁸ with a focus on the steady-state density distribution.

In the present work, we investigate the motion of a charged ABP under resetting and the effect of Lorentz force. The particle is stochastically reset to the line $x = 0$ at a constant rate. In addition, the system is periodic in the y direction. We start with a generalized coarse-grained Fokker–Planck equation and analytically determine the density, flux, and polarization, first for a uniform magnetic field and then for a spatially inhomogeneous magnetic field. We show that whereas for a uniform magnetic field the properties of the stationary state of the active system can be obtained from its passive counterpart, novel features emerge in the case of an inhomogeneous magnetic field which have no counterpart in passive systems. In particular, there exists an activity-dependent threshold rate such that for smaller resetting rates, the density distribution of active particles becomes non-monotonic. We also study the Mean First-Passage Time (MFPT) to the x axis and find a surprising result: it takes an active particle more time to reach the target from any given point for the case when the magnetic field increases away from the axis.

The paper continues as follows. In Section 2, we define the model and provide a description of the methods used to analyze the system. In Section 3, we study the system in the presence of a constant magnetic field. We then consider a space-dependent magnetic field and derive expressions for the density, flux, and polarization in the system in Section 4. In Section 5, we obtain the MFPT for the active and passive systems. The conclusion of the paper is presented in Section 6.

2 Model and theory

We consider a single self-propelled, charged Brownian particle of mass m and charge q subjected to an external magnetic field $\vec{B}(\vec{r})$ of strength $B(\vec{r})$, whose direction is along the z axis where the Lorentz force does not influence the motion of the particle. As a consequence, we study the dynamics of the particle in the xy plane with $\vec{r} = (x, y)$. The particle is stochastically reset to the line $x = 0$ at a constant rate μ . The generalized Fokker–Planck equation for the probability density of finding the particle at position \vec{r} with orientation $\vec{u} = (u_x, u_y)$ at time t given that the particle started its motion at the origin, $P(\vec{r}, \vec{u}; t)$ is given as

$$\begin{aligned} \frac{\partial}{\partial t} P(\vec{r}, \vec{u}; t) = & \nabla \cdot [\Gamma^{-1}(\vec{r}) \cdot (D_t \nabla - v_0 \vec{u}) P(\vec{r}, \vec{u}; t)] \\ & + D_r \vec{\mathcal{R}}^2 P(\vec{r}, \vec{u}; t) + \Phi_l + \Phi_g, \end{aligned} \quad (1)$$

where $\nabla = (\partial_x, \partial_y)$ and

$$\Phi_l = -\mu P(\vec{r}, \vec{u}; t), \quad (2)$$

is the loss of the probability from the position \vec{r} due to resetting while

$$\Phi_g = \mu \delta(x) \int P(x', y, u_x, u_y; t) dx', \quad (3)$$

is the gain of the probability at the point $(0, y)$ on the x -axis. Although eqn (1) is not of the form of a continuity equation, the total probability is conserved. Here $v_0 = f/\gamma$ is the self-propulsion speed where γ is the friction coefficient and f is the magnitude of the self-propulsion force that drives the particle into the direction of its (unit) orientation vector \vec{u} . In addition, $\vec{\mathcal{R}} = \vec{u} \times \nabla_{\vec{u}}$ is the rotation operator, $D_t = k_B T / \gamma$ with k_B being the Boltzmann constant that is the translational diffusion coefficient and D_r is the rotational diffusion coefficient. The matrix Γ is defined as $\mathbb{I} + \kappa(\vec{r}) \mathbb{M}$ where \mathbb{I} is the identity matrix, the dimensionless parameter $\kappa(\vec{r}) = qB(\vec{r})/\gamma$ quantifies the strength of the Lorentz force relative to the frictional force and \mathbb{M} is the matrix with elements $M_{ij} = -\epsilon_{ijk} n_k$ where ϵ_{ijk} is the antisymmetric Levi–Civita symbol in two dimensions and n_k is the k component of the unit vector \vec{n} along which the magnetic field is pointed. The inverse of Γ reads

$$\Gamma^{-1}(\vec{r}) = \mathbb{I} - \frac{\kappa(\vec{r})}{1 + \kappa^2(\vec{r})} \mathbb{M} + \frac{\kappa(\vec{r})^2}{1 + \kappa^2(\vec{r})} \mathbb{M}^2. \quad (4)$$

Note that the orientation of the particle remains unchanged under resetting; the particle restarts its motion with the orientation that it had at the time of resetting.

We also perform Brownian dynamics simulations to validate our theoretical predictions. The dynamics of the particle can be described by the following Langevin equations

$$\frac{m}{\gamma} \dot{\vec{v}}(t) = \Gamma(\vec{r}) \cdot \vec{v} + v_0 \vec{u}(t) + \vec{\xi}(t), \quad (5)$$

$$\dot{\vec{u}}(t) = \vec{v}(t), \quad \text{and} \quad \dot{\vec{u}}(t) = \vec{\eta}(t) \times \vec{u}(t), \quad (6)$$

where the dot over the vectors denotes the time derivative and the stochastic forces $\vec{\xi}(t)$ and $\vec{\eta}(t)$ satisfy the properties of Gaussian white noise with a zero mean value and correlation functions $\langle \vec{\xi}(t) \vec{\xi}^T(t') \rangle = 2D_t \mathbb{I} \delta(t - t')$ and $\langle \vec{\eta}(t) \vec{\eta}^T(t') \rangle = 2D_r \mathbb{I} \delta(t - t')$. As the resetting mechanism we consider Poisson distribution for the resetting time which gives the probability of the number of resets to the line $x = 0$ in a small interval of time with a constant rate μ (see Fig. 1). The Fokker–Planck equation in (1) is the small-mass limit of the Langevin equations in (5) and (6) which corresponds to overdamped dynamics of the system. We numerically integrate the set of equations in (5) and (6) with a small mass $m = 0.002$ and the integration time step $dt = 10^{-6} \tau$ where $\tau = \gamma/k_B T$ is the time the particle takes to diffuse over a unit distance. We also fix $k_B = \gamma = 1.0$, the self-propulsion force $f = 10.0$, and $D_r = 20.0$. The particle starts its motion at the origin with the initial velocity $(v_{0x}, v_{0y}) = (1.0, 1.0)$ and initial orientation $(u_{0x}, u_{0y}) = (1.0, 0.0)$. The choice of the parameters holds throughout the paper.

The Fokker–Planck equation in (1) provides a full statistical description of the position and orientation of an ABP under




Fig. 1 Schematic of a charged active Brownian particle which is stochastically reset to the line $x = 0$ at a constant rate μ . The self-propulsion velocity is shown by an arrow inside the disc. Between any two consecutive resetting events the particle undergoes Brownian motion and self-propulsion. Immediately after a resetting event, the orientation of the ABP remains unchanged. The system is subjected to an external magnetic field, $B(\vec{r})$ in the z direction.

stochastic resetting. However, it is a formidable task to obtain an exact solution of this equation. To theoretically analyze the system we make the following assumptions: (1) $D_r \gg \mu$ and (2) the gradients in the system are small on the length scale of persistence length of the ABP. Under these assumptions, one can integrate out the orientational degrees of freedom by a gradient expansion (see Appendix A for details) to yield an equation for the (marginal) probability density as a function of time and position degrees of freedom alone.³⁶ The final equation for the coarse-grained density reads as

$$\frac{\partial \rho(\vec{r}; t)}{\partial t} = \nabla \cdot [\Gamma^{-1}(\vec{r}) \cdot (D_t \nabla + v_0 \pi(\vec{r}; t)) \rho(\vec{r}; t)] + \phi_l + \phi_g, \quad (7)$$

where

$$\phi_l = -\mu \rho(\vec{r}; t), \quad (8)$$

$$\phi_g = \mu \delta(x) \int \rho(x', y; t) dx', \quad (9)$$

are the loss and gain of probabilities and $\rho(\vec{r}; t)$ is the (marginal) probability density of finding the particle at position \vec{r} at time t given that the particle started its motion at the origin. The first and second terms in the square brackets are the (negative) probability fluxes stemming from the thermal fluctuations and activity, respectively. The polarization, $\pi(\vec{r}; t)$, defined as the average orientation per particle, is given as

$$\pi(\vec{r}; t) = -\frac{l_p}{2\rho(\vec{r}; t)} \nabla \cdot [\Gamma^{-1}(\vec{r}) \rho(\vec{r}; t)], \quad (10)$$

where $l_p = v_0/(D_r + \mu)$ denotes the modified persistence length of the ABP. An alternative approach to the above derivation is to treat activity as a perturbation and use the linear response theory for ABPs as outlined in ref. 59 and 60. The gradient expansion approach, in contrast, does not require the activity to be small but only that the gradients be small compared to the persistence length of the ABP. It therefore allows one to consider an active system in which the activity dominates over thermal fluctuations.

By resetting the particles to the x axis we confine the motion of the particles in the x direction. Now we impose periodicity in the y direction over a length scale of L which results in the normalisation condition $\int_{-\infty}^{\infty} dx \int_0^L dy \rho(x, y; t) = 1$. Since the motion is translationally invariant in the y direction the above normalisation condition can be written as $L \int_{-\infty}^{\infty} g(x; t) = 1$ where $g(x; t)$ is the probability density of finding the particle at position x at time t given that its initial position was at $x = 0$. The Fokker-Planck equation for $g(x; t)$ reads

$$\frac{\partial g(x; t)}{\partial t} = -\nabla \cdot [\vec{j}(x; t) + \vec{j}^a(x; t)] - \mu g(x; t) + \frac{\mu}{L} \delta(x), \quad (11)$$

where the flux due to thermal fluctuations is

$$\vec{j}(x; t) = -D_t \Gamma^{-1}(x) \nabla g(x; t), \quad (12a)$$

and

$$\vec{j}^a(x; t) = -v_0 \Gamma^{-1}(x) \vec{p}(x; t) g(x; t). \quad (12b)$$

is the flux due to activity where $\nabla g(x; t) = (\partial_x g(x; t), 0)^T$, and

$$\vec{p}(x; t) = -\frac{l_p}{2g(x; t)} \nabla \cdot [\Gamma^{-1}(x) g(x; t)], \quad (13)$$

is the polarization. Note that since there is no variation in the y direction all the derivatives with respect to y are zero resulting in the reduction of ∇ in eqn (11) and (13) to simply the derivative with respect to x .

To highlight the new features which emerge in the system of ABPs we make a comparison between the active system and its passive counterpart. As the active system we consider the motion of the particle purely due to the activity by ignoring the thermal term (*i.e.*, $D_t = 0$) in eqn (5), (6) and (12a). We compare the active system with the passive one wherein the motion of the particle is due to the thermal fluctuations. The governing Langevin equations and the corresponding Fokker-Planck equation of the passive system can be obtained by setting the self-propulsion velocity, v_0 to zero in eqn (5), (6) and (12b).

3 Uniform magnetic field

We first consider the system subjected to a uniform magnetic field $\kappa(x) \equiv \kappa$. For the active system, the stationary probability density, denoted by $g^a(x)$, can be easily obtained by plugging eqn (12b) into eqn (11) and setting $\partial_t g(x; t) = 0$. The solution can be written as

$$g^a(x) = \frac{\alpha}{2L} \exp(-\alpha|x|), \quad (14)$$

where $\alpha = \sqrt{1 + \kappa^2} \alpha_a$ with $\alpha_a = \sqrt{\mu/D_a}$ and $D_a = v_0^2/2(D_r + \mu)$ being the modified active diffusivity. The stationary solution in (14) is the same as that of the passive system wherein D_a is replaced by D_t .³⁷

The polarization can be obtained by plugging eqn (14) in eqn (13), which in the x direction can be written as

$$p_x(x) = \frac{l_p \alpha_a}{2(1 + \kappa^2)} \text{sign}(x), \quad (15a)$$




Fig. 2 Density, flux in the x direction, and orientations in the x and y directions are shown in (a) to (d), respectively. An ABP is subjected to a constant magnetic field such that $\kappa = 3.0$ and is stochastically reset to the line $x = 0$ at the rate $\mu = 1.0$ with $L = 10.0$. Despite the translational invariant motion in the y direction the magnetic field induces polarization in the y direction. However, the y component of the stationary flux is zero due to the cancellation of fluxes arising from the polarization in the x and y directions. The solid lines show the analytical solutions from eqn (14)–(16) and the circles depict the results from Brownian dynamics simulations.

and in spite of the translational invariance in the y direction there exists polarization which is given as

$$p_y(x) = \kappa p_x(x), \quad (15b)$$

However, the substitution of the polarization into eqn (12b) gives zero fluxes in the y direction and

$$j_x^a(x) = \frac{\mu}{2L} \text{sign}(x) \exp(-\alpha|x|), \quad (16)$$

in the x direction where $\text{sign}(\cdot)$ denotes the sign function. Note that the stationary polarization and fluxes in the y direction are zero in the absence of the magnetic field.

In Fig. 2(a–d) we show the density, stationary flux and the x and y components of the orientation, respectively. Note that despite the translational invariant motion in the y direction there exists polarization in this direction. However, the y component of the stationary flux is zero due to the cancellation of fluxes arising from the polarization in the x and y directions.

4 Inhomogeneous magnetic field

In this section, we show that novel features emerge in the case of an inhomogeneous magnetic field which have no counterpart in passive systems. We consider a system subjected to an exponentially varying magnetic field such that $\kappa(x) = \sqrt{e^{2\lambda|x|} - 1}$ where λ is a constant. With this choice of the magnetic field, the Fokker–Planck equation in (11) can be solved exactly. The stationary probability density is given as

$$g^a(x) = \frac{\alpha_a}{2L} \exp\left[\frac{\lambda|x|}{2} - \frac{2\alpha_a}{\lambda} \left(\exp\left(\frac{\lambda|x|}{2}\right) - 1\right)\right]. \quad (17)$$

Using eqn (14), the polarization in the x direction is given as

$$p_x(x) = \frac{l_p \text{sign}(x)}{2} \left[\frac{\lambda}{2} \exp(-\lambda|x|) + \alpha_a \exp\left(\frac{-\lambda|x|}{2}\right) \right], \quad (18a)$$

and similarly

$$p_y(x) = \frac{l_p \lambda \text{sign}(x)}{4\sqrt{\exp(\lambda|x|) - 1}} \left[\frac{4\alpha_a}{\lambda} \sinh\left(\frac{\lambda|x|}{2}\right) - \exp(-\lambda|x|) \right], \quad (18b)$$

is the polarization in the y direction. The x and y components of the stationary flux can be obtained using eqn (12b), which read

$$j_x^a(x) = \alpha_a D_a \text{sign}(x) \exp\left(\frac{-\lambda|x|}{2}\right) g^a(x), \quad (19a)$$

$$j_y^a(x) = \frac{\lambda D_a \exp(-\lambda|x|)}{2\sqrt{\exp(\lambda|x|) - 1}} \text{sign}(x) g^a(x). \quad (19b)$$

Note that the stationary polarization and stationary flux in the y direction cease to exist in the absence of the magnetic field.

We also consider a passive system under resetting and subjected to the same magnetic field as in the active system. The governing Fokker–Planck equation for the system can be easily written by setting $v_0 = 0$ in eqn (12b) and substituting eqn (12a) into eqn (11). The stationary solution, $g^p(x)$, of the resulting equation is

$$g^p(x) = \frac{\alpha_p}{2LK_0 \left(\frac{2\alpha_p}{\lambda}\right)} \exp\left(\frac{\lambda|x|}{2}\right) K_1\left(\frac{2\alpha_p}{\lambda} \exp\left(\frac{\lambda|x|}{2}\right)\right), \quad (20)$$

where $\alpha_p = \sqrt{\mu/D_t}$ and K_0 and K_1 are the modified Bessel functions of the second kind of order 0 and 1, respectively. The x component of the stationary flux can be written as

$$j_x^p(x) = \frac{D\alpha_p^2}{2LK_0 \left(\frac{2\alpha_p}{\lambda}\right)} \text{sign}(x) K_0\left(\frac{2\alpha_p}{\lambda} \exp\left(\frac{\lambda|x|}{2}\right)\right), \quad (21)$$

and similarly

$$j_y^p(x) = -\sqrt{e^{2\lambda|x|} - 1} j_x^p(x). \quad (22)$$

is the flux in the y direction.

Fig. 3(a) and (b) show the dynamical density distribution in the active and passive systems without resetting, respectively. The density distribution in the active system evolves from a delta function at time $t = 0$ to a non-monotonic distribution at later times. In contrast, the density distribution in the passive system decays monotonically at all times. The non-monotonic behaviour can be qualitatively understood as follows: It has been shown that one of the effects of a spatially varying magnetic field on active particles is similar to subjecting them to an inhomogeneous activity field, for which it is well known that (i) ABPs exhibit a net polarization anti-parallel to the activity gradient and (ii) the steady-state density distribution scales inversely with the active force.³⁶ For an increasing magnetic field away from the origin, the polarization flux takes ABPs away from the x axis where due to the large magnetic field, they accumulate. This gives rise to non-monotonic density distribution, shown in Fig. 3(a). Since there are no polarization effects in passive particles, the density distribution is always monotonically




Fig. 3 Density distributions in (a) the active system and (b) the passive system without resetting are shown at different times. The systems are subjected to a spatially inhomogeneous magnetic field such that $\kappa(x) = \sqrt{e^{\lambda|x|} - 1}$ with $\lambda = 2.0$ and $L = 10.0$. For the passive system, the translational diffusivity, D_t has the same value as the active diffusivity, D_a . The density distribution in the active system evolves, according to eqn (11) by setting $D_t = 0$, from a delta function at time $t = 0$ to a non-monotonic distribution at later times. For the passive system the distribution evolves according to eqn (11) by setting $v_0 = 0$. In contrast to the active system the density distribution decays monotonically at all times. By stochastically resetting the particles one renews these dynamics on the time scale of resetting for which we show density distribution in (c) the active system and (d) the passive system for different values of μ . While for the passive system the accumulation of particles is in a vicinity of $x = 0$, in the active system it is non-monotonic with local maxima at $x = \pm(2/\lambda)\ln(\lambda/2\alpha_a)$ for $\mu < \lambda^2 v_0^2 / (8D_t)$. The lines show the theoretical results from eqn (17) and (20) and the symbols depict simulation results.

decaying. By stochastically resetting the particles one renews these dynamics on the time scale of resetting for which we show the probability density in Fig. 3(c) for the active system and in Fig. 3(d) for the passive system. Clearly the steady-state distributions carry the signature of the underlying dynamics. For the active system there exists an activity-dependent threshold rate such that for smaller resetting rates, the density distribution of the particles becomes non-monotonic. Below this threshold rate, $\mu < \lambda^2 v_0^2 / (8D_t)$, the ABPs accumulate in a vicinity of positions given by $x = \pm(2/\lambda)\ln(\lambda/2\alpha_a)$.




Fig. 4 The x and y components of the flux and orientation are shown in (a) to (d), respectively. An active particle is stochastically reset to the line $x = 0$ at the rate $\mu = 2.0$ with $L = 10.0$. The particle is subjected to the magnetic field $\kappa(x) = \sqrt{e^{\lambda|x|} - 1}$ with $\lambda = 2.0$. The solid lines show the analytical solutions from eqn (18a) to eqn (19b) and the circles depict the results from Brownian dynamics simulations. Note that the polarization and flux in the y direction cease to exist in the absence of the magnetic field.




Fig. 5 Flux in the x and y directions is shown in (a) and (b), respectively. A passive particle is stochastically reset to the line $x = 0$ at the rate $\mu = 2.0$ with $L = 10.0$. The particle is subjected to the magnetic field $\kappa(x) = \sqrt{e^{\lambda|x|} - 1}$ with $\lambda = 2.0$. The translational diffusivity, D_t , has the same value as the active diffusivity, D_a . The solid lines show the analytical solutions from eqn (21) and (22) and the circles depict the results from Brownian dynamics simulations.

In Fig. 4 we use eqn (18a) to eqn (19b) to plot the fluxes and the polarization in the active system. While in the case of a constant magnetic field there is no flux in the y direction, inhomogeneity in the magnetic field gives rise to orientation which results in fluxes in the x and y directions. Note that the polarization and flux in the y direction cease to exist in the absence of the magnetic field. Fig. 5 shows the x and y components of the stationary flux in the passive system. As can be seen from eqn (21) and (22), the theoretical results are in good agreement with the simulation results.

Transport properties of Brownian particles have been usually studied by considering systems which are restricted within the confines of structured and inhomogeneous environments. While in many cases, such structured environments can be viewed as confined channels with different boundaries and properties,^{61–65} directed transport can be obtained *via* spatial control of activity.^{66,67} Here we show that the Lorentz force can result in directed transport with no need for structured geometries. We consider an ABP under resetting, subjected to the magnetic field $\kappa(x) = \sqrt{e^{-\lambda x} - 1}$ if $x < 0$ and $\kappa(x) = -\sqrt{e^{\lambda x} - 1}$ otherwise. The system is invariant to translation in the y direction. However, for a better visualisation of the directed transport, we show the flux and density in the system in two dimensions. Fig. 6(a) depicts a vector plot of the stationary flux in the system which clearly shows the particle transport along the y axis. In Fig. 6(b) we show a surface plot of the stationary probability density in which the arrows show the direction of the particle transport.

5 Mean first-passage time

We now study the first-passage properties of the system in the case of a fixed target at the origin. The searching particle is subjected to the magnetic field $\kappa(x) = \sqrt{e^{\lambda|x|} - 1}$ with $\lambda = 2.0$ and is stochastically reset to its initial position x_0 to be fixed. The backward Fokker-Planck equation for the survival probability, $G(x;t)$ – the probability that the searching particle




Fig. 6 A spatial control of Lorentz force (or self-propulsion speed) can direct transport with no need for structured geometries. (a) A vector plot of the stationary flux whose direction is shown by the arrows and the magnitude is color coded and (b) a surface plot of the stationary probability density to which the flux is attached. An ABP under resetting to the line $x = 0$ at the rate $\mu = 1.0$ is subjected to the magnetic field such that $\kappa(x) = \sqrt{e^{-\lambda x} - 1}$ if $x < 0$ and $\kappa(x) = -\sqrt{e^{\lambda x} - 1}$ otherwise with $\lambda = 2.0$. The system is invariant to translation in the y direction. However, for a better visualisation of the directed transport, we show the figures in two dimensions.

starting at x at $t = 0$ has not reached the target in time t – can be written as

$$\frac{\partial G(x; t)}{\partial t} = A(x) \frac{\partial^2 G(x; t)}{\partial x^2} + B(x) \frac{\partial G(x; t)}{\partial x} - \mu G(x; t) + \mu G(x_0; t), \quad (23)$$

where the initial and boundary conditions are $G(x; 0) = 1$ and $G(0; t) = 0$, respectively. While the coefficients $A(x)$ and $B(x)$ for the active system are $D_a e^{-\lambda x}$ and $-D_a \lambda e^{-\lambda x}/2$, those for the passive one are $D e^{-\lambda x}$ and $-D \lambda e^{-\lambda x}$, respectively. We first solve eqn (23) and then set x to x_0 to find the MFPT (see Appendix B for details). The Laplace transform of the backward Fokker-Planck equation in (23) reads

$$B(x) \frac{\partial^2 \tilde{G}(x; s)}{\partial x^2} + A(x) \frac{\partial \tilde{G}(x; s)}{\partial x} - (\mu + s) \tilde{G}(x; s) = -1 - \mu \tilde{G}(x_0; s), \quad (24)$$

where $\tilde{G}(x; s) = \int_0^\infty dt e^{-st} G(x; t)$ is the Laplace transform of the survival probability. Solving eqn (24) and setting $x = x_0$ we obtain the expressions for the survival probability for the active and passive systems in the Laplace space, which when evaluated at $s = 0$ gives the MFPT as

$$T^a(x_0) = \frac{1}{\mu} \left[\exp \left(\frac{2\alpha_a}{\lambda} \left(\exp \left(\frac{\lambda x_0}{2} \right) - 1 \right) \right) - 1 \right], \quad (25)$$

for the active system, and

$$T^p(x_0) = \frac{1}{\mu} \left[\frac{K_1 \left(\frac{2\alpha_p}{\lambda} \right) \exp \left(-\frac{\lambda x_0}{2} \right)}{K_1 \left(\frac{2\alpha_p}{\lambda} \right) \exp \left(\frac{\lambda x_0}{2} \right)} - 1 \right], \quad (26)$$

for the passive system. Note that the MFPTs of the systems




Fig. 7 Mean first-passage time in the active and passive systems is shown in red and blue, respectively. The systems are subjected to the magnetic field such that $\kappa(x) = \sqrt{e^{2|x|} - 1}$ with $\lambda = 2.0$. The initial position of the particles is located at $x_0 = 2$. For the passive system, the translational diffusivity, D_i has the same value as the active diffusivity, D_a . The solid lines show the theoretical predictions from eqn (25) and (26) and the symbols depict the results from Brownian dynamics simulations. The inset shows the optimal resetting rate with respect to the initial position x_0 . The numerical solution of eqn (25) is compared with the simulation results.

diverge as $\mu \rightarrow 0$ or $\mu \rightarrow \infty$. This implies that there exists an optimal rate at which the MFPT becomes minimum.

In Fig. 7 we show the MFPT with respect to the stochastic rate for the active and passive systems. We compare the results from the theory, given by eqn (25) and (26) and those from Brownian dynamics simulations. It is clear that there exists an optimal resetting rate, μ^* that minimizes the time for the searcher to reach the target. The optimal resetting rate decreases exponentially with increasing starting point x_0 due to inhomogeneity in the magnetic field. The inset shows how the optimal resetting rate varies with increasing initial position of the particle in the active system. The optimal resetting rate is obtained by numerically solving eqn (25) and verified by the simulation results.




Fig. 8 The ratio of the MFPT of the active particle to the passive one for different values of μ . The systems are subjected to the magnetic field such that $\kappa(x) = \sqrt{e^{2\lambda|x|} - 1}$ with $\lambda = 2.0$. For the passive system, the translational diffusivity, D_t has the same value as the active diffusivity, D_a . The solid lines show the theoretical predictions from eqn (25) and (26) and the symbols depict the results from Brownian dynamics simulations. For a fixed target at the origin, the active particle is slower than the passive one. The relative slowness increases as $x_0 \rightarrow 0$ or $x_0 \rightarrow \infty$. It implies that there exists a position, x_0^* where if the particles start from, they reach the target with a minimum time difference. In the limit of large x_0 the MFPT for the active system to find the target is exponentially longer than the passive one and scales as $\sim e^{\lambda x_0/4}$, which is shown by dashed lines. The inset depicts the simulation results of the ratio of the MFPTs for a searcher whose initial position is the origin and the target is fixed at x_0 . In this case, either an active or passive searcher can be faster. There is also a point at which if the particles started from, they would have the same MFPT which occurs in the case of a constant magnetic field (e.g. $\lambda = 0$) as well.

Fig. 8 shows the ratio of the MFPT of the active system to its passive counterpart. Interestingly, the active particle is slower than its passive counterpart to reach the target. The relative slowness increases as $x_0 \rightarrow 0$ or $x_0 \rightarrow \infty$. It implies that there exists a position, x_0^* where if the particles start from, they reach the target with a minimum time difference. In the limit of large x_0 the MFPT for the active system to find the target is exponentially longer than the passive one and scales as $\sim e^{\lambda x_0/4}$, which is shown by dashed lines. The inset depicts the simulation results of the ratio of the MFPTs for the searcher starting at the origin and the target is set at x_0 . In this case, either an active or passive searcher can be faster. There is also a point at which if the particles started from, they would have the same MFPT which occurs in the case of a constant magnetic field (e.g. $\lambda = 0$) as well.

6 Discussion and conclusion

In this paper, we studied the motion of a charged ABP under resetting and the effect of Lorentz force. We showed that whereas for a uniform magnetic field the properties of the stationary state of the active system can be obtained from its passive counterpart, novel features emerge in the case of an inhomogeneous magnetic field which have no counterpart in passive systems. In particular, there exists an activity-dependent threshold rate such that for smaller resetting rates, the density distribution of active particles becomes non-monotonic. Moreover, somewhat counter intuitively, it may take an active particle much longer to reach a fixed target than its passive counterpart in an

inhomogeneous magnetic field. We also showed that the Lorentz force can result in directed transport with no need for structured geometries.

The regime in which the rotational relaxation occurs slowly in comparison to the resetting time scale can be a problem of interest. In this regime, where $D_r < \mu$, the effects of the persistent motion of ABPs would be more pronounced. However, as long as one only resets the position of the particle without changing the orientation, we do not expect any new features to emerge in addition to those already reported in the paper. For instance, even in this new regime, the density distribution will show a peak away from the x axis for sufficiently small resetting rates. This would occur simply due to the persistence of the ABPs which takes them away from the x axis following a resetting event. More drastic signatures of activity would be seen if the orientation of the ABP is also reset together with its position. This has been recently investigated in ref. 58 in the regime of $D_r < \mu$ where the density distribution can be highly anisotropic depending on the chosen protocol.

We would like to emphasize that the choice of the magnetic field is motivated by the mathematical convenience, which allows us to theoretically analyse the system. The qualitative behaviour of the system will remain unaffected by other choices of the magnetic field. We note that an ABP in an inhomogeneous activity field and subjected to a constant magnetic field will give rise to the same phenomenology as presented in this study. A possible experimental realization is to reset the particle in a rotating frame of reference using optical tweezers. By rotating the reference frame one can induce a Coriolis force which acts similarly as the Lorentz force arising from an external magnetic field.⁶⁸ From a future perspective, it would be interesting to investigate the effect of stochastic resetting on inertial ABPs.^{12,69}

Conflicts of interest

There are no conflicts to declare.

A Elimination of orientational degrees of freedom

We use a gradient-expansion approach to integrate the orientational degrees of freedom from the probability density. To do this, we recall the Fokker-Planck equation for the probability density, $P(\vec{r}, \vec{u}; t)$ as

$$\begin{aligned} \frac{\partial}{\partial t} P(\vec{r}, \vec{u}; t) = & \nabla \cdot [\Gamma^{-1}(\vec{r}) \cdot (D_t \nabla - v_0 \vec{u}) P(\vec{r}, \vec{u}; t)] \\ & + D_r \vec{\mathcal{R}}^2 P(\vec{r}, \vec{u}; t) + \Phi_1 + \Phi_g, \end{aligned} \quad (27)$$

where

$$\Phi_1 = -\mu P(\vec{r}, \vec{u}; t), \quad (28)$$

is the loss of the probability from the position \vec{r} due to resetting while

$$\Phi_g = \mu \delta(x) \int P(x', y, u_x, u_y; t) dx', \quad (29)$$

is the gain of the probability at the point $(0, y)$ on the x -axis.

The probability density, $P(\vec{r}; t)$, can be projected on spherical harmonics and consequently written as an expansion. By projecting the probability density on the zeroth and first spherical harmonics we find

$$P(\vec{r}; t) = \rho(\vec{r}; t) + \vec{\sigma}(\vec{r}; t) \cdot \vec{u} + \Xi \quad (30)$$

where $\rho(\vec{r}; t)$ is the positional probability density and the vector $\vec{\sigma}(\vec{r}; t)$ is the polar order parameter. Note that Ξ denotes higher-order contributions which are ignored in this study. Plugging this expansion in eqn (27) and then integrating the orientational degrees of freedom, we find

$$\frac{\partial \rho(\vec{r}; t)}{\partial t} = -\nabla \cdot [\vec{J}(\vec{r}; t) + \vec{J}^a(\vec{r}; t)] + \phi_1 + \phi_g, \quad (31)$$

where

$$\phi_1 = -\mu \rho(\vec{r}; t), \quad (32)$$

$$\phi_g = \mu \delta(x) \int \rho(x', y; t) dx', \quad (33)$$

and

$$\vec{J}(\vec{r}; t) = -D_t \Gamma^{-1}(\vec{r}) \nabla \rho(\vec{r}; t), \quad (34a)$$

$$\vec{J}^a(\vec{r}; t) = -\frac{v_0}{2} \Gamma^{-1}(\vec{r}) \vec{\sigma}(\vec{r}; t). \quad (34b)$$

To calculate the polar order parameter, $\vec{\sigma}(\vec{r}; t)$ we multiply eqn (27) by \vec{p} and integrate over the orientational degrees of freedom. This gives rise to an equation for $\vec{\sigma}(\vec{r}; t)$ as

$$\begin{aligned} \frac{\partial \vec{\sigma}(\vec{r}; t)}{\partial t} = & \nabla \cdot [\Gamma^{-1}(\vec{r}) (D_t \nabla \vec{\sigma}(\vec{r}; t) - v_0 \rho(\vec{r}; t))] \\ & - (D_r + \mu) \vec{\sigma}(\vec{r}; t) + \mu \delta(x) \int \vec{\sigma}(x', y; t) dx', \end{aligned} \quad (35)$$

This is the point where we make an assumption in which the density is the slowest mode in the system. In this limit, the gradients in the system are small in comparison to the modified persistence length of the active particle, $l_p = v_0/(D_r + \mu)$ with $D_r \gg \mu$. With this approximation the time derivative of $\vec{\sigma}(\vec{r}; t)$ and the first term on the right hand side of eqn (35), sitting in the bracket, is negligible. In addition, due to the symmetry in the system the integral over the polar order, given as the last term in eqn (35), is zero. In this limit, eqn (35) gives

$$\vec{\sigma}(\vec{r}; t) = -\frac{v_0}{(D_r + \mu)} \nabla \cdot [\Gamma^{-1}(\vec{r}) \rho(\vec{r}; t)]. \quad (36)$$

The substitution of eqn (36) into eqn (34b) and then the resulting equation together with eqn (34a) into eqn (31) gives

$$\begin{aligned} \frac{\partial \rho(\vec{r}; t)}{\partial t} = & \nabla \cdot [\Gamma^{-1}(\vec{r}) \cdot (D_t \nabla + v_0 \pi(\vec{r}; t)) \rho(\vec{r}; t)] + \phi_1 \\ & + \phi_g, \end{aligned} \quad (37)$$

where (in two dimensions) the polarization, $\pi(\vec{r}; t)$ is related to the polar parameter, $\vec{\sigma}(\vec{r}; t)$ through

$$\pi(\vec{r}; t) = \frac{\vec{\sigma}(\vec{r}; t)}{2\rho(\vec{r}; t)}. \quad (38)$$

B Derivation of the MFPT

Here we present the derivation of the MFPT of the active system. A similar approach can be used to derive the MFPT of the passive system. Using eqn (23) in the main text, the backward Fokker-Planck equation for the active system can be written as

$$\begin{aligned} \partial_t G(x; t) = & D_a e^{-\lambda x} \left[\frac{\partial}{\partial x^2} G(x; t) - \frac{\lambda}{2} \frac{\partial}{\partial x} G(x; t) \right] \\ & - \mu G(x; t) + \mu G(x_0; t), \end{aligned} \quad (39)$$

where $G(x_0; t)$ is the probability that the particle, started at x_0 , is not absorbed by the target. The initial and boundary conditions are given as

$$\begin{cases} G(x; t) = 0, & \text{at } t = 0, \\ G(x; t) = 1, & \text{at } x = 0. \end{cases} \quad (40)$$

Using the Laplace transform of $G(x; t)$, which is defined as

$$\tilde{G}(x; s) = \int_0^\infty e^{-st} G(x; t) dt, \quad (41)$$

the transformed backward equation can be written as

$$\begin{aligned} D_a e^{-\lambda x} \left[\frac{\partial}{\partial x^2} \tilde{G}(x; s) - \frac{\lambda}{2} \frac{\partial}{\partial x} \tilde{G}(x; s) \right] - (\mu + s) \tilde{G}(x; s) \\ = -1 - \mu \tilde{G}(x_0; s) \end{aligned} \quad (42)$$

To solve eqn (42), we first obtain the solution to the equation without the reinjection flux (the RHS terms), denoted by $\tilde{G}_0(x; s)$, i.e.

$$e^{-\lambda x} \left[\frac{\partial}{\partial x^2} \tilde{G}_0(x; s) - \frac{\lambda}{2} \frac{\partial}{\partial x} \tilde{G}_0(x; s) \right] - \alpha_a^2(s) \tilde{G}_0(x; s) = 0, \quad (43)$$

where $\alpha_a(s) = \sqrt{(\mu + s)/D_a}$. The solution to eqn (43) is

$$\tilde{G}_0(x; s) = A \exp \left[\frac{2\alpha_a(s)}{\lambda} \exp \left(\frac{\lambda x}{2} \right) \right] + B \exp \left[-\frac{2\alpha_a(s)}{\lambda} \exp \left(\frac{\lambda x}{2} \right) \right], \quad (44)$$

where A and B are constants. The condition that the probability density is finite as $x \rightarrow \infty$ implies that $A = 0$. Thus, the solution to eqn (42) can be written as

$$\tilde{G}(x; s) = B \exp \left[-\frac{2\alpha_a(s)}{\lambda} \exp \left(\frac{\lambda x}{2} \right) \right] + \frac{1 + \mu \tilde{G}(x_0; s)}{\mu + s}, \quad (45)$$

where the constant B can be calculated using the boundary condition in eqn (40), which reads

$$B = -\frac{1 + \mu \tilde{G}(x_0; s)}{\mu + s} \exp \left(\frac{2\alpha_a(s)}{\lambda} \right). \quad (46)$$

Plugging eqn (46) into eqn (45) gives

$$\begin{aligned}\tilde{G}(x; s) = & -\frac{1 + \mu\tilde{G}(x_0; s)}{\mu + s} \exp\left[-\frac{2\alpha_a(s)}{\lambda} \left(\exp\left(\frac{\lambda x}{2}\right) - 1\right)\right] \\ & + \frac{1 + \mu\tilde{G}(x_0; s)}{\mu + s}.\end{aligned}\quad (47)$$

Finally, by setting $x = x_0$ we obtain the survival probability, which can be written as

$$\tilde{G}(x_0; s) = \frac{1 - \exp\left[-\frac{2\alpha_a(s)}{\lambda} \left(\exp\left(\frac{\lambda x_0}{2}\right) - 1\right)\right]}{s + \exp\left[-\frac{2\alpha_a(s)}{\lambda} \left(\exp\left(\frac{\lambda x_0}{2}\right) - 1\right)\right]}, \quad (48)$$

which when evaluated at $s = 0$ gives the MFPT. This yields

$$T^a(x_0) = \frac{1}{\mu} \left[\exp\left(\frac{2\alpha_a}{\lambda} \left(\exp\left(\frac{\lambda x_0}{2}\right) - 1\right)\right) - 1 \right], \quad (49)$$

where $\alpha_a \equiv \alpha_a(0)$. Note that we obtained the MFPT for the active system. However, a similar method can be used to derive the MFPT for the passive system, as well.

Acknowledgements

A. Sharma acknowledges the support by the Deutsche Forschungsgemeinschaft (DFG) within the project SH 1275/3-1.

Notes and references

- 1 A. Fiasconaro, W. Ebeling and E. Gudowska-Nowak, *Eur. Phys. J. B*, 2008, **65**, 403–414.
- 2 P. Romanczuk, M. Bär, W. Ebeling, B. Lindner and L. Schimansky-Geier, *Eur. Phys. J.: Spec. Top.*, 2012, **202**, 1–162.
- 3 H. H. Wensink, H. Löwen, M. Marechal, A. Härtel, R. Wittkowski, U. Zimmermann, A. Kaiser and A. M. Menzel, *Eur. Phys. J.: Spec. Top.*, 2013, **222**, 3023–3037.
- 4 A. Walther and A. H. Muller, *Chem. Rev.*, 2013, **113**, 5194–5261.
- 5 J. Wang, *Nanomachines: fundamentals and applications*, John Wiley & Sons, 2013.
- 6 G. Volpe, S. Gigan and G. Volpe, *Am. J. Phys.*, 2014, **82**, 659–664.
- 7 J. Elgeti, R. G. Winkler and G. Gompper, *Rep. Prog. Phys.*, 2015, **78**, 056601.
- 8 M. E. Cates, *Rep. Prog. Phys.*, 2012, **75**, 042601.
- 9 É. Fodor, C. Nardini, M. E. Cates, J. Tailleur, P. Visco and F. van Wijland, *Phys. Rev. Lett.*, 2016, **117**, 038103.
- 10 M. E. Cates and J. Tailleur, *Annu. Rev. Condens. Matter Phys.*, 2015, **6**, 219–244.
- 11 P. Digregorio, D. Levis, A. Suma, L. F. Cugliandolo, G. Gonnella and I. Pagonabarraga, *Phys. Rev. Lett.*, 2018, **121**, 098003.
- 12 S. Mandal, B. Liebchen and H. Löwen, *Phys. Rev. Lett.*, 2019, **123**, 228001.
- 13 H. Löwen, *J. Chem. Phys.*, 2020, **152**, 040901.
- 14 J. Singh and A. A. Kumar, *Phys. Rev. E*, 2020, **101**, 022606.
- 15 A. D. Sezer, *Smart Drug Delivery System*, BoD-Books on Demand, 2016.
- 16 Z. Yang, A. H. Muller, C. Xu, P. S. Doyle, J. M. DeSimone, J. Lahann, F. Sciortino, S. Glotzer, L. Hong and D. A. Aarts *et al.*, *Janus particle synthesis, self-assembly and applications*, Royal Society of Chemistry, 2012.
- 17 W. F. Paxton, K. C. Kistler, C. C. Olmeda, A. Sen, S. K. St. Angelo, Y. Cao, T. E. Mallouk, P. E. Lammert and V. H. Crespi, *J. Am. Chem. Soc.*, 2004, **126**, 13424–13431.
- 18 S. Fournier-Bidoz, A. C. Arsenault, I. Manners and G. A. Ozin, *Chem. Commun.*, 2005, 441–443.
- 19 J. R. Howse, R. A. Jones, A. J. Ryan, T. Gough, R. Vafabakhsh and R. Golestanian, *Phys. Rev. Lett.*, 2007, **99**, 048102.
- 20 S. J. Ebbens and J. R. Howse, *Soft Matter*, 2010, **6**, 726–738.
- 21 W. Poon, *Proc. Int. Sch. Phys. “Enrico Fermi”*, 2013, **184**, 317–386.
- 22 T. Ekeh, M. E. Cates and É. Fodor, 2020, arXiv preprint arXiv:2002.05932.
- 23 W. Gao, A. Pei, R. Dong and J. Wang, *J. Am. Chem. Soc.*, 2014, **136**, 2276–2279.
- 24 P. K. Ghosh, Y. Li, F. Marchesoni and F. Nori, *Phys. Rev. E: Stat., Nonlinear, Soft Matter Phys.*, 2015, **92**, 012114.
- 25 H. D. Vuijk, A. Sharma, D. Mondal, J. U. Sommer and H. Merlitz, *Phys. Rev. E*, 2018, **97**, 042612.
- 26 H. Merlitz, H. D. Vuijk, R. Wittmann, A. Sharma and J.-U. Sommer, *PLoS One*, 2020, **15**, e0230873.
- 27 H. D. Vuijk, H. Merlitz, M. Lang, A. Sharma and J.-U. Sommer, 2020, arXiv preprint arXiv:2009.09060.
- 28 R. Filliger and P. Reimann, *EPL*, 2007, **77**, 30008.
- 29 V. Balakrishnan, *Elements of Nonequilibrium Statistical Mechanics*, Ane Books, India, 2008.
- 30 P. Friz, P. Gassiat and T. Lyons, *Transactions of the American Mathematical Society*, 2015, **367**, 7939–7955.
- 31 H.-M. Chun, X. Durang and J. D. Noh, *Phys. Rev. E*, 2018, **97**, 032117.
- 32 H. D. Vuijk, J. M. Brader and A. Sharma, *Soft Matter*, 2019, **15**, 1319–1326.
- 33 H. D. Vuijk, J. M. Brader and A. Sharma, *J. Stat. Mech.: Theory Exp.*, 2019, **2019**, 063203.
- 34 H.-M. Chun, L. P. Fischer and U. Seifert, *Phys. Rev. E*, 2019, **99**, 042128.
- 35 I. Abdoli, H. D. Vuijk, J.-U. Sommer, J. M. Brader and A. Sharma, *Phys. Rev. E*, 2020, **101**, 012120.
- 36 H. D. Vuijk, J.-U. Sommer, H. Merlitz, J. M. Brader and A. Sharma, *Phys. Rev. Res.*, 2020, **2**, 013320.
- 37 I. Abdoli, H. Vuijk, R. Wittmann, J. Sommer, J. Brader and A. Sharma, *Phys. Rev. Res.*, 2020, **2**, 023381.
- 38 I. Abdoli, E. Kalz, H. D. Vuijk, R. Wittmann, J.-U. Sommer, J. M. Brader and A. Sharma, *New J. Phys.*, 2020, **22**, 093057.
- 39 S. Cerrai, J. Wehr and Y. Zhu, 2020, arXiv preprint arXiv:2003.02330.
- 40 H.-J. Van Leeuwen, *J. Phys. Radium*, 1921, **2**, 361–377.
- 41 M. R. Evans and S. N. Majumdar, *Phys. Rev. Lett.*, 2011, **106**, 160601.

- 42 M. R. Evans and S. N. Majumdar, *J. Phys. A: Math. Theor.*, 2011, **44**, 435001.
- 43 M. R. Evans, S. N. Majumdar and K. Mallick, *J. Phys. A: Math. Theor.*, 2013, **46**, 185001.
- 44 X. Durang, M. Henkel and H. Park, *J. Phys. A: Math. Theor.*, 2014, **47**, 045002.
- 45 S. N. Majumdar, S. Sabhapandit and G. Schehr, *Phys. Rev. E: Stat., Nonlinear, Soft Matter Phys.*, 2015, **91**, 052131.
- 46 M. R. Evans and S. N. Majumdar, *J. Phys. A: Math. Theor.*, 2018, **51**, 475003.
- 47 J. Masoliver, *Phys. Rev. E*, 2019, **99**, 012121.
- 48 A. Pal, Ł. Kuśmierz and S. Reuveni, *Phys. Rev. E*, 2019, **100**, 040101.
- 49 T. T. da Silva and M. D. Fragoso, *J. Phys. A: Math. Theor.*, 2019, **52**, 495001.
- 50 M. R. Evans, S. N. Majumdar and G. Schehr, *J. Phys. A: Math. Theor.*, 2020, **53**, 193001.
- 51 M. Magoni, S. N. Majumdar and G. Schehr, 2020, arXiv preprint arXiv:2002.04867.
- 52 D. Gupta, C. A. Plata and A. Pal, *Phys. Rev. Lett.*, 2020, **124**, 110608.
- 53 S. Belan, *Phys. Rev. Res.*, 2020, **2**, 013243.
- 54 O. Tal-Friedman, A. Pal, A. Sekhon, S. Reuveni and Y. Roichman, 2020, arXiv preprint arXiv:2003.03096.
- 55 B. Besga, A. Bovon, A. Petrosyan, S. N. Majumdar and S. Ciliberto, *Phys. Rev. Res.*, 2020, **2**, 032029.
- 56 A. Scacchi and A. Sharma, *Mol. Phys.*, 2018, **116**, 460–464.
- 57 I. Santra, U. Basu and S. Sabhapandit, 2020, arXiv preprint arXiv:2009.09891.
- 58 V. Kumar, O. Sadekar and U. Basu, 2020, arXiv preprint arXiv:2008.03294.
- 59 A. Sharma and J. M. Brader, *J. Chem. Phys.*, 2016, **145**, 161101.
- 60 H. Merlitz, H. D. Vuijk, J. Brader, A. Sharma and J.-U. Sommer, *J. Chem. Phys.*, 2018, **148**, 194116.
- 61 P. Maggaretti, M. Janssen, I. Pagonabarraga and J. M. Rubi, *J. Chem. Phys.*, 2019, **151**, 084902.
- 62 B.-Q. Ai, W.-J. Zhu and J.-J. Liao, *New J. Phys.*, 2019, **21**, 093041.
- 63 P. Maggaretti, G. Oshanin and J. Talbot, *J. Phys.: Condens. Matter*, 2019, **31**, 270201.
- 64 Y. Li, R. Mei, Y. Xu, J. Kurths, J. Duan and R. Metzler, *New J. Phys.*, 2020, **22**, 053016.
- 65 P. C. Bressloff, *J. Phys. A: Math. Theor.*, 2020, **53**, 355001.
- 66 J. Stenhammar, R. Wittkowski, D. Marenduzzo and M. E. Cates, *Sci. Adv.*, 2016, **2**, e1501850.
- 67 A. Sharma and J. M. Brader, *Phys. Rev. E*, 2017, **96**, 032604.
- 68 H. Kählert, J. Carstensen, M. Bonitz, H. Löwen, F. Greiner and A. Piel, *Phys. Rev. Lett.*, 2012, **109**, 155003.
- 69 L. Caprini and U. M. B. Marconi, 2020, arXiv preprint arXiv:2009.14032.