

## Expression of the alpha8 integrin chain facilitates phagocytosis by renal mesangial cells

Ines Marek, Robert Becker, Fabian Fahlbusch, Carlos Menendez-Castro, Wolfgang Rascher, Christoph Daniel, Gudrun Volkert, Andrea Hartner

### Angaben zur Veröffentlichung / Publication details:

Marek, Ines, Robert Becker, Fabian Fahlbusch, Carlos Menendez-Castro, Wolfgang Rascher, Christoph Daniel, Gudrun Volkert, and Andrea Hartner. 2018. "Expression of the alpha8 integrin chain facilitates phagocytosis by renal mesangial cells." *Cellular Physiology and Biochemistry* 45 (6): 2161–73. <https://doi.org/10.1159/000488160>.

Original Paper

# Expression of the Alpha8 Integrin Chain Facilitates Phagocytosis by Renal Mesangial Cells

Ines Marek<sup>a</sup> Robert Becker<sup>b</sup> Fabian B. Fahlbusch<sup>a</sup> Carlos Menendez-Castro<sup>a</sup>  
Wolfgang Rascher<sup>a</sup> Christoph Daniel<sup>b</sup> Gudrun Volkert<sup>a</sup> Andrea Hartner<sup>a</sup>

<sup>a</sup>Department of Pediatrics and Adolescent Medicine, <sup>b</sup>Institute of Nephropathology, University Hospital of Erlangen, Erlangen, Germany

## Key Words

Alpha8 integrin • *Itga8* • Phagocytosis • Mesangial cells • Extracellular matrix resolution • Cytoskeleton • Glomerulonephritis

## Abstract

**Background/Aims:** Healing of mesangioproliferative glomerulonephritis involves degradation of excess extracellular matrix, resolution of hypercellularity by apoptosis and phagocytosis of apoptotic cells. Integrin receptors participate in the regulation of phagocytosis. In mice deficient for alpha8 integrin (*Itga8*<sup>-/-</sup>) healing of glomerulonephritis is delayed. As *Itga8* is abundant in mesangial cells (MC) which are non-professional phagocytes, we hypothesized that *Itga8* facilitates phagocytosis of apoptotic cells and matrix components by MC. **Methods:** MC were isolated from wild type (WT) and *Itga8*<sup>-/-</sup> mice. Latex beads were coated with matrix components. Apoptosis was induced by cisplatin in macrophages and in DiI-stained MC. After coincubation of latex beads or apoptotic cells with MC, the phagocytosis rate was detected in WT and *Itga8*<sup>-/-</sup> MC via fluorescence microscopy and FACS analysis. **Results:** *Itga8*<sup>-/-</sup> MC showed reduced phagocytosis of matrix-coated beads and apoptotic cells compared to WT MC. Reduction of stress fibers was observed in *Itga8*<sup>-/-</sup> compared to WT MC. Inhibition of cytoskeletal reorganization by inhibition of Rac1 or ROCK during phagocytosis significantly decreased the rate of phagocytosis by WT MC but not by *Itga8*<sup>-/-</sup> MC. **Conclusion:** The expression of *Itga8* facilitates phagocytosis in MC, likely mediated by *Itga8*-cytoskeleton interactions. An impairment of MC phagocytosis might thus contribute to a delayed glomerular regeneration in *Itga8*<sup>-/-</sup> mice.

© 2018 The Author(s)  
Published by S. Karger AG, Basel

G. Volkert and A. Hartner contributed equally to this work.

Ines Marek

Department of Pediatrics and Adolescent Medicine, University Hospital of Erlangen  
Loschgestrasse 15, Erlangen (Germany)  
Tel. +49-9131/8533118, Fax +49-9131/8539202, E-Mail Ines.Marek@uk-erlangen.de

KARGER

## Introduction

Glomerulonephritis is characterized by invasion of inflammatory cells, cell proliferation and extracellular matrix expansion. In acute forms of glomerulonephritis, regenerative mechanisms contribute to the resolution of glomerular inflammation, hypercellularity and fibrosis, thereby restoring the balance between proliferation, matrix deposition, apoptosis, phagocytosis of inflammatory cells, surplus mesangial cells and extracellular matrix components [1]. The latter involves the engulfment of large particles through formation of vesicles called phagosomes. Phagocytosis is not limited to “professional” phagocytes such as macrophages, but also performed by “non-professional” phagocytes such as epithelial, endothelial, neuronal and mesenchymal cells [2]. Phagocytosis by non-professional phagocytes is an important mechanism to maintain tissue homeostasis which eliminates apoptotic cells and matrix components from tissue [3, 4]. In experimental glomerulonephritis, phagocytosis of apoptotic mesangial cells by neighbouring healthy mesangial cells leads to the resolution of hypercellularity [5] and thereby contributes to restoring the integrity of the glomerular tuft.

A wide variety of receptor-ligand interactions and signaling pathways has been implicated in phagocytosis, summarized in [6–8]. Recognition and binding of apoptotic cells commonly lead to the intracellular activation of Rho-GTPases and subsequent remodeling of the actin cytoskeleton [2]. Signaling via Rac1 is described to be active in the phagocytic cup [9], and in FCyR-dependent phagocytosis, whereas signaling via RhoA is essential for  $\alpha\text{M}\beta 2$  integrin-mediated phagocytosis [10]. Integrins can promote phagocytosis in different cell types, such as epithelial, endothelial and mesenchymal cells dependent on their lineage [2, 11]. In immune cells, e.g.  $\alpha\text{M}\beta 2$ ,  $\alpha\text{V}\beta 3$ ,  $\alpha\text{V}\beta 5$  and  $\alpha 6\beta 1$  integrins are described as mediators of phagocytosis (summarized in [2]). In macrophages,  $\alpha\text{M}\beta 5$  integrin is described as a pro-phagocytotic integrin which activates RhoA. RhoA activation results in ROCK-mediated myosin II phosphorylation which is necessary for F-actin turnover to seal engulfed particles [12].  $\alpha\text{V}\beta 5$  integrin is able to bind and internalize apoptotic cells via Rac1 activation [13, 14]. In trabecular meshwork cells,  $\alpha\text{V}\beta 5$  integrin mediates phagocytosis via FAK activation, while  $\alpha\text{V}\beta 3$  reduces phagocytotic activity in these cells [15]. In contrast,  $\alpha\text{V}\beta 3$  serves to internalize apoptotic neutrophils in macrophages [13]. In human breast cancer cells,  $\alpha 3\beta 1$  mediates phagocytosis of extracellular matrix [16]. In microglia,  $\alpha 6\beta 1$  integrin is described to be a component of the complex that can internalize fibrillary  $\beta$ -amyloids [17]. Fibroblasts are able to phagocyte collagens via  $\alpha 2\beta 1$  integrin after integrin clustering and Rac1 activated F-actin reassembly [18, 19]. In mesangial cells,  $\alpha\text{V}\beta 3$  seems to be involved in phagocytosis as internalization of apoptotic cells by mesangial cells was found to be reduced following inhibition of the  $\alpha\text{V}\beta 3$ /thrombospondin/CD36 pathway [20, 21].

Alpha8 integrin (*Itga8*) is an RGD binding integrin which is expressed on mesangial cells and regulates their biological properties including adhesion, proliferation, migration and apoptosis by interacting with extracellular ligands such as fibronectin, vitronectin or osteopontin [22, 23]. Its role for phagocytosis is not yet known. However, previous *in vivo* studies in *Itga8*-deficient mice indicated a contribution of *Itga8* to the resolution of glomerular injury after acute mesangioproliferative glomerulonephritis [24]. Healing of glomerular damage was considerably delayed in *Itga8*-deficient mice [24]. Thus, we hypothesized that *Itga8* might contribute to phagocytosis of apoptotic mesangial cells, inflammatory cells and degraded extracellular matrix components in glomeruli after an inflammatory insult. It was further assumed that in mesangial cells with a deficiency for *Itga8* phagocytosis might be impaired. We performed *in vitro* studies using wild type and *Itga8*-deficient mesangial cells to analyze the contribution of *Itga8* to phagocytosis of apoptotic cells and extracellular matrix components.

## Materials and Methods

### Cell culture

Mesangial cells (MC) were isolated from kidneys of healthy wild type (WT) or *Itga8*-deficient (*Itga8*<sup>-/-</sup>) mice (obtained from U. Müller, Scripps Institute, LaJolla, USA) by the sieving method and characterized as described before [23, 25]. Rat MC were isolated from Sprague-Dawley rats as described in [26]. MC were grown in Dulbecco's modified Eagle's Medium (DMEM with L-Glutamine; PAA Laboratories GmbH, Pasching, Austria) containing 10% FCS, 1% penicillin-streptomycin, 0.1% insulin (Sigma, Deisenhofen, Germany) in a 95% air – 5% CO<sub>2</sub> humidified atmosphere at 37°C. MC were used for experiments in passages 5-18. A stably GFP-transfected murine macrophage cell line (GFP-RAW) (gift from B. Wielockx and J. Kalucka, Dresden, Germany [27]) was cultivated in DMEM with L-Glutamine containing 10% FCS, 1% penicillin-streptomycin in a 95% air – 5% CO<sub>2</sub> humidified atmosphere at 37°C. A stably *Itga8* overexpressing HEK cell line (*Itga8*<sup>+/+</sup> HEK) (described in [23]) was cultivated in DMEM with L-Glutamine containing 10% FCS, 1% penicillin-streptomycin and 0.4mg/ml G418 (PAA Laboratories GmbH, Pasching, Austria). Wild type HEK cells were used as a control (co HEK).

### Generation of apoptotic cells

Apoptosis was induced *in vitro* by incubation of either GFP-RAW cells or vybrant DiI (Molecular Probes Life Technologies, Eugene, OR, USA) stained MC with 50µM cisplatin (cis-diamineplatinum (II) dichloride from Sigma-Aldrich, Munich, Germany) for 24 hours at 37°C similar as described before [24]. For vybrant DiI dye staining, MC were seeded in culture dishes and were allowed to adhere overnight. Then, vybrant DiI dye was added (5µl/ml DMEM) for 10min at 37°C. Apoptosis was evaluated as described in [24]. Hoechst staining was used to analyze apoptosis via counting of chromatin-condensed cells. Trypan blue exclusion was used to determine viability (in average 80%).

### Phagocytosis of apoptotic cells


Phagocytosis was analyzed by performing co-culture of apoptotic cells (GFP-RAW, MC) with non-apoptotic MC at different time points. Apoptotic GFP-RAW macrophages and native, adherent MC were co-incubated at a ratio of 10:1 (apoptotic RAW to native MC) for 2, 4, 6 and 24 hours. For phagocytosis of apoptotic MC, apoptotic DiI stained MC and native, unstained adherent MC were co-incubated at a ratio of 2.5:1 (apoptotic to native MC) for 4 and 24 hours. The phagocytosis rate of apoptotic cells was compared in WT and *Itga8*<sup>-/-</sup> MC via fluorescence microscopy or FACS analysis.

### Phagocytosis of matrix-coated beads

A latex bead model [28-30] was employed to study phagocytosis of extracellular matrix components (2µm, latex, carboxylate-modified polystyrene, yellow-green fluorescent beads from Sigma-Aldrich) in MC. Latex beads were coated with the *Itga8* ligand fibronectin (10µg/ml in PBS) (Sigma-Aldrich), collagen type I, which is not a ligand for *Itga8* (0.3µg/ml in PBS) (Collaborative Biomedical Products, Bedford, MA, USA), or mouse serum (gift from C. Warnecke, Erlangen, Germany) at 37°C for 1 hour. Mouse serum was used as an unspecific coating control (similar to [31]). MC (WT: 10, 000 cells/well, *Itga8*<sup>-/-</sup> 15, 000 cells/well) were seeded on glass 8-well chamber slides coated with fibronectin (10µg/ml). Cells were allowed to attach overnight. MC were layered with coated beads in a ratio of 1:50 (cells to beads). The optimal matrix coating concentration and the optimal incubation time for phagocytosis was assessed in pilot studies, similar as in [29, 30, 32-35] (see Fig. 1). After 24 hours of incubation at 37°C, cells were washed with saline and fixed in 4% paraformaldehyde. Nuclei were stained with DaPi (1:2000; Sigma-Aldrich) and actin filaments with rhodamine/phalloidin (5mM; Molecular Probes, Eugene, OR, USA), as described before [22]. The phagocytosis rate of latex beads was assessed via fluorescence microscopy.

For analysis of phagocytosis of fibronectin-coated latex beads in *Itga8* overexpressing cells, *Itga8*<sup>+/+</sup> and co HEK cells (30, 000 cells/well) were seeded on glass 8-well chamber slides coated with fibronectin (10µg/ml). The next day, HEK cells were layered with fibronectin-coated (10µg/ml) beads in the ratio of 1:50 (cells to beads). For HEK cell experiments, latex beads (carboxylate-modified polystyrene, yellow-green fluorescent beads from Sigma-Aldrich) with a size of 1µm were chosen as described in [36, 37] and phagocytosis was analyzed as described above for MC after 3 and 24 hours by fluorescence microscopy.

**Fig. 1.** Testing conditions for phagocytosis of beads. A, Comparison of the phagocytosis of serum-coated beads by  $\alpha 8$  integrin-deficient (Itga8<sup>-/-</sup>) and wild type (WT) mesangial cells (MC) after 3 and 24 hours (3h, 24h), cell-to-bead ratio = 1:50, N=4, \*=p<0.05 24h versus 3h. B, Comparison of the phagocytosis of matrix-coated beads, using different coating concentrations, by Itga8<sup>-/-</sup> and wild type MC after 24 hours, cell-to-bead ratio = 1:50, FN= Fibronectin, COL= Collagen I, N=3, \*=p<0.05 Itga8<sup>-/-</sup> versus wild type MC. C, Phagocytosis of serum-coated beads in Itga8<sup>-/-</sup> and wild type MC in dependence of the cell-to-bead ratio (1:5 versus 1:50) after 24 hours, N=3, \*=p<0.05 1:5 versus 1:50 cell-to-bead ratio. D, Inhibition of the ingestion of serum-coated beads by cytochalasin D (cyto D) in Itga8<sup>-/-</sup> and wild type MC in dependence of the cyto D concentration after 24 hours. Results are representative for three similar experiments. co=solvent control. \*= no phagocytosis detected, all Itga8<sup>-/-</sup> MC dead. Phagocytosis in untreated wild type MC is set to 100%.




#### Flow cytometry

For FACS analysis (Canto II, BD Biosciences, Heidelberg, Germany), FITC-channel (530nm wave length) was used to detect the phagocytosis of apoptotic GFP-RAW cells. PE-channel (585nm wave length) was used to detect phagocytosis of apoptotic red fluorescent DiI stained MC. FSC (forward scatter channel) was used to quantify the size and SSC (side scatter channel) to detect the granularity of the cells (see Fig. 2).

#### Fluorescence microscopy

Internalization of apoptotic GFP-RAW cells by MC was analyzed after co-incubation for 2, 4, 6 and 24 hours. Cells were fixed in 4% paraformaldehyde before microscopic evaluation (NIKON Eclipse 80i, Düsseldorf) in bright field and fluorescence microscopy. For phagocytosis of apoptotic DiI stained MC by MC, cells were fixed after 4 and 24 hours and evaluated in the same manner. Bead internalization was confirmed with fluorescence microscopy (NIKON Eclipse 80i) and inverted microscopy with z-stack images (Keyence, Neu-Isenburg, Germany), (see Fig. 3). The quantitative analysis of the phagocytosis rate was



**Fig. 2.** FACS analysis of phagocytosis. A, B, Phagocytosis of apoptotic DiI-stained  $\alpha 8$  integrin-deficient (Itga8<sup>-/-</sup>) mesangial cells (MC) after 4 hours. Apoptotic, DiI-stained MC correspond to green dots in (A), green arrow points to DiI-stained apoptotic MC in (B). Native, not phagocytosing MC are shown by red dots (A) and red arrow (B). Phagocytosing MC correspond to blue dots (A), blue arrow points to phagocytosing MC (B). C, D, Phagocytosis of apoptotic GFP-RAW by wild type MC (C) and by Itga8<sup>-/-</sup> MC (D) after 4 hours of co-culture. Apoptotic GFP-RAW are depicted in green (P1), not phagocytosing MC in blue (P3), phagocytosing MC in purple dots (P4).



calculated as the ratio of the number of phagocytosing cells to the total cell number, similar to [28, 38]. In inhibition assays, the phagocytosis rate of matrix-coated beads by WT MC seeded on fibronectin was normalized to 100% as baseline level.

#### Selective integrin blocking

Blocking antibodies to  $\alpha 6$  and  $\alpha 2$  integrin chains (both BD Pharmingen, Heidelberg, Germany) were used at a concentration of  $2\mu\text{g}/\text{ml}$ ,  $\alpha\text{V}\beta 3$  integrin blocking RGD peptides and the negative peptide control (cRAD) (both from Bachem, Bubendorf, Switzerland) at a concentration of 5 and  $50\mu\text{M}$ , as described in [39]. MC were incubated with blocking antibodies or with blocking RGD peptides for 1 hour at  $37^\circ\text{C}$  before starting phagocytosis assays.

#### Inhibition of actin polymerization

MC were either treated with medium containing Cytochalasin D (0.5, 1.0 and  $10\mu\text{M}$ ) for 30 minutes to inhibit actin polymerization or treated with an equal amount of the solvent dimethylsulfoxide (DMSO) only, see Fig. 1D (similar as described [31, 40, 41]). Then, latex beads were added to start phagocytosis, as described above.

#### Inhibition of Rho GTPase-dependent actin remodeling


ROCK inhibitor Y27632 (Calbiochem, La Jolla, CA, USA) is a selective inhibitor of the Rho-associated kinase p160 ROCK (an effector kinase of RhoA), NSC23766 (Abcam, Cambridge, UK) is an inhibitor of Rac1. Before starting phagocytosis assays with latex beads, as described above, MC were incubated with Y27632 at a concentration of  $10\mu\text{M}$  for 15 minutes at  $37^\circ\text{C}$ , as described in [42]. NSC23766 was used at two different concentrations (10 and  $50\mu\text{M}$ ) for incubation of MC for 30 minutes at  $37^\circ\text{C}$ , as described in [43].

#### Silencing of *Itga8* in rat MC

Gene silencing of *Itga8* was achieved by transfection of rat MC with *Itga8* siRNA (si *Itga8*) according to the “fast protocol” from the HiPerFect Transfection Reagent Handbook (Qiagen, Hilden, Germany). MC were transfected in cell suspension with a final concentration of  $5\text{nM}$  si *Itga8* (sense: GAC CUC CUC AGG AUG AAA UdT dT, antisense: AUU UCA UCC UGA GGA GGU CdT dT, Qiagen) before seeding to allow silencing for 72 hours. A nonsilencing siRNA (si co) (sense: UUC UCC GAA CGU GUC ACG UdT dT, antisense: ACG UGA CAC GUU CGG AGA AdTdT, Qiagen) was used as control. To control for effective silencing, real-time PCR for *Itga8* expression levels was performed, as described in [26]. Moreover, western blot analysis was used to confirm down regulation of *Itga8* on protein level.

#### Isolation of mRNA and Real-time PCR

To evaluate mRNA expression levels of *Itga8* and *Acta2* ( $\alpha$ -smooth muscle actin) after silencing of *Itga8* with siRNA, total RNA was obtained from rat MC by extraction with RNeasy® Mini columns (Qiagen, Hilden, Germany). First-strand cDNA was synthesized with TaqMan reverse transcription reagents



**Fig. 3.** Visualization of bead localization. Examples of fluorescence microscopic images with z-stacks to verify intracellular bead localization in wild type MC. Left: intracellular bead localization (white arrow). Right: extracellular bead localization (white arrow). Nuclei are stained with DaPi, actin filaments with rhodamine/phalloidin.

(Applied Biosystems, Weiterstadt, Germany) using random hexamers as primers. Final RNA concentration in the reaction mixture was adjusted to 100ng/μL. Reactions without MultiScribe Reverse Transcriptase (Applied Biosystems) were used as negative controls for genomic DNA contamination. PCR was performed with an ABI PRISM 7000 Sequence Detector System and SYBR Green (Applied Biosystems) according to the manufacturer's instructions. The relative amount of the specific mRNA was normalized with respect to *Rn18s*. Primers were created with Primer3 Output program ([http://biotools.umassmed.edu/bioapps/primer3\\_www.cgi](http://biotools.umassmed.edu/bioapps/primer3_www.cgi)). Primer sequences (MWG Biotech, Ebersberg, Germany) used for *Rn18s* were forward 5'-TTGATTAAGTCCCTGCCCTTTGT-3', reverse 5'-CGATCCGAGGGCCTCACTA-3', for *Itga8* forward 5'-TCCAAATCAGAAGCTCCAACAA-3', reverse 5'-CGCTCACGAAATTGCTGTCA-3' and for *Acta2* forward 5'-TCCTGACCCTGAAGTATCCGATA-3', reverse 5'-GGTGCCAGATCTTTTCCATGTC-3'. All samples were run in triplicates.

## Detection of F-actin and Itga8 in cultured cells

For F-actin staining, MC were seeded on glass 8-well chamber slides blocked with 2% BSA. Cells were allowed to adhere for 24 hours. Then, adherent cells were rinsed three times with PBS and fixed in 4% paraformaldehyde (PFA) for 10 minutes. Cells were permeabilized using 1% Triton X-100 and nonspecific binding was blocked using 100% FCS. F-actin was visualized with rhodamin/phalloidin from Molecular Probes (Leiden, The Netherlands). Cells were embedded in Tris-buffered Mowiol, pH 8.6 (Hoechst, Frankfurt, Germany). For staining of *Itga8* in HEK cells, *Itga8*<sup>+/+</sup> HEK and co HEK were seeded on fibronectin (10μg/ml) coated 8-well chamber slides blocked with 2% BSA. Cells were allowed to adhere overnight. Then, cells were prefixed with 4% PFA (0.3ml 4% PFA added to 0.5ml medium/well) for 2 minutes at room temperature, washed carefully with PBS and fixed with 4% PFA for 20 minutes. After washing twice with PBS, cells were permeabilized using 0.2% Triton X-100/PBS. After rinsing twice with PBS, cells were stained with an *Itga8* antibody (polyclonal goat anti-mouse/rat IgG, AF4076, R&D Systems, Minneapolis, USA) 1:50 in PBS overnight at 4°C. After washing three times with PBS, a cy<sup>TM</sup>3-conjugated donkey anti-goat IgG secondary antibody (Jackson ImmunoResearch Laboratories, Nr 705-166-147, West Grove, PA, USA) was applied 1:300 in PBS for 1 hour at 37°C. Finally, after washing with PBS, chamber slides were embedded with Fluoromount<sup>TM</sup> (Sigma Aldrich).

## Western Blot Analysis

Protein was isolated from MC with cell lysis buffer (50mM HEPES pH 7.4, 150mM NaCl, 1% Triton X-100, 1mM EDTA, Complete Protease-Inhibitor Cocktail (Roche, Mannheim) by incubating harvested cells on ice for 30 minutes with intermittent vortexing, followed by centrifugation for 10 minutes at 13,000 rpm. Protein concentration of cell lysate supernatants was determined using a protein assay kit (Pierce, Rockford, IL). Protein samples containing 30μg – 50μg total protein were denatured by boiling for five minutes and separated on an 8% denaturing SDS-PAGE gel. After electrophoresis, the gels were electro blotted onto PVDF membranes (Pall Filtron, Karlsruhe, Germany), blocked with Roti-Block (Roth, Karlsruhe, Germany) for α-smooth muscle actin (SMA), *Itga8* and β-actin detection, or 5% low fat milk in TBS-T (Humana Milichunion eG, Herford, Germany) for β-tubulin detection, for 1 hour and incubated with the primary antibody overnight. A mouse monoclonal antibody to SMA (DAKO Diagnostika, Hamburg, Germany) was used at a dilution of 1:1000 in 1x Roti-Block. A goat anti-mouse antibody to *Itga8* (R&D Systems) was used at a dilution of 1:1000 in 1x Roti-Block. As loading controls, a rabbit polyclonal antibody to β-tubulin (Abcam, Cambridge, UK) was used at a dilution of 1:5000 in 1% FCS/PBS and a mouse monoclonal antibody to β-actin (Sigma Aldrich) was used at a dilution of 1:20,000 in Roti-Block. Horseradish peroxidase-conjugated anti-mouse IgG antibody and anti-rabbit IgG antibody (Santa Cruz Biotechnology, Heidelberg, Germany), both diluted 1:10,000 in TBS-T, and anti-goat IgG antibody (Santa Cruz Biotechnology), diluted 1:50,000 in TBS-T were used as secondary antibodies, respectively. Immunoreactivity was visualized with a secondary horseradish peroxidase-conjugated IgG antibody, using the ECL system according to the manufacturer's instructions (Amersham, Braunschweig, Germany). Densitometric evaluation was performed using AIDA image analyzer 4.15 (Raytest, Berlin).

## Statistics

A two-tailed student's t-test was used to test significance of differences between groups. A p-value of <0.05 was considered significant. Values are displayed as means±standard deviation.

## Results

### Phagocytosis of latex beads by MC


The number of MC performing phagocytosis of extracellular matrix-coated beads was significantly higher in wild type MC compared to *Itga8*<sup>-/-</sup> MC (Fig. 4) with a higher affinity of both cell types to fibronectin-coated than collagen-coated beads (Fig. 4). 24 hours after incubation with fibronectin-coated beads, phagocytosis was detected in nearly 80% of wild type MC, but only in about 45% of *Itga8*<sup>-/-</sup> MC. At the same time, nearly 40% of wild type MC but only 10% of *Itga8*<sup>-/-</sup> MC internalized collagen-coated beads.

To gain more insight in the pathways involved in phagocytosis of wild type versus *Itga8*<sup>-/-</sup> MC, we performed inhibition assays for certain integrins and regulators of cytoskeletal organization.


Functional blocking of the RGD-binding  $\alpha$ V integrin, an integrin known to be mainly involved in phagocytosis in MC [20], resulted in a significant dose-dependent reduction of phagocytosis of fibronectin-coated beads in wild type MC (Fig. 5A), while there was only a moderate, yet insignificant reduction of phagocytosis in *Itga8*<sup>-/-</sup> MC (Fig. 5B).

Previous studies showed an increase in the expression of  $\alpha$ 2 and  $\alpha$ 6 integrins in *Itga8*<sup>-/-</sup> MC [22]. Therefore, we hypothesized that these integrins might partly compensate the loss of *Itga8* for phagocytosis. The use of inhibitory antibodies to  $\alpha$ 2 and  $\alpha$ 6 integrins revealed a decreased rate of phagocytosis of fibronectin-coated latex beads in wild type MC to 88.9% after blocking of  $\alpha$ 2 integrin, to 85.5% after blocking of  $\alpha$ 6 integrin and to 78% after simultaneous blocking of both  $\alpha$ 2 and  $\alpha$ 6 integrin compared to untreated controls (Fig. 5C). In contrast, no significant reduction of phagocytosis after blocking  $\alpha$ 2 and  $\alpha$ 6 integrin was observed in *Itga8*<sup>-/-</sup> MC arguing against a compensatory role for these integrins (Fig. 5D).

Inhibition of actin remodeling was achieved by ROCK and Rac1 inhibitors. ROCK inhibition showed a significant reduction of phagocytosis of fibronectin-coated latex beads in wild type MC, but not in *Itga8*<sup>-/-</sup> MC (Fig. 6A and B, respectively). Rac1 inhibition dose-dependently reduced phagocytosis of fibronectin-coated beads in wild type MC, with the strongest effect at a concentration of 50  $\mu$ M (Fig. 6C). In *Itga8*<sup>-/-</sup> MC, no significant effect of the Rac1 inhibitor on phagocytosis was detected (Fig. 6D).




**Fig. 4.** Phagocytosis of matrix-coated beads. A, Comparison of the uptake of extracellular matrix-coated latex beads in  $\alpha$ 8 integrin-deficient (*Itga8*<sup>-/-</sup>) and in wild type (WT) mesangial cells (MC). B, exemplary photomicrographs of WT and *Itga8*<sup>-/-</sup> MC during phagocytic uptake of matrix-coated beads. Nuclei were stained with DAPI, actin filaments with rhodamine/phalloidin. FN= Fibronectin, COL= Collagen I, N=3, \* =  $p < 0.05$  *Itga8*<sup>-/-</sup> versus WT MC.




**Fig. 5.** Inhibition of integrin chains ( $\alpha$ V,  $\alpha$ 2,  $\alpha$ 6) during phagocytosis. Inhibition of the uptake of fibronectin-coated beads in wild type (A) and  $\alpha$ 8 integrin-deficient (B) mesangial cells (MC) by cRGD. RAD=control non-blocking peptide, cRGD=  $\alpha$ V integrin-blocking peptide. N=3, \* =  $p < 0.05$  versus unstimulated control (co). Inhibition of phagocytosis of fibronectin-coated beads by  $\alpha$ 2 or  $\alpha$ 6 integrin-blocking antibodies (anti  $\alpha$ 2, anti  $\alpha$ 6) and by both antibodies applied together (anti  $\alpha$ 2 $\alpha$ 6) in wild type (C) and  $\alpha$ 8 integrin-deficient (D) mesangial cells (MC). N=3, \* =  $p < 0.05$  versus unstimulated control (co).



**Fig. 6.** Inhibition of actin remodeling during phagocytosis. Inhibition of phagocytosis of fibronectin-coated beads by a ROCK inhibitor (ROCK-Inh) in wild type (A) and  $\alpha 8$  integrin-deficient (B) mesangial cells (MC). Inhibition of phagocytosis of fibronectin-coated beads by a RAC1 inhibitor (Rac1-Inh) in wild type (C) and  $\alpha 8$  integrin-deficient (D) mesangial cells (MC). N=6, \*=  $p < 0.05$  versus unstimulated control (co).



**Fig. 7.** Phagocytosis of fibronectin-coated latex beads in HEK cells. A, uptake of fibronectin-coated beads (green fluorescence) by Itga8-overexpressing HEK cells (Itga8+/+, C+E) and wildtype HEK cells (co, B, D+F) after 3 and 24 hours, exemplary photomicrographs representative for 3 similar independent experiments. \*=  $p < 0.05$  Itga8+/+ versus co. E, F, control staining of  $\alpha 8$  integrin in Itga8+/+ (E) and co HEK cells (F).




Inhibition of actin polymerization resulted in a significant reduction of phagocytosing cells. Pretreatment with Cytochalasin D reduced the number of phagocytosing MC, both of the wild type and the *Itga8*<sup>-/-</sup> phenotype, in a dose dependent manner (Fig. 1D).

#### Phagocytosis of latex beads by HEK cells

After 3 hours of phagocytosis of fibronectin-coated beads, *Itga8* overexpressing HEK cells (*Itga8*<sup>+/+</sup> HEK) showed a higher phagocytosis rate (34.9% phagocytosing cells) compared to wild type HEK cells (co HEK, 12.1% phagocytosing cells). After 24 hours, the phagocytosis rate was not significantly different in both cell types anymore (91.2% in *Itga8*<sup>+/+</sup> HEK cells versus 87.8% in co HEK, see Fig. 7).

#### Phagocytosis of apoptotic RAW cells by MC

Cisplatin was used to induce apoptosis in GFP-RAW. 50μM cisplatin induced an apoptotic rate of 82.8% compared to 4.6% in unstimulated controls (Fig. 8A). Phagocytosis of apoptotic RAW cells was significantly reduced in *Itga8*<sup>-/-</sup> MC compared to wild type MC after 2, 4 and 6 hours of coincubation. After 24



**Fig. 8.** Phagocytosis of GFP-labelled RAW cells (GFP-RAW). A, Induction of apoptosis in GFP-RAW by 50μM cisplatin (CP). N=3. \*=  $p < 0.05$  versus unstimulated control (co). B, Uptake of apoptotic GFP-RAW in wild type (WT) and  $\alpha 8$  integrin-deficient (*Itga8*<sup>-/-</sup>) mesangial cells (MC). N=3, \*=  $p < 0.05$  *Itga8*<sup>-/-</sup> versus WT MC. C, exemplary photomicrographs of MC phagocytosing apoptotic GFP-RAW after 6 hours.

**Fig. 9.** Phagocytosis of Dil-labelled apoptotic MC. A, Uptake of apoptotic Dil-stained wild type (WT) and  $\alpha 8$  integrin deficient (*Itga8*<sup>-/-</sup>) mesangial cells (MC) by native MC after 4 and 24 hours. N=3, \* $p < 0.05$  *Itga8*<sup>-/-</sup> versus WT. B, exemplary photomicrographs of MC phagocytosing apoptotic MC after 4 hours.

hours, phagocytosis did not differ significantly in both cell types any more (Fig. 8B). Fig. 8C shows exemplary photomicrographs after 6 hours of coincubation.

#### Phagocytosis of apoptotic MC by MC


Apoptosis was induced in MC using 50  $\mu$ M cisplatin as described before [24]. Phagocytosis of apoptotic MC differed significantly after 4 hours with a higher rate of uptake in wild type MC (40.8% versus 23.7% *Itga8*<sup>-/-</sup> MC). After 24 hours, the rate of phagocytosis of apoptotic MC by *Itga8*<sup>-/-</sup> MC and wild type MC was comparable (Fig. 9).

#### Changes in the actin cytoskeleton after loss of *Itga8*

*Itga8*<sup>-/-</sup> MC acquire a different cell shape compared to wild type MC (Fig. 10A). Stainings for F-actin revealed that typical stress fibers and the fibroblast-like phenotype of wild type MC are no longer present in *Itga8*<sup>-/-</sup> MC (Fig. 10A). Similar observations were made after a knockdown of *Itga8* using si RNA to *Itga8*, which also results in a change of the arrangement of actin filaments in the cell boundary close to the cell membrane and a more compact morphology (Fig. 10B). Moreover, like already described for *Itga8*<sup>-/-</sup> MC [22], in si *Itga8* MC  $\alpha$ -smooth muscle actin is downregulated after a transient knock down of *Itga8* (Fig. 10C, D, E). Thus, underexpression of *Itga8* results in profound changes in the cytoskeletal structure.

#### Discussion

Mesangial cells can serve as non-professional phagocytes, thereby supporting healing of the glomerular tuft after glomerulonephritis [1]. *Itga8* is an integrin which is specifically expressed on mesangial cells in the glomerulus [25]. Based on our *in vivo* findings of delayed glomerular regeneration after glomerulonephritis in *Itga8*-deficient mice [24], phagocytosis of apoptotic cells and matrix components was analyzed in *Itga8*-deficient mesangial cells *in vitro* as a possible mechanism affecting the speed of tissue regeneration. In mesangial cells, the



**Fig. 10.** Changes in the actin cytoskeleton after the loss of  $\alpha 8$  integrin. A, F-actin staining in  $\alpha 8$  integrin-deficient (*Itga8*<sup>-/-</sup>) and wild type (WT) mesangial cells (MC). B, F-actin staining after silencing of *Itga8* in MC (si *Itga8*). C,  $\alpha$ -smooth muscle actin (SMA) mRNA expression in si *Itga8* MC. N=3. \* $p < 0.05$  versus non-silencing control (co). D, representative western blots for *Itga8* and SMA after silencing of *Itga8* in MC (si *Itga8*),  $\beta$ -tubulin,  $\beta$ -actin= loading control. E, statistical analysis after densitometric evaluation of western blots for *Itga8* and SMA after silencing of *Itga8* in MC (si *Itga8*). N=3, \* $p < 0.05$  versus control (co).

functional relevance of  $\alpha V\beta 3$  integrin as a mediator of phagocytosis was described by Cortes-Hernandez et al [20]. The role of *Itga8*, however, for phagocytosis in glomerular regeneration or healing has not yet been elucidated. In our study, we investigated phagocytosis of matrix components by using matrix-coated beads and phagocytosis of apoptotic cells by using apoptotic GFP-transfected RAW cells or apoptotic DiI-stained mesangial cells in wild type and *Itga8*-deficient mesangial cells.

Our results show that the presence of *Itga8* facilitates phagocytosis of matrix components and apoptotic cells by mesangial cells. Thus, the expression of *Itga8* on mesangial cells could be especially beneficial after acute glomerulonephritis. Glomerulonephritis leads to an influx of inflammatory cells, mesangial cell hyperplasia and excessive deposition of extracellular matrix (fibrosis). Resolution of glomerulonephritis comprises apoptosis of surplus cells, which are removed via phagocytosis conducted by mesangial cells in addition to macrophages [1]. It is therefore conceivable that an ineffective mesangial phagocytosis, e.g. secondary to *Itga8* deficiency, could result in a retardation of glomerular healing as seen in *Itga8*-deficient mice after induction of acute Habu nephritis [24].

The amount of phagocytosing cells was significantly reduced in *Itga8*-deficient compared to wild type mesangial cells. This was independent of the ingested particles. A deficiency in *Itga8* led to reduced phagocytosis of apoptotic mesangial cells, apoptotic inflammatory cells and matrix components. Thus, *Itga8* seems to play an important role for phagocytosis by mesangial cells. However, there is still a considerable amount of *Itga8*-deficient mesangial cells capable of engulfing and ingesting particles, which implies that there are additional factors conveying phagocytosis by mesangial cells. The RGD-binding  $\alpha V$  integrin is abundant on mesangial cells. This integrin is known to be critically involved in the phagocytosis of apoptotic cells by mesangial cells [20]. The inhibition of  $\alpha V$  integrin in *Itga8*-deficient cells, however, did not significantly reduce phagocytosis of coated beads by *Itga8*-deficient mesangial cells, arguing against a decisive contribution of  $\alpha V$  integrin to phagocytosis by *Itga8*-deficient mesangial cells. Previous studies have shown that *Itga8*-deficient mesangial cells express more  $\alpha 2$  and  $\alpha 6$  integrins than wild type mesangial cells [24, 44]. A contribution to collagen internalization was described for  $\alpha 2\beta 1$  integrin in fibroblasts [18]. Therefore, we analyzed the phagocytosis capacity of *Itga8*-deficient mesangial cells after blocking of  $\alpha 2$  and  $\alpha 6$  integrins. Although blocking of  $\alpha 2$  and  $\alpha 6$  integrins readily reduced the ingestion of matrix-coated beads by wild type mesangial cells, it had no effect on phagocytosis by *Itga8*-deficient mesangial cells.

Moreover, the reduction of the rate of phagocytosis in mesangial cells lacking *Itga8* seemed to be independent from the matrix protein used for coating. Fibronectin is a ligand for *Itga8*, while collagen I is not, but ingestion of both matrix components was reduced in *Itga8*-deficient mesangial cells. This argues against an effect mediated by direct integrin receptor-ligand binding.

In line with this observation, phagocytosis by both wild type and *Itga8*-deficient mesangial cells is strongly dependent on a functional cytoskeleton. Actin rearrangement is a crucial prerequisite for the uptake of particles and apoptotic cells by phagocytes [45]. Thus, inhibition of actin cytoskeleton rearrangement results in inhibition of phagocytosis. As important regulators of actin rearrangement Rho GTPases are effective modulators of phagocytosis in various cell types [45]. Rac1 is a Rho GTPase, which is involved in cell motility and other cell functions by controlling the function of lamellipodia. Activation of Rac1 is important for the engulfment of apoptotic cells by the phagocytes' lamellipodia [46]. Similarly, the Rho-associated protein kinase (ROCK) is able to support phagocytosis as described in macrophages [47]. In accordance with these data, inhibition of Rac1 or ROCK attenuated phagocytosis of wild type mesangial cells, while in *Itga8*-deficient cells inhibition of Rac1 or ROCK was less effective on phagocytosis. This further argues for *Itga8*-mediated differences in the organization of the cytoskeleton in wild type and *Itga8*-deficient mesangial cells [22]. Phagocytosis by *Itga8*-deficient mesangial cells is less effective, but is also less dependent on the ability of the cell to perform actin rearrangements. While wild type mesangial cells show the typical mesenchymal phenotype, *Itga8*-deficient mesangial cells are characterized by a

more epithelial rather than a mesenchymal phenotype [22]. This observation was also made in mesangial cells with a transient knock-down of *Itga8* which similarly results in a loss of the fibroblast-like phenotype with a different, more compact arrangement of actin filaments close to the cell membrane. In mesangial cells with a transient knock-down of *Itga8*, however, the actin cytoskeleton changes are somewhat milder than in mesangial cells with a genetic *itga8* knock-out, which might be due to an incomplete blockade of *Itga8* expression. Thus, the process of phagocytosis might be somewhat different from wild type mesangial cells. In mammary epithelial cells for instance, E-cadherin seems to contribute to phagocytosis [48]. This is in contrast to the phagocytosis in mesenchymal cells, where E-cadherin is commonly not found. Moreover, changes of the cell shape including cell rounding can result in reduced phagocytosis capacity as described in macrophages [49-51].

## Conclusion

Taken together, we show that a loss of *Itga8* leads to a reduced phagocytosis capacity of matrix coated beads and of apoptotic cells by mesangial cells. Thus, *Itga8* seems to contribute to an effective mesangial cell phagocytosis *in vitro*. This does not seem to be due to a direct involvement of *Itga8* in the process of phagocytosis. Instead, reduced phagocytosis might arise from a change in cytoskeletal organization, secondary to the lack of *Itga8*. We speculate that the attenuation of phagocytosis as a consequence of an underexpression of *Itga8* could contribute to the observed retardation in glomerular regeneration after renal damage in a model of mesangioproliferative glomerulonephritis in *Itga8*-deficient mice. However, our study is limited in that the significance of our *in vitro* findings for the *in vivo* situation remains to be shown.

## Acknowledgements

This study was supported by a grant from the ELAN fund, University Hospital of Erlangen and an educational grant from the Interdisciplinary Center for Clinical Research of the Faculty of Medicine of the Friedrich-Alexander University Erlangen-Nürnberg to IM and by a grant from the Deutsche Forschungsgemeinschaft SFB 423 (A2) to AH. The authors thank Dr. Ulrich Muller (Scripps Institute, LaJolla, USA) for providing the *Itga8*-deficient mouse strain and Prof. Ben Wielockx and Dr. Joanna Kalucka (Dresden, Germany) for providing the stably GFP-transfected RAW cells. The authors thank Mirosława Kupraszewicz-Hutzler for technical assistance.

We acknowledge support by Deutsche Forschungsgemeinschaft and Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU) within the funding programme Open Access Publishing.

## Disclosure Statement

The authors declare that they have no competing interests.

## References

- 1 Watson S, Cailhier JF, Hughes J, Savill J: Apoptosis and glomerulonephritis. *Curr Dir Autoimmun* 2006;9:188-204.
- 2 Sayedyahosseini S, Dagnino L: Integrins and small GTPases as modulators of phagocytosis. *Int Rev Cell Mol Biol* 2013;302:321-354.
- 3 Goodridge HS, Underhill DM, Touret N: Mechanisms of Fc receptor and dectin-1 activation for phagocytosis. *Traffic* 2012;13:1062-1071.
- 4 Flannagan RS, Jaumouille V, Grinstein S: The cell biology of phagocytosis. *Ann Rev Pathol* 2012;7:61-98.



- 5 Baker AJ, Mooney A, Hughes J, Lombardi D, Johnson RJ, Savill J: Mesangial cell apoptosis: the major mechanism for resolution of glomerular hypercellularity in experimental mesangial proliferative nephritis. *J Clin Invest* 1994;94:2105-2116.
- 6 Krieser RJ, White K: Engulfment mechanism of apoptotic cells. *Curr Opin Cell Biol* 2002;14:734-738.
- 7 Underhill DM, Ozinsky A: Phagocytosis of microbes: complexity in action. *Ann Rev Immunol* 2002;20:825-852.
- 8 Lauber K, Blumenthal SG, Waibel M, Wesselborg S: Clearance of apoptotic cells: getting rid of the corpses. *Mol Cell* 2004;14:277-287.
- 9 Hoppe AD, Swanson JA: Cdc42, Rac1, and Rac2 display distinct patterns of activation during phagocytosis. *Mol Biol Cell* 2004;15:3509-3519.
- 10 Dupuy AG, Caron E: Integrin-dependent phagocytosis: spreading from microadhesion to new concepts. *J Cell Sci* 2008;121:1773-1783.
- 11 Sayedyahosseini S, Nini L, Irvine TS, Dagnino L: Essential role of integrin-linked kinase in regulation of phagocytosis in keratinocytes. *FASEB J* 2012;26:4218-4229.
- 12 Olazabal IM, Caron E, May RC, Schilling K, Knecht DA, Machesky LM: Rho-kinase and myosin-II control phagocytic cup formation during CR, but not FcγR, phagocytosis. *Curr Biol* 2002;12:1413-1418.
- 13 Albert ML, Kim JI, Birge RB: alphavbeta5 integrin recruits the CrkII-Dock180-rac1 complex for phagocytosis of apoptotic cells. *Nat Cell Biol* 2000;2:899-905.
- 14 Mao Y, Finnemann SC: Essential diurnal Rac1 activation during retinal phagocytosis requires alphavbeta5 integrin but not tyrosine kinases focal adhesion kinase or Mer tyrosine kinase. *Mol Biol Cell* 2012;23:1104-1114.
- 15 Gagen D, Filla MS, Clark R, Liton P, Peters DM: Activated alphavbeta3 integrin regulates alphavbeta5 integrin-mediated phagocytosis in trabecular meshwork cells. *Invest Ophthalmol Vis Sci* 2013;54:5000-5011.
- 16 Coopman PJ, Thomas DM, Gehlsen KR, Mueller SC: Integrin alpha 3 beta 1 participates in the phagocytosis of extracellular matrix molecules by human breast cancer cells. *Mol Biol Cell* 1996;7:1789-1804.
- 17 Koenigsknecht J, Landreth G: Microglial phagocytosis of fibrillar beta-amyloid through a beta1 integrin-dependent mechanism. *J Neurosci* 2004;24:9838-9846.
- 18 Arora PD, Marignani PA, McCulloch CA: Collagen phagocytosis is regulated by the guanine nucleotide exchange factor Vav2. *Am J Physiol Cell Physiol* 2008;295:C130-137.
- 19 Arora PD, Manolson MF, Downey GP, Sodek J, McCulloch CA: A novel model system for characterization of phagosomal maturation, acidification, and intracellular collagen degradation in fibroblasts. *J Biol Chem* 2000;275:35432-35441.
- 20 Cortes-Hernandez J, Fossati-Jimack L, Carugati A, Potter PK, Walport MJ, Cook HT, Botto M: Murine glomerular mesangial cell uptake of apoptotic cells is inefficient and involves serum-mediated but complement-independent mechanisms. *Clin Exp Immunol* 2002;130:459-466.
- 21 Hughes J, Liu Y, Van Damme J, Savill J: Human glomerular mesangial cell phagocytosis of apoptotic neutrophils: mediation by a novel CD36-independent vitronectin receptor/thrombospondin recognition mechanism that is uncoupled from chemokine secretion. *J Immunol* 1997;158:4389-4397.
- 22 Marek I, Volkert G, Jahn A, Fahlbusch F, Zurn C, Ozcan Z, Goppelt-Strube M, Hilgers KF, Rascher W, Hartner A: Lack of alpha8 integrin leads to morphological changes in renal mesangial cells, but not in vascular smooth muscle cells. *BMC Cell Biol* 2010;11:102.
- 23 Bieritz B, Spessotto P, Colombatti A, Jahn A, Prols F, Hartner A: Role of alpha8 integrin in mesangial cell adhesion, migration, and proliferation. *Kidney Int* 2003;64:119-127.
- 24 Hartner A, Marek I, Cordasic N, Haas C, Schocklmann H, Hulsmann-Volkert G, Plasa I, Rascher W, Hilgers KF, Amann K: Glomerular regeneration is delayed in nephritic alpha 8-integrin-deficient mice: contribution of alpha 8-integrin to the regulation of mesangial cell apoptosis. *Am J Nephrol* 2008;28:168-178.
- 25 Hartner A, Schocklmann H, Prols F, Muller U, Sterzel RB: Alpha8 integrin in glomerular mesangial cells and in experimental glomerulonephritis. *Kidney Int* 1999;56:1468-1480.
- 26 Volkert G, Jahn A, Dinkel C, Fahlbusch F, Zurn C, Hilgers KF, Rascher W, Hartner A, Marek I: Contribution of the alpha8 integrin chain to the expression of extracellular matrix components. *Cell Commun Adhes* 2014;21:89-98.
- 27 Mamlouk S, Kalucka J, Singh RP, Franke K, Muschter A, Langer A, Jakob C, Gassmann M, Baretton GB, Wielockx B: Loss of prolyl hydroxylase-2 in myeloid cells and T-lymphocytes impairs tumor development. *Int J Cancer* 2014;134:849-858.

- 28 McCulloch CA, Knowles GC: Deficiencies in collagen phagocytosis by human fibroblasts *in vitro*: a mechanism for fibrosis? J Cell Physiol 1993;155:461-471.
- 29 Knowles GC, McKeown M, Sodek J, McCulloch CA: Mechanism of collagen phagocytosis by human gingival fibroblasts: importance of collagen structure in cell recognition and internalization. J Cell Sci 1991;98:551-558.
- 30 van der Pauw MT, Van den Bos T, Everts V, Beertsen W: Phagocytosis of fibronectin and collagens type I, III, and V by human gingival and periodontal ligament fibroblasts *in vitro*. J Periodontol 2001;72:1340-1347.
- 31 Segal G, Lee W, Arora PD, McKee M, Downey G, McCulloch CA: Involvement of actin filaments and integrins in the binding step in collagen phagocytosis by human fibroblasts. J Cell Sci 2001;114:119-129.
- 32 Cosio FG, Sedmak DD, Nahman NS, Jr.: Cellular receptors for matrix proteins in normal human kidney and human mesangial cells. Kidney Int 1990;38:886-895.
- 33 Blystone SD, Graham IL, Lindberg FP, Brown EJ: Integrin alpha v beta 3 differentially regulates adhesive and phagocytic functions of the fibronectin receptor alpha 5 beta 1 J Cell Biol 1994;127:1129-1137.
- 34 Zhao MW, Jin ML, He S, Spee C, Ryan SJ, Hinton DR: A distinct integrin-mediated phagocytic pathway for extracellular matrix remodeling by RPE cells. Invest Ophthalmol Vis Sci 1999;40:2713-2723.
- 35 McKeown M, Knowles G, McCulloch CA: Role of the cellular attachment domain of fibronectin in the phagocytosis of beads by human gingival fibroblasts *in vitro*. Cell Tissue Res 1990;262:523-530.
- 36 Gu BJ, Saunders BM, Jursik C, Wiley JS: The P2X7-nonmuscle myosin membrane complex regulates phagocytosis of nonopsonized particles and bacteria by a pathway attenuated by extracellular ATP. Blood 2010;115:1621-1631.
- 37 Gu BJ, Saunders BM, Petrou S, Wiley JS: P2X(7) is a scavenger receptor for apoptotic cells in the absence of its ligand, extracellular ATP. J Immunol 2011;187:2365-2375.
- 38 Savill J, Smith J, Sarraf C, Ren Y, Abbott F, Rees A: Glomerular mesangial cells and inflammatory macrophages ingest neutrophils undergoing apoptosis. Kidney Int 1992;42:924-936.
- 39 Amann K, Haas CS, Schussler J, Daniel C, Hartner A, Schocklmann HO: Beneficial effects of integrin alphavbeta3-blocking RGD peptides in early but not late phase of experimental glomerulonephritis. Nephrol Dial Transplant 2012;27:1755-1768.
- 40 Takahashi N, Kobayashi M, Takaki T, Takano K, Miyata M, Okamatsu Y, Hasegawa K, Nishihara T, Yamamoto M: Actinobacillus actinomycetemcomitans lipopolysaccharide stimulates collagen phagocytosis by human gingival fibroblasts. Oral Microbiol Immunol 2008;23:259-264.
- 41 Tiwari V, Shukla D: Nonprofessional phagocytosis can facilitate herpesvirus entry into ocular cells. Clin Dev Immunol 2012;2012:651691.
- 42 Rojas AM, Fuentes G, Rausell A, Valencia A: The Ras protein superfamily: evolutionary tree and role of conserved amino acids. J Cell Biol 2012;196:189-201.
- 43 Levay M, Krobert KA, Wittig K, Voigt N, Bermudez M, Wolber G, Dobrev D, Levy FO, Wieland T: NSC23766, a widely used inhibitor of Rac1 activation, additionally acts as a competitive antagonist at muscarinic acetylcholine receptors. J Pharmacol Exp Therapeut 2013;347:69-79.
- 44 Haas CS, Amann K, Schittny J, Blaser B, Muller U, Hartner A: Glomerular and renal vascular structural changes in alpha8 integrin-deficient mice. J Am Soc Nephrol 2003;14:2288-2296.
- 45 Mao Y, Finnemann SC: Regulation of phagocytosis by Rho GTPases. Small GTPases 2015;6:89-99.
- 46 Nakaya M, Kitano M, Matsuda M, Nagata S: Spatiotemporal activation of Rac1 for engulfment of apoptotic cells. Proc Nat Acad Sci USA 2008;105:9198-9203.
- 47 Liu Y, Tejpal N, You J, Li XC, Ghobrial RM, Kloc M: ROCK inhibition impedes macrophage polarity and functions. Cell Immunol 2016;300:54-62.
- 48 Fornetti J, Flanders KC, Henson PM, Tan AC, Borges VF, Schedin P: Mammary epithelial cell phagocytosis downstream of TGF-beta3 is characterized by adherens junction reorganization. Cell Death Diff 2016;23:185-196.
- 49 Akei H, Whitsett JA, Buroker M, Ninomiya T, Tatsumi H, Weaver TE, Ikegami M: Surface tension influences cell shape and phagocytosis in alveolar macrophages. Am J Physiol Lung Cell Mol Physiol 2006;291:L572-579.
- 50 Haussler S, Rohde M, von Neuhoff N, Nimtz M, Steinmetz I: Structural and functional cellular changes induced by Burkholderia pseudomallei rhamnolipid. Infect Immun 2003;71:2970-2975.
- 51 Rossi AG, McCutcheon JC, Roy N, Chilvers ER, Haslett C, Dransfield I: Regulation of macrophage phagocytosis of apoptotic cells by cAMP. J Immunol 1998;160:3562-3568.