

## The potential of prediction models of functioning remains to be fully exploited: a scoping review in the field of spinal cord injury rehabilitation

Jsabel Hodel, Gerold Stucki, Birgit Prodingen

### Angaben zur Veröffentlichung / Publication details:

Hodel, Jsabel, Gerold Stucki, and Birgit Prodingen. 2021. "The potential of prediction models of functioning remains to be fully exploited: a scoping review in the field of spinal cord injury rehabilitation." *Journal of Clinical Epidemiology* 139: 177–90.  
<https://doi.org/10.1016/j.jclinepi.2021.07.015>.

**REVIEW**

# The potential of prediction models of functioning remains to be fully exploited: A scoping review in the field of spinal cord injury rehabilitation

Jsabel Hodel<sup>a,b,\*</sup>, Gerold Stucki<sup>a,c</sup>, Birgit Prodinger<sup>a,b,d</sup><sup>a</sup>Swiss Paraplegic Research, Guido A. Zäch Strasse 4, 6207 Nottwil, Switzerland<sup>b</sup>Department of Health Sciences and Medicine, University of Lucerne, Frohburgstrasse 3, 6002 Lucerne, Switzerland<sup>c</sup>Center for Rehabilitation in Global Health Systems, Department of Health Sciences and Medicine, University of Lucerne, Frohburgstrasse 3, 6002 Lucerne, Switzerland<sup>d</sup>Faculty of Applied Health and Social Sciences, Technical University of Applied Sciences Rosenheim, Hochschulstrasse 1, 83024 Rosenheim, Germany

Accepted 22 July 2021; Available online 27 July 2021

**Abstract****Objective:** The study aimed to explore existing prediction models of functioning in spinal cord injury (SCI).**Study Design and Setting:** The databases *PubMed*, *EBSCOhost* *CINAHL Complete*, and *IEEE Xplore* were searched for relevant literature. The search strategy included published search filters for prediction model and impact studies, index terms and keywords for SCI, and relevant outcome measures able to assess functioning as reflected in the International Classification of Functioning, Disability and Health (ICF). The search was completed in October 2020.**Results:** We identified seven prediction model studies reporting twelve prediction models of functioning. The identified prediction models were mainly envisioned to be used for rehabilitation planning, however, also other possible applications were stated. The method predominantly used was regression analysis and the investigated predictors covered mainly the ICF components of *body functions* and *activities and participation*, next to characteristics of the health condition and health interventions.**Conclusion:** Findings suggest that the development of prediction models of functioning for use in clinical practice remains to be fully exploited. By providing a comprehensive overview of what has been done, this review informs future research on prediction models of functioning in SCI and contributes to an efficient use of research evidence. © 2021 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)**Keywords:** Spinal Cord Injuries; Rehabilitation; Prognosis; Diagnosis; Clinical Decision Rules; Forecasting**1. Introduction**

Spinal cord injury (SCI) is a chronic health condition devastatingly affecting a person's life in a variety of ways. The structural damage to the spinal cord and the resulting loss of neurologic functions adversely affects the ability of a person to perform simple and complex activities and to participate in community and major life areas [1]. After

the injury, persons with SCI go through an extensive rehabilitation process to live independently with the health condition: from intensive care and inpatient rehabilitation to outpatient specialized care after returning to the community. The World Health Organization (WHO) refers to the lived experience of a health condition as 'functioning' [2]. The concept of functioning, as described in WHO's International Classification of Functioning, Disability and

Competing interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Presentation of this material: None.

Funding: The study was funded by the Swiss Paraplegic Research.

Ethics approval: Not applicable.

Author contributions: **JH:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Software, Validation, Visualization, Writing - original draft, Writing - review & editing. **GS:** Conceptualization, Funding acquisition, Supervision, Writing - review & editing. **BP:** Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Resources, Supervision, Validation, Writing - review & editing.

\* Correspondence: Tel.: +41 41 939 66 32; Fax: +41 41 939 66 40

E-mail address: [jsabel.hodel@paraplegie.ch](mailto:jsabel.hodel@paraplegie.ch) (J. Hodel).

Health (ICF), includes different components – *body functions* and *body structures* as well as *activities and participation* – which interact with each other and are outcomes of the interaction between a health condition and *environmental* and *personal contextual factors*. Against this background, the objective of rehabilitation after SCI can be formulated as the optimization and maintenance of a person's functioning [3]. In order to achieve this objective, comprehensive and relevant functioning information is essential to guide rehabilitation planning and management, individual clinical care and decision making.

Prediction research aims to enhance individual health and health care practice by investigating and improving the diagnosis or prognosis of a specific health condition [4–6]. For the purpose of this review, roughly three types of prediction research can be distinguished: [4,7] (1) predictor finding studies, (2) prediction model studies, and (3) impact studies. Predictor finding studies generally aim to explore or identify which variables within a set of candidate predictors are independently associated with a specific outcome. Prediction model studies aim to develop and/or externally validate (with or without updating) a multivariable prediction model for use in medical or clinical practice. Impact studies build on a developed and validated prediction model and aim to assess the impact of the use of such a model in a specific context or setting compared to not using it. Prediction model development, validation and impact studies correspond with the phases, which prediction models for use in practice usually have to undergo in their development process [8–11]. The development of prediction models has gained increasing attention by the recognition of evidence-based health care and the uptake of new statistical methods in the health sciences and clinical epidemiology.

In rehabilitation research, the role of functioning as key health indicator complementing mortality and morbidity [12] poses the question of how prediction research, and specifically prediction models, can improve the use of functioning information for practice. In SCI literature, various efforts have been undertaken to develop and/or validate prediction models for outcomes related to specific aspects of functioning, such as ambulation, [13–20] or bladder and bowel outcomes [21–23]. Predictor finding studies for several functioning outcomes have already been reviewed and synthesized [24–27]. What remains to be investigated is how functioning, as a multidimensional concept is reflected in current prediction models across the corresponding development phases depicted by development, validation and impact studies in the field of SCI rehabilitation. Therefore, the objective of this scoping review is to explore existing prediction models of functioning in SCI. Specifically, the review aims to (1) identify prediction models of functioning in SCI, (2) examine their content by using the ICF as a reference language, (3) examine their use from a systems perspective, and (4) document which methods were used to develop them. The

scoping review will shed light on current research gaps as well as on promising directions for future developments and improvements of prediction models of functioning for SCI.

## What is new?

### Key findings

- Identification of seven prediction model studies reporting twelve prediction models of functioning in SCI; no impact study was identified.

### What this adds to what is known?

- The development of prediction models of functioning in SCI is still in its infancy. This review highlights potential future directions in the development of prediction models in the field of SCI rehabilitation with regards to content, use and methods.

### What is the implication, what should change now?

- Functioning, as outcome of the identified models, was measured with the FIM<sup>TM</sup> or the SCIM. The investigated predictors covered mainly body functions, activities and participation, characteristics of the health condition or health interventions. The integration of a broad range of potential predictors including imaging, biomarkers, and genetics, as well as predictors covering body structures and contextual factors remains to be investigated.
- The method predominantly used was linear regression analysis. The application and usefulness of other methods such as machine learning techniques need to be further investigated and its potential merit compared to current methods.
- The identified prediction models were intended to be used for guidance in rehabilitation planning, patient counselling, financial aspects related to the reduction of costs by guided management strategies, and improvements in clinical trial designs. To delineate the value of prediction models for the field of SCI rehabilitation in detail, further research is needed related to validation and impact assessment of prediction models.

## 2. Methods

The scoping review followed the methodological framework of Arksey and O'Malley [28] and incorporate recent experiences of the application of the framework [29–31] as well as the guidance for the conduction of systematic scoping reviews developed by Peters *et al.* [32]. An unpublished review protocol was developed and agreed upon by all authors prior to conducting the review and is avail-

able from the authors on request. The reporting followed the *Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews* [33] and the corresponding checklist can be found in the Supplemental Table 1.

### 2.1. Searching for relevant literature

The following three databases were searched for relevant literature: *PubMed*, [34] *EBSCOhost CINAHL Complete*, [35] and *IEEE Xplore* [36]. The databases were chosen to cover literature from a broad spectrum of rehabilitation research topics including clinical and biomedical sciences, nursing and allied health, as well as biomechanical and engineering sciences. We did not explicitly search for grey literature.

The search strategy was defined in an iterative fashion [37] and included the following components: 1) The *Haynes Broad Search Strategy* for prediction studies, [38] which is available on PubMed via the search filters for "Clinical Queries", 2) an update to the strategy in step one in the form of the *Teljeur/Murphy Inclusion Filter* introduced by Keogh *et al.* [39] and adapted by the authors of this study, 3) index terms and keywords for SCI, and 4) relevant outcome measures able to assess the lived experience of health in persons with SCI as operationalized by functioning. The latter were identified by the development of an initial list based on literature [40–50] and feedback by scholars in the field about the most important measures to consider, given the scope of this study. Included languages were German and English, no limits were chosen with regards to the publication date. The search strategy was developed using PubMed and afterwards translated and adapted to the particularities of the identified other databases. The full search strategy for all databases can be found in the Supplemental Table 2. The search was completed on October 12th 2020.

### 2.2. Study selection

Eligibility was formulated according to in- and exclusion criteria for title/abstract and full-text screening separately (see Table 1). Underlying the eligibility criteria are the different types of prediction research explained in the introduction. Prediction models are thereby understood as “tools that combine multiple predictors by assigning relative weights to each predictor to obtain a risk or probability” [5]. Other notions include (clinical) prediction rules, probability assessments, decision rules or risk scores. In accordance with the objective of this review, only models were included that predicted functioning: Outcome variables included in the studies had to reflect different domains of functioning (classified as chapters in the ICF), but at least two chapters of *activities and participation*. Published conference proceedings in the biomechanical and

engineering sciences were considered as original publications.

After database searching and removing of duplicates, [51] we followed the approach applied by Maritz *et al.* [52,53] for screening of titles and abstracts. A random sample incorporating 50 articles of the records were screened independently by two reviewers (JH, BP) in light of the eligibility criteria to determine whether an article is relevant. If the agreement in decisions for article in- or exclusion of the reviewers was acceptable (>90%), one reviewer continued to screen the remaining articles (JH). Otherwise, a new random sample of the same size was screened independently by the two reviewers. Disagreement was solved by discussions and the procedure was repeated until an acceptable agreement was reached.

Before starting the full-text screening, the eligibility criteria were revisited and further detailed by the study team. Subsequently, full-texts were screened by one reviewer (JH) and in the case of ambiguity, discussed with a second reviewer (BP). After full-text screening, an additional hand search was conducted. The database findings, screening and references were organized with EndNote [54].

### 2.3. Data extraction and results charting

The extraction fields presented by Peters *et al.* [32] were entered into a Microsoft Excel sheet and complemented by elements of the checklists for *Critical Appraisal and Data Extraction for Systematic Reviews of Prediction Modelling Studies* (CHARMS) [55] and *Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis* (TRIPOD) [5] in order to document the identified prediction models of functioning in SCI (see Supplemental Table 3).

To examine the content of identified prediction models, the established linking method developed by Cieza *et al.* [56] was applied. This method allows to link the content of outcomes or predictors included in the respective prediction models to the ICF as a reference model, and thus enables the comparison of outcomes and predictors contained in different prediction models. The linking process entails the linking at the conceptual and the classification level. For the purpose of this review, outcomes and predictors reported in the identified studies were extracted and linked if possible at chapter-level of the ICF. The ICF Research Branch (<https://www.icf-research-branch.org>) was contacted to request existing linking results of specific outcomes and predictors. To examine the envisioned use and implications of the identified prediction models, micro (patient-provider interaction), meso (service provision and payment) and macro (policies and programs) system levels were used as framework of reference. To document the methods used to develop the identified prediction models, the respective author's description used within the article were extracted together with the stated

**Table 1.** Eligibility criteria according to title/abstract screening and full-text screening**Inclusion and exclusion criteria for title/abstract screening***Inclusion criteria:*

- Primary study
- Prediction model study or impact study
- Study includes at least one variable (predictor and/or outcome) assessed with a measure of the lived experience of health as operationalised by functioning, which reflects two or more chapters of activities and participation as described in the ICF
- Study population includes males and/or females with SCI (traumatic and/or non-traumatic)
- Publication language is English or German

*Exclusion criteria:*

- Animal study
- Paediatric study
- Predictor finding study
- Prediction model study or impact study with mixed-diagnosis populations
- Study population includes SCI as a complication
- Study includes mortality as solely outcome

**Inclusion and exclusion criteria for full-text screening***Inclusion criteria:*

- Study includes measure of functioning as outcome variable

*Exclusion criteria:*

- Study includes measure of functioning as predictor variable only
- Study includes as outcome variable only single items or subscales of a measure of functioning, which no longer reflect two or more chapters of activities and participation as described in the ICF
- Study with outcome assessed/evaluated within the acute rehabilitation setting

Abbreviations. ICF, International Classification of Functioning, Disability and Health; SCI, spinal cord injury.

argumentation for its use, as well as stated advantages and disadvantages.

The data extraction was performed by one reviewer (JH) and cross-checked by a second reviewer (BP). The results of the scoping review were arranged in tabular format and discussed narratively.

### 3. Results

#### 3.1. Study identification

In total, 2378 articles were retrieved through database searching and after screening the titles and abstracts of 1851 articles and the full-texts of 234 articles, seven eligible studies were identified for inclusion in the scoping review [57–63]. The corresponding flow diagram of the screening process is presented in Figure 1.

#### 3.2. Screening and study selection process

For the title and abstract screening, in total three random samples were screened independently by the two reviewers until acceptable agreement was reached. The specific agreement levels reached for each sample were 78%, 86%, and 94%, respectively. Main reason for disagreement was the challenging distinction between predictor finding studies and prediction model studies. Following the framework of Kent *et al.* [7] the distinction should be based on the study aim. However, often authors did not clearly state the study aim, which was also reported by authors who

conducted reviews on prediction models previously [4]. If a study aim was not clearly stated or unsure, studies were nevertheless included for full-text screening if they described a functioning outcome, or mentioned some form of model performance or accuracy assessment.

As the eligibility criteria for the full-text screening were revisited, for prediction model development studies the criteria, that studies need to include an internal validation of the prediction models to be eligible for this review, was decided. This decision was based on the recommendation of the TRIPOD statement for prediction model development studies to include some form of internal validation. In addition, this decision enhanced the consistency in the distinction between prediction model and predictor finding studies.

In the hand search we applied the following criteria: 1) publications based on identified SCI cohorts, trials or research projects (European Multicenter Study about Spinal Cord Injury, Rick Hansen Spinal Cord Injury Registry, Spinal Cord Injury Model System, SCIRehab) were specifically searched for in *PubMed*, and 2) the identified eligible studies were checked for updates using the 'Cited-by'-function of *PubMed*.

#### 3.3. Characteristics of the included studies

The basic characteristics of the seven included prediction model studies are shown in Table 2. Six studies [57–60,62,63] described model development and included inter-

**Table 2.** Overview of included prediction model studies.

Study	Population												Location	Data handling	Modelling		
Authors	Sample size	Mean age (SD) in years	Sex (%)		Aetiology (%)		Level of injury (%)		Severity of injury according to AIS grade (%) <sup>a</sup>				Country, centres	Approaches to handle missing observations	Methods	Predictor selection procedure	Validation approach
			Male	Female	Traumatic	Non-traumatic	Para-plegia (T1-S5)	Tetra-plegia (C1-C8)	A	B	C	D					
Ariji et al.	137	60 (16)	80	20	100	0	17	83	36	14	32	18	Japan, singe-centre	complete case analysis	linear regression	backward stepwise	internal, bootstrap
Facchinello et al.	172	49 (18)	NA	NA	100	0	34 <sup>b</sup>	66 <sup>c</sup>	40	10	15	36	Canada, single-centre	complete case analysis	machine learning	literature	internal, cross-validation
Harrington et al.	417	56 ±28 <sup>d</sup>	66	31	75	NA	40	57	25	11	35	26	UK, single-centre	median imputation, LOCF, NOCB	linear regression, generalized linear regression	significance, elastic net penaliza-tion	internal, cross-validation
Kaminski et al.	76	43 (18)	76	24	100	0	54	46	53	11	9	27	Canada, single-centre	multiple imputation analysis	linear regression	forward stepwise	internal, bootstrap
Tomioaka et al.	31	59 (19)	87	13	100	0	16	84	19	3	52	26	Japan, single-centre	no missing observations reported	logarithmic equation	not applicable	external, extrapolation
Wilson et al.	376	43 (17)	78	NA	100	0	NA	NA	36	17	15	32	Canada/USA, multi-centre	multiple imputation analysis	linear regression, logistic regression	no selection procedure performed	internal, bootstrap
Zariffa et al.	14	44 (18)	93	7	100	0	0	100	NA	NA	NA	NA	Canada/ Switzerland, multi-centre	no missing observations reported	linear regression	cross-validation	internal, cross-validation

Abbreviations. AIS, American Spinal Injury Association Impairment Scale; LOCF, last observation carried forwards; NA, not available; NOCB, next observation carried backwards; SD, standard deviation; UK, United Kingdom; USA, United States of America.


Note. Estimates and percentages have been rounded to zero decimal places for the purpose of this review.

<sup>a</sup> If AIS grade was reported at several time points, the earliest was chosen for this overview;

<sup>b</sup> Paraplegia: T2-L2;

<sup>c</sup> Tetraplegia: C1-T1;

<sup>d</sup> Median ± interquartile range.



**Figure 1.** Flow diagram of the scoping review. Note that the reasons for full-text exclusion are not mutually exclusive. Figure adapted from Moher *et al.* 2009 [64].

nal validation approaches either based on cross-validation or bootstrap procedure, one study [61] described an external validation of a prediction model originally developed in stroke [65] and extrapolated to SCI. Only two studies included data from multiple institutions [62,63]. The mean age of the study populations under investigation ranged from 43 (SD=18) to 60 (SD=16) years and the population samples focused on traumatic aetiology and tend to include predominantly men and persons with tetraplegia. No impact studies were found.

In total, the seven included articles described 12 prediction models of functioning. Table 3 shows the identified models, their specific outcomes, investigated predictors and the corresponding linking to the ICF. The functioning outcome variables used in the prediction models all related to the two instruments Spinal Cord Independence Measure (SCIM) and Functional Independence Measure (FIM<sup>TM</sup>), which both are assessing functional independence of a person in daily life, specifically focusing on self-care, mobility, and bladder and bowel management. The time scope for prediction ranged up to one year after

injury. Predictors were assessed during early acute phase and up to one month after injury. Investigated predictor variables described concepts covered by the ICF components *body functions*, and *activities and participation*. Predictors that could not be linked to the ICF mainly described characteristics of the health condition or health interventions. With regards to their intended or envisioned use, all prediction models were assigned to the micro system level (e.g. guidance in rehabilitation planning, goal setting and patient care) [57–63] and some also to the meso system level (e.g. determination of appropriate length of stay, diminishing costs by guided management strategies) [60,61]. Some studies explicitly stated in addition a potential application for research purposes (e.g. improving clinical trial designs) [62,63] and for patient counselling (e.g. informing patients and relatives about expectations and relieving from psychological uncertainty) [58–60,62]. The reported statistical methods for the development of the prediction models were mostly regression analyses (linear and logistic), one study reported the use of machine learning methods, specifically regression tree analysis [58].

**Table 3.** Overview of outcome and predictor variables of included prediction model studies

Study	Final model(s)							Linking to ICF components						
Authors	No.	Variable specification						b	s	d	e	pf	nc/nd	
		<i>Outcome</i>	<i>Predictors</i>	<i>Prediction time frame</i>	<i>Measurement time point</i>	Included in final model?								
						1	2	3	4					
Ariji et al.	1	<b>SCIM III, total score</b>	<b>6 months after injury</b>	X	-	-	-	-	X		X			
		Age at injury	NA	X										nd
		ASIA key motor muscle items <sup>a</sup>	1 month after injury	X					X					
		ASIA key sensory point items <sup>b</sup>	1 month after injury						X					
		SCIM III items <sup>c</sup>	1 month after injury	X					X		X			
		WISCI II	1 month after injury	X							X			
Facchinello et al.	2	<b>SCIM III, total score</b>	<b>6/12 MT after injury</b>	X	X	-	-	-	X		X			
		Age at injury	Acute care hospitalization	X	X									nd
		ASIA impairment scale	Acute care hospitalization	X	X				X					nc_hc
		Delay from the injury to surgery	Acute care hospitalization	X										nc_ICHI
		Early spasticity	Acute care hospitalization	X					X					
		Energy associated with injury	Acute care hospitalization	X	X									nc_hc
		ISS	Acute care hospitalization	X										nc_hc
		Mechanism of injury	Acute care hospitalization	X										nc_hc
		Neurological level of the injury	Acute care hospitalization	X	X									nc_hc
		Pneumonia	Acute care hospitalization	X										nc_hc
		Pressure ulcers	Acute care hospitalization	X										nc_hc
		Urinary tract infection	Acute care hospitalization	X										nc_hc
Harrington et al.	4 <sup>d,i</sup>	<b>SCIM III, total score</b>	<b>Discharge</b>	X		X			X		X			
		<b>SCIM III, total score</b>	<b>12 months after injury</b>		X		X		X		X			
		Age at injury	NA			X	X							nd
		ASIA impairment scale, grade B	Rehabilitation admission			X			X					nc_hc
		ASIA impairment scale, grade C	Rehabilitation admission						X					nc_hc
		ASIA impairment scale, grade D	Rehabilitation admission			X			X					nc_hc
		ASIA light touch score	Rehabilitation admission						X					
		ASIA motor score	Rehabilitation admission	X	X	X	X		X					
		ASIA pin prick score	Rehabilitation admission	X	X				X					

(continued on next page)

Table 3 (continued)

Study	Final model(s)						Linking to ICF components						
Authors	No.	Variable specification				b	s	d	e	pf	nc/nd		
		Outcome /Predictors	Prediction time frame /Measurement time point	Included in final model?									
				1	2							3	4
		Alanine transaminase	Time of blood test <sup>e</sup>	X	X			X					
		Albumin	Time of blood test	X				X					
		Alkaline phosphatase	Time of blood test	X				X					
		C-reactive protein	Time of blood test					X					
		Creatinine	Time of blood test	X		X		X					
		Drinking status	NA	X						X			
		Fracture	NA	X							nc_hc		
		Gamma glutamyl transferase	Time of blood test	X				X					
		Hematocrit	Time of blood test					X					
		Hemoglobin	Time of blood test					X					
		Lumbar injury	NA								nc_hc		
		Mean cell hemoglobin	Time of blood test					X					
		Mean cell volume	Time of blood test	X	X			X					
		Monocytes	Time of blood test	X				X					
		Neurological level of injury, traumatic	NA	X							nc_hc		
		Platelets	Time of blood test	X				X					
		Potassium	Time of blood test					X					
		SCIM III, total score	Rehabilitation admission	X	X	X	X	X		X			
		Sex	NA	X	X		X				nd		
		Smoker status known	NA							X			
		Smoker status unknown	NA			X				X			
		Surgery	NA	X							nc_ICHI		
		Time to first blood test	Time of blood test			X	X				nc_ICHI		
		Total bilirubin	Time of blood test					X					
		Total protein	Time of blood test	X				X					
		Type 1 diabetes	NA	X							nc_hc		
		Type 2 diabetes	NA			X					nc_hc		
		Urea	Time of blood test			X		X					
		White blood count	Time of blood test	X				X					
Kaminski et al.	1	<i>SCIM III, total score</i>	<i>12 months follow-up</i>	<b>X</b>	-	-	-	<b>X</b>		<b>X</b>			
		Age	Acute phase after injury								nd		
		ASIA impairment scale	Acute phase after injury	X				X			nc_hc		
		ASIA light touch score	Acute phase after injury	X				X					
		ASIA motor score	Acute phase after injury	X				X					

(continued on next page)

Table 3 (continued)

Study Authors	Final model(s)		Linking to ICF components					
	No.	Variable specification						
			Outcome /Predictors	Prediction time frame /Measurement time point	Included in final model?			
					1	2	3	4
		ASIA pin prick score	Acute phase after injury				X	
		Comorbidity	Acute phase after injury					nc_hc
		Delay to surgery	Acute phase after injury					nc_ICHI
		ISS	Acute phase after injury	X				nc_hc
		Level of injury	Acute phase after injury					nc_hc
		Sex	Acute phase after injury					nd
		TBI	Acute phase after injury					nc_hc
		Type of injury	Acute phase after injury					nc_hc
Tomioka et al.	1	<b>SCIM III, total score</b>	<b>Day X after injury</b>	<b>X</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>X</b>
		SCIM III, total score at day A	First assessment of SCIM III in days after injury <sup>f</sup>	X			X	X
		SCIM III, total score at day B	Third assessment of SCIM III in days after injury <sup>g</sup>	X			X	X
		Day A	First assessment of SCIM III in days after injury	X				nc_ICHI
		Day B	Third assessment of SCIM III in days after injury	X				nc_ICHI
		Day X	Assessment of SCIM X days after injury	X				nc_ICHI
Wilson et al.	2 <sup>h</sup>	<b>FIM<sup>TM</sup>, motor score</b>	<b>6/12 months follow-up</b>	<b>X</b>	<b>X</b>	<b>-</b>	<b>-</b>	<b>X</b>
		Age at injury	NA	X	X			nd
		ASIA impairment scale	Within 3 days after injury	X	X		X	nc_hc
		ASIA motor score	Within 3 days after injury	X	X		X	
		MRI intramedullary signal characteristics	Within 3 days after injury	X	X			nc_hc
Zariffa et al.	1 <sup>d</sup>	<b>SCIM III, total score</b>	<b>Inpatient rehabilitation</b>	<b>X</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>X</b>
		Hand range of motion, x direction	All predictor variables were assessed within two weeks of the SCIM III assessment (before or after)	X			X	

(continued on next page)

Table 3 (continued)

Study Authors	Final model(s)				Linking to ICF components					
	No.	Variable specification			b	s	d	e	pf	nc/nd
		Outcome	Predictors	Prediction time frame Measurement time point						
				Included in final model?						
				1	2	3	4			
		Hand range of motion, y direction						X		
		Hand range of motion, z direction		X				X		
		Joint range of motion, angle 1						X		
		Joint range of motion, angle 2						X		
		Joint range of motion, angle 3						X		
		Joint range of motion, angle 4						X		
		Joint range of motion, angle 5						X		
		Movement mean jerk over task duration						X		
		Movement mean velocity over task duration						X		
		Number of changes in hand's trajectory direction, normalized by task length						X		
		Range of grip pressure		X				X		
		Ratio of mean to maximum velocity over task duration						X		
		Skewness of grip pressure		X				X		

Abbreviations. ASIA, American Spinal Injury Association examination according to the International Standards for Neurological Classification of Spinal Cord Injury; b, body functions; d, activities and participation; e, environmental factors; FIM<sup>TM</sup>, Functional Independence Measure; ICF, International Classification of Functioning, Disability and Health; ISS, Injury Severity Score; MRI, magnetic resonance imaging; nc, not covered in the ICF; nc\_hc, not covered in the ICF, health condition; nc\_ICHI, not covered in the ICF, health intervention (International Classification of Health Interventions); NA, not available; nd, not defined; pf, personal factors (not classified in the ICF); s, body structures; SCIM III, Spinal Cord Independence Measure version three; TBI, traumatic brain injury; WISCI II, Walking Index for Spinal Cord Injury version two.

<sup>a</sup> In total, 20 variables were tested, of which 3 entered the final model;

<sup>b</sup> In total, 112 variables were tested, of which none entered the final model;

<sup>c</sup> In total, 19 variables were tested, of which 1 entered the final model;

<sup>d</sup> Only prediction models of functioning outcomes are reported for the purpose of this review;

<sup>e</sup> Mean time of blood test was 31 days (SD = 30 days) post-injury;

<sup>f</sup> Mean assessment time of SCIM III was 69.8 days (SD = 55.6 days) from admission, and mean time between injury and admission was 45.2 days (SD = 60.8);

<sup>g</sup> Mean assessment time of SCIM III was 123.4 days (SD = 58.2 days) from admission, and mean time between injury and admission was 45.2 days (SD = 60.8);

<sup>h</sup> The two models differ according to the used coding scheme of FIM<sup>TM</sup> and corresponding regression method (discrete score and linear regression model vs. dichotomization according to the achievement of a score of at least 6 in all FIM<sup>TM</sup> motor score items and logistic regression model);

<sup>i</sup> The respective models differ according to the used regression method and predictor selection (linear regression and significance criteria used for models 3 and 4 vs. generalized linear regression and elastic net penalization used for models 1 and 2).

#### 4. Discussion

We identified seven prediction model studies reporting twelve prediction models of functioning. No corresponding impact studies were found. This suggests that the development of prediction models of functioning and their use in practice is not fully exploited. In order to improve prediction models in SCI, it might be helpful to contrast current models with recent suggestions and examples from other health conditions.

All functioning outcome variables used in the identified prediction models related either to SCIM or FIM<sup>TM</sup>. Predictor variables covered the ICF components *body functions* (e.g. assessed by the American Spinal Injury Association examination), and *activities and participation* (e.g. assessed by SCIM). Other predictors described characteristics of the health condition (e.g. level of injury, complications) or of health interventions (e.g. delay to surgery). Only few studies investigated predictors such as blood measures, [59] magnetic resonance imaging, [62] and sensor data [63]. These findings are in line with Wingbermühle *et al.* [66] and Wartenberg *et al.*, [67] which both identified gaps in the investigation of a broad range of possible predictors including biological and physical, as well as psychosocial measures, and especially in the use of directly observable predictors such as imaging, biomarkers, and genetics. In terms of covered ICF components, the integration of *body structures* and *contextual factors* in prediction models remains scarce. Despite the use of the ICF as a frame of reference in the study and the consistency of using FIM<sup>TM</sup> and SCIM as outcomes, the comparability of the findings with regards to selected predictors is limited due to the application of different variable coding schemes such as dichotomized, discrete, or interval scores. Moreover, the comparability of the identified prediction models is further hampered by the heterogeneity of the study populations and settings, as well as by the different time points of predictor and outcome measurements. Further information standards are needed to enhance the interoperability of functioning outcomes or existing standards, such as the ICF or the SCI Data Set actually used in research and practice.

The method most often used in these identified prediction models was linear regression analysis. Only two identified studies were multi-centre studies and the respective population samples focused on traumatic aetiology and tend to include predominantly men and persons with tetraplegia, which limits the generalizability of the developed prediction models. Due to the complex and multidimensional nature of functioning in SCI, prediction models based on new methods such as machine learning techniques are promising and may allow a dynamic and real time modelling of interactions among a variety of predictors [66]. Beyond the findings of our review, also other methods are deployed in SCI prediction research, such as artificial neural network analysis [68] or individual growth

curve models [69]. However, the applicability and usefulness of these methods needs to be further investigated [70]. To do so, large data sets, ideally designed specifically for prediction research, including a broad variety of predictors and appropriately reflecting the population under study are needed [66].

The identified prediction models were intended for clinical purposes including guidance in individual rehabilitation planning, financial aspects related to the reduction of costs by guided management strategies, patient counselling, as well as for research purposes including the improvement of clinical trial designs, which are in line with other prediction research studies in SCI [13,14,22,23,69]. To delineate the value of prediction models for the field of SCI rehabilitation in detail, validation and impact assessment of prediction models require further research.

##### 4.1. Limitations

There are some limitations to this review. Firstly, scoping reviews aim to give an overview of existing evidence on a given topic, regardless of the quality of the reviewed literature [32]. Since we did not assess the quality of the included studies, we are not able to make any statement about the performances, the usefulness or applicability of the presented prediction models for practice. Secondly, the search strategy specifically included common instruments assessing functioning and used in SCI. We do not claim this list to be complete and it might be the case that prediction model studies were missed because their instruments were not included in our search strategy. Thirdly, although our search strategy based on published search filters for prediction model and impact studies, these filters have been shown to low perform for the search of impact studies [71]. Furthermore, despite the absence of relevant impact studies, prediction models of functioning might be developed and implemented locally and not published internationally. Fourthly, the eligibility criteria understand functioning outcomes as variables covering at least two chapters of the ICF component *activities and participation*. Fifthly, the present review only includes prediction model studies which performed at least some kind of internal validation. Although internal validation is strongly recommended in prediction model development, this eligibility criterion lead to the exclusion of studies [69,72,73] about prediction model development which did not intend or failed for some reason to perform an internal validation. Such studies might also include valuable details to inform the development of prediction models in the future. For example, they might include information on potentially important predictor variables to consider in the development of prediction models of functioning, such as different neurophysiological variables as investigated by Hupp *et al.* [72]. Sixthly, we considered conference proceedings from the engineering sciences as original publications. However, these proceedings were often shorter than ordinary journal

articles and thus, provided less information for the full-text screening and the categorization of excluded articles. Lastly, the authors had primarily expertise in the field of health sciences and less so in biomechanical and engineering sciences.

## 5. Conclusion

This scoping review sheds light on existing prediction models of functioning in SCI and highlights their content, use cases, and development methods. Findings suggest that the development of prediction models of functioning for use in clinical practice remains to be fully exploited. However, we believe that SCI with its many different functioning aspects concerned and its life-long perspective and requirement for health and social services across the entire continuum of care is an excellent learning example for the development of prediction models of functioning. By providing a comprehensive overview of what has been done, we hope to inform future research on prediction models of functioning in SCI, including the development of new prediction models for specific purposes or the external validation and improvement of existing ones, and contribute to an efficient and meaningful synthesis and use of research evidence.

## Acknowledgements

The authors would like to thank Jerome Bickenbach for his valuable feedback on the manuscript, and Hildegard Oswald for her support in the search strategy development and its translation between the different databases.

This review is part of the cumulative dissertation of Jsabel Hodel which was conducted within the Swiss National Science Foundation's National Research Programme "Smarter Health Care" (NRP74) and the project "Enhancing continuous quality improvement and supported clinical decision making by standardized reporting of functioning".

## Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:[10.1016/j.jclinepi.2021.07.015](https://doi.org/10.1016/j.jclinepi.2021.07.015).

## References

- [1] World Health Organization. *The International Spinal Cord Society-International Perspectives on Spinal Cord Injury*. Geneva: World Health Organization; 2013.
- [2] World Health Organization. *International Classification of Functioning, Disability and Health (ICF)*. Geneva: World Health Organization; 2001.
- [3] Meyer T, Gutenbrunner C, Bickenbach J, Cieza A, Melvin J, Stucki G. Towards a conceptual description of rehabilitation as a health strategy. *J Rehabil Med* 2011;43(9):765–9. doi:[10.2340/16501977-0865](https://doi.org/10.2340/16501977-0865).
- [4] Bouwmeester W, Zuithoff NP, Mallett S, Geerlings MI, Vergouwe Y, Steyerberg EW, et al. Reporting and methods in clinical prediction research: a systematic review. *PLoS Med* 2012;9(5):1–12. doi:[10.1371/journal.pmed.1001221](https://doi.org/10.1371/journal.pmed.1001221).
- [5] Collins GS, Reitsma JB, Altman DG, Moons KG. Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis (TRIPOD). *Ann Intern Med* 2015;162(10):735–6. doi:[10.7326/L15-5093-2](https://doi.org/10.7326/L15-5093-2).
- [6] Hemingway H, Croft P, Perel P, Hayden JA, Abrams K, Timmis A, et al. Prognosis research strategy (PROGRESS) 1: a framework for researching clinical outcomes. *BMJ* 2013;346:e5595. doi:[10.1136/bmj.e5595](https://doi.org/10.1136/bmj.e5595).
- [7] Kent P, Cancelliere C, Boyle E, Cassidy JD, Kongsted A. A conceptual framework for prognostic research. *BMC Med Res Methodol* 2020;20(1):172. doi:[10.1186/s12874-020-01050-7](https://doi.org/10.1186/s12874-020-01050-7).
- [8] Moons KG, Royston P, Vergouwe Y, Grobbee DE, Altman DG. Prognosis and prognostic research: what, why, and how? *BMJ* 2009;338:b375. doi:[10.1136/bmj.b375](https://doi.org/10.1136/bmj.b375).
- [9] Royston P, Moons KG, Altman DG, Vergouwe Y. Prognosis and prognostic research: Developing a prognostic model. *BMJ* 2009;338:b604. doi:[10.1136/bmj.b604](https://doi.org/10.1136/bmj.b604).
- [10] Altman DG, Vergouwe Y, Royston P, Moons KG. Prognosis and prognostic research: validating a prognostic model. *BMJ* 2009;338:b605. doi:[10.1136/bmj.b605](https://doi.org/10.1136/bmj.b605).
- [11] Moons KG, Altman DG, Vergouwe Y, Royston P. Prognosis and prognostic research: application and impact of prognostic models in clinical practice. *BMJ* 2009;338:b606. doi:[10.1136/bmj.b606](https://doi.org/10.1136/bmj.b606).
- [12] Stucki G, Bickenbach J. Functioning: the third health indicator in the health system and the key indicator for rehabilitation. *Eur J Phys Rehabil Med* 2017;53(1):134–8. doi:[10.23736/S1973-9087.17.04565-8](https://doi.org/10.23736/S1973-9087.17.04565-8).
- [13] Zörner B, Blanckenhorn WU, Dietz V, EM-SCI Study Group, Curt A. Clinical algorithm for improved prediction of ambulation and patient stratification after incomplete spinal cord injury. *J Neurotrauma* 2010;27(1):241–52. doi:[10.1089/neu.2009.0901](https://doi.org/10.1089/neu.2009.0901).
- [14] van Middendorp JJ, Hosman AJ, Donders AR, Pouw MH, Dittunio JF Jr, Curt A, et al. A clinical prediction rule for ambulation outcomes after traumatic spinal cord injury: a longitudinal cohort study. *Lancet* 2011;377(9770):1004–10. doi:[10.1016/S0140-6736\(10\)62276-3](https://doi.org/10.1016/S0140-6736(10)62276-3).
- [15] van Silfhout L, Peters AE, Graco M, Schembri R, Nunn AK, Berlowitz DJ. Validation of the Dutch clinical prediction rule for ambulation outcomes in an inpatient setting following traumatic spinal cord injury. *Spinal Cord* 2016;54(8):614–18. doi:[10.1038/sc.2015.201](https://doi.org/10.1038/sc.2015.201).
- [16] Hicks KE, Zhao Y, Fallah N, Rivers CS, Noonan VK, Plashkes T, et al. A simplified clinical prediction rule for prognosticating independent walking after spinal cord injury: a prospective study from a Canadian multicenter spinal cord injury registry. *Spine J* 2017;17(10):1383–92. doi:[10.1016/j.spinee.2017.05.031](https://doi.org/10.1016/j.spinee.2017.05.031).
- [17] Phan P, Budhram B, Zhang Q, Rivers CS, Noonan VK, Plashkes T, et al. Highlighting discrepancies in walking prediction accuracy for patients with traumatic spinal cord injury: an evaluation of validated prediction models using a Canadian Multicenter Spinal Cord Injury Registry. *Spine J* 2019;19(4):703–10. doi:[10.1016/j.spinee.2018.08.016](https://doi.org/10.1016/j.spinee.2018.08.016).
- [18] Sturt R, Hill B, Holland A, New PW, Bevans C. Validation of a clinical prediction rule for ambulation outcome after non-traumatic spinal cord injury. *Spinal Cord* 2020;58(5):609–15. doi:[10.1038/s41393-019-0386-x](https://doi.org/10.1038/s41393-019-0386-x).
- [19] DeVries Z, Hoda M, Rivers CS, Maher A, Wai E, Moravek D, et al. Development of an unsupervised machine learning algorithm for the prognostication of walking ability in spinal cord injury patients. *Spine J* 2020;20(2):213–24. doi:[10.1016/j.spinee.2019.09.007](https://doi.org/10.1016/j.spinee.2019.09.007).
- [20] Engel-Haber E, Zeilig G, Haber S, Worobey L, Kirshblum S. The effect of age and injury severity on clinical prediction rules

- for ambulation among individuals with spinal cord injury. *Spine J* 2020;20(10):1666–75. doi:10.1016/j.spinee.2020.05.551.
- [21] Scivoletto G, Pavese C, Bachmann LM, Schubert M, Curt A, Finazzi Agro E, et al. Prediction of bladder outcomes after ischemic spinal cord injury: A longitudinal cohort study from the European multicenter study about spinal cord injury. *Neurourol Urodyn* 2018;37(5):1779–84. doi:10.1002/nau.23521.
- [22] Pavese C, Schneider MP, Schubert M, Curt A, Scivoletto G, Finazzi Agro E, et al. Prediction of Bladder Outcomes after Traumatic Spinal Cord Injury: A Longitudinal Cohort Study. *PLoS Med* 2016;13(6):e1002041. doi:10.1371/journal.pmed.1002041.
- [23] Pavese C, Bachmann LM, Schubert M, Curt A, Mehnert U, Schneider MP, et al. Bowel Outcome Prediction After Traumatic Spinal Cord Injury: Longitudinal Cohort Study. *Neurorehabil Neural Repair* 2019;33(11):902–10. doi:10.1177/1545968319868722.
- [24] AlHuthaifi F, Krzak J, Hanke T, Vogel LC. Predictors of functional outcomes in adults with traumatic spinal cord injury following inpatient rehabilitation: A systematic review. *J Spinal Cord Med* 2017;40(3):282–94. doi:10.1080/10790268.2016.1238184.
- [25] Richard-Denis A, Beauséjour M, Thompson C, Nguyen BH, Mac-Thiong JM. Early Predictors of Global Functional Outcome after Traumatic Spinal Cord Injury: A Systematic Review. *J Neurotrauma* 2018;35(15):1705–25. doi:10.1089/neu.2017.5403.
- [26] Wilson JR, Cadotte DW, Fehlings MG. Clinical predictors of neurological outcome, functional status, and survival after traumatic spinal cord injury: a systematic review. *J Neurosurg Spine* 2012;17(1 Suppl):11–26 Suppl. doi:10.3171/2012.4.Aospine1245.
- [27] Al-Habib AF, Attabi N, Ball J, Bajammal S, Casha S, Hurlbert RJ. Clinical predictors of recovery after blunt spinal cord trauma: systematic review. *J Neurotrauma* 2011;28(8):1431–43. doi:10.1089/neu.2009.1157.
- [28] Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005;8(1):19–32. doi:10.1080/1364557032000119616.
- [29] Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci* 2010;5:69. doi:10.1186/1748-5908-5-69.
- [30] O'Brien KK, Colquhoun H, Levac D, Baxter L, Tricco AC, Straus S, et al. Advancing scoping study methodology: a web-based survey and consultation of perceptions on terminology, definition and methodological steps. *BMC Health Serv Res* 2016;16:305. doi:10.1186/s12913-016-1579-z.
- [31] Tricco AC, Lillie E, Zarin W, O'Brien K, Colquhoun H, Kastner M, et al. A scoping review on the conduct and reporting of scoping reviews. *BMC Med Res Methodol* 2016;16:15. doi:10.1186/s12874-016-0116-4.
- [32] Peters MD, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB. Guidance for conducting systematic scoping reviews. *Int J Evid Based Healthc* 2015;13(3):141–6. doi:10.1097/XEB.000000000000050.
- [33] Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018;169(7):467–73. doi:10.7326/M18-0850.
- [34] National Library of Medicine. PubMed. Available at: <https://pubmed.ncbi.nlm.nih.gov/>. Accessed October 12 2020.
- [35] EBSCOhost. CINAHL Complete. Available at: <http://web.b.ebscohost.com/ehost/search/advanced?vid=8&sid=ed9eea5c-13f9-4e68-9c0c-30dbf885fec1%40pdc-v-sessmgr03>. Accessed October 12 2020.
- [36] Institute of Electrical and Electronics Engineers. IEEE Xplore. Available at: <https://ieeexplore.ieee.org/Xplore/home.jsp>. Accessed October 12 2020.
- [37] Aromataris E, Riitano D. Constructing a search strategy and searching for evidence. A guide to the literature search for a systematic review. *Am J Nurs* 2014;114(5):49–56. doi:10.1097/01.NAJ.0000446779.99522.f6.
- [38] Wong SS, Wilczynski NL, Haynes RB, Ramkissoonsingh R for the Hedges Team. Developing optimal search strategies for detecting sound clinical prediction studies in MEDLINE. *AMIA Annu Symp Proc* 2003;728–32.
- [39] Keogh C, Wallace E, O'Brien KK, Murphy PJ, Teljeur C, McGrath B, et al. Optimized retrieval of primary care clinical prediction rules from MEDLINE to establish a Web-based register. *J Clin Epidemiol* 2011;64(8):848–60. doi:10.1016/j.jclinepi.2010.11.011.
- [40] The Spinal Cord Injury Research Evidence (SCIRE) Project. Outcome Measures. Available at: <http://scireproject.com/outcome-measures/alphabetical/>. Accessed September 15, 2020.
- [41] Alexander MS, Anderson KD, Biering-Sorensen F, Blight AR, Brannon R, Bryce TN, et al. Outcome measures in spinal cord injury: recent assessments and recommendations for future directions. *Spinal Cord* 2009;47(8):582–91. doi:10.1038/sc.2009.18.
- [42] Anderson K, Aito S, Atkins M, Biering-Sorensen F, Charlifue S, Curt A, et al. Functional recovery measures for spinal cord injury: an evidence-based review for clinical practice and research. *J Spinal Cord Med* 2008;31(2):133–44. doi:10.1080/10790268.2008.11760704.
- [43] Dawson J, Shamley D, Jamous MA. A structured review of outcome measures used for the assessment of rehabilitation interventions for spinal cord injury. *Spinal Cord* 2008;46(12):768–80. doi:10.1038/sc.2008.50.
- [44] Furlan JC, Noonan V, Singh A, Fehlings MG. Assessment of disability in patients with acute traumatic spinal cord injury: a systematic review of the literature. *J Neurotrauma* 2011;28(8):1413–30. doi:10.1089/neu.2009.1148.
- [45] Jackson AB, Carnel CT, Ditunno JF, Read MS, Boninger ML, Schmeler MR, et al. Outcome measures for gait and ambulation in the spinal cord injury population. *J Spinal Cord Med* 2008;31(5):487–99. doi:10.1080/10790268.2008.11753644.
- [46] Lam T, Noonan VK, Eng JJ the SCIRE Research Team. A systematic review of functional ambulation outcome measures in spinal cord injury. *Spinal Cord* 2008;46(4):246–54. doi:10.1038/sj.sc.3102134.
- [47] Magasi SR, Heinemann AW, Whiteneck GG Quality of Life/Participation Committee. Participation following traumatic spinal cord injury: an evidence-based review for research. *J Spinal Cord Med* 2008;31(2):145–56. doi:10.1080/10790268.2008.11760705.
- [48] Noonan VK, Miller WC, Noreau L the SCIRE Research Team. A review of instruments assessing participation in persons with spinal cord injury. *Spinal Cord* 2009;47(6):435–46. doi:10.1038/sc.2008.171.
- [49] Tomaschek R, Gemperli A, Rupp R, Geng V, Scheel-Sailer A German-speaking Medical SCI Society (DMGP) Ergebniserhebung Guideline Development Group. A systematic review of outcome measures in initial rehabilitation of individuals with newly acquired spinal cord injury: providing evidence for clinical practice guidelines. *Eur J Phys Rehabil Med* 2019;55(5):605–17. doi:10.23736/S1973-9087.19.05676-4.
- [50] Ballert CS, Hopfe M, Kus S, Mader L, Prodinger B. Using the refined ICF Linking Rules to compare the content of existing instruments and assessments: a systematic review and exemplary analysis of instruments measuring participation. *Disabil Rehabil* 2019;41(5):584–600. doi:10.1080/09638288.2016.1198433.
- [51] Bramer WM, Giustini D, de Jonge GB, Holland L, Bekhuis T. De-duplication of database search results for systematic reviews in EndNote. *J Med Libr Assoc* 2016;104(3):240–3. doi:10.3163/1536-5050.104.3.014.
- [52] Maritz R, Aronsky D, Prodinger B. The International Classification of Functioning, Disability and Health (ICF) in Electronic Health Records. A Systematic Literature Review. *Appl Clin Inform* 2017;8(3):964–80. doi:10.4338/aci-2017050078.
- [53] Maritz R, Scheel-Sailer A, Schmitt K, Prodinger B. Overview of quality management models for inpatient healthcare settings. A scop-

- ing review. *Int J Qual Health Care* 2019;31(6):404–10. doi:[10.1093/intqhc/mzy180](https://doi.org/10.1093/intqhc/mzy180).
- [54] Bramer WM, Milic J, Mast F. Reviewing retrieved references for inclusion in systematic reviews using EndNote. *J Med Libr Assoc* 2017;105(1):84–7. doi:[10.5195/jmla.2017.111](https://doi.org/10.5195/jmla.2017.111).
- [55] Moons KG, de Groot JA, Bouwmeester W, Vergouwe Y, Mallett S, Altman DG, et al. Critical appraisal and data extraction for systematic reviews of prediction modelling studies: the CHARMS checklist. *PLoS Med* 2014;11(10):e1001744. doi:[10.1371/journal.pmed.1001744](https://doi.org/10.1371/journal.pmed.1001744).
- [56] Cieza A, Fayed N, Bickenbach J, Prodinger B. Refinements of the ICF Linking Rules to strengthen their potential for establishing comparability of health information. *Disabil Rehabil* 2019;41(5):574–83. doi:[10.3109/09638288.2016.1145258](https://doi.org/10.3109/09638288.2016.1145258).
- [57] Arijji Y, Hayashi T, Ideta R, Koga R, Murai S, Towatari F, et al. A prediction model of functional outcome at 6 months using clinical findings of a person with traumatic spinal cord injury at 1 month after injury. *Spinal Cord* 2020;58(11):1158–65. doi:[10.1038/s41393-020-0488-5](https://doi.org/10.1038/s41393-020-0488-5).
- [58] Facchinello Y, Beausejour M, Richard-Denis A, Thompson C, Mac-Thiong JM. Use of Regression Tree Analysis for Predicting the Functional Outcome after Traumatic Spinal Cord Injury. *J Neurotrauma* 2021;38(9):1285–91. doi:[10.1089/neu.2017.5321](https://doi.org/10.1089/neu.2017.5321).
- [59] Harrington GMB, Cool P, Hulme C, Osman A, Chowdhury JR, Kumar N, et al. Routinely Measured Hematological Markers Can Help to Predict American Spinal Injury Association Impairment Scale Scores after Spinal Cord Injury. *J Neurotrauma* 2021;38(3):301–8. doi:[10.1089/neu.2020.7144](https://doi.org/10.1089/neu.2020.7144).
- [60] Kaminski L, Cordemans V, Cernat E, M'Bra KI, Mac-Thiong JM. Functional Outcome Prediction after Traumatic Spinal Cord Injury Based on Acute Clinical Factors. *J Neurotrauma* 2017;34(12):2027–33. doi:[10.1089/neu.2016.4955](https://doi.org/10.1089/neu.2016.4955).
- [61] Tomioka Y, Uemura O, Ishii R, Liu M. Using a logarithmic model to predict functional independence after spinal cord injury: a retrospective study. *Spinal Cord* 2019;57(12):1048–56. doi:[10.1038/s41393-019-0315-z](https://doi.org/10.1038/s41393-019-0315-z).
- [62] Wilson JR, Grossman RG, Frankowski RF, Kiss A, Davis AM, Kulkarni AV, et al. A clinical prediction model for long-term functional outcome after traumatic spinal cord injury based on acute clinical and imaging factors. *J Neurotrauma* 2012;29(13):2263–71. doi:[10.1089/neu.2012.2417](https://doi.org/10.1089/neu.2012.2417).
- [63] Zariffa J, Kapadia N, Kramer JL, Taylor P, Alizadeh-Meghbrazi M, Zivanovic V, et al. Relationship between clinical assessments of function and measurements from an upper-limb robotic rehabilitation device in cervical spinal cord injury. *IEEE Trans Neural Syst Rehabil Eng* 2012;20(3):341–50. doi:[10.1109/TNSRE.2011.2181537](https://doi.org/10.1109/TNSRE.2011.2181537).
- [64] Moher D, Liberati A, Tetzlaff J, Altman DG the PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6(7):e1000097. doi:[10.1371/journal.pmed.1000097](https://doi.org/10.1371/journal.pmed.1000097).
- [65] Koyama T, Matsumoto K, Okuno T, Domen K. A new method for predicting functional recovery of stroke patients with hemiplegia: logarithmic modelling. *Clin Rehabil* 2005;19(7):779–89. doi:[10.1191/0269215505cr876oa](https://doi.org/10.1191/0269215505cr876oa).
- [66] Wingbermühle RW, Chiarotto A, Koes B, Heymans MW, van Trijffel E. Challenges and solutions in prognostic prediction models in spinal disorders. *J Clin Epidemiol* 2021. doi:[10.1016/j.jclinepi.2020.12.017](https://doi.org/10.1016/j.jclinepi.2020.12.017).
- [67] Wartenberg KE, Hwang DY, Haeusler KG, Muehlschlegel S, Sakowitz OW, Madzar D, et al. Gap Analysis Regarding Prognostication in Neurocritical Care: A Joint Statement from the German Neurocritical Care Society and the Neurocritical Care Society. *Neurocrit Care* 2019;31(2):231–44. doi:[10.1007/s12028-019-00769-6](https://doi.org/10.1007/s12028-019-00769-6).
- [68] Belliveau T, Jette AM, Seetharama S, Axt J, Rosenblum D, Larose D, et al. Developing Artificial Neural Network Models to Predict Functioning One Year After Traumatic Spinal Cord Injury. *Arch Phys Med Rehabil* 2016;97(10):1663–1668.e3. doi:[10.1016/j.apmr.2016.04.014](https://doi.org/10.1016/j.apmr.2016.04.014).
- [69] Pretz CR, Kozłowski AJ, Charlifue S, Chen Y, Charlifue S, Heineemann AW. Using Rasch motor FIM individual growth curves to inform clinical decisions for persons with paraplegia. *Spinal Cord* 2014;52(9):671–6. doi:[10.1038/sc.2014.94](https://doi.org/10.1038/sc.2014.94).
- [70] Gravestijn BY, Nieboer D, Ercole A, Lingsma HF, Nelson D, van Calster B, et al. Machine learning algorithms performed no better than regression models for prognostication in traumatic brain injury. *J Clin Epidemiol* 2020;122:95–107. doi:[10.1016/j.jclinepi.2020.03.005](https://doi.org/10.1016/j.jclinepi.2020.03.005).
- [71] Geersing GJ, Bouwmeester W, Zuithoff P, Spijker R, Leeflang M, Moons KG. Search filters for finding prognostic and diagnostic prediction studies in Medline to enhance systematic reviews. *PLoS One* 2012;7(2):e32844. doi:[10.1371/journal.pone.0032844](https://doi.org/10.1371/journal.pone.0032844).
- [72] Hupp M, Pavese C, Bachmann LM, Koller R, EMSCI Study Group, Schubert M, Group ES. Electrophysiological Multimodal Assessments Improve Outcome Prediction in Traumatic Cervical Spinal Cord Injury. *J Neurotrauma* 2018;35(24):2916–23. doi:[10.1089/neu.2017.5576](https://doi.org/10.1089/neu.2017.5576).
- [73] Ribeiro Neto F, Gomes Costa RR, Tanhoffer RA, Leal JC, Bottaro M, Carregaro RL. Muscle Strength Cutoff Points for Functional Independence and Wheelchair Ability in Men With Spinal Cord Injury. *Arch Phys Med Rehabil* 2020;101(6):985–93. doi:[10.1016/j.apmr.2020.01.010](https://doi.org/10.1016/j.apmr.2020.01.010).