

Tools and drugs for uracil nucleotide-activated P2Y receptors

Muhammad Rafehi, Christa E. Müller

Angaben zur Veröffentlichung / Publication details:

Rafehi, Muhammad, and Christa E. Müller. 2018. "Tools and drugs for uracil nucleotide-activated P2Y receptors." *Pharmacology & Therapeutics* 190: 24–80.
<https://doi.org/10.1016/j.pharmthera.2018.04.002>.

Tools and drugs for uracil nucleotide-activated P2Y receptors

Muhammad Rafehi, Christa E. Müller*

Pharmaceutical Chemistry I, Pharmaceutical Institute, Pharmaceutical Sciences Bonn, University of Bonn, Germany

Contents

1. Introduction	25
2. The P2Y ₂ receptor	34
3. The P2Y ₄ receptor	58
4. The P2Y ₆ receptor	67
5. The P2Y ₁₄ receptor	69
6. Conclusions	73
Conflict of Interest	74
References	74

Abbreviations: ADP, Adenosine 5'-diphosphate; AP₄A, P¹,P⁴-Di(adenosine-5')-tetraphosphate; AR, Adenosine receptor; AR-C118925, 5-[[5-(2,8-Dimethyl-5H-dibenzo[a,d]cyclohepten-5-yl)-3,4-dihydro-2-oxo-4-thioxo-1(2H)-pyrimidinyl)methyl]-N-(1H-tetrazol-5-yl)-2-furancarboxamide; ATP, Adenosine 5'-triphosphate; ATPγS, Adenosine 5'-(γ-thio)-triphosphate; BODIPY®, Boron-dipyrromethene; cAMP, 3',5'-Cyclic adenosine monophosphate; CDP, Cytidine 5'-diphosphate; CFTR, Cystic fibrosis transmembrane conductance regulator; CNS, Central nervous system; CTP, Cytidine 5'-triphosphate; DAG, Diacylglycerol; DTPA, Diethylenetriaminepentaacetic acid; GDP, Guanosine 5'-diphosphate; GI tract, Gastrointestinal tract; GPCR, G protein-coupled receptor; GTP, Guanosine 5'-triphosphate; HEK, Human embryonic kidney; IDP, Inosine 5'-diphosphate; IMP, Inosine 5'-monophosphate; INS48823, P¹-(2-Benzyl-1,3-dioxolo-4-yl)uridine-5'-P³-(uridine-5'-)-triphosphate; monobenzylacetal-Up₃U; IP₃, Inositol 1,4,5-trisphosphate; ITP, Inosine 5'-triphosphate; Me, Methyl; MeO, Methoxy; MeS, Methylthio; MRS2567, 1,2-Di-(4-isothiocyantophenyl)ethane; MRS2578, N,N"-1,4-Butanediyl-bis[N'-(3-isothiocyantophenyl)thiourea]; MRS2957, N⁴-MeO-Cp₃U; P¹-(uridine 5'-)-P³-(N⁴-methoxycytidine-5'-)-triphosphate; n.d., No data was found; NECA, 5'-(N-Ethylcarboxamido)adenosine; P2YR, P2Y receptor; PAMAM, Polyamidoamine; PIP₂, Phosphatidylinositol 4,5-bisphosphate; PPADS, Pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid; PPTN, 4-(4-(Piperidin-4-yl)-phenyl)-7-(4-(trifluoromethyl)-phenyl)-2-naphthoic acid; PSB-16133, Sodium 1-amino-4-[4-(2,4-dimethylphenylthio)]-9,10-dioxo-9,10-dihydroanthracene-2-sulfonate; PSB-716, 1-Amino-4-(2-methoxyphenyl)-2-sulfoanthraquinone; RB-2, Reactive Blue 2; SAR, Structure-activity relationship; TDP, Thymidine 5'-diphosphate; TIM-38, 3-Nitro-2-(trifluoromethyl)-2H-chromene; TMPS, Thymidine 5'-O-monophosphorothioate; TTP, Thymidine 5'-triphosphate; UDP, Uridine 5'-diphosphate; UDPβS, Uridine 5'-(β-thio)-diphosphate; UMP, Uridine 5'-monophosphate; UMPS, Uridine 5'-O-monophosphorothioate; UP₄U, P¹,P⁴-Di(uridine-5')-tetraphosphate; UTP, Uridine 5'-triphosphate; UTPγS, Uridine 5'-(γ-thio)-triphosphate.

* Corresponding author at: Pharmazeutisches Institut, Pharmazeutische Chemie I, An der Immenburg 4, D-53121 Bonn, Germany.

E-mail address: christa.mueller@uni-bonn.de. (C.E. Müller).

1. Introduction

1.1. Purinergic and pyrimidinergic receptors

The first reported observation that purines are capable of evoking cellular responses was published in 1929, when Drury and Szent-Györgyi investigated the effects of adenosine and adenosine 5'-monophosphate on the mammalian heart (Drury & Szent-Györgyi, 1929). However, it was not until 1976 that the existence of a distinct family of purinergic receptors was proposed (Burnstock, 1976). Shortly thereafter, the distinction into P1 (adenosine receptors, xanthine-sensitive) and P2 subfamilies (nucleotide receptors, not blocked by xanthine derivatives) was made. A third class, P0 (activated by the nucleobase adenine), was later discovered in rodents through reverse pharmacology (Bender et al., 2002; Brunschweiler & Müller, 2006). Both P0 and P1 receptors are G protein-coupled receptors (GPCRs). The P2 family is subdivided into the metabotropic P2Y receptors (P2YRs) and the ligand-gated ion channels known as P2X receptors (Abbracchio & Burnstock, 1994). The latter group consists of seven subunits, termed P2X1-7, which form homo- or heterotrimeric ligand-gated cation channels (Aschrafi, Sadtler, Niculescu, Rettinger, & Schmalzing, 2004; Kaczmarek-Hájek, Lorinczi, Hausmann, & Nicke, 2012). They share little sequence homology with other proteins and form, besides Cys-loop and glutamate-gated channels, a third major family of ionotropic receptors (North, 1996). P2X channels are cation-selective (mainly Ca^{2+} and Na^+ , but also K^+) and are activated by the nucleotide adenosine 5'-triphosphate (ATP) (Coddou, Yan, Obsil, Huidobro-Toro, & Stojilkovic, 2011; Kaczmarek-Hájek et al., 2012).

1.2. P2Y receptors

P2YRs belong to the δ -branch of class A rhodopsin-like GPCRs. They have an extracellular, glycosylated N-terminus, the characteristic seven hydrophobic transmembrane domains, three intra- and three extracellular loops, and an intracellular C-terminus that possesses consensus binding motifs for protein kinases. The mammalian P2YR family consists of eight subtypes: P2Y₁, P2Y₂, P2Y₄, P2Y₆, P2Y₁₁, P2Y₁₂, P2Y₁₃, and P2Y₁₄. The numbers missing in the sequence (p2y3, p2y5, p2y7-10) account for non-mammalian receptors or for GPCRs that have a certain degree of sequence homology but do not appear to bind nucleotides. Examples include the chicken p2y3 receptor and the *Xenopus laevis* p2y8 receptor, which may be homologs of the mammalian P2Y₆R and the P2Y₄R, respectively (Bogdanov, Dale, King, Whittock, & Burnstock, 1997; Li, Olesky, Palmer, Harden, & Nicholas, 1998; Webb et al., 1996). The p2y7 and the p2y9 receptor are a leukotriene B₄ and a lysophosphatidic acid receptor (Akbar et al., 1996; Herold, Li, Schachter, Harden, & Nicholas, 1997; Noguchi, Ishii, & Shimizu, 2003). It is possible that further human P2YR subtypes have not been identified yet, as several orphan receptors share some sequence similarity with current P2YR members. It has to be noted that sequence homology between the current members of the P2YR family is moderate to low (21-48 % sequence identity) (Abbracchio et al., 2006).

The human P2YRs can be further categorized into two groups based on sequence homology and the type of G protein they are primarily coupled to (Fig. 1) (Costanzi, Mamedova, Gao, & Jacobson, 2004). The P2Y₁, P2Y₂, P2Y₄, P2Y₆, and P2Y₁₁ receptors signal mainly through G_{q/11} with subsequent activation of phospholipase C β . This enzyme in turn hydrolyzes phosphatidylinositol 4,5-bisphosphate (PIP₂), producing the second messengers inositol 1,4,5-trisphosphate (IP₃) and diacylglycerol (DAG). IP₃ and DAG mediate the release of calcium ions from intracellular stores and activate protein kinase C, respectively. The P2Y₁₂, P2Y₁₃, and P2Y₁₄ receptors signal preferentially through G_{i/o}, thereby inhibiting adenylate cyclase and reducing 3',5'-cyclic adenosine monophosphate (cAMP) levels (Abbracchio et al., 2006).

However, P2YR signaling is more complex and coupling to other second messenger systems has been described: P2Y₂ was reported to

couple to G₀ and G₁₆, P2Y₁₁ to G_s, and P2Y₂, P2Y₆, and P2Y₁₂ were found to activate Rho through G_{12/13} (Bagchi et al., 2005; Baltensperger & Porzig, 1997; Liao, Seye, Weisman, & Erb, 2007; Nishida et al., 2008; Sauzeau et al., 2000). Furthermore, P2YRs may form homo- or hetero-oligomers, as is the case for other GPCRs (a database of GPCR oligomers can be found at www.gpcr-okb.org (Khelashvili et al., 2010; Skrabanek et al., 2007)). This may alter their pharmacological, signaling, desensitization, and trafficking properties. The formation of P2Y₆R homodimers was described (Kawashita, Tsuji, Kanno, Tsuchida, & Itoh, 2016). The P2Y₁R and the P2Y₂R were observed to coimmunoprecipitate with the A₁ adenosine receptor (A₁AR) in cotransfected human embryonic kidney (HEK) 293T cells (Suzuki, Namba, Tsuga, & Nakata, 2006; Yoshioka, Saitoh, & Nakata, 2001). The results suggested that the A₁AR can form heteromers with either one, the P2Y₁R and the P2Y₂R. The formation of A₁AR/P2Y₁R heteromers was also observed *in vivo* in colocalization and coimmunoprecipitation studies in rat brain tissue (Yoshioka, Hosoda, Kuroda, & Nakata, 2002). The heteromeric receptors were shown to possess novel pharmacological properties: A₁AR/P2Y₁R heteromers behaved like A₁ARs with respect to signaling but showed the agonist profile of P2Y₁Rs. The P2Y₁R agonist adenosine 5'- β -thio-diphosphate (ADP β S) displaced the adenosine receptor agonist [³H]5'-(N-ethylcarboxamido)adenosine ([³H]NECA) from its binding site on the A₁AR and induced G_{i/o}-mediated signaling, which is typical for A₁ARs (Yoshioka et al., 2001). However, the selective allosteric P2Y₁R antagonist N⁶-methyl-2'-deoxyadenosine 3',5'-bisphosphate (MRS2179) failed to compete with [³H]NECA for its binding site on the A₁/P2Y₁ heteromeric receptor (Yoshioka et al., 2001). In A₁AR/P2Y₂R heteromers, binding of [³H]NECA and of the selective A₁AR agonist [³H]2-chloro-N⁶-cyclopentyladenosine ([³H]CCPA) was inhibited by the P2Y₂R agonists ATP and uridine 5'-triphosphate (UTP) (Suzuki et al., 2006). UTP prevented the G_{i/o}-mediated inhibition of adenylate cyclase in A₁AR/P2Y₂R heteromers that is normally observed as a result of A₁AR activation by an A₁AR agonist. P2Y₂R-mediated intracellular Ca^{2+} release *via* G_{q/11} was synergistically enhanced by the simultaneous activation of the A₁AR/P2Y₂R heteromers by the P2Y₂R agonist UTP and the A₁AR agonist NECA (Suzuki et al., 2006). Heteromerization within the P2YR family appears to exist as well. Coimmunoprecipitation and Förster resonance energy transfer (FRET) experiments provided evidence for P2Y₁R/P2Y₂R, P2Y₁R/P2Y₄R, and P2Y₁R/P2Y₁₁R heteromers (Ecke et al., 2008; Ribeiro-Filho et al., 2016). The physiological implications of P2YR heteromerization needs to be further elucidated, but it appears likely that it leads to allosteric modulation altering agonist affinity and specificity, and impacting intracellular signaling pathways.

1.3. P2Y receptor ligands

Another approach to categorizing the human P2YR family is based on their preference for natural ligands. The P2Y₁, P2Y₁₁, P2Y₁₂, and P2Y₁₃ receptors are activated by adenine di- or triphosphates. In contrast, uracil nucleotides are the endogenous agonists for the P2Y₄, P2Y₆, and P2Y₁₄ receptors. The P2Y₂R responds to both ATP and UTP. The P2 receptor family was initially named P2 purinoceptors. With the discovery of the uracil nucleotide-activated subtypes, the inadequacy of this nomenclature became evident and the receptor family was renamed accordingly (purine and pyrimidine receptors). The pronounced differences in agonist preferences, which is uncommon in other GPCR families, likely results from the relatively low sequence homology shared between members of the P2YR family. Yet, it has proven difficult with most P2YR subtypes to develop potent agonists and antagonists that are highly selective over other subtypes. A major issue with many current P2YR ligands – mainly concerning agonists but also some classes of antagonists – is their low oral bioavailability due to several negative charges (e.g. phosphate or sulfonate groups that are deprotonated at physiological pH). Nucleotides are subject to degradation by ectonucleotidases, which results in the production of nucleoside di- or monophosphates and eventually leads to the




Fig. 1. Overview of the P2YR subtypes categorized into two groups based on sequence homology and main signaling cascades.

formation of the nucleosides adenosine or uridine (Harden, Lazarowski, & Boucher, 1997; also see reviews by Zimmermann, 2016 and Zimmermann, Zebisch, & Sträter, 2012). Some of these metabolites in turn can activate other receptors and thereby cause ambiguity in pharmacological experiments. Moreover, ectokinases may phosphorylate nucleosides and nucleotides, further causing difficulties in the correct interpretation of the measured biological responses (Harden et al., 1997; Lazarowski, Boucher, & Harden, 2000; Nicholas, Watt, Lazarowski, Li, & Harden, 1996). Additional complicating factors in cell-based experiments could also be the result of nucleotide release via connexins, pannexins, maxi-channels, ATP-binding cassette transporters, exocytosis of secretory granules, or vesicular transport. This may occur in response to stress experienced by the cells, for example through physical movement (e.g. shaking), hypoxia, or changes in pH or nutrient medium (Lazarowski, Boucher, & Harden, 2003; Lazarowski, Watt, Stutts, Boucher, & Harden, 1995; for more information, refer to the recent review by Burnstock & Knight, 2017). In addition, nucleotides are released from cells that undergo necrosis or apoptosis, acting as signalling molecules to promote phagocytic clearance of the dying cells (Elliott et al., 2009; Idzko et al., 2007). To complicate matters even further, some commercial preparations of nucleoside

diphosphates were found to contain significant amounts of the corresponding nucleoside triphosphates, and vice versa. This had previously led to incorrect conclusions regarding pharmacological properties of P2Y receptors (Nicholas et al., 1996).

1.4. Scope and structure of this review

Given the significant clinical potential of the P2YR family, substantial research efforts directed at developing P2YR ligands for use as pharmacological tools and drugs have led to the discovery and development of a significant number of agonists but so far only a moderate number of antagonists. An exception is the P2Y₁₂R, for which antagonists are already marketed as antithrombotic drugs. The present review summarizes the current state of ligand development for the class of uracil nucleotide-activated P2Y receptors – the P2Y₂R, P2Y₄R, P2Y₆R, and P2Y₁₄R – and discusses the compounds' advantages and limitations. In addition, a series of tables provide a comprehensive overview of the available receptor ligands, their structures and their properties, to allow for direct comparison. For simplicity, compounds are represented as free acids or free bases. It should be noted, however, that they are synthesized, purchased and tested in their more stable salt forms. The

Table 1
Overview of the human uracil nucleotide-activated P2YR subtypes.

Receptor	Entrez gene ID	Uniprot ID ^a	Amino acid length	Main physiological agonists	Main tissue distribution ^b	Therapeutic potential ^c
P2Y ₂	5029	P41231	377	ATP, UTP, (Ap ₄ A, Up ₄ U)	Endocrine tissue, immune system, skeletal & cardiac muscle, lung, GI tract, male & female reproductive tract, bladder, kidney, skin ^c Prominent expression in epithelial & glandular cells ^c	Agonists: Neurodegenerative disorders, dry eyes, ocular hypertension, retinal degeneration, cystic fibrosis, myocardial infarction Antagonists: Atherosclerosis, nephrogenic diabetes insipidus, psoriasis, osteoporosis, cancer, inflammation, pain Agonists: Neurodegenerative disorders, cystic fibrosis Antagonists: Neurodegenerative disorders, myocardial infarction, constipation, diarrhea, cancer Agonists: Neurodegenerative disorders, ocular hypertension, glaucoma, cystic fibrosis, blood pressure, cancer, diabetes, infectious diseases Antagonists: Neurodegenerative disorders, cerebral vasospasms, cardiac hypertrophy, cardiac fibrosis, atherosclerosis, inflammation, pain, cancer, obesity Agonists: Autoimmune diseases, diabetes Antagonists: Inflammation, diabetes, osteoporosis
P2Y ₄	5030	P51582	365	UTP, (Up ₄ U)	GI tract; lower expression in CNS, lung, heart, prostate, skin, adipose tissue, skeletal muscle, spleen & immune cells ^d	
P2Y ₆	5031	Q15077	328	UDP, (Up ₃ U)	Spleen, placenta, kidney; lower expression in CNS, lung, heart, GI tract, adipose tissue, bone ^d	
P2Y ₁₄	9934	Q15391	338	UDP, UDP-glucose	CNS, endocrine tissue, immune system, muscle (skeletal, cardiac, smooth), lung, pancreas, GI tract, kidney, bladder, male & female reproductive tract, skin ^c	

^a UniProt Consortium, 2015.

^b Moore et al., 2001; Uhlén et al., 2015.

^c Detected at both protein and mRNA level.

^d Detected at mRNA level.

^e Refer to the text for more details and for references.

most important compounds listed in these tables will be discussed in more detail. Agonists are collected in Tables 2 to 9, categorized according to structural modifications. These include physiological mononucleotides (Fig. 2 and Table 2) and synthetic derivatives thereof, with substitutions at the nucleobase (Fig. 3 and Table 3), the ribose moiety (Fig. 4 and Table 4), or the phosphate chain (Fig. 5 and Table 5). Nucleotide derivatives with combined substitutions at different sites are depicted in Fig. 6 and Table 6. Physiological dinucleotides and synthetic analogues are listed in Fig. 7 and Table 7. Nucleotide sugars are summarized in Fig. 8 and Table 8, while non-nucleotide agonists reported very recently are shown in Fig. 9 and Table 9. For each compound, published EC₅₀ values and, where available, K_i or K_B values at the P2Y₂R, P2Y₄R,

P2Y₆R, and P2Y₁₄R are presented. These potencies are given as a range if more than two different values for a compound were found in the literature. Since EC₅₀ values of GPCRs are influenced by receptor expression and can vary significantly between different test systems (Fujioka & Omori, 2012), a direct comparison is facilitated by means of a correlation factor given in brackets. This factor expresses the multiplicity by which the potency of the compound directly compares to that of the endogenous agonists ATP (superscript a), UTP (superscript b), UDP (superscript c), or UDP-glucose (superscript d) assessed in the same assay system. The correlation factor for a compound that is n-fold more potent (i.e. smaller EC₅₀, K_i, or K_B) is shown as (n^{mx})^{a,b,c, or d}, while compounds of n-fold lower potency (i.e. larger EC₅₀, K_i, or K_B) are described




Fig. 2. Structures of physiological mononucleotides (for data, refer to Table 2).

as (\downarrow nx)^{a,b,c, or d}. In cases where large series of structural analogues were published, only the most important compounds are shown.

The development of antagonists for the uracil nucleotide-activated P2YRs has yielded significantly fewer compounds so far as compared to the number of published agonists, but they display

greater structural variability (Fig. 10 and Table 10). The antagonists were organized according to receptor selectivity. Antagonist potencies are mostly given as IC₅₀ values; in some cases they are shown as pA₂, K_i, K_B values, or as the percentage of inhibition at a given concentration.

Table 2

Physiological mononucleotides for the uracil-activated P2YRs. Shown are the EC₅₀ or, in some cases, K_i and K_B values in μ M at the human receptor, unless stated otherwise. The correlation factor by which the potency is n-fold higher (\uparrow nx) or lower (\downarrow nx) compared to that of the endogenous agonist in the same test system is given in brackets. Particularly interesting values are shown in bold.

Physiological mononucleotides							
No.	Name	EC ₅₀ in μM (potency higher (↑) or lower (↓) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
2-01	ATP	0.015 - 0.230 (1x-↓4x) ^b	antagonist (see Table 10)	>3 mM (↓>10,000x) ^c	0.252 (↓100x ^c / ↓115x ^d) inactive at 1 μM (n.d. ^c / ↓>10x ^d)		Bogdanov et al., 1998; Chambers et al., 2000; Chang et al., 1995; Chen, Krull, Xu, Levy, & Lightman, 1996; Communi, Parmentier et al., 1996; Erb, Lustig, Sullivan, Turner, & Weisman, 1993; Filippov, Webb, Barnard, & Brown, 1999; Hamel et al., 2011; Ivanov, Fricks, Kendall Harden, & Jacobson, 2007; Jacobson et al., 2006; Janssens, Paindavoine, Parmentier, & Boeynaems, 1999; Kennedy et al., 2000; Knoblauch et al., 1999; Lazarowski et al., 1995; Lazarowski et al., 2001; Lazarowski & Harden, 1994; Lustig et al., 1993; Nicholas et al., 1996; Sakuma et al., 2017; Shaver et al., 2005; Shen et al., 2004; Suarez-Huerta, Pouillon, Boeynaems, & Robaye, 2001; Webb, Henderson, Roberts, & Barnard, 1998; Wildman, Unwin, & King, 2003; Zambon et al., 2000
		Mouse: 0.7 - 17.7 (↑2x-↓14x) ^b	Mouse: 0.435 (1x ^a / ↓2x ^b) 0.7 (1x ^a / ↓2x ^b)	Mouse: inactive at 100 μM (↓>2300x) ^c			
		Rat: 2.7 (1x) ^b 0.2 (1x) ^b	Rat: 0.51 - 1.8 (1x ^a / 1x-↓3x ^b)	Rat: 500 (n.d.) ^c inactive (n.d.) ^c			
		Dog: ≈0.2 (1x) ^b ≈2 (1x) ^b					
		Pig: 2.7 (↓5x) ^b 80 % efficacy					

(continued on next page)

Table 2 (continued)

Physiological mononucleotides						
No.	Name	EC ₅₀ in μ M (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄	
2-02	ADP	inactive at 100 μ M (\downarrow 500 - >6600x ^a / \downarrow 1000 - >6600x ^b)	inactive at 100 μ M (\downarrow >500x) ^b partial agonist with 15 % efficacy	30 - 100 (\downarrow 100-500x) ^c inactive at 100 μ M (\downarrow >200x) ^c	0.0139 (\downarrow 6x) ^{c,d} inactive at 1 μ M (n.d. ^c / \downarrow >10x ^d)	Bogdanov et al., 1998; Chambers et al., 2000; Chang et al., 1995; Chen et al., 1998; Communi, Parmentier et al., 1998; Filippov et al., 1999; Hamel et al., 2011; Lazarowski et al., 2001; Lazarowski & Harden, 1994; Lin, Lustig, Sportiello, Weisman, & Sun, 1993; Lustig et al., 1993; Nguyen et al., 1995; Nicholas et al., 1998; Robaye et al., 1997; Shaver et al., 2005; Shen et al., 2004; Webb et al., 1998; Zamboni et al., 2000
		Mouse: \approx 100 (n.d.) ^{a,b} >100 (\downarrow >140x ^a / \downarrow >90x ^b) partial agonist	Mouse: inactive at 100 μ M (\downarrow >230x ^a / \downarrow >390x ^b) Rat: partial agonist with 34 % efficacy inactive at 100 μ M (\downarrow >95x ^a / \downarrow >160x ^b)	Mouse: >1 (\downarrow >24x) ^c Rat: 50 (n.d.) ^c inactive (\downarrow >130x) ^c		
2-03	GTP	2.64 (\downarrow 31x ^a / \downarrow 54x ^b) 12.3 (\downarrow 53x ^a / \downarrow 88x ^b)	6.59 (\downarrow 12x) ^b inactive at 100 μ M (\downarrow >500x) ^b	n.d.	0.814 (\downarrow 340x ^c / \downarrow 370x ^d)	Chen et al., 1996; Erb et al., 1993; Hamel et al., 2011; Jacobson et al., 2006; Kennedy et al., 2000; Lazarowski et al., 1995; Lazarowski et al., 2001; Lazarowski & Harden, 1994; Lin et al., 1993; Shen et al., 2004; Wildman et al., 2003
		Mouse: >100 (n.d.) ^{a,b} inactive at 1 mM (\downarrow >670x ^a / \downarrow >1100x ^b)	Mouse: 7 (\downarrow 16x ^a / \downarrow 27x ^b) Rat: 1.4 (1x) ^{a,b} 2.28 (\downarrow 5x ^a / \downarrow 11x ^b)	Rat: inactive at 300 μ M (\downarrow >130x) ^c		

Table 2 (continued)

Physiological mononucleotides							
No.	Name	EC ₅₀ in μ M (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
		Pig: 1.3 (\uparrow 2x ^a / \downarrow 3x ^b) 63 % efficacy					
2-04	GDP	66.0 (\downarrow 290x ^a / \downarrow 470x ^b) Rat: inactive at 100 μ M (\downarrow >500x) ^{a,b} Pig: inactive at 100 μ M (\downarrow >37x ^a / \downarrow >190x ^b)	n.d.	44.6 (\downarrow 150x) ^c Rat: inactive at 300 μ M (\downarrow >130x) ^c Mouse: >1 (\downarrow >24x) ^c	0.0800 (\downarrow 33x ^c / \downarrow 36x ^d) \approx 30 (\downarrow 400x ^c / \downarrow 93x ^d)		Carter et al., 2009; Chen et al., 1996; Hamel et al., 2011; Lazarowski et al., 1995; Lazarowski et al., 2001; Lazarowski & Harden, 1994; Robaye et al., 1997; Shen et al., 2004
2-05	ITP	\approx 10 (n.d.) ^{a,b} Mouse: ca. 100 (n.d.) ^{a,b} Rat: 20.9 (\downarrow 8x ^a / \downarrow 6x ^b) Pig: 1.7 (\uparrow 2x ^a / \downarrow 3x ^b)	7.38 - 32.8 (\downarrow 13x) ^b >1000 (\downarrow >5000x) ^b Mouse: 2 (\downarrow 5x ^a / \downarrow 8x ^b) Rat: 1.4 - 1.82 (1x- \downarrow 4x ^a / 1x- \downarrow 9x ^b)	n.d.	n.d.		Bogdanov et al., 1998; Brunschweiler & Müller, 2006; Communi, Motte et al., 1996; Kennedy et al., 2000; Lazarowski et al., 2001; Lin et al., 1993; Nguyen et al., 1995; Shen et al., 2004; Wildman et al., 2003
2-06	IDP	n.d. Pig: 35.7 (\downarrow 13x ^a / \downarrow 67x ^b)	n.d.	34.4 (\downarrow 120x) ^c	0.010 (\downarrow 4x ^c / \downarrow 5x ^d) inactive at 1 μ M (1x ^c / \downarrow >10x ^d)		Chambers et al., 2000; Hamel et al., 2011; Robaye et al., 1997; Shen et al., 2004
2-07	IMP	n.d.	n.d.	n.d.	5.43 (\downarrow 2300x ^c / \downarrow 2500x ^d)		Hamel et al., 2011

(continued on next page)

Table 2 (continued)

Physiological mononucleotides							
No.	Name	EC ₅₀ in μM (potency higher (↑) or lower (↓) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
2-08	UTP (INS316)	0.014 - 0.64 (↑4x-1x) ^a	0.069 - 2.5	6 (↓20x) ^c >10 (↓>100x) ^c	0.0175 - 0.381 (↓7-54x ^c / ↓3-16x ^d) inactive at 1 μM (n.d. ^a / ↓>10x ^d) K _i = 0.009 (↑2x) ^{c,d} K _i = 0.251 (↓2x ^c / ↓4x ^d)	Investigated in clinical studies to stimulate mucociliary clearance by targeting the P2Y ₂ R in patients with mild chronic bronchitis*	Bogdanov et al., 1998; Chambers et al., 2000; Chang et al., 1995; Charlton et al., 1996b; Chen et al., 1996; Communi, Motte et al., 1996; Communi, Parmentier et al., 1996; Eliahu et al., 2009; El-Tayeb et al., 2006; Erb et al., 1993; Filippov et al.,
		Mouse: 0.9 - 1.25 (↑14x- ↓2x) ^a	Mouse: 0.260 (↑2x) ^a 0.4 (↑2x) ^a	Mouse: ≈0.8 (↓20x) ^c	Mouse: 0.049 - 0.0303 (↓2-3x ^c / ↓4-6x ^d)		1999, 1997, 1998; WO002009066298A1, 2008; WO002012073237A1, 2011; WO002012032513A1, 2011; Fricks et al., 2008; Guille et al., 2001; Hamel et al., 2011; Ivanov, Ko et al., 2007; Jacobson et al., 2006; Janssens et al., 1999; US020090148850A1, 2008; Kennedy et al., 2000; Kim et al., 2002; Knoblauch et al., 1999; Ko et al., 2007; Ko et al., 2008; Lazarowski et al., 1995; Lazarowski et al., 2001; Lazarowski & Harden, 1994; Lustig et al., 1993; Marucka et al., 2011; Nguyen et al., 1995; Nicholas et al., 1996; Patel et al., 2001; Pendergast et al., 2001; Sakuma et al., 2017; Shaver et al., 2005; Shen et al., 2004; Suarez-Huerta et al., 2001; Webb et al., 1998; Wildman et al., 2003; Zambon et al., 2000
		Rat: 0.50 - 3.6 (1x) ^a 0.2 (1x) ^a	Rat: 0.20 - 2.6 (1x-↑3x) ^a	Rat: 0.020 - >100 (↓25->500x) ^c	Rat: competitive antagonist		
		Dog: ≈2 (1x) ^a			Chimpanzee: 0.0246 - 0.0445 (↓6-8x ^c / ↓4-6x ^d) K _i = 0.011 (1x ^c / ↑2x ^d)		
		Pig: 0.53 (↑5x) ^a					*Johnson, Donchue, & Shaffer, 2002

Table 2 (continued)

Physiological mononucleotides							
No.	Name	EC ₅₀ in μM (potency higher (↑) or lower (↓) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
2-09	UDP	4.20 (↓280x) ^{a,b} 16.5 (↓72x ^a / ↓120x ^b) inactive at 100 (↓500x ^a / ↓1000x ^b) <hr/> Mouse: 3.21 (↑6x ^a / ↓3x ^b)	9.48 - 19.5 (↓8-70x) ^b inactive at 100 μM (↓>130x) ^b <hr/> Mouse: inactive at 100 μM	0.007 - 0.530 <hr/> Mouse: 0.0259 - 0.042	0.0024 - 0.160 (1x-↑15x) ^d <hr/> Mouse: 0.0084 - 0.018		Besada et al., 2006; Bogdanov et al., 1998; WO002007002945A2, 2006; WO002007002945A2, 2006; Br��ser et al., 2017; Carter et al., 2009; Chambers et al., 2000; Chen et al., 1996; Communi, Motte et al., 1996; Communi, Parmentier
		partial agonist with 61 % efficacy inactive at 1 mM (↓>670x ^a / ↓>1100x ^b) <hr/> Rat: 16 (↓80x) ^{a,b} <hr/> Dog: ≈10 (↓≈50x ^a / n.d. ^b) <hr/> Pig: 1.5 (↑2x ^a / ↓3x ^b) 35 % efficacy	(↓>230x ^a / ↓>390x ^b) <hr/> Rat: 6.3 (↓4x ^a / ↓2x ^b) partial agonist inactive at 100 μM (↓>95x ^a / ↓>160x ^b)	 <hr/> Rat: 0.0059 - 2.3	(↓2x) ^d <hr/> Rat: 5.23 (1x) ^d partial agonist <hr/> Chimpanzee: 0.0031 - 0.0068 (1x-↑2x) ^d <		

(continued on next page)

Table 2 (continued)

Physiological mononucleotides							
No.	Name	EC ₅₀ in μM (potency higher (↑) or lower (↓) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
2-10	UMP	inactive at 100 μM (n.d. ^a / ↓>2300x ^b)	inactive at 100 μM (↓>1150x) ^c	inactive at 100 μM (↓>2400x) ^c	0.0540 - 0.703 (↓14-100x ^c / ↓7-25x ^d) inactive (n.d.) ^{c,d} K _i = 0.695 (↓50x ^c / ↓35x ^d)		Chen et al., 1996; Communi, Motte et al., 1996; El-Tayeb et al., 2006; Hamel et al., 2011; US020090148850A1, 2008; Ko et al., 2007; Lazarowski & Harden, 1994; Zambon et al., 2000
		Rat: inactive at 100 μM (↓>500x) ^{a,b}		Rat: inactive at 300 μM (↓>130x) ^c	Mouse: 0.0634 - 0.082 (↓5-8x ^c / ↓7-13x ^d)		
		Dog: inactive at 100 μM (↓>500x ^a / n.d. ^b)			Chimpanzee: 0.169 - 0.289 (↓43-54x ^c / ↓27-45x ^d) K _i = 1.15 (↓88x ^c / ↓50x ^d) K _i = 5.10 (↓31x ^c / ↓84x ^d)		
2-11	CTP	5.63 (↓>66x ^a / ↓>120x ^b) Mouse: almost inactive at 100 μM (n.d.) ^{a,b} inactive at 1 mM (↓>670x ^a / ↓>1100x ^b) Rat: 6.8 (↓3x ^a / ↓2x ^b) >100 (↓>500x) ^{a,b} Pig: 8.4 (↓3x ^a / ↓16x ^b)	antagonist inactive at 100 μM (↓>500x) ^b Mouse: 25 (↓58x ^a / ↓96x ^b) Rat: 1.2 (1x) ^{a,b} 7.24 (↓14x ^a / ↓36x ^b)	n.d. Rat: inactive at 300 μM (↓>130x) ^c	3.7 (↓1500x ^c / ↓1700x ^d) inactive at 1 μM (n.d. ^c / ↓>10x ^d)		Chambers et al., 2000; Chen et al., 1996; Erb et al., 1993; Hamel et al., 2011; Jacobson et al., 2006; Kennedy et al., 2000; Lazarowski et al., 2001; Lazarowski & Harden, 1994; Lin et al., 1993; Nguyen et al., 1995; Shen et al., 2004; Wildman et al., 2003

Table 2 (continued)

Physiological mononucleotides						
No.	Name	EC ₅₀ in μM (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄	
2-12	CDP	n.d.	n.d.	88.0 ($\downarrow 290x^c$) partial agonist	0.319 ($\downarrow 130x^c$ / $\downarrow 150x^d$) ≈ 50 ($\downarrow 680x^c$ / $\downarrow 160x^d$) inactive at 1 μM ($1x^c$ / $\downarrow >10x^d$)	Carter et al., 2009; Chambers et al., 2000; Chen et al., 1996; Hamel et al., 2011; Lazarowski et al., 2001; Lazarowski & Harden, 1994; Robaye et al., 1997; Shen et al., 2004
		Pig: 87.1 ($\downarrow 32x^a$ / $\downarrow 160x^b$)		Mouse: inactive at 100 μM ($\downarrow >2400x^c$)		
		Rat: inactive at 100 μM ($\downarrow >500x^{a,b}$)		Rat: inactive at 300 μM ($\downarrow >130x^c$)		
2-13	TTP	n.d.	n.d.	n.d.	n.d.	Shen et al., 2004
		Pig: 8.6 ($\downarrow 3x^a$ / $\downarrow 16x^b$) 74 % efficacy				
2-14	TDP	n.d.	n.d.	7.7 ($\downarrow 26x^c$)	2.17 ($\downarrow 900x^c$ / $\downarrow 960x^d$)	Hamel et al., 2011; Robaye et al., 1997; Shen et al., 2004
		Pig: 31.4 ($\downarrow 12x^a$ / $\downarrow 59x^b$)				

^arelative to ATP; ^brelative to UTP; ^crelative to UDP; ^drelative to UDP-glucose.

2. The P2Y₂ receptor

The mouse P2Y₂R, formerly referred to as the P_{2U} receptor, was first cloned in 1993 from a cDNA library obtained from NG108-15 cells, a mouse N18TG2 neuroblastoma x rat C6 glioma hybrid cell line (Lustig, Shiao, Brake, & Julius, 1993). The human receptor was cloned shortly thereafter (Parr et al., 1994). P2Y₂R mRNA is predominantly expressed in the gastrointestinal (GI) tract and the immune system, in muscle (mostly skeletal and cardiac muscle), and in endocrine tissue (thyroid, parathyroid, and adrenal glands). Lower P2Y₂R mRNA expression was also detected in the CNS, lungs, liver and gallbladder, kidney and bladder, male and female reproductive organs, placenta, skin, bone, and adipose tissue (Moore et al., 2001; Uhlén et al., 2015). At protein level, the P2Y₂R was mostly detected in epithelial and glandular cells of the lungs (nasopharynx and bronchi), gastrointestinal tract (salivary glands, oral mucosa, esophagus), tonsils, breasts, male and female reproductive tracts (prostate, epididymis, fallopian tubes, to a lower extent also in vagina, cervix, uterus, endometrium), and in placenta trophoblasts. Low P2Y₂R protein expression was also observed in immune tissue (bone marrow, spleen, lung macrophages, lymphoid tissue of the appendix) and skin keratinocytes (Uhlén et al., 2015). An overview of the tissue distribution and therapeutic significance of the P2Y₂R and

the other uracil nucleotide-activated P2YRs is provided in Table 1 (for discussion, see below).

A relatively large number of agonists – mostly derivatives of the endogenous ligand UTP – and a few antagonists have been described so far. Those of greater potential are discussed in the subsequent sections, while a comprehensive overview is provided in Tables 2 to 10.

2.1. P2Y₂ receptor agonists

2.1.1. Therapeutic potential of P2Y₂ receptor agonists

P2Y₂R activation bears significant therapeutic potential for different applications, including cystic fibrosis. The P2Y₂R is involved in epithelial ion transport. Na⁺ absorption by the airway epithelium is reduced as a result of P2Y₂R activation, and knockout mice were found to exhibit defective Cl⁻ secretion. Thus, P2Y₂R agonism was expected to compensate for the malfunctioning of the Cl⁻ channel cystic fibrosis transmembrane conductance regulator (CFTR) in cystic fibrosis patients (Cressman et al., 1999; Kunzelmann & Mall, 2003). In the eye, P2Y₂Rs mediate tear production and reduce intraocular pressure. P2Y₂R agonists are therefore useful for the treatment of dry eyes, and may also be beneficial for treating ocular hypertension and retinal degeneration (Jacobson & Civan, 2016; Lau, Samarawickrama, & Skaliky, 2014; Pintor et al.,




Fig. 3. Structures of agonists at uracil nucleotide-activated P2YRs with substitutions at the nucleobase (for data, refer to Table 3).

Table 3

Base-modified nucleotides as agonists for the uracil-activated P2YRs. Shown are the EC₅₀ or, in some cases, K_i and K_B values in μM at the human receptor, unless stated otherwise. The correlation factor by which the potency is n-fold higher (↑*nx*) or lower (↓*nx*) compared to that of the endogenous agonist in the same test system is given in brackets. Particularly interesting values are shown in bold.

Base-substituted mononucleotides							
No.	Name	EC ₅₀ in μM				Comments	References
		(potency higher (↑) or lower (↓) relative to endogenous agonist)					
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
3-01	2-S-UTP	≈0.007 - 0.050 (↑2x ^a / 1x ^b)	1.77 (↓20x) ^b 0.35 (↓5x) ^b	≈1.5 (↓36x) ^c	n.d.	10-fold selective for P2Y ₂ R Commercially available	El-Tayeb et al., 2006; Jacobson et al., 2006; Sakuma et al., 2017
3-02	2-S-UDP	≈50 (n.d. ^a / ↓1200x ^b)	≈40 (↓460x) ^b	0.239 (↓6x) ^c 0.447 (1x) ^c	0.00192 (↑83x ^c / ↑210x ^d) 0.002 (↑37x ^c / ↑160x ^d)	125-fold P2Y ₁₄ R selective	Carter et al., 2009; Das et al., 2010; El-Tayeb et al., 2006
3-03	2-S-UDP-glucuronic acid	n.d.	n.d.	n.d.	0.42 (n.d. ^c / 1x ^d)		Ko et al., 2007
3-04	2-Phenylethyl thio-UDP	0.544 (n.d. ^a / ↓30x ^b)	≥100 (↓≥1100x) ^b	≈2.5 (↓52x) ^c	n.d.		El-Tayeb et al., 2011
3-05	2-Phenylethyl thio-UMP	1.32 (n.d. ^a / ↓74x ^b)	>100 (↓>1100x) ^b	>100 (↓>2100x) ^c	n.d.		El-Tayeb et al., 2011
3-06	2-Amino-UDP (iso-CDP)	0.604 (n.d. ^a / ↓34x ^b)	≥100 (↓≥1100x) ^b	≥100 (↓≥2100x) ^c	n.d.	170-fold P2Y ₂ R selective	El-Tayeb et al., 2011
3-07	2-Amino-UMP (iso-CMP)	>100 (n.d. ^a / ↓>5600x ^b)	4.98 (↓56x) ^b	>100 (↓>2100x) ^c	n.d.	>20-fold P2Y ₄ R selective 2-(Me-amino)-UMP inactive at P2Y ₄ R	El-Tayeb et al., 2011
3-08	2-S-4-MeS-UTP	0.91 (n.d. ^a / ↓15x ^b)	5.35 (↓59x) ^b	>10 (↓>33x) ^c	n.d.		Ko et al., 2008
3-09	3-Me-UTP	0.564 (n.d. ^a / ↓13x ^b) 1.20 (↓14x ^a / ↓25x ^b)	2.92 (↓34x) ^b 3.40 (↓47x) ^b	≈8 (↓5300x) ^c	n.d.		El-Tayeb et al., 2006; Jacobson et al., 2006
3-10	3-Me-UDP	n.d.	n.d.	3.3 (↓250x) ^c	n.d.		Besada et al., 2006
3-11	3-Phenacyl-UDP (PSB-0474)	≈40 (n.d. ^a / ↓930x ^b)	>100 (↓>1200x) ^b	0.070 (↓2x) ^c	n.d. <div>Rat: inactive at 10 μM (↓>2x^c / ↓>2x^d)</div>	570-fold P2Y ₆ R selective Commercially available	El-Tayeb et al., 2006; Gao et al., 2010

(continued on next page)

Table 3 (continued)

Base-substituted mononucleotides							
No.	Name	EC ₅₀ in μ M (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
						Potency of 3-phenacyl-UTP in the μ M range at P2Y ₂ R, P2Y ₄ R, and P2Y ₆ R	
3-12	4-S-UTP	0.026 - 0.71 (\uparrow 3x ^a / \uparrow 2x- \downarrow 6x ^b)	0.023 (\uparrow 3x) ^b	n.d.	n.d.		Brunschweiler & Müller, 2006; Jacobson et al., 2006; Shaver et al., 1997
3-13	4-S-UDP	n.d.	n.d.	0.08 (\downarrow 6x) ^c 2.36 (\downarrow 5x) ^c	0.320 (\downarrow 2x ^c / 1x ^d)		Besada et al., 2006; Das et al., 2010
3-14	Zebularine triphosphate	8.9 (\downarrow 100x ^a / \downarrow 180x ^b)	inactive at 10 μ M (\downarrow >140x) ^b	n.d.	n.d.		Jacobson et al., 2006
3-15	N ⁴ -phenylpropoxy -CTP (MRS4062)	0.640 (n.d. ^a / \downarrow 12x ^b)	0.023 (\uparrow 4x) ^b	0.740 (n.d.) ^c	n.d.	28-fold selective for P2Y ₄ R Commercially available	Jayasekara et al., 2013; Maruoka et al., 2011
3-16	N ⁴ -phenylethoxy- CTP	1.20 (n.d. ^a / \downarrow 22x ^b)	0.073 (1x) ^b	1.21 (n.d.) ^c	n.d.	16-fold selective for P2Y ₄ R	Jayasekara et al., 2013; Maruoka et al., 2011
3-17	N ⁴ -benzyloxy- CDP (MRS2964)	2.13 (n.d. ^a / \downarrow 36x ^b)	1.15 (\downarrow 13x) ^b	0.026 (\uparrow 12x) ^c	n.d.	44-fold selective for P2Y ₆ R	Jayasekara et al., 2013; Maruoka et al., 2010
3-18	N ⁴ -MeO-CTP	0.05 (n.d. ^a / 1x ^b)	0.05 (\uparrow 2x) ^b	0.08 (\uparrow 4x) ^c	n.d.		Jayasekara et al., 2013; Maruoka et al., 2010
3-19	N ⁴ -MeO-CDP	3.60 (n.d. ^a / \downarrow 60x ^b)	6.45 (\downarrow 72x) ^b	0.070 (\uparrow 4x) ^c	3.32 (\downarrow 21x ^c / \downarrow 8x ^d)	4-fold more potent than UDP at P2Y ₆ R 47-fold selective for P2Y ₆ R	Das et al., 2010; Jayasekara et al., 2013; Maruoka et al., 2010
3-20	MRS4141	6.20 (n.d.) ^{a,b}	>10 (n.d.) ^b	0.039 (n.d.) ^c	n.d.		Jayasekara et al., 2013
3-21	N ⁴ -(3-(4- methoxyphenyl)- propyl)oxy-UTP	0.047 (n.d.) ^{a,b}	0.023 (\uparrow 4x) ^b	0.277 (n.d.) ^c	n.d.		Jayasekara et al., 2014
3-22	4-(Hexyl-S)-UTP	0.84 (n.d. ^a / \downarrow 7x ^b)	n.d.	n.d.	n.d.		Shaver et al., 1997

Table 3 (continued)

Base-substituted mononucleotides							
No.	Name	EC ₅₀ in μ M (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
3-23	5-Br-UTP	0.347 - 2.06 (\downarrow 9x ^a / \downarrow 8x-15 ^b)	2.1 - 49 (\downarrow 14-75x) ^b	0.291 (\downarrow 7x) ^c 0.800 (\downarrow 3x) ^c <hr/> Rat: 9 (\downarrow 4x) ^c inactive at 100 μ M (\downarrow >530x) ^c	n.d.	Patented	Communi, Parmentier et al., 1996; El-Tayeb et al., 2006; Jacobson et al., 2006; US000005620676A, 1995; Lazarowski et al., 1995; Lazarowski & Harden, 1994; Nguyen et al., 1995; Nicholas et al., 1996
3-24	5-Br-UDP	3.67 (n.d. ^a / \downarrow 85x ^b)	7.20 (\downarrow 83x) ^b inactive at 1 mM (\downarrow >1300x) ^b	0.151 (\downarrow 4x) ^c 0.800 (\downarrow 3x) ^c <hr/> Rat: 0.13 (\uparrow 2x) ^c	n.d.		Communi, Parmentier et al., 1996; El-Tayeb et al., 2006; Nicholas et al., 1996
3-25	5-I-UTP	0.83 (\downarrow 10x ^a / \downarrow 17x ^b)	4.0 (\downarrow 55x) ^b	n.d.	n.d.		Jacobson et al., 2006
3-26	5-I-UDP (MRS2693)	n.d.	n.d.	0.015 (1x) ^c	n.d.	Commercially available	Besada et al., 2006
3-27	5-F-UTP	6 (n.d. ^a / \downarrow 60x ^b)	0.6 (1x) ^b	>100 (\downarrow >710x) ^c	n.d.	Possibly a partial agonist Patented	Ginsburg-Shmuel et al., 2010; US000005620676A, 1995
3-28	5-F-UDP	2 (n.d. ^a / \downarrow 20x ^b)	3.5 (\downarrow 7x) ^b	10 (\downarrow 71x) ^c	n.d.	Possibly a partial agonist	Ginsburg-Shmuel et al., 2010
3-29	5-Methyl-UTP	0.48 (\downarrow 6x ^a / \downarrow 10x ^b)	3.9 (\downarrow 53x) ^b	n.d.	n.d.		Jacobson et al., 2006
3-30	5-Ethyl-UTP	n.d. <hr/> Mouse: 99 (\downarrow 6x ^a / \downarrow 79x ^b)	n.d.	n.d.	n.d.		Knoblauch et al., 1999
3-31	5-MeO-UDP	inactive (n.d.) ^{a,b}	\geq 20 (\downarrow >40x) ^b	0.08 (\uparrow 2x) ^c	n.d.		Ginsburg-Shmuel et al., 2010
3-32	5-Amino-UDP	n.d.	n.d.	0.61 (\downarrow 2x) ^c	n.d.		Ko et al., 2008
3-33	6-Aza-UTP	8.6 (\downarrow 100x ^a / \downarrow 180x ^b)	>10 (\downarrow >140x) ^b	n.d.	n.d.		Jacobson et al., 2006
3-34	Pseudouridine 5'-triphosphate	0.78 (\downarrow 9x ^a / \downarrow 16x ^b)	3.0 (\downarrow 41x) ^b	n.d.	n.d.		Jacobson et al., 2006

(continued on next page)

Table 3 (continued)

Base-substituted mononucleotides							
No.	Name	EC ₅₀ in μ M (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
3-35	Nucleotide triphosphate with 'unnatural' bicyclic aromatic base substitution	0.003 (n.d. ^a / \uparrow 2x ^b)	>20 (\downarrow >510x) ^b	>20 (n.d.) ^c	n.d.	>6600-fold P2Y ₂ R selective Patented	Brookings et al., 2007; Davenport et al., 2007; WO002002062816A1, 2002; WO002003011885A1, 2002
3-36	2-Cl-ATP	2.30 (\downarrow 10x ^a / \downarrow 16x ^b) "low activity"	n.d.	n.d.	n.d.	Patented	Guile et al., 2001; US0000005620676A, 1995; Lazarowski et al., 1995
3-37	2-MeSATP	inactive (n.d.) ^{a b} Rat: 9.4 (\downarrow 4x ^a / \downarrow 3x ^b) partial agonist with 34 % efficacy >100 (\downarrow >500x) ^{a b} Mouse: \approx 100 (n.d.) ^{a b} \approx 1000 (\downarrow 1400x ^a / \downarrow 900x ^b) Dog: >10 (\downarrow >5x ^a / \downarrow >5x ^b) Pig: inactive at 100 μ M (\downarrow 37x ^a / \downarrow 190x ^b)	inactive at 1 mM (\downarrow >1300x) ^b Rat: 1.4 (1x) ^{a b} partial agonist with 24 % efficacy 2.1 (1x) ^{a b} partial agonist inactive at 100 μ M (\downarrow >95x ^a / \downarrow >160x ^b)	100 (\downarrow 330x) ^c Rat: 50 (n.d.) ^c \approx 1000 (\downarrow 5300x) ^c inactive at 300 μ M (\downarrow >130x) ^c	n.d.	Patented	Bogdanov et al., 1998; Chang et al., 1995; Chen et al., 1998; Communi, Parmentier et al., 1996; Guile et al., 2001; US0000005620676A, 1995; Lazarowski & Harden, 1994; Lin et al., 1993; Lustig et al., 1993; Nicholas et al., 1996; Shen et al., 2004; Webb et al., 1998; Wildman et al., 2003; Zamboni et al., 2000

Table 3 (continued)

Base-substituted mononucleotides							
No.	Name	EC ₅₀ in μ M (potency higher (↑) or lower (↓) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
3-38	8-Br-ATP	23.0 (↓100x ^a / ↓160x ^b)	n.d.	0.800 (↓3x) ^c	n.d.	Patented	Communi, Parmentier et al., 1996; US000005620676A, 1995; Lazarowski et al., 1995
3-39	MRS4160	0.109 (n.d.) ^{a,b}	0.040 (↑2x) ^b	0.183 (n.d.) ^c	n.d.	Suitable for chain extension by click tethering Derivative of 3-21	Jayasekara et al., 2014
3-40	MRS4162	0.066 (n.d.) ^{a,b}	0.070 (1x) ^b	0.023 (n.d.) ^c	n.d.	Fluorescent probe Derivative of 3-21	Jayasekara et al., 2014
3-41	MRS4063	3.9 (n.d.) ^{a,b}	0.952 (n.d.) ^b	0.100 (↑3x) ^c	n.d.	Suitable for chain extension using click tethering	Jayasekara et al., 2013
3-42	MRS4129	2.5 (n.d.) ^{a,b}	>10 (n.d.) ^b	0.009 (↑33x) ^c	n.d.	Fluorescent probe	Jayasekara et al., 2013

^arelative to ATP; ^brelative to UTP; ^crelative to UDP; ^drelative to UDP-glucose.

2003; Pintor, Carracedo, Alonso, Bautista, & Peral, 2002). The concept of using P2Y₂R agonists to promote secretion and mucus clearance for the treatment of dry eyes, ocular hypertension, and retinal degeneration has been patented, as well as their use in the treatment of sinusitis, ear infections, bronchitis, pneumonia, gastrointestinal tract disorders, vaginal dryness, and for joint lubrication (Cowlen, Yerxa, Jones, & Brown, 2002; Drutz, Rideout, & Jacobus, 1997; Jacobus & Leighton, 1997; Pendergast, Rideout, Siddiqi, & Yerxa, 1998; Pendergast, Shaver, Drutz, & Rideout, 1999; Peterson, 2001; Peterson & Yerxa, 2002; Shaffer, Boucher, Rideout, & Jacobus, 1997; Yerxa et al., 2000; Yerxa, Peterson, Rideout, & Pendergast, 2009). P2Y₂R agonism may possibly also find application in cardiovascular incidents, as it was shown to reduce post-ischemic myocardial damage *in vivo* and protect cardiomyocytes from hypoxia *in vitro* (Cohen et al., 2011; Hochhauser et al., 2013). Moreover, a neuroprotective role was postulated for the P2Y₂R via the induction of α -secretase-dependent amyloid precursor protein processing in astrocytoma cells, microglia-mediated clearance of β -amyloid, and modulation of the ubiquitin-proteasome system (Ajit et al., 2014; Camden et al., 2005; Diego-García et al., 2017; Kim et al., 2012). In addition, a reduced P2Y₂R expression was observed post mortem in the parietal cortex of patients who had suffered from Alzheimer's disease (Lai et al., 2008).

2.1.2. Mononucleotides and analogues

Almost all of the current P2Y₂R agonists are nucleotide derivatives, which are generally not ideal for drug development or *in vivo* studies due to their susceptibility to enzymatic degradation and their lack of oral bioavailability as a result of the several negative charges at physiological pH.

The endogenous agonists for the P2Y₂R are ATP (2-01, compound 01 in Fig. 2 and Table 2) and UTP (2-08, compound 08 in Fig. 2 and Table 2)

with similar potencies. Many commercial preparations of uridine 5'-diphosphate (UDP, 2-09, Fig. 2 and Table 2) were found to contain significant amounts of UTP that can activate the P2Y₂R. Furthermore, UDP can be phosphorylated to UTP by nucleoside diphosphokinase. These effects initially caused misinterpretations regarding the potency of UDP. Following purification of the test compounds, and in experimental conditions that excluded enzymatic conversion, both UDP and adenosine 5'-diphosphate (ADP, 2-02, Fig. 2 and Table 2) were found inactive at the human P2Y₂R (Lazarowski et al., 1995; Nicholas et al., 1996). Adenosine 5'-(γ -thio)-triphosphate (ATP γ S, 5-03, compound 03 in Fig. 5 and Table 5) and uridine 5'-(γ -thio)-triphosphate (UTP γ S, 5-05, Fig. 5 and Table 5) are also full agonists. The γ -thiophosphate group increases stability towards ectonucleotidases (Lazarowski et al., 1995; Malmström et al., 2000). A thio-substitution at the α -phosphate group caused a reduction in potency (Jacobson et al., 2006).

UTP (2-08, compound 08 in Fig. 2 and Table 2) is not only an agonist for the P2Y₂R but activates the related P2Y₄R as well. Relatively simple nucleobase modifications (Fig. 3 and Table 3) to UTP may enhance the selectivity for the P2Y₂R: 2-thio-UTP (3-01) was described as a P2Y₂R agonist with potency similar to UTP (EC₅₀ of 35–50 nM) and 10- to 30-fold selectivity versus the related P2Y₄ and P2Y₆ receptors (El-Tayeb, Qi, & Müller, 2006; Jacobson et al., 2006). Moreover, 2-amino-UDP (3-06), although 34-fold less potent (EC₅₀ of 604 nM), is at least 170-fold selective (El-Tayeb, Qi, Nicholas, & Müller, 2011). Further modifications, not only on the pyrimidine ring but additionally through substitutions at the ribose moiety, led to one of the most potent and selective P2Y₂R agonists known to date: 2'-amino-2'-deoxy-2-thio-UTP (MRS2698, 6-01, Fig. 6 and Table 6; EC₅₀ of 8 nM). It is 6-fold more potent than UTP and >300-fold selective versus the P2Y₄ and P2Y₆ receptors (Ivanov et al., 2007). Several thousandfold improvement in selectivity with potencies similar to UTP could be achieved by




Fig. 4. Structures of agonists at uracil nucleotide-activated P2Y₂R with substitutions at the ribose ring (for data, refer to Table 4).

replacing the nucleobase with various 'unnatural' bicyclic aromatic residues, as in compound **3-35** (Fig. 3 and Table 3) (Brookings et al., 2007). These structures could serve as templates for the development of more selective compounds than the uracil-derived P2Y₂R agonists.

Improved ectonucleotidase stability was achieved in a series of nucleobase-substituted nucleotide derivatives through β,γ -dichloro- and β,γ -difluoromethylene substitutions (El-Tayeb et al., 2011). The most promising compound of this series is 4-thio- β,γ -difluoromethylene-UTP (PSB-1114, **6-05**, Fig. 6 and Table 6) with >50-fold selectivity versus the P2Y₄ and P2Y₆ receptors. PSB-1114, which is available from commercial sources, is with an EC₅₀ of 134 nM approx. 8-fold less potent than UTP (El-Tayeb et al., 2011).

Masking of the terminal phosphate group may also be a strategy to reduce the susceptibility of nucleotides to ectonucleotidase hydrolysis.

Up δ -3-chlorophenyl phosphoester (**5-17**, Fig. 5 and Table 5) exhibited an EC₅₀ value of 840 nM at the P2Y₂R and is 15-fold less potent than UTP (Maruoka et al., 2011). It showed no effect on the P2Y₄R at 10 μ M, but the selectivity versus the P2Y₆R was merely 4-fold (Maruoka et al., 2011). Up δ - δ -phenyl ester (MRS2768, **5-02**, Fig. 5 and Table 5) also displayed some selectivity for the P2Y₂R, as no effect was observed at 10 μ M for the P2Y₄R and P2Y₆R subtypes. However, the potency is with an EC₅₀ of 1.89 μ M at the P2Y₂R lower compared to many other agonists (Ko et al., 2008). Moreover, the stability of such esters in biological media may be limited.

2.1.3. Dinucleotides

Nucleoside triphosphates can be mimicked by dinucleotide tetraphosphates, as both classes of compounds bear four negative

Table 4

Mononucleotides with modifications at the ribose ring as agonists for the uracil-activated P2YRs. Shown are the EC₅₀ or, in some cases, K_i and K_B values in μM at the human receptor, unless stated otherwise. The correlation factor by which the potency is n-fold higher ($\uparrow\text{nx}$) or lower ($\downarrow\text{nx}$) compared to that of the endogenous agonist in the same test system is given in brackets. Particularly interesting values are shown in bold.

Ribose-modified mononucleotides							
No.	Name	EC ₅₀ in μM (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
4-01	2'-Me-UTP	1.45 (n.d. ^a / $\downarrow\text{24x}^b$)	1.26 ($\downarrow\text{14x}^b$)	inactive at 10 μM ($\downarrow\text{>33x}^c$)	n.d.		Ko et al., 2008
4-02	2'-Deoxy-UTP	1.08 ($\downarrow\text{13x}^a$ / $\downarrow\text{22x}^b$)	1.9 ($\downarrow\text{26x}^b$)	n.d. Rat: inactive at 300 μM ($\downarrow\text{>130x}^c$)	n.d.		Jacobson et al., 2006; Lazarowski & Harden, 1994
4-03	2'-Deoxy-UDP	n.d.	n.d.	1.72 ($\downarrow\text{130x}^c$) Rat: ≈ 1000 ($\downarrow\text{440x}^c$)	n.d.		Besada et al., 2006; Costanzi et al., 2005; Lazarowski & Harden, 1994
4-04	2'-Amino-2'-deoxy-UTP	0.062 (1x) ^{a,b}	1.2 ($\downarrow\text{16x}^b$)	inactive at 10 μM ($\downarrow\text{>33x}^c$)	n.d.		Jacobson et al., 2006; Ko et al., 2008
4-05	2'-Amino-2'-deoxy-UDP	n.d.	n.d.	3.9 ($\downarrow\text{300x}^c$)	n.d.		Besada et al., 2006
4-06	2'-Azido-2'-deoxy-UTP	5.0 ($\downarrow\text{59x}^a$ / $\downarrow\text{100x}^b$)	1.1 ($\downarrow\text{15x}^b$)	n.d.	n.d.		Jacobson et al., 2006
4-07	2'-Azido-2'-deoxy-UDP	n.d.	n.d.	1.5 ($\downarrow\text{120x}^c$)	n.d.		Besada et al., 2006
4-08	2'-Deoxy-2'-F-UTP	0.78 ($\downarrow\text{9x}^a$ / $\downarrow\text{16x}^b$)	0.54 ($\downarrow\text{7x}^b$)	n.d.	n.d.		Jacobson et al., 2006
4-09	2'-Deoxy-2'-MeO-UTP	14.3 ($\downarrow\text{170x}^a$ / $\downarrow\text{290x}^b$)	8.2 ($\downarrow\text{110x}^b$)	n.d.	n.d.		Jacobson et al., 2006
4-10	2'-Deoxy-2'-ureido-UTP	1.74 (n.d. ^a / $\downarrow\text{29x}^b$)	4.64 ($\downarrow\text{52x}^b$)	>10 ($\downarrow\text{>33x}^c$)	n.d.		Ko et al., 2008
4-11	2'-Deoxy-2'-ureido-UDP	n.d.	n.d.	4.70 ($\downarrow\text{16x}^c$)	n.d.		Ko et al., 2008
4-12	2'-F-2'-deoxyara-UDP	n.d.	n.d.	5.5 ($\downarrow\text{423x}^c$)	n.d.		Besada et al., 2006
4-13	Arabino-UTP	0.087 (1x ^a / $\downarrow\text{2x}^b$)	0.71 (1x) ^b	n.d.	n.d.		Jacobson et al., 2006
4-14	2'-Deoxyarabino-2'-F-UTP	0.52 ($\downarrow\text{6x}^a$ / $\downarrow\text{11x}^b$)	0.52 ($\downarrow\text{7x}^b$)	n.d.	n.d.		Jacobson et al., 2006
4-15	3'-Deoxy-UDP	n.d.	n.d.	2.5 ($\downarrow\text{192x}^c$)	n.d.		Besada et al., 2006
4-16	3',4'-Cyclopropyl-UDP	n.d.	n.d.	3.5 ($\downarrow\text{269x}^c$)	n.d.		Besada et al., 2006
4-17	3'-O-(4-benzoyl)-benzoyl-ATP (BzATP)	n.d. Rat:	n.d. Rat:	n.d.	n.d.	ATP analogue used as photoaffinity label	Erb et al., 1993; Lin et al., 1993; Wildman et al., 2003

(continued on next page)

Table 4 (continued)

Ribose-modified mononucleotides							
No.	Name	EC ₅₀ in μ M (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
		4.7 ($\downarrow 2x^a$ / $1x^b$) Mouse: 104 % receptor activation at 100 μ M	antagonist (see Table 10)				
4-18	(N)-methano-carba-UTP	0.0159 ($\downarrow 5x^a$ / $\downarrow 2x^b$)	0.085 ($\downarrow 2x$) ^b	inactive at 100 μ M ($\downarrow >6700x$) ^c	n.d.		Kim et al., 2002
4-19	(N)-methano-carba-ATP	0.091 ($1x^a$ / $\downarrow 11x^b$)	>10 ($\downarrow >200x$) ^b	n.d.	n.d.	Methanocarba-adenosine analogues patented	WO002001051490A1, 2001; Kim et al., 2002
4-20	(S)-methano-carba-UTP	0.08 (n.d. ^a / $1x^b$)	0.30 ($\downarrow 3x$) ^b	1.37 ($\downarrow 5x$) ^c	n.d.		Maruoka et al., 2010
4-21	(S)-methano-carba-ATP	3.7 ($\downarrow 44x^a$ / $\downarrow 460x^b$)	inactive at 100 μ M ($\downarrow >2000x$) ^b	n.d.	n.d.		Kim et al., 2002
4-22	(S)-methano-carba-UDP (MRS2795)	n.d.	n.d.	0.042 ($\uparrow 2x$) ^c	inactive at 10 μ M ($\downarrow >19x$) ^c	(N)-methano-carba-UDP inactive at P2Y ₆ R	Besada et al., 2006; Das et al., 2010; Maruoka et al., 2010
4-23	2'-Deoxy-(S)-methanocarba-UDP (MRS2633)	n.d.	n.d.	0.230 ($\downarrow 18x$) ^c	n.d.		Besada et al., 2006; Costanzi et al., 2005
4-24	L- α -threo furanosyl-UTP (MRS2488)	9.9 (n.d. ^a / $\downarrow 1200x^b$)	26 ($\downarrow 530x$) ^b	inactive at 10 μ M (n.d.) ^c	n.d.		Ohno et al., 2004

^arelative to ATP; ^brelative to UTP; ^crelative to UDP; ^drelative to UDP-glucose.

charges at physiological pH (Brunschweiler & Müller, 2006). The naturally occurring dinucleotides P^1, P^4 -di(adenosine-5')-tetrphosphate (AP₄A, **7-01**, Fig. 7 and Table 7) and P^1, P^4 -di(uridine-5')-tetrphosphate (Up₄U, **7-03**, Fig. 7 and Table 7) are potent P2Y₂R agonists, albeit non-selective versus the P2Y₄R. Up₄U is also known as diquafosol or INS365, and is marketed since 2010 under the trade name of Diquas® in Japan, Korea, Thailand, and Vietnam as a treatment for dry eye syndrome (see reviews by Lau et al., 2014; Keating, 2015, and Nichols, Yerxa, & Kellerman, 2004). It is an analogue of the diadenosine polyphosphates naturally found in human tears that were shown to promote tear secretion (Mundasat et al., 2001; Pintor et al., 2002; Tauber et al., 2004; Yamane et al., 2015). AP₄A was shown to induce controlled hypotension in patients and exert modulating effects on blood pressure during anesthesia (Kikuta, Ohiwa, Okada, Watanabe, & Haruki, 1999). Dinucleotides are less susceptible to enzymatic hydrolysis *in vivo* than mononucleotides with a free terminal phosphate group. Replacement of one of the two uridine moieties with 2'-deoxycytidine

to form Up₄dC (denufosol, INS37217, **7-07**, Fig. 7 and Table 7) further enhanced ectonucleotidase stability. Denufosol was shown to significantly enhance tracheal mucus transport for more than 8 hours in an animal model (Deterding et al., 2007; Yerxa et al., 2002). It was subsequently evaluated in two phase III clinical trials as a therapy for cystic fibrosis patients. In the first trial (TIGER-1), it was found to significantly improve lung function in cystic fibrosis patients with normal to mildly impaired lung function. However, this could not be confirmed in the second phase III clinical trial: in the TIGER-2 study, denufosol failed to meet the primary endpoint, a significant change in baseline FEV1 (forced expiratory volume in one second) at week 48 (Accurso et al., 2011; Kellerman et al., 2008; Ratjen et al., 2012; for a detailed account on the denufosol clinical trials, refer to the commentary by Moss, 2013). A series of further dinucleotides with potencies in the high nanomolar or low micromolar range at the P2Y₂R is described in the patent literature (Pendergast et al., 1998; Rideout, Yerxa, Shaver, & Douglass III, 2003).




Fig. 5. Structures of agonists at uracil nucleotide-activated P2YRs with substitutions at the phosphate chain (for data, refer to Table 5).

2.1.4. An allosteric P2Y₂ receptor partial agonist

A carbon-phosphorus bond cannot be hydrolyzed by ectonucleotidases. Thus, in an attempt to enhance metabolic stability, the α-phosphate group of UTP was replaced by an isosteric phosphonate (Cosyn et al., 2009). It was subsequently discovered that the entire triphosphate group could be replaced by a smaller 5'-methylenephosphonate moiety. This modification also enhanced the compounds' selectivity for the P2Y₂R over the P2Y₄R (van Poecke et al., 2012). For a series of uridine 5'-methylenephosphonates with varying substitutions in the 5-position of the uracil base the potencies were in the high nanomolar to low micromolar range. The most potent compound, SVP333 (**6-11**, Fig. 6 and Table 6), displayed an EC₅₀ value of 400 nM (van Poecke et al., 2012). The maximal receptor stimulation elicited by these compounds was not higher than half of that observed with UTP, suggesting that they act as partial agonists. High concentrations (100 μM) did not affect UTP-mediated receptor activation, as would otherwise be expected from orthosteric partial agonists. They were thus described as allosteric partial agonists (van Poecke et al., 2012). Replacing the 5-(4-fluorophenyl)uracil ring of SVP333 (**6-11**, Fig. 6 and Table 6) with quinazoline-2,4-dione led to a reduction in potency (Song et al., 2014).

2.1.5. Discovery of a non-nucleotide allosteric P2Y₂ receptor modulator

Very recently, the discovery of a weakly potent non-nucleotide agonist for the P2Y₂R was reported (Sakuma, Nakagawa, Oikawa, Noda, & Ikeda, 2017). The 4(1H)-quinoline derivative 2-((ethyl(4-fluorobenzyl)amino)methyl)-7,8-dimethylquinolin-4(1H)-one (**9-01**, Fig. 9 and Table 9), referred to as "Compound 89", exhibited an EC₅₀ value of 10.5 μM at the human P2Y₂R, and of 2.7 μM at the mouse P2Y₂R in calcium mobilization assays using recombinant 1321N1 astrocytoma cells. It was described as a partial agonist with 66 % (human P2Y₂R) and 51 % (mouse P2Y₂R) efficacy. However, these maximum efficacy values were calculated by extrapolation and could not be determined experimentally due to the low potency and limited solubility. **9-01** was inactive at 30 μM at the human P2Y₁R, P2Y₄R, P2Y₆R, P2Y₁₁R, and P2Y₁₂R, and was thus reported to be selective for the

P2Y₂R (Sakuma et al., 2017). In recombinant 1321N1 astrocytoma cells, it activated the P2Y₂R in the absence of an orthosteric agonist but also exhibited positive allosteric modulation in combination with the agonists ATP, UTP, and AP₄A; it thus appears to function as an ago-allosteric agonist (for more information on allosteric modulation of GPCRs, see Müller, Schiedel, & Baqi, 2012 and Christopoulos et al., 2014). In neonatal rat cardiomyocytes, **9-01** behaved as a positive allosteric modulator only (Sakuma et al., 2017). Being the only non-nucleotide P2Y₂R agonist described in the literature so far, **9-01** might serve as a lead structure for the development of analogues with higher potency that could be useful tool compounds with the potential to become therapeutics.

2.1.6. N⁴-Substituted cytidine 5'-triphosphate (CTP) derivatives

Recently, a series of CTP derivatives with varying (3-arylpropyl)oxy substitutions on the cytosine amino group (N⁴) was published (Jayasekara et al., 2014). These compounds revealed that sterically demanding groups in this region of the nucleotide were tolerated by the P2Y₂R, P2Y₄R, and P2Y₆R. The potencies and selectivity profiles could be modulated by structural changes of the N⁴-substituent. Among these compounds showed N⁴-(3-(4-methoxyphenyl)propyl)oxy-UTP (**3-21**, Fig. 3 and Table 3) relatively high potencies on the P2Y₂R (EC₅₀ of 47 nM), the P2Y₄R (EC₅₀ of 23 nM), and the P2Y₆R (EC₅₀ of 277 nM) (Jayasekara et al., 2014). Since steric bulk in this region does not significantly impede binding to those receptors, **3-21** was selected for the attachment of fluorophores and other reporter moieties as prosthetic groups. MRS4162 (**3-40**, Fig. 3 and Table 3) is a fluorescent derivative that features a boron-dipyrromethene (BODIPY®) 630/650 moiety (Jayasekara et al., 2014). MRS4162 exhibited potencies in the mid-nanomolar range on the P2Y₂R, P2Y₄R, and P2Y₆R. It was shown in a preliminary feasibility study using flow cytometry to label 1321N1 astrocytoma cells that express P2Y₆Rs but not those without P2YR expression (Jayasekara et al., 2014). MRS4162 is thus a useful tool for the labeling and quantification of the P2Y₆R (and most likely also the P2Y₂R and P2Y₄R) in living cells, similar to the development of fluorescent ligands for adenosine receptors (Dale, Hill, & Kellam, 2012; Kozma et al., 2012). However, it will be difficult to differentiate

Table 5

Nucleotide derivatives with substitutions at the phosphate groups as agonists for the uracil-activated P2Y_{Rs}. Shown are the EC₅₀ or, in some cases, K_i and K_B values in μM, unless stated otherwise. The correlation factor by which the potency is n-fold higher (↑**nx**) or lower (↓**nx**) compared to that of the endogenous agonist in the same test system is given in brackets. Particularly interesting values are shown in bold.

Modifications at the phosphate groups						
No.	Name	EC ₅₀ in μM				Comments
		(potency higher (↑) or lower (↓) relative to endogenous agonist)				References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄	
5-01	Up ₄	2.61 (n.d. ^a / ↓ 44x^b)	4.64 (↓ 52x^b)	7.56 (↓ 25x^c)	n.d.	Ko et al., 2008
5-02	Up ₄ -δ-phenyl ester (MRS2768)	1.89 (n.d. ^a / ↓ 32x^b)	>10 (↓> 110x^b)	>10 (↓> 33x^c)	n.d.	Commercially available Ko et al., 2008
5-03	ATPyS	0.570 - 1.72 (↓ 7-8x^a / ↓ 12-29x^b) Rat: 10.5 (↓ 4x^a / ↓ 3x^b) Mouse: 94 % receptor activation at 100 μM 7.9 (↓ 11x^a / ↓ 7x^b) 10.4 (↓ 7x^a / ↓ 12x^b) Dog: ≈1 (↓≈ 5x^a / n.d. ^b) Pig: 1.0 (↑ 3x^a / ↓ 2x^b) 54 % efficacy	inactive (n.d.) ^b Rat: 2.1 (1x) ^{a,b} partial agonist 5.4 (↓ 4x^a / ↓ 3x^b) partial agonist with 24 % efficacy	n.d. Rat: inactive at 300 μM (↓> 130x^c)	n.d.	Bogdanov et al., 1998; Communi, Motte et al., 1996; Erb et al., 1993; Janssens et al., 1999; Lazarowski et al., 1995; Lazarowski & Harden, 1994; Lin et al., 1993; Lustig et al., 1993; Shen et al., 2004; Wildman et al., 2003; Zambon et al., 2000
5-04	GTPyS	26.5 (↓ 120x^a / ↓ 190x^b) Pig: 1.2 (↑ 3x^a / ↓ 2x^b) 80 % efficacy	n.d.	n.d.	n.d.	Lazarowski et al., 1995; Shen et al., 2004
5-05	UTPyS	0.240 (1x ^a / ↓ 2x^b)	1.6 (↓ 2x^b)	n.d. Rat: weak agonist	n.d.	α-Thiophosphate unfavorable for P2Y ₂ R & P2Y ₄ R Erb et al., 1993; Jacobson et al., 2006; Lazarowski et al., 1996; Nicholas et al., 1996
5-06	UDPβS	inactive at 10 μM (n.d.) ^{a,b}	inactive at 10 μM (n.d.) ^b	0.028 (↑ 2x^c)	0.026 (↓ 3x^c / ↓ 12x^d)	Initially thought to be a competitive Carter et al., 2009; Fricks et al., 2008;

Table 5 (continued)

Modifications at the phosphate groups							
No.	Name	EC ₅₀ in μ M				Comments	References
		(potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)					
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
						antagonist at P2Y ₁₄ R	Hou et al., 2002
5-07	UMPS	n.d.	n.d.	62 % receptor activation at 10 μ M	81 % receptor activation at 10 μ M	AMPS & CMPS are also agonists with EC ₅₀ >10 μ M at P2Y ₁₄ R; TMPS is an antagonist	Gendaszewska-Darmach et al., 2016; Gendaszewska-Darmach & Szustak, 2016
5-08	TMPS	inactive at 1 mM (n.d.) ^{a,b}	inactive at 1 mM (n.d.) ^b	partial agonist with 50 % efficacy	antagonist (see Table 10)	AMPS, UMPS & CMPS are also P2Y ₁₄ R agonists	Gendaszewska-Darmach et al., 2016; Gendaszewska-Darmach & Szustak, 2016
5-09	R _p - α -S-UTP	5.4 (\downarrow 64x ^a / \downarrow 110x ^b)	27 (\downarrow 370x) ^b	n.d.	n.d.	2'-Deoxy-analogues less potent or inactive at P2Y ₂ R & P2Y ₄ R	Jacobson et al., 2006
5-10	β , γ -Imido-ATP (AppNHp)	5.66 (\downarrow 25x ^a / \downarrow 40x ^b) Mouse: inactive	n.d.	n.d.	n.d.		Erb et al., 1993; Lazarowski et al., 1995
5-11	β , γ -Imido-UTP (UppNHp)	1.45 (n.d.) ^{a,b}	n.d.	n.d.	n.d.		Brunschweiler & Müller, 2006
5-12	β , γ -Methylene-UTP (UppCH ₂ p)	73.3 (n.d.) ^{a,b}	n.d.	n.d.	n.d.		Brunschweiler & Müller, 2006
5-13	α , β -Methylene-UDP (UpCH ₂ p)	n.d.	n.d.	0.339 (\uparrow 2x) ^c 0.66 (\downarrow 2x) ^c	0.011 (\uparrow 15x ^c / \uparrow 36x ^d)		Das et al., 2010; Ko et al., 2008
5-14	β , γ -Difluoro methylene-UTP (UppCF ₂ p)	4.9 (n.d. ^a / \downarrow 270x ^b)	n.d.	n.d.	n.d.		El-Tayeb et al., 2011
5-15	α , β -Difluoro methylene-UDP (MRS2802)	n.d.	n.d.	inactive at 10 (\downarrow >33x) ^c	0.063 (\uparrow 3x ^c / \uparrow 6x ^d)		Carter et al., 2009; Das et al., 2010; Ko et al., 2008
5-16	Up ₂ - β -Me-phosphonate	n.d.	n.d.	8.0 (\downarrow 15x) ^c	4.58 (\downarrow 29x ^c / \downarrow 12x ^d)		Das et al., 2010
5-17	Up α - δ -3-chlorophenyl phosphoester	0.840 (n.d. ^a / \downarrow 15x ^b)	inactive at 10 μ M (\downarrow >130x) ^b	3.69 (n.d.) ^c	n.d.	4-fold selective for P2Y ₂ R	Maruoka et al., 2011

^arelative to ATP; ^brelative to UTP; ^crelative to UDP; ^drelative to UDP-glucose.




Fig. 6. Structures of agonists at uracil nucleotide-activated P2YRs with combined substitutions at the base, ribose ring, and/or phosphate chain (for data, refer to Table 6).

the response for cell lines where different P2YRs are simultaneously expressed. Developing analogues with improved selectivity may be a next step, as would be the development of assays for routine application

of this fluorescent probe. Comprehensive information on fluorescent probes for P2YRs (as well as adenosine receptors) and their use were published by Ciruela, Fernández-Duenas, & Jacobson, 2015.

Table 6

Mononucleotide derivatives with combined modifications at the base, ribose ring, and/or phosphate chain as agonists for the uracil-activated P2YRs. Shown are the EC₅₀ or, in some cases, K_i and K_B values in μ M, unless stated otherwise. The correlation factor by which the potency is n-fold higher (\uparrow nx) or lower (\downarrow nx) compared to that of the endogenous agonist in the same test system is given in brackets. Particularly interesting values are shown in bold.

Mononucleotide derivatives with combined substitutions							
No.	Name	EC ₅₀ in μM				Comments	References
		(potency higher (↑) or lower (↓) relative to endogenous agonist)					
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
6-01	2'-Amino-2'-deoxy-2-S-UTP (MRS2698)	0.008 (↑11x ^a / ↑6x ^b)	2.4 (↓33x) ^b	inactive at 10 μM (↓>440x) ^c	n.d.	300-fold P2Y ₂ R selective	Ivanov, Ko et al., 2007; Ko et al., 2008
6-02	2-S-arabino-UTP	0.14 (n.d. ^a / ↓2x ^b)	7.93 (↓88x) ^b	inactive at 10 μM (↓>33x) ^c	n.d.		Ko et al., 2008
6-03	2-S-β,γ-dichloromethylene-UTP	2.51 (n.d. ^a / ↓42x ^b)	inactive at 10 μM (↓>110x) ^b	inactive at 10 μM (↓>33x) ^c	n.d.		Ko et al., 2008
6-04	2-S-β,γ-difluoromethylene-UTP	1.63 (n.d. ^a / ↓27x ^b)	8.11 (↓90x) ^b	5.15 (↓17x) ^c	n.d.		Ko et al., 2008
6-05	4-S-β,γ-difluoromethylene-UTP (PSB-1114)	0.134 (n.d. ^a / ↓8x ^b)	9.3 (↓110x) ^b	7.0 (↓150x) ^c	n.d.	50-fold selective for P2Y ₂ R Enhanced ectonucleotidase stability Commercially available	El-Tayeb et al., 2011
6-06	N3-phenacyl-β,γ-dichloromethylene-UTP	0.826 (n.d. ^a / ↓46x ^b)	7.3 (↓83x) ^b	0.142 (↓3x) ^c	n.d.		El-Tayeb et al., 2011
6-07	β,γ-Dichloromethylene-5-Br-UTP	0.354 (n.d. ^a / ↓20x ^b)	3.99 (↓45x) ^b	0.120 (↓3x) ^c	n.d.		El-Tayeb et al., 2006
6-08	α,β-Methylene-2-S-UDP (MRS2905)	n.d.	n.d.	1.99 (↓4x) ^c	0.00092 (↑170x ^c / ↑440x ^d)	Commercially available α,β-Methylene-UDP 12-fold less potent at P2Y ₁₄ R	Das et al., 2010
6-09	α,β-Methylene-5-I-UDP	n.d.	n.d.	0.13 (↑2x) ^c	n.d.		Maruoka et al., 2010
6-10	α,β-Difluoromethylene-5-I-UDP	n.d.	n.d.	0.127 (↑4x) ^c	0.142 (1x ^c / ↑3x ^d)		Das et al., 2010
6-11	SVP333	0.4 (n.d. ^a / ↓20x ^b)	>100 (n.d.) ^b	>100 (n.d.) ^c	n.d.	Allosteric partial agonist	van Poecke et al., 2012
6-12	5-MeO-uridine 5'-	inactive at 100 μM	inactive at 100 μM	0.008 (↑19x) ^c	n.d.	>12,500-fold	WO002012073237A1,

(continued on next page)

Table 6 (continued)

Mononucleotide derivatives with combined substitutions							
No.	Name	EC ₅₀ in μM				Comments	References
		(potency higher (↑) or lower (↓) relative to endogenous agonist)					
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
	O-(α-borano diphosphate), R _p isomer	(n.d. ^a / ↓>710x ^b)	(n.d. ^a / ↓110x ^b)			P2Y ₆ R selective	2011; Ginsburg-Shmuel et al., 2012
						Patented	
6-13	α,β-Methylene-N ⁴ -MeO-CDP	>10 (n.d. ^a / ↓>170x ^b)	inactive at 10 μM (↓110x) ^b	0.678 (↓2x) ^c	n.d.		Maruoka et al., 2010
6-14	2-S-Up ₅	0.57 (n.d. ^a / ↓10x ^b)	5.27 (↓59x) ^b	7.33 (↓24x) ^c	n.d.		Ko et al., 2008
6-15	2-S-Up ₄	0.60 (n.d. ^a / ↓10x ^b)	5.52 (↓61x) ^b	6.83 (↓23x) ^c	n.d.		Ko et al., 2008
6-16	4-S-Up ₄	0.070 (n.d. ^a / 1x ^b)	0.28 (↓3x) ^b	6.46 (↓22x) ^c	n.d.		Ko et al., 2008
6-17	2-S-Up ₂ -OMe	n.d.	n.d.	inactive at 10 μM (↓19x) ^c	0.056 (↑3x ^c / ↑7x ^d)	4-S-analogue of 6-17 inactive at P2Y ₁₄ R	Das et al., 2010
6-18	2-S-Up ₂ -OEt (MRS2906)	n.d.	n.d.	inactive at 10 μM (↓19x) ^c	0.039 (↑4x ^c / ↑10x ^d)		Das et al., 2010
6-19	2-S-Up ₂ -OPr (MRS2907)	n.d.	n.d.	9.1 (↓17x) ^c	0.040 (↑4x ^c / ↑10x ^d)		Das et al., 2010; Fricks et al., 2009
6-20	2-S-Up ₂ -OC(CH ₃) ₃	n.d.	n.d.	2.04 (↓4x) ^c	0.032 (↑5x ^c / ↑13x ^d)		Das et al., 2010
6-21	2-S-Up ₂ - O(CH ₂) ₂ C≡CH	n.d.	n.d.	inactive at 10 μM (↓19x) ^c	0.011 (↑15x ^c / ↑36x ^d)		Das et al., 2010





^arelative to ATP; ^brelative to UTP; ^crelative to UDP; ^drelative to UDP-glucose.

2.2. P2Y₂ receptor antagonists



2.2.1. Therapeutic potential of P2Y₂ receptor antagonists

P2Y₂R antagonists could function, among other possible indications, as anti-metastatic cancer therapeutics. This hypothesis is based in part on the observation that P2Y₂Rs expressed on the endothelium open the endothelial barrier in response to ATP. The nucleotide is released from platelets following induction by tumor cells, thereby facilitating tumor cell extravasation from the blood stream into surrounding tissue to form secondary tumor sites (Schumacher, Strilic, Sivaraj, Wettschureck, & Offermanns, 2013). In addition, the proliferation and migration of different tumor and non-tumor cells, as well as the induction of cell cycle progression in vascular smooth muscle cells was attributed to P2Y₂R activation (Greig, Linge, Cambrey, & Burnstock, 2003; Malam-Souley et al., 1996; Martínez-Ramírez, Garay, García-Carrancá, & Vázquez-Cuevas, 2016; Miyagi et al., 1996; Muscella, Elia, Greco, Storelli, & Marsigliante, 2003; Qiu et al., 2018; Robles-Martinez et al., 2017; Schafer, Sedehizade, Welte, & Reiser, 2003; Shen et al., 2004; Tu et al., 2000; Wilden, Agazie, Kaufman, & Halenda, 1998; Zhang et al., 2017). Another therapeutic application for P2Y₂R antagonists may be psoriasis, as P2Y₂R activation was shown to induce keratinocyte proliferation and neutrophil migration (Chen et al., 2006; Dixon et al., 1999). P2Y₂R antagonists may in fact be useful for a range of excessive inflammatory reactions, including atherosclerosis. This is because the P2Y₂R was found to activate cytosolic phospholipase A₂, which in turn promotes the release of arachidonic acid and the subsequent synthesis of prostaglandins and leukotrienes (Seye et al., 2002; Welch, Carlson, Shi, Myatt, &


Kishore, 2003; Xing, Post, Ostrom, Samardzija, & Insel, 1999; Xu et al., 2002; Xu et al., 2003). The potential of P2Y₂R blockade in the treatment of atherosclerosis is further emphasized by the following observations: the receptor appears to mediate the uptake of low-density lipoprotein in vascular smooth muscle cells and contribute to vascular inflammation by increasing vascular cell adhesion molecule-1 expression. Moreover, plaques were more stable in mice with an endothelial cell-specific deficiency of P2Y₂Rs (Chen et al., 2017; Dissmore et al., 2016). P2Y₂R function was also linked to the development of neointimal hyperplasia and pulmonary fibrosis (Agca, Qian, Agca, & Seye, 2016). In addition, P2Y₂R activation in human esophageal epithelial cells led to the release of the pro-inflammatory chemokine interleukin-8, which is believed to contribute to mucosal inflammation and gastroesophageal reflux disease (Wu, Oshima, Fukui, Watari, & Miwa, 2016). Furthermore, the P2Y₂R is expressed in collecting ducts of the kidney, where it opposes the actions of antidiuretic hormone (arginine vasopressin) and, consequently, reduces water reuptake into the blood (Kishore et al., 2015). Thus, patients suffering from nephrogenic diabetes insipidus acquired, for example, through chronic use of lithium in a bipolar disorder therapy, may benefit from P2Y₂R antagonists; a concept that has been patented (Kishore, Carlson, & Zhang, 2012). P2Y₂R antagonists could further prove useful in the treatment of osteoporosis and pain. They could prevent the P2Y₂R-mediated inhibition of bone formation and its involvement in thermal nociception (Hoebertz, Mahendran, Burnstock, & Arnett, 2002; Malin et al., 2008; Orriss et al., 2017; Shi et al., 2017). Refer to Table 1 for an overview of the possible indications for P2Y₂R modulation.

A


- 7-01 Ap₄A R = R¹, n = 3
 7-02 Ap_nA R = R¹, n = 1, 2, 4, or 5
 7-03 Up₄U Diquafosol R = R², n = 3
 7-04 Up₃U R = R², n = 2

7-05 Ap₄U


- 7-06 Up₄C X = OH
 7-07 Denufosol X = H




7-08 MRS2957

B


7-09



7-10



7-11 INS48823



7-12

Fig. 7. Structures of dinucleotide agonists at uracil nucleotide-activated P2Y₂Rs (for data, refer to Table 7).

2.2.2. AR-C118925

In comparison to the relatively large number of agonists reported for the P2Y₂R, only few antagonists have been described to date. The most potent antagonist reported in the literature is the thiouracil derivative 5-[[5-(2,8-dimethyl-5H-dibenzo[*a,d*]cyclohepten-5-yl)-3,4-dihydro-2-oxo-4-thioxo-1(2H)-pyrimidinyl]methyl]-N-(1H-tetrazol-5-yl)-2-furancarboxamide (AR-C118925, **10-01**, compound 01 in Fig. 10 and Table 10). For AR-C118925, IC₅₀ values in the mid-nanomolar to low

micromolar range were reported (Kemp, Sugar, & Jackson, 2004; Rafehi, Burbiel, Attah, Abdelrahman, & Müller, 2017). The selectivity of this compound for the P2Y₂R over other P2Y₂R subtypes is approx. 50- to 500-fold. It is particularly high versus the P2Y₄R (500-fold), which is the P2Y₂R subtype that is structurally and pharmacologically most closely related to the P2Y₂R (Rafehi, Burbiel, et al., 2017). Being a structural analogue of the endogenous agonist UTP (refer to Kindon et al., 2017 for an account on the systematic development of AR-C118925

Table 7

Physiological and synthetic dinucleotide polyphosphate compounds and analogues as agonists for the uracil-activated P2YRs. Shown are the EC₅₀ or, in some cases, K_i and K_B values in μ M, unless stated otherwise. The correlation factor by which the potency is n-fold higher (\uparrow **nx**) or lower (\downarrow **nx**) compared to that of the endogenous agonist in the same test system is given in brackets. Particularly interesting values are shown in bold.

Dinucleotides							
No.	Name	EC ₅₀ in μM				Comments	References
		(potency higher (↑) or lower (↓) relative to endogenous agonist)					
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
7-01	Ap ₄ A	0.054 - 0.720 (↓3-4x ^a / 1x-↓13x ^b)	inactive at 100 μM (↓1800-14300x) ^b low activity at 100 μM (n.d.) ^b	low activity at 100 μM (n.d.) ^c	inactive at 1 μM (n.d. ^c / ↓10x ^d)	>140-fold P2Y ₂ R selective Boranophosphate analogue patented	Bogdanov et al., 1998; Chambers et al., 2000; Communi et al., 1995; US020060287271A1, 2006; Janssens et al., 1999; Kennedy et al., 2000; Lazarowski et al., 1995; Patel et al., 2001; Sakuma et al., 2017; Shaver et al., 2005; Webb et al., 1998; Wildman et al., 2003
7-02	Ap _n A, where n = 2, 3, 5, or 6	12.6 - 39.8 (n.d. ^a / ↓160-500x ^b) 30 % efficacy of Ap ₃ A	>1000 (n.d.) ^b low activity at 100 μM (n.d.) ^b	Only Ap ₃ A and Ap ₅ A active (n.d.) ^c	inactive at 1 μM (n.d. ^c / ↓10x ^d)	Boranophosphate analogues patented	Chambers et al., 2000; Communi et al., 1995; US020060287271A1, 2006; Janssens et al., 1999; Patel et al., 2001; Shaver et al., 2005; Wildman et al., 2003
7-03	Up ₄ U (Diquafosol, INS365)	0.06 - 0.210 (↓4x ^a / ↓3-4x ^b)	0.130 - 0.20 (↓2-4x) ^b	1.16 - 24.8 (↓50-200x) ^c	inactive (n.d.) ^{c,d}	Diquafosol tetrasodium (Diquas®) approved in Japan, Korea, Vietnam, and Thailand as 3 % ophthalmic solution for dry eye disease Boranophosphate analogues patented	US020060287271A1, 2006; Ivanov, Fricks et al., 2007; Keating, 2015; Ko et al., 2007; Lau et al., 2014; Maruoka et al., 2011; Pendergast et al., 2001; Shaver et al., 2005

Table 7 (continued)

7-04	Up ₃ U	1.31 - 22 (n.d. ^a / ↓22-730x ^b)	0.87 (1x) ^b >100 (↓>1000x) ^b	0.2 - 0.92 (↓2x) ^c	n.d.	Boranophosphate analogues patented	WO002007002945A2, 2006; US020060287271A1, 2006; Maruoka et al., 2010; Pendergast et al., 2001; Shaver et al., 2005
7-05	Ap ₄ U	0.35 (↓23x ^a / ↓23x ^b)	1.45 (↓21x) ^b	>100 (↓>200x) ^c	n.d.		
7-06	Up ₄ C	0.45 (n.d.) ^{a,b} 0.46 (↓3x) ^{a,b}	0.65 (n.d.) ^b 0.85 (↓12x) ^b	3.5 (n.d.) ^c 12.1 (↓24x) ^c	n.d.	Patented	Shaver et al., 2005; Yerxa et al., 2002; WO001999061012A2, 2000
7-07	Up ₄ dC (Denufosol, INS37217)	0.22 (n.d.) ^{a,b} 0.27 (↓18x ^a / ↓18x ^b)	0.8 (n.d.) ^b 1.22 (↓17x) ^b	≈70 % at 100 μM 16.0 (↓32x) ^c	n.d.		1999
7-08	N ⁴ -MeO-Cp ₃ U (MRS2957)	0.17 (n.d. ^a / ↓3x ^b)	0.79 (↓9x) ^b	0.012 (↑25x) ^c	K _i = 5.18 (↓8x ^c / ↓2x ^d)	Commercially available	Kiselev et al., 2015; Maruoka et al., 2010
7-09	4-S-Up ₄ U	0.04 (↓3x) ^{a,b}	0.17 (↓2x) ^b	190 (↓380x) ^c	n.d.	Patented	WO001998034942A2, 1998; Shaver et al., 2005
7-10	P ¹ ,P ⁴ -di(3,N ⁴ - ethenocytidine 5'-)tetrphosphate	0.46 (n.d.) ^{a,b}	19.8 (n.d.) ^b	inactive (n.d.) ^c	n.d.		
7-11	Monobenzylnacetal- Up ₃ U (INS48823)	inactive (n.d.) ^{a,b}	inactive (n.d.) ^b	≈0.125 (1x) ^c 0.140 (↑4x) ^c	n.d.	Patented	WO002007002945A2, 2006; Ginsburg- Shmuel et al., 2012; Jayasekara et al., 2013; Korcok et al., 2005; Maruoka et al., 2010
7-12	α,β-Methylene- Up ₃ U	>10 (n.d. ^a / ↓>170x ^b)	inactive at 10 μM (↓110x) ^b	1.30 (↓4x) ^c	n.d.	Patented	WO002007002945A2, 2006; Maruoka et al., 2010

^arelative to ATP; ^brelative to UTP; ^crelative to UDP; ^drelative to UDP-glucose.

from UTP), AR-C118925 acts as a competitive antagonist (Rafehi, Burbiel, et al., 2017). In the absence of an X-ray structure, a homology model supported by mutagenesis data was recently developed that shows a likely binding pose and interactions of AR-C118925 with residues in the putative orthosteric site (Rafehi et al., 2017). This is so far the only reported P2Y₂R homology model that was developed based on the recently published P2Y₁R and P2Y₁₂R X-ray structures (Zhang et al., 2014; Zhang et al., 2014; Zhang et al., 2015), while previous models were based on the distantly related bovine rhodopsin.

In addition to the relatively high degree of selectivity for the P2Y₂R, AR-C118925 (10-01, Fig. 10 and Table 10) has a number of desirable physicochemical and pharmacokinetic properties, including high solubility in phosphate-buffered saline and a very high metabolic stability in human and mouse liver microsomes. It has reasonable rat *in vivo* pharmacokinetic properties, with intravenous clearance of 75 mL/min/kg, V_{ss} of 4.34 L/kg and a half-life of 2.12 h. AR-C118925 (10-01) was shown to inhibit CYP2C8 but not eight other cytochrome P450 subtypes (including CYP3A4) at 1 μM. However, its bioavailability following oral administration is very low (Conroy, Kindon, Kellam, & Stocks, 2016; Kindon et al., 2017; Rafehi, Burbiel, et al., 2017). The compound, developed by AstraZeneca, was assessed in pre-clinical trials as a topical skin treatment for chronic psoriasis, but was found to be inefficacious (Conroy et al., 2016; Kindon, Meghani, & Thom, 1998). One reason for this might be its high polarity, which may prevent it from

penetrating into deeper layers of the skin. AR-C118925 is nevertheless a valuable pharmacological tool to study P2Y₂R function and has been utilized in several *in vivo* and *in vitro* studies (Cosentino et al., 2012; Hochhauser et al., 2013; Kemp et al., 2004; Magni, Merli, Verderio, Abbracchio, & Ceruti, 2015; Önnheim et al., 2014; Wang et al., 2015).

2.2.3. Suramin and Reactive Blue 2

AR-C118925 (10-01, Fig. 10 and Table 10) became available through commercial sources only recently and was therefore inaccessible for most research groups. Thus, the antagonists used most widely to study P2Y₂R function were suramin (10-04, Fig. 10 and Table 10) and the dye Reactive Blue 2 (RB-2, 10-07, Fig. 10 and Table 10). RB-2 is a large (molecular weight of 840 g/mol), polysulfonated, and polyaromatic anthraquinone derivative with an IC₅₀ value in the micromolar range at the P2Y₂R (Baqi et al., 2010; Rafehi, Neumann, et al., 2017; Weyler et al., 2008). However, smaller derivatives that retain the anthraquinone core are similarly potent at the P2Y₂R (Weyler et al., 2008). RB-2 is more potent at the P2Y₄R than at the P2Y₂R; it will consequently be dealt with in more detail in Section 3.2.2.

Suramin (10-04, Fig. 10 and Table 10) was developed by the pharmaceutical company Bayer in 1920 as an anti-parasitic drug and is still in use for the treatment of sleeping sickness caused by trypanosomes (see reviews by Singh Grewal, Pandita, Bhardwaj, & Lather, 2016; Voogd, Vansterkenburg, Wilting, & Janssen, 1993). With a molecular




Table 8

Physiological and synthetic nucleotide-sugar conjugates as agonists for the uracil-activated P2YRs. Shown are the EC₅₀ or, in some cases, K_i and K_B values in μM , unless stated otherwise. The correlation factor by which the potency is n-fold higher ($\uparrow\text{nx}$) or lower ($\downarrow\text{nx}$) compared to that of the endogenous agonist in the same test system is given in brackets. Particularly interesting values are shown in bold.

Nucleotide sugars							
No.	Name	EC ₅₀ in μM (potency higher (↑) or lower (↓) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
8-01	UDP-glucose	10 (n.d.) ^{a,b}	n.d.	16 (↓30) ^c	0.0022 - 0.400 (1x-↓15x) ^c K _i = 0.020 - 2.23 (↑2x-↓4x) ^c	UDP-β- [1'']glucose half as potent at P2Y ₁₄ R UDP-[6'']glucose of similar potency Fluorinated deoxy-derivatives published	Carter et al., 2009; Chambers et al., 2000; Das et al., 2009; Das et al., 2010; Freeman et al., 2001; Fricks et al., 2009; Gao et al., 2010; Hamel et al., 2011; Ivanov, Fricks et al., 2007; US020090148850A1, 2008; Kiselev et al., 2015; Ko et al., 2007; Ko et al., 2009; Lazarowski & Harden, 1994; Scrivens & Dickenson, 2005
				Rat: >1000 (↓>440x) ^c	Rat: 0.026 (n.d.) ^c 0.189 - 5.98 (1x) ^c Mouse: 0.0028 - 0.022 (↑2x) ^c Chimpanzee: 0.0044 - 0.0106 (1x-↓2x) ^c K _i = 0.023 (↓2x) ^c		
8-02	UDP-galactose	n.d.	n.d.	n.d.	0.0028 - 0.671 (1x-↓32x ^c / 1x-↓4x ^d) partial agonist Rat: 0.077 (n.d. ^c / ↓3x ^d) Mouse: 0.0028 - 0.063 (↓4x ^c / 1x-↓6x ^d) Chimpanzee: 0.0056 - 0.0272 (↓2-4x ^c / 1x-↓3x ^d)	Freeman et al., 2001; Hamel et al., 2011; Ivanov, Fricks et al., 2007; US020090148850A1, 2008; Ko et al., 2007; Ko et al., 2009; Scrivens & Dickenson, 2005	

(continued on next page)

Table 8 (continued)

Nucleotide sugars							
No.	Name	EC ₅₀ in μ M (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
8-03	UDP-glucuronic acid	n.d.	n.d.	n.d.	0.006 - 0.576 (\downarrow 2-82x ^c / 1x- \downarrow 6x ^d) partial agonist K _i = 5.3 (\downarrow 8x ^c / \downarrow 2x ^d) Rat: 0.102 (n.d. ^c / \downarrow 4x ^d) Mouse: 0.0096 - 0.0915 (\downarrow 2-5x ^c / \downarrow 3-8x ^d) Chimpanzee: 0.0054 - 0.0762 (1- \downarrow 11x ^c / \downarrow 2-7x ^d)	Derivatives of less potency published as well Dendrimer analogues also published	Chambers et al., 2000; Das et al., 2009; Freeman et al., 2001; Fricks et al., 2009; Hamel et al., 2011; Ivanov, Fricks et al., 2007; US020090148850A1, 2008; Kiselev et al., 2015; Ko et al., 2007; Ko et al., 2009; Scrivens & Dickenson, 2005
8-04	UDP-N-acetylglucosamine	n.d.	n.d.	n.d.	0.0417 - 4.38 (\downarrow 14-180x ^c / \downarrow 2-19x ^d) partial agonist Rat: 0.170 (n.d. ^c / \downarrow 7x ^d) Mouse: 0.0257 - 0.101 (\downarrow 3x ^c / \downarrow 5-9x ^d) Chimpanzee: 0.079 - 0.149 (\downarrow 17-22x ^c / \downarrow 12-16x ^d)		Chambers et al., 2000; Freeman et al., 2001; Fricks et al., 2009; Hamel et al., 2011; Ivanov, Fricks et al., 2007; US020090148850A1, 2008; Ko et al., 2007; Ko et al., 2009; Scrivens & Dickenson, 2005
8-05	UDP-N-acetylglactosamine	n.d.	n.d.	n.d.	0.810 (n.d. ^c / \downarrow 2x ^d)		Ko et al., 2007; Ko et al., 2009
8-06	UDP-inositol	n.d.	n.d.	n.d.	1.88 (n.d. ^c / \downarrow 5x ^d)		Ko et al., 2007; Ko et al., 2009
8-07	UDP-[1"]fructose	n.d.	n.d.	n.d.	0.880 (n.d. ^c / \downarrow 3x ^d)		Ko et al., 2007; Ko et al., 2009
8-08	UDP-[6"]fructose	inactive at 10 μ M (n.d.) ^{a,b}	n.d.	n.d.	0.323 (n.d. ^c / 1x ^d)		Ko et al., 2009

Table 8 (continued)

Nucleotide sugars							
No.	Name	EC ₅₀ in μ M (potency higher (\uparrow) or lower (\downarrow) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
8-09	UDP-fucose	n.d.	n.d.	n.d.	0.562 (n.d. ^c / \downarrow 2x ^d)		Ko et al., 2009
8-10	UDP-[1 ^{'''}]mannose	n.d.	n.d.	n.d.	0.910 (n.d. ^c / \downarrow 3x ^d)		Ko et al., 2007; Ko et al., 2009
8-11	UDP-[6 ^{'''}]mannose	n.d.	n.d.	n.d.	0.658 (n.d. ^c / \downarrow 3x ^d)		Ko et al., 2007; Ko et al., 2009
8-12	UDP-[5 ^{'''}]ribose (MRS2738)	n.d.	n.d.	n.d.	0.238 (n.d. ^c / 1x ^d)		Ko et al., 2009
8-13	UDP-[5 ^{'''}]arabinose	n.d.	n.d.	n.d.	0.460 (n.d. ^c / \downarrow 2x ^d)		Ko et al., 2009
8-14	GDP-glucose	n.d.	n.d.	n.d.	0.141 (\downarrow 59x ^c / \downarrow 64x ^d)		Hamel et al., 2011
8-15	2-S-UDP-[1 ^{'''}]glucose (MRS2690)	inactive at 10 μ M (n.d.) ^{a,b}	n.d.	>10 (\downarrow >19x) ^c	0.011 (\uparrow 15x ^c / \uparrow 36x ^d) 0.049 (n.d. ^c / \uparrow 7x ^d) K_i = 0.34 (\uparrow 2x ^c / \uparrow 7x ^d) <hr/> Rat: 0.0081 - 0.538 (\uparrow 10x ^c / \uparrow 11-23x ^d)	Commercially available	Das et al., 2010; Gao et al., 2010; Kiselev et al., 2015; Ko et al., 2007; Ko et al., 2009
8-16	4-S-UDP-[1 ^{'''}]glucose (MRS2670)	n.d.	n.d.	n.d.	0.29 (n.d. ^c / 1x ^d)		Ko et al., 2007
8-17	N ⁴ -MeO-CTP-glucose	>10 (n.d. ^a / \downarrow >170x ^b)	inactive at 10 μ M (\downarrow >110x) ^b	0.18 (\uparrow 2x) ^c	n.d.		Maruoka et al., 2010
8-18	(S)-methanocarba-UTP-glucose	inactive at 10 μ M (n.d. ^a / \downarrow >170x ^b)	inactive at 10 μ M (\downarrow >110x) ^b	2.47 (\downarrow 8x) ^c	n.d.		Maruoka et al., 2010
8-19	Up ₄ -[1 ^{'''}]glucose (MRS2732)	0.30 (n.d. ^a / \downarrow 5x ^b)	2.06 (\downarrow 23x) ^b	7.83 (\downarrow 26x) ^c	n.d.	6-fold selective for P2Y ₂ R	Ko et al., 2008
8-20	Up ₄ -[1 ^{'''}]3'-deoxy-3'-fluoroglucose (MRS2927)	0.710 (n.d. ^a / \downarrow 13x ^b)	0.062 (1x) ^b	0.950 (n.d.) ^c	n.d.	11-fold selective for P2Y ₄ R	Maruoka et al., 2011
8-21	UDP-glucuronic acid-ethylenediamine (MRS2892)	n.d.	n.d.	n.d.	2.59 (n.d. ^c / \downarrow 10x ^d)	Amino-functionalized P2Y ₁₄ R agonist	Das et al., 2009
8-22	UDP-glucuronic acid-(N-acetyl-ethylenediamine)	n.d.	n.d.	n.d.	0.496 (n.d. ^c / \downarrow 2x ^d)		Das et al., 2009

(continued on next page)

Table 8 (continued)

Nucleotide sugars							
No.	Name	EC ₅₀ in μM (potency higher (↑) or lower (↓) relative to endogenous agonist)				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
8-23	MRS4183	n.d.	n.d.	n.d.	0.00096 (↑170x ^c / ↑270x ^d)	Analogue with shorter linker published (EC ₅₀ = 0.091 μM at P2Y ₁₄ R)	Kiselev et al., 2015
8-24	G6-(UDP-glucuronic acid) ₁₄₇	n.d.	n.d.	n.d.	0.0008 (n.d. ^c / ↑330x ^d)	Dendrimer-conjugate	Das et al., 2009
8-25	G3-(UDP-glucuronic acid) _{20.1} -(biotin) _{4.9}	n.d.	n.d.	n.d.	0.0034 (n.d. ^c / ↑77x ^d)	Dendrimer-conjugate coupled to biotin for avidin complexation	Das et al., 2009
8-26	G3-(UDP-glucuronic acid) _{20.1} -(Alexa Fluor® 488) _{2.3}	n.d.	n.d.	n.d.	0.0398 (n.d. ^c / ↑7x ^d)	Dendrimer-conjugate coupled to fluorophore	Das et al., 2009
8-27	G3-(UDP-glucuronic acid) _{20.1} -(DTPA) _{4.5}	n.d.	n.d.	n.d.	0.0041 (n.d. ^c / ↑64x ^d)	Dendrimer-conjugate coupled to metal-chelating group	Das et al., 2009
8-28	G3-(UDP-glucuronic acid) _{20.1} -(DTPA-Gd(III)) _{4.5}	n.d.	n.d.	n.d.	0.421 (n.d. ^c / ↓2x ^d)	Dendrimer-conjugate coupled to metal-chelating group and complexed with Gadolinium	Das et al., 2009
8-29	MRS5259	1.39 (n.d.) ^{a,b}	n.d.	n.d.	0.00224 (n.d.) ^{c,d}	UDP-glucuronic acid coupled to A ₃ AR agonist (MRS3558) via PAMAM dendrimer Patented	WO002011068978A1, 2010

^arelative to ATP; ^brelative to UTP; ^crelative to UDP; ^drelative to UDP-glucose.

weight of 1297 g/mol, it is a comparatively large, polyaromatic, and polysulfonated naphthalene derivative. Suramin has a symmetrical structure with a urea group at the center. It caused a parallel rightward shift of UTP dose-response curves at the human P2Y₂R. The corresponding Schild plot yielded a slope of 1.57 and an apparent pA₂ value of 4.32

(48 μM) (Charlton et al., 1996a). Suramin can therefore be classified as a very weak competitive P2Y₂R antagonist.

Both RB-2 and suramin are partially selective antagonists but also inhibit ectonucleotidases. In light of insufficient alternatives, both have been in use as pharmacological tools for P2R characterization for




Fig. 9. Structures of non-nucleotide agonists at uracil nucleotide-activated P2YRs (for data, refer to Table 9).

a long time. However, the negative charge on both RB-2 and suramin that prevents the penetration of cell membranes as well as the low degree of potency and, in particular, of selectivity are serious limitations to their usefulness.

2.2.4. Flavonoids

Flavonoids are secondary plant metabolites found in fruits, vegetables, bark, and flowers of many higher plants, including those that are used in traditional herbal medicine. Over 5000 different flavonoids with diverse pharmacological properties are known. A series of 40 flavonoids was assessed for P2Y₂R antagonism. Tangeretin (**10-02**, Fig. 10 and Table 10), isolated from a tangerine fruit peel extract (*Citrus reticulata* ssp., Rutaceae), was found to be the most potent of these, with an IC₅₀ value of 12 μM at the mouse P2Y₂R (Kaulich, Streicher, Mayer, Müller, & Müller, 2003). The structure-activity relationships (SARs) were found to be rather complex, but flavone derivatives were generally more potent than flavanones, indicating that a flat aromatic ring system was preferred. The maximal receptor activation of agonist (UTP) concentration-response curves was reduced with increasing concentrations of tangeretin, while the EC₅₀ value of UTP remained unaffected. These compounds, therefore, appeared to act as non-competitive, allosteric antagonists (Kaulich et al., 2003).

2.2.5. Acyclic nucleotide derivatives

Acyclic derivatives of uracil nucleotides, in which the ribose ring is replaced by aliphatic residues, were published as P2Y₂R antagonists (Sauer, El-Tayeb, Kaulich, & Müller, 2009). The phosphate groups were replaced by phosphonates to enhance chemical and metabolic stability. The most potent P2Y₂R antagonist was diphosphoric 5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)pentylphosphonic anhydride (**10-05**, Fig. 10 and Table 10), which had an IC₅₀ of 92 μM at the mouse P2Y₂R (Sauer et al., 2009). Thus, their suitability as pharmacological tools appears to be limited at present due to their low potency, but further optimization may improve their properties.

3. The P2Y₄ receptor

The existence of specific pyrimidineric receptors activated by uracil nucleotides but not adenine nucleotides was postulated several years prior to the actual cloning of the first uracil nucleotide-selective receptor, the P2Y₄R, in 1995 (Communi, Pirotton, Parmentier, & Boeynaems, 1995; Nguyen et al., 1995; Seifert & Schultz, 1989). The P2Y₄R was discovered with the aid of degenerate oligonucleotide primers. These were designed and synthesized based on the most conserved segments of the

Table 9

Non-nucleotide agonists for the uracil-activated P2YRs. Shown are the EC₅₀ values in μM, unless stated otherwise. The correlation factor by which the potency is n-fold higher (↑nx) or lower (↓nx) compared to that of the endogenous agonist in the same test system is given in brackets. Particularly interesting values are shown in bold.

Non-nucleotide agonists							
No.	Name	EC ₅₀ in μM				Comments	References
		(potency higher (↑) or lower (↓) relative to endogenous agonist)					
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
9-01	Compound 89	10.5 (↓≈500x ^a / n.d. ^b)	inactive at 30 μM	inactive at 30 μM	n.d.	Ago-allosteric agonist with 66 % (human P2Y ₂ R) and 51 % (mouse P2Y ₂ R) efficacy calculated by extrapolation	Sakuma et al., 2017
		53.9 (n.d.) ^{a,b}	(↓>600x) ^b	(↓>600x) ^c			
		<hr/> Mouse: 2.7 (↓≈15x ^a / n.d. ^b)					
9-02	Prostaglandin E ₂ glyceryl ester	n.d.	n.d.	0.2 - 1.2 pM	n.d.		Brüser et al., 2017
				(↑6000-65,000x) ^c			
				<hr/> Mouse: 0.8 pM (↑48,000x) ^c			

^arelative to ATP; ^brelative to UTP; ^crelative to UDP.

chick P2Y₁R and the mouse P2Y₂R sequences (Communi et al., 1995). Polymerase chain reactions on human genomic DNA using these primers led to the localization of a 712-base pair sequence, which was subsequently used as a probe to screen a human genomic DNA library. The P2Y₄R was eventually characterized as an intronless 1095-base pair open reading frame sequence (Communi et al., 1995). The receptor is both structurally and pharmacologically closely related to the P2Y₂R. Accordingly, it also signals primarily via $G\alpha_{q/11}$. The residues His262,

Arg265, Lys289, and Arg292, that are believed to be involved in nucleotide binding, are conserved between these two receptors (Communi et al., 1995). In fact, the charged amino acids Arg265 and Arg292 were proposed to be directly responsible for the binding of the negatively charged phosphate groups of nucleotides. Mutating Lys289 to arginine in the human P2Y₄R resulted in a shift in agonist preferences from UTP to UDP (Communi et al., 1995). A P2Y₄R homology model was developed based on the human P2Y₁R X-ray structure, and molecular




Fig. 10. Structures of antagonists at uracil nucleotide-activated P2YRs (for data, refer to Table 10).




Fig. 10 (continued).

docking studies were performed (Rafehi et al., 2017; Zhang et al., 2015). According to these, Arg265 interacts with the uracil base, Arg292 with the α - and β -phosphate groups, and Lys289 with the γ -phosphate group. His262 is in close proximity (approx. 3 Å) to the uracil base and interaction is likely (Rafehi, Malik, et al., 2017). No *N*-glycosylation consensus sequence is found in the putative exofacial region of the human P2Y₄R, which is rare but not unique in GPCRs (Communi et al., 1995). In contrast, the rat ortholog contains an Asn in position 179 and a Thr in position 181 that creates a consensus *N*-linked glycosylation site (Herold, Qi, Harden, & Nicholas, 2004).

P2Y₄R expression appears to be more restricted than of the P2Y₂R; it is mostly found (at mRNA level) in the GI tract and to a lower extent in the CNS, lungs, heart, prostate, skin, adipose tissue, skeletal muscle, spleen and lymphocytes (Table 1) (Moore et al., 2001; Uhlén et al., 2015). A possible physiological function of the P2Y₄R is to mediate Cl⁻ secretion in the gastrointestinal tract (DuBose, Wolff, Qi, Naruszewicz, & Nicholas, 2013; Ghanem et al., 2005; Matos, Robaye, Boeynaems, Beauwens, & Leipziger, 2005). Another potentially important role is in

the cochlea, where it regulates Na⁺ homeostasis in the endolymph. It does this by inhibiting epithelial Na⁺ channels of Reissner's membrane, and by K⁺ secretion across strial marginal cell epithelial cells in the apical membrane (Kim et al., 2010; Marcus et al., 2005). Furthermore, the P2Y₄R appears to be important for postnatal cardiac development, since P2Y₄R-null mice showed microcardia (Horckmans et al., 2012).

Relatively few compounds that target the P2Y₄R have been developed in comparison to the other pyrimidineric P2YR subtypes. An overview of the ligands and pharmacological tools currently available for the P2Y₄R is provided in Tables 2 to 10. Those of greatest potential are further discussed below.

3.1. P2Y₄ receptor agonists

3.1.1. Therapeutic potential of P2Y₄ receptor agonists

P2Y₄Rs were shown to be involved in Cl⁻ secretion in the intestinal epithelium (Ghanem et al., 2005). The Cl⁻ transport across the epithelium of both the jejunum and the colon was found to be abolished in

Table 10Antagonists for the uracil-activated P2YRs. Shown are the IC_{50} or, in some cases, pA_2 , K_i and K_B values in μM . Particularly interesting values are shown in bold

Antagonists							
No.	Name	IC_{50} in μM				Comments	References
		P2Y ₂	P2Y ₄	P2Y ₆	P2Y ₁₄		
10-01	AR-C118925	0.0574 - 1	37.1	30.4	>3	Patented Smaller derivative AR-C126313 has pA_2 of 6.9	Kemp et al., 2004; Kindon et al., 2017; WO00199902501A1, 1998; Rafehi, Burbiel et al., 2017
10-02	Tangeretin	n.d. Mouse: 12	n.d.	n.d.	n.d.	Non-competitive antagonist	Kaulich et al., 2003
10-03	PSB-716	9.82 Mouse: 9	n.d.	n.d.	n.d.	Derivative of RB-2	Hillmann et al., 2009; Rafehi, Malik et al., 2017; Weyler et al., 2008
10-04	Suramin	48 $pA_2 = 4.32$ Rat: 8.9 competitive antagonist with $pA_2 = 5.40$ 77 % inhibition at 100 μM Mouse: 31	inactive at 300 μM Rat: 1027 10 % inhibition at 100 μM Mouse: inactive at 100 μM	27 % inhibition at 100 μM Rat: competitive antagonist	n.d.		Bogdanov et al., 1998; Charlton et al., 1996b; Chen et al., 1996; Communi, Parmentier et al., 1996; Kaulich et al., 2003; Lazarowski & Harden, 1994; Müller, 2002; Robaye et al., 1997; Suarez-Huerta et al., 2001; Wildman et al., 2003
10-05	Diphosphoric 5-(2,4-dioxo-3,4-dihydro pyrimidin-1(2H)-yl) pentylphosphonic anhydride	n.d. Mouse: 92	n.d.	n.d.	n.d.		Sauer et al., 2009
10-06	PSB-16133	8.54 allosteric antagonist	0.233	12.5	n.d.	37-fold selective for P2Y₄R Derivative of RB-2	Rafehi, Malik et al., 2017

Table 10 (continued)

10-07	Reactive Blue 2 (RB-2)	1.85 22.1	0.625 (purified) 1.14 9.79 33 % inhibition at 100 μ M	31 (87 % inhibition at 100 μ M)	n.d.	Lead structure for development of 10-03 & 10-06	Baqi et al., 2010; Bogdanov et al., 1998; Communi, Motte et al., 1996; Kaulich et al., 2003; Kim et al., 2010; Lazarowski & Harden, 1994; Rafehi, Malik et al., 2017; Rafehi, Neumann et al., 2017;
		Rat: 10,000	Rat: 18.5 21.1 competitive antagonist with pA_2 = 6.43	Rat: 4.34 non-competitive antagonist			Robaye et al., 1997; Suarez-Huerta et al., 2001; Weyler et al., 2008; Wildman et al., 2003
		Mouse: 5.0 5.5	Mouse: 47 competitive antagonist				
			Gerbil: 4.9				
2-01	ATP	agonist (see Table 2)	competitive antagonist with pA_2 =	almost inactive as agonist (see Table 2)	agonist (see Table 2)		Communi, Motte et al., 1996; Kennedy et al., 2000; Nguyen et al., 1995
			6.15 Rat: agonist (see Table 2)				
4-17	3'-O-(4-benzoyl)-benzoyl-ATP (BzATP)	agonist (see Table 4)	159	n.d.	n.d.	ATP analogue used as photoaffinity label	Erb et al., 1993; Lin et al., 1993; Wildman et al., 2003
10-08	MRS2578	inactive at 10 μ M	inactive at 10 μ M	0.037 Rat: 0.098	no binding at 10 μ M	Non-competitive, insurmountable antagonist	Kiselev et al., 2015; Mamedova et al., 2004
10-09	MRS2567	inactive at 10 μ M	inactive at 10 μ M	0.126 Rat: 0.101	n.d.	10-08 is commercially	

(continued on next page)

Table 10 (continued)

10-10	MRS2575	inactive at 10 μ M	inactive at 10 μ M	0.155	n.d.	available	
				Rat: inactive			
10-11	TIM-38	inactive at 50 μ M	inactive at 50 μ M	4.3	n.d.		Ito et al., 2017
10-12	PPADS	inactive	15 (73 % inhibition at 100 μ M) >100 Rat: >10,000 Dog: inactive	69 % inhibition at 100 μ M Rat: 25 - >1000 non-competitive antagonist Mouse: 45 non-competitive antagonist	n.d.		Bogdanov et al., 1998; Brown, Tanna, & Boarder, 1995; Charlton et al., 1996b, 1996a; Communi, Motte et al., 1996; Lambrecht, 1996; Müller, 2002; Robaye et al., 1997; Suarez-Huerta et al., 2001; Vigne et al., 1996; Wildman et al., 2003; Zambon et al., 2000
10-13	Uridylol phosphosulfate	inactive	inactive	112	n.d.		Meltzer et al., 2015
10-14	PPTN	inactive at 1 μ M	inactive at 1 μ M	inactive at 1 μ M	0.00043 0.006 $K_i = 0.0003$ $K_i = 0.0019$ $K_B = 0.000434$ Chimpanzee: 0.0022 $K_i = 0.0019$	Competitive antagonist >2300-fold selective for P2Y₁₄R Low oral bioavailability due to zwitterionic structure but ester-prodrug published Patented	Barrett et al., 2013; WO2009070873A1, 2008; Junker et al., 2016; Kiselev et al., 2014; Kiselev et al., 2015; Robichaud et al., 2011
10-	Ester-prodrug of PPTN	n.d.	n.d.	n.d.	n.d.		Robichaud et al., 2011
15							

Table 10 (continued)

10-16	4,7-Disubstituted naphthoic acid derivative	n.d.	n.d.	n.d.	n.d.	Competitive antagonist 67 % oral bioavailability in mice, low intrinsic clearance (1.6 ml/min/kg) but > 99% bound to human serum albumin Patented	WO2009070873A1, 2008; Gauthier et al., 2011; Robichaud et al., 2011
					Mouse: 0.008		
					Chimpanzee: 0.001 $K_i = 0.004$		
10-17	4,7-Disubstituted naphthoic acid derivative	n.d.	n.d.	n.d.	n.d.	Reversible, competitive antagonist	Gauthier et al., 2011
					Mouse: 3.5		
					Chimpanzee: 3.5 $K_i = 0.16$		
10-18	MRS4174	n.d.	n.d.	n.d.	$K_i = 0.000080$	Analogue of PPTN coupled to a fluorophore Precursors published	Kiselev et al., 2014
10-19	Triazole analogue of PPTN	inactive at 10 μ M	inactive at 10 μ M	inactive at 10 μ M	0.032		Junker et al., 2016
10-20	Alkyne analogue of PPTN	inactive at 10 μ M	inactive at 10 μ M	n.d.	5.69		Junker et al., 2016
10-21	3,4-Methylenedioxyphenyl derivative with dihydropyridopyrimidine core	n.d.	n.d.	n.d.	n.d.	Non-competitive antagonist SARs and	Guay et al., 2011; WO002009000087A1, 2008
					Mouse: 0.010		
					Chimpanzee: 0.081 0.0885		

(continued on next page)

Table 10 (continued)

10-22	Compound A (phosphonate derivative)	n.d.	n.d.	n.d.	2.3	Hamel et al., 2011; US020090148850A1, 2008
					8.58	
					$K_i = 1.20$	
5-08	TMPS	inactive	inactive	partial agonist (see Table 5)	Chimpanzee: $K_i = 1.52$	AMPS, UMPS, and CMPS are P2Y ₁₄ R agonists
					antagonist	
						Gendaszewska-Darmach et al., 2016; Gendaszewska-Darmach & Szustak, 2016

P2Y₄R-null mice, which showed an otherwise normal behavior, growth, and reproduction (Ghanem et al., 2005; Robaye et al., 2003). It was proposed that the P2Y₄R is responsible for Cl⁻ secretion in response to UTP in both the small and large intestines, and, together with the P2Y₂R, at the basolateral side of the jejunum (Ghanem et al., 2005). Cystic fibrosis is a disorder that results from a mutation in the Cl⁻ ion channel CFTR. Thus, it was hypothesized that stimulation of the P2Y₄R could compensate for the defective CFTR in the intestine (in analogy to P2Y₂R activation in the airway epithelium). This is supported by observations in mice homozygous for the $\Delta F508$ mutation of CFTR, the most common cause for cystic fibrosis. The jejunum of these mice was responsive to UTP, although the magnitude of the response was smaller than in wild-type mice (Ghanem et al., 2005). P2Y₄R agonists are therefore considered to be potential drugs for patients suffering from cystic fibrosis. P2Y₄Rs were also found to be involved in the growth and migration of cardiac endothelial cells, and in the secretion of platelet-derived growth factor B (Horckmans et al., 2012). P2Y₄R agonists may modulate angiogenesis, cardiac remodeling, and post-ischemic revascularization. Moreover, activation of the P2Y₄R in rat microglia was observed to promote pinocytosis of soluble β -amyloid, the major constituent of amyloid plaques observed in the brains of patients suffering from Alzheimer's disease (Li et al., 2013). Consequently, P2Y₄R agonists may serve as therapies for this debilitating and as yet incurable disorder.

On the other hand, P2Y₄R antagonism has also been suggested to be potentially beneficial in Alzheimer's disease treatment (see Section 3.2.1): The P2Y₄R contributes to the release of amyloid precursor protein, which can be converted to β -amyloid that forms the characteristic plaques (Tran, 2011). This indicates a complex involvement of the P2Y₄R and demonstrates the necessity of suitable, brain-permeable pharmacological tools for target validation studies.

3.1.2. Physiological ligands of the P2Y₄ receptor

The endogenous agonist of the P2Y₄R is UTP (2-08, compound 08 of Fig. 2 and Table 2) (Nicholas et al., 1996). Other nucleotide triphosphates, such as guanosine 5'-triphosphate (GTP, 2-03, Fig. 2 and Table 2) and inosine 5'-triphosphate (ITP, 2-05, Fig. 2 and Table 2), are less potent than UTP at the human P2Y₄R receptor (Communi, Motte, Boeynaems, & Pirotton, 1996; Kennedy, Qi, Herold, Harden, & Nicholas, 2000; Nguyen et al., 1995). ATP (2-01, Fig. 2 and Table 2) was observed to competitively antagonize the human P2Y₄R with moderate potency (Communi, Motte, et al., 1996; Kennedy et al., 2000). ADP (2-02, Fig. 2 and Table 2) is inactive at the human P2Y₄R. UDP (2-09, Fig. 2 and Table 2) exhibited EC₅₀ values in the mid-micromolar range

but was found inactive when UTP contamination and enzymatic conversion to UTP was excluded (Communi et al., 1995; Communi, Motte, et al., 1996; Nguyen et al., 1995; Nicholas et al., 1996). Up₄U (7-03, Fig. 7 and Table 7) is also a physiological P2Y₄R agonist with an EC₅₀ value in the high nanomolar range (Maruoka et al., 2011; Pendergast et al., 2001; Shaver et al., 2005). Diuridine polyphosphates with different numbers of phosphate groups are significantly less potent. Ap₄A (7-01, Fig. 7 and Table 7) is inactive (Kennedy et al., 2000; Patel et al., 2001; Shaver et al., 2005). The agonist profile of the rat P2Y₄R is strikingly different from that of the human ortholog and more closely resembles that of the P2Y₂R (Bogdanov, Wildman, Clements, King, & Burnstock, 1998). It is thus likely that some of the effects attributed to the functionally defined "P_{2U} receptor" in native rodent-derived cells were not (solely) mediated by the P2Y₂R but possibly by the rodent P2Y₄R as well. At the rat P2Y₄R, ATP acts as an agonist with similar potency as UTP, Ap₄A, and ITP (see Tables 2 and 7) (Bogdanov et al., 1998; Kennedy et al., 2000). UDP, ADP, ATP γ S (5-03, Fig. 5 and Table 5), and 2-methylthio-ATP (2-MeSATP, 3-37, Fig. 3 and Table 3) were described as partial agonists at the rat receptor (Bogdanov et al., 1998). The sequence identity between the human and the rat orthologs is 83% in total, and 90% in the transmembrane regions and extracellular loops (Bogdanov et al., 1998; Kennedy et al., 2000). Human/rat P2Y₄R chimera studies showed that whether ATP behaves as an agonist or an antagonist on the P2Y₄R was mainly dependent on the second extracellular loop, and to a lesser extent on the N-terminus (Herold et al., 2004). Mutating three residues within the second extracellular loop of the human receptor to the corresponding residues of the rat ortholog (S177N, V183I, and R190L) imparted ATP sensitivity onto the human receptor (Herold et al., 2004).

3.1.3. Synthetic nucleotide derivatives

The P2Y₄R is often expressed in the same tissues as the more abundant P2Y₂R and, as for example in epithelial cells of the intestine, sometimes even in the same cells. As a result, it is challenging to distinguish pharmacologically between these two structurally related receptor subtypes, especially given the similarities in their pharmacological profiles. While extensive efforts have led to the development of several selective agonists for the P2Y₂R, the availability of P2Y₄R-selective agonists has been more limited. The currently most selective compounds described in the literature are the CTP derivatives N⁴-(phenylpropoxy)-CTP (MRS4062, 3-15, Fig. 3 and Table 3) and N⁴-(phenylethoxy)-CTP (3-16, Fig. 3 and Table 3) (Maruoka et al., 2011). The commercially available MRS4062 was twice as potent as UTP and displayed 27- and

32-fold selectivity for the P2Y₄R over the P2Y₂R and P2Y₆R, respectively. Compound **3-16** displayed a potency in the mid-nanomolar range, similar to UTP, and was 16-fold selective for the P2Y₄R versus the other two subtypes (Maruoka et al., 2011). These two P2Y₄R agonists are part of a series of synthetic compounds that were initially designed with the intention to develop improved P2Y₂R agonists through combining different substitutions of the pyrimidine moiety and the phosphate groups (Maruoka et al., 2011).

Among the first reported P2Y₄R agonists with good selectivity (>20-fold versus the P2Y₂R and P2Y₆R) was an analogue of uridine 5'-monophosphate (UMP) with the 2-keto group replaced by an amino-function. 2-Amino-UMP (iso-CMP, **3-07**, Fig. 3 and Table 3), showed micromolar potency (EC₅₀ of 4.98 μ M) (El-Tayeb et al., 2011). It is worth mentioning that the 2-amino substitution conferred agonist activity at the P2Y₄R to the otherwise inactive UMP. Certain substitutions in the uracil moiety of UTP, including 2-thio- (**3-01**, Fig. 3 and Table 3), 3-methyl- (**3-09**), 5-bromo- (**3-23**), 5-iodo- (**3-25**), and 5-methyl- (**3-29**), were tolerated but caused a reduction in potency (Communi, Parmentier, & Boeynaems, 1996; El-Tayeb et al., 2006; Jacobson et al., 2006; Nguyen et al., 1995; Nicholas et al., 1996). Several compounds with substitutions in 6-position were found to be completely inactive (Ginsburg-Shmuel et al., 2010). In contrast, thio-substitution in the 4-position did not cause a decrease in potency but appeared to be favorable instead. An EC₅₀ value of 23 nM was reported for 4-thio-UTP (**3-12**, Fig. 3 and Table 3), which was thus 3-fold more potent than UTP at the P2Y₄R (but unselective versus the P2Y₂R) (Jacobson et al., 2006). As was observed with the P2Y₂R, even larger substituents in the 4-position were well-tolerated. The fluorescent probe MRS4162 (**3-40**, Fig. 3 and Table 3), discussed previously in Section 2.1.5., and its analogues with varying substitutions in the 4-position all showed potencies in the nanomolar range at the P2Y₄R (Jayasekara et al., 2014). The fluorescent probe MRS4162 and its precursor MRS4063 (**3-41**, Fig. 3 and Table 3) that can be extended, e.g. by click tethering, are versatile pharmacological tools. The fact that they did not only show activity at the P2Y₂R but also at the P2Y₄R expands the possibilities for further characterization of this receptor. A UTP analogue with a 2'-azido-2'-deoxy substitution at the ribose ring (**4-06**, Fig. 4 and Table 4; EC₅₀ of 1.1 μ M) was selective for the P2Y₄R, albeit only 5-fold versus the P2Y₂R (Jacobson et al., 2006). Not only substitutions at the ribose ring were tolerated by the P2Y₄R, but also replacement of the ribose by a methanocarba (bicyclo[3.1.0]hexane) ring, which includes a ribose fixed in the (N)-conformation, without significant loss in potency (Jacobson et al., 2006; Kim et al., 2002; for more information on methanocarba compounds and their syntheses, see review article by Tosh & Jacobson, 2013). Related methanocarba analogues of adenosine were patented (Jacobson & Marquez, 2001). (N)-Methanocarba substitution is also tolerated by the P2Y₂R, which is another example of the large overlap of the pharmacological profiles of these two receptors (Kim et al., 2002). Another similarity to the P2Y₂R is that a thio-substitution at the γ -phosphate group is tolerated while α -thiophosphate is unfavorable (Jacobson et al., 2006).

Substitutions of one uridine moiety of Up₄U (**7-03**, Fig. 7 and Table 7) with various sugars as well as aliphatic or aromatic alcohol moieties led to the development of Up₄-[1'']3'-deoxy-3'-fluoroglucose (MRS2927, **8-20**, Fig. 8 and Table 8). It showed similar potency as UTP in the mid-nanomolar range, 11-fold selectivity over the P2Y₂R, and 15-fold selectivity versus the P2Y₆R. The δ -phenyl phosphoesters but not the δ -glucosylphosphoesters appeared to be enzymatically more stable than UTP (Maruoka et al., 2011).

3.2. P2Y₄ receptor antagonists

3.2.1. Therapeutic potential of P2Y₄ receptor antagonists

As P2Y₄R activation promotes epithelial Cl⁻ secretion in the intestine (refer to Section 3.1.1), blockade of this receptor could be beneficial for treating diarrhea. However, contrary to this, the P2Y₄R was found to

inhibit the contraction of mouse ileum longitudinal muscle (Zizzo, Mastropaolo, Grahlert, Mule, & Serio, 2012). Antagonists may induce contractions and have therefore also been proposed for treating constipation. It was further observed that the P2Y₄R (as well as the related P2Y₂R) is involved in the production and release of amyloid precursor protein in rat cortical astrocytes (Tran, 2011). Amyloid precursor protein can be cleaved to β -amyloid, and aggregates have been associated with Alzheimer's disease. P2Y₄R blockade may reduce the amount of amyloid precursor protein and, as a consequence, that of β -amyloid as well. It is thus postulated that P2Y₄R antagonists could prevent the formation of amyloid plaques. However, in rat microglia, P2Y₄R activation promoted pinocytosis of β -amyloid (Li et al., 2013). Therefore, the P2Y₄R appears to be a complex target in relation to neurodegenerative diseases, and better pharmacological tools would be required to further elucidate its role. Another possible indication for P2Y₄R antagonists might be myocardial infarction. The P2Y₄R appears to be an important mediator of the inflammatory response following cardiac ischemia. P2Y₄R-null mice displayed smaller infarcts in the left descending artery ligation model as well as reduced neutrophil infiltration and fibrosis, indicating that antagonists may be beneficial (Horckmans et al., 2015). The P2Y₄R (as well as the P2Y₂R) was found to be significantly overexpressed in different colon cancer cell lines, and antagonizing these two receptors may be beneficial for the treatment of certain cancers (Delbro, Nylund, & Nordgren, 2005; Nylund, Hultman, Nordgren, & Delbro, 2007). Moreover, P2Y₄R expression was shown to be upregulated under hypoglycemic conditions and linked to neuronal cell death. P2Y₄R blockade could thus have neuroprotective effects in diabetes (Cavaliere et al., 2004).

3.2.2. Anthraquinone derivatives

Potent and selective antagonists for the P2Y₄R are scarce. In light of insufficient alternatives, RB-2 (**10-07**, compound **07** in Fig. 10 and Table 10) has been the primary antagonist used for studying P2Y₄Rs. The antagonistic potential of RB-2 was already discovered in 1979, when the field of purinergic signaling was still in its early days. In fact, pharmacological studies involving RB-2 assisted in the differentiation and characterization of purinergic receptors (Kerr & Krantis, 1979).

RB-2 exhibited an IC₅₀ value of 1.14 μ M at the human P2Y₄R and was approximately 20-fold selective versus the P2Y₂R subtype, at which it also acted as a weak antagonist (Rafehi, Malik, et al., 2017; Rafehi, Neumann, et al., 2017). It is a dye that can be obtained from different commercial sources. Commercially available RB-2 is a mixture of isomers with differing identity and purity (Glänzel, Bültmann, Starke, & Frahm, 2003). This causes difficulties in obtaining reliable pharmacological results. For example, a certain batch obtained from Sigma-Aldrich contained only 54 % of RB-2, 12 % of its dichlorotriazinyl precursor, and 34 % of inorganic salts (NaCl and/or Na₃PO₄) (Rafehi, Malik, et al., 2017). As to be expected, purification of the commercially obtained compound by reversed phase-18 flash column chromatography yielded a pure product with increased potency at the P2Y₄R (IC₅₀ value of 0.625 μ M) (Rafehi, Malik, et al., 2017). A recently published review article provides comprehensive information on RB-2 and related anthraquinone derivatives, including a discussion on their safety and potential toxicity (Malik & Müller, 2016).

RB-2 (**10-07**, Fig. 10 and Table 10) has served as a lead structure for the extensive development of antagonists for different P2 receptors. Examples include the potent and selective competitive P2Y₁₂R antagonist 1-amino-4-[4-phenylamino-3-sulfophenylamino]-9,10-dioxo-9,10-dihydroanthracene-2-sulfonate (PSB-0739; A₂ value of 0.158 nM) and disodium 1-amino-4-[3-(4,6-dichloro[1,3,5]triazine-2-ylamino)-4-sulfophenylamino]-9,10-dioxo-9,10-dihydroanthracene-2-sulfonate (PSB-1011; IC₅₀ of 79 nM), a competitive inhibitor of the rat P2X₂ receptor (Baqi et al., 2011; Baqi, Atzler, Köse, Glänzel, & Müller, 2009; Hoffmann et al., 2009). Although several P2 receptors are blocked by RB-2 derivatives, SARs are significantly different for each receptor (Malik & Müller, 2016).

Recently, the synthesis of a series of P2Y₄R antagonists that are smaller-size analogues of RB-2 was reported (Rafehi, Malik, et al., 2017). While the anthraquinone scaffold was retained, the three aromatic rings in the 4-position of the anthraquinone core of RB-2 were reduced to one or two, and the substituents were varied. This was generally well-tolerated by the P2Y₄R. The most potent compound of this series, 1-amino-4-[4-(2,4-dimethylphenylthio)]-9,10-dioxo-9,10-dihydroanthracene-2-sulfonate (PSB-16133, **10-06**, Fig. 10 and Table 10), showed an IC₅₀ value of 233 nM (Rafehi, Malik, et al., 2017). While its improvement in potency compared to RB-2 was moderate (5-fold), its selectivity towards other P2Y_R subtypes was significantly enhanced: PSB-16133 is 37-fold and 54-fold selective for the P2Y₄R versus the P2Y₂ and P2Y₆ receptors, respectively. In comparison, the selectivity of RB-2 for the P2Y₄R is 19-fold and merely 2-fold versus the P2Y₂R and P2Y₆R, respectively (Rafehi, Malik, et al., 2017; Rafehi, Neumann, et al., 2017). Based on pharmacological assessments and molecular docking to a homology model of the P2Y₄R, the anthraquinone derivatives were described as non-competitive antagonists binding to a site located close to the extracellular surface above the putative orthosteric binding site (Rafehi, Malik, et al., 2017).

4. The P2Y₆ receptor

The rat P2Y₆R was first recombinantly expressed in 1995 using cDNA isolated from a rat aortic smooth muscle cell library (Chang, Hanaoka, Kumada, & Takuwa, 1995). Cloning of the human ortholog succeeded a year later following a search for further P2Y_R subtypes by screening of a human placenta cDNA library using a P2Y₄R probe (Communi, Parmentier, & Boeynaems, 1996). P2Y₆R mRNA is strongly expressed in the spleen, placenta, and kidney. It was also detected at lower levels in the CNS, lung, heart, GI tract, adipose tissue, bone, and other parts of the immune system (Table 1) (Moore et al., 2001; Uhlén et al., 2015).

4.1. P2Y₆ receptor agonists

4.1.1. Therapeutic potential of P2Y₆ receptor agonists

A possible physiological function of the P2Y₆R may be, similar to the P2Y₂R and P2Y₄R, to mediate electrolyte transport in the airway epithelium and the colon (Dulong, Bernard, & Ehrenfeld, 2007; Köttgen et al., 2003; Schreiber & Kunzelmann, 2005). The P2Y₆R may therefore be another possible target for cystic fibrosis. Furthermore, P2Y₆R knockout mice revealed an involvement in the contraction and endothelium-dependent relaxation of the aorta (Bar et al., 2008). Thus, the P2Y₆R could additionally become a target for controlling blood pressure. The receptor was further found to be involved in insulin secretion and is thereby of potential interest for the treatment of diabetes (Balasubramanian, Ruiz de Azua, Wess, & Jacobson, 2010; Ohtani, Suzuki, Jacobson, & Oka, 2008; Parandeh, Abaraviciene, Amisten, Erlinge, & Salehi, 2008). In analogy to P2Y₂R agonists, P2Y₆R activation could become a new strategy for treating ocular hypertension and glaucoma. Topical application of the P2Y₆R agonist UDP led to a reduction in intraocular pressure in mice and rabbits, and P2Y₆R knockout mice showed hypertensive glaucoma-like optic neuropathy (Jacobson & Civan, 2016; Markovskaya et al., 2008; Shinozaki et al., 2017). Furthermore, P2Y₆R activation was observed to have an anti-apoptotic function in different cell types by preventing the activation of caspases 3 and 8 following tumor necrosis factor- α exposure (Haas, Ginsburg-Shmuel, Fischer, & Reiser, 2014; Kim et al., 2003; Kim et al., 2003). This effect was also observed in mouse C2C12 skeletal muscle cells, and P2Y₆R agonism protected against skeletal muscle ischemia/reperfusion injury in mouse hindleg muscle. Thus, P2Y₆R agonists could be useful in skeletal muscle injuries (Mamedova, Wang, Besada, Liang, & Jacobson, 2008). The P2Y₆R was further shown to protect cells from γ -rays by promoting cellular repair responses (Ide, Nishimaki, Tsukimoto, & Kojima, 2014). Anti-proliferative effects on gastric cancer cells were associated with P2Y₆R activation (Wan et al., 2017). Moreover, P2Y₆R activation

induced recruitment of monocytes and macrophages in response to bacterial invasion. It also triggered an antiviral immune response following infection with vesicular stomatitis virus. These mechanisms could be employed in the therapy of infectious diseases (Li et al., 2014; Zhang et al., 2011). Patients suffering from neurodegenerative diseases may also benefit from P2Y₆R agonists. This is based on the observation that the P2Y₆R functions as a sensor for phagocytosis on microglia. P2Y₆R agonists could thus promote the clearance of dead cells or harmful debris in the brain (Inoue, Koizumi, Kataoka, Tozaki-Saitoh, & Tsuda, 2009; Koizumi et al., 2007). Altogether, it is evident that P2Y₆R agonists (Tables 2 to 10) are in high demand for a wide range of research fields and clinical applications.

4.1.2. Physiological mononucleotides

The endogenous agonist for the P2Y₆R is UDP (**2-09**, compound **09** in Fig. 2 and Table 2) (Communi, Parmentier, & Boeynaems, 1996). However, since it also activates the P2Y₁₄R, it is not selective (Carter et al., 2009). UDP can be phosphorylated to UTP by nucleoside diphosphokinase, and commercial preparations of UDP may contain impurities of UTP, which should be taken into consideration in pharmacological experiments (Lazarowski et al., 2000; Nicholas et al., 1996). Other physiological mononucleotides, e.g. UTP (**2-08**), GDP (**2-04**), and thymidine 5'-diphosphate (TDP, **2-14**), can also activate the P2Y₆R at high concentrations (Table 2). Uracil nucleotides are preferred over adenine nucleotides, and diphosphates are more potent than triphosphates. Accordingly, ATP (**2-01**, Fig. 2 and Table 2) is almost inactive (Communi, Parmentier, & Boeynaems, 1996; Nicholas et al., 1996).

4.1.3. Prostaglandin E₂ glyceryl ester

Very recently, prostaglandin E₂ glyceryl ester (**9-02**, Fig. 9 and Table 9) was proposed as another physiological P2Y₆R agonist with (sub-)picomolar potency (Brüser et al., 2017). Such a high potency is rarely seen with endogenous ligands but appears physiologically reasonable, as **9-02** is produced in low concentrations by cyclooxygenase 2 and disappears rapidly due to hydrolysis. It had previously been shown to induce calcium mobilization and activate protein kinase C as well as extracellular signal-regulated kinase 1/2 (Brüser et al., 2017). To identify the receptor responsible for these effects, mRNA encoding for GPCRs from different cell lines was sequenced. The GPCR expression profiles of cell lines responsive to **9-02** were compared with those that could not be stimulated by **9-02**. Six GPCRs were found to be expressed exclusively in the responsive cell lines, and thus were considered as potential candidates for a putative prostaglandin E₂ glyceryl ester receptor. The P2Y₆R was eventually discovered as the target of **9-02** using calcium mobilization, IP₃, and ERK1/2 phosphorylation assays in recombinant and native cells (Brüser et al., 2017). Based on a P2Y₆R homology model and mutagenesis studies, the binding pocket of **9-02** was proposed to partially overlap with that of UDP. Several mutations resulted in a loss of response to **9-02** but not to UDP, and *vice versa*. This indicated a direct activation of the P2Y₆R by **9-02** as opposed to a paracrine or autocrine mechanism in which **9-02** would induce UDP release leading to P2Y₆R activation (Brüser et al., 2017).

It would not be unusual for the P2Y₆R to be activated by both nucleotides and lipids in light of the fact that the phylogenetically related cysteinyl leukotriene receptors 1 and 2 (CysLT1 and CysLT2) respond to lipids (cysteinyl leukotrienes) as well as to the nucleotide UDP (Costanzi et al., 2004; Mellor et al., 2003; Mellor, Maekawa, Austen, & Boyce, 2001). So far, **9-02** has not been assessed at other P2Y_R subtypes. Should these recent findings be confirmed, the non-nucleotide lipid agonist with extraordinarily high potency would likely inspire new compound or perhaps even drug development efforts for the P2Y₆R and the other P2Y_R subtypes. The low stability of prostaglandin E₂ derivatives under physiological conditions, however, constitutes a major challenge (Kozak et al., 2001).

4.1.4. Dinucleotides and analogues

The larger dinucleotide tetraphosphates (Fig. 7 and Table 7) are less potent at the P2Y₆R compared to the nucleotide triphosphate-preferring receptors P2Y₂ and P2Y₄. Up₄U (**7-03**) has a potency in the micromolar range, while Ap₄A (**7-01**) showed only low activity at 100 μM (Maruoka et al., 2011; Pendergast et al., 2001; Shaver et al., 2005). Up₃U (**7-04**) is of higher potency (EC₅₀ of 200–920 nM) and similarly potent as UDP (Maruoka et al., 2010; Pendergast et al., 2001). This matches the pattern observed for the triphosphate nucleotide-preferring P2YR subtypes, where dinucleotide tetraphosphates mimic the activity of the corresponding triphosphates (Brunschweiler & Müller, 2006; Pendergast et al., 2001).

P¹-(Uridine-5'-)-P³-(N⁴-methoxycytidine-5'-)-triphosphate (N⁴-(MeO)-Cp₃U, MRS2957, **7-08**, Fig. 7 and Table 7) was 25-fold more potent than UDP and exhibited 14-fold and 66-fold selectivity versus the P2Y₂R and P2Y₄R, respectively (Maruoka et al., 2010). The stable analogue of Up₃U, P¹-(2-benzyl-1,3-dioxolo-4-yl)uridine-5'-)-P³-(uridine-5'-)-triphosphate (INS48823, monobenzylacetal-Up₃U, **7-11**, Fig. 7 and Table 7), is another P2Y₆R-selective agonist with an EC₅₀ value of 125 nM and no appreciable activity at the P2Y₂R and P2Y₄R (Korcok, Raimundo, Du, Sims, & Dixon, 2005).

4.1.5. Boranophosphate analogues of nucleotides

Among the most promising P2Y₆R agonists are boranophosphate analogues of UDP (**2-09**). The R_p isomer of 5-MeO-uridine 5'-O-(α-boranodiphosphate) (**6-12**, Fig. 6 and Table 6) showed an EC₅₀ value of 8 nM, and was thus 19-fold more potent than UDP (Ginsburg-Shmuel et al., 2012). It is the most potent P2Y₆R agonist described to date. Since it exhibited no activity at the P2Y₂R and P2Y₄R, it is also the most selective one. Furthermore, it was found to be very stable in acidic conditions (e.g. in gastric juice) as well as in blood serum (Ginsburg-Shmuel et al., 2012). However, the negative charges of this compound could be an impediment to its usefulness *in vivo*, as is the case for all nucleotide analogues, which are therefore not orally bio-available. Compound **6-12** was found to protect 1321N1 astrocytoma cells transfected with the human P2Y₆R from apoptosis induced by tumor necrosis factor-α; it did so more prominently than UDP (Haas et al., 2014). The agonist was patented and its therapeutic potential is under further investigation (Fischer, Pintor, Elyahu, & Ginsburg-Shmuel, 2011). Compound **6-12** is an analogue of 5-MeO-UDP (**3-31**), which was developed two years earlier through a data mining analysis of binding interactions in 44 protein-uridine nucleoside or nucleotide complexes (Ginsburg-Shmuel et al., 2010). Although dinucleotide derivatives were less active than mononucleotides, the introduction of an α-borano-group enhanced their potency (Ginsburg-Shmuel et al., 2012; also see patent by Fischer & Nahum, 2006). Interestingly, the P2Y₂R does not appear to tolerate the presence of an α-borano-substitution; α-borano-substituted ATP and 5-MeO-UTP are less potent at the P2Y₂R than their non-substituted analogues, which explains their selectivity for the P2Y₆R over the P2Y₂R (Ginsburg-Shmuel et al., 2012; Tulapurkar, Laubinger, Nahum, Fischer, & Reiser, 2004).

4.1.6. Phosphorothioate derivatives of nucleotides

Unsurprisingly, uridine 5'-(β-thio)-diphosphate (UDPβS, **5-06**, Fig. 5 and Table 5) potently (EC₅₀ of 28 nM, similar to UDP) activates the P2Y₆R. This is analogous to UTPγS (**5-05**, Fig. 5 and Table 5) being an agonist at the P2Y₂R and the P2Y₄R, and ATPγS (**5-03**, Fig. 5 and Table 5) at the P2Y₂R (Hou et al., 2002; Lazarowski et al., 1995; Lazarowski et al., 1996; Nicholas et al., 1996). UDPβS was equally potent at the P2Y₁₄R, which is not astonishing in light of the fact that UDP functions as a potent P2Y₁₄R agonist as well (Carter et al., 2009). Although UMP (**2-10**, Fig. 2 and Table 2) was inactive at the P2Y₆R, uridine 5'-O-monophosphorothioate (UMPS, **5-07**, Fig. 5 and Table 5) showed some activity (62 % activation at 10 μM, relative to the response induced by 1 μM UDP). Its structural analogue thymidine 5'-O-monophosphorothioate (TMPS, **5-08**, Fig. 5 and Table 5) was described

as a selective partial agonist at the P2Y₆R. At 1 mM concentration, it elicited 50 % receptor activation relative to the effect observed with 1 μM UDP (Gendaszewska-Darmach & Szustak, 2016). At the P2Y₁₄R, TMPS appeared to have antagonistic effects (Gendaszewska-Darmach, Weglowska, Walczak-Drzewiecka, & Karas, 2016). The terminal thiophosphate group increases stability towards different ectonucleotidases (Gendaszewska-Darmach et al., 2016; Gendaszewska-Darmach & Szustak, 2016).

4.1.7. Nucleobase-modified nucleotide analogues

Large substituents in the N3-position of the uracil base were well tolerated by the P2Y₆R, which distinguishes it from the P2Y₂R and the P2Y₄R (El-Tayeb et al., 2006). As a result, 3-phenacyl-UDP (PSB-0474, **3-11**, Fig. 3 and Table 3) was 570-fold selective for the P2Y₆R versus the P2Y₂R and the P2Y₄R. With an EC₅₀ of 70 nM, it is among the most potent P2Y₆R agonists described to date, and has been made commercially available (El-Tayeb et al., 2006). Also interesting are the following cytidine 5'-diphosphate (CDP) derivatives with mid-nanomolar potencies (Fig. 3 and Table 3): N⁴-(benzyloxy)-CDP (MRS2964, **3-17**; 82- and 44-fold selectivity versus the P2Y₂R and P2Y₄R, respectively) and N⁴-(MeO)-CDP (**3-19**; 51-, 92-, and 47-fold selectivity versus the P2Y₂, P2Y₄, and P2Y₁₄ receptors). An analogue of **3-19**, α,β-methylene-N⁴-MeO-CDP (**6-13**, Fig. 6 and Table 6), exhibited an EC₅₀ value of 678 nM and 15-fold selectivity (Maruoka et al., 2010). Different nucleobase modifications led to further potent and selective P2Y₆R agonists, including 5-MeO-UDP (**3-31**, Fig. 3 and Table 3; EC₅₀ of 80 nM, 250-fold selectivity) and 5-iodo-UDP (MRS2693, **3-26**, Fig. 3 and Table 3; EC₅₀ of 15 nM) (Besada et al., 2006; Ginsburg-Shmuel et al., 2010). Substitutions in the 6-position appeared to be unfavorable for the P2Y₆R, as was the case for the P2Y₂R and P2Y₄R (Ginsburg-Shmuel et al., 2010). Although different substitutions at the uracil base were tolerated by the P2Y₆R, complete replacement by other nucleobases was not (Robaye, Boeynaems, & Communi, 1997).

4.1.8. Ribose-modified nucleotide analogues

Replacement of the ribose moiety by a constrained methanocarba (bicyclo[3.1.0]hexane) ring was tolerated by the P2Y₆R, but only if the methanocarba ring was in the (S)-configuration (Kim et al., 2002). This is in contrast to the other G_q-coupled human P2YRs, all of which show a preference for the (N)-configuration (Costanzi et al., 2005; Kim et al., 2012). The rigid methanocarba analogues of UDP in the (S)-configuration, (S)-methanocarba-UDP (MRS2795, **4-22**, Fig. 4 and Table 4) and 2'-deoxy-(S)-methanocarba-UDP (**4-23**), both displayed a 7-fold higher potency than the flexible UDP and 2'-deoxy-UDP, respectively (Costanzi et al., 2005; Maruoka et al., 2010). Interestingly, **4-23** had been developed following a rational design using molecular modeling (Costanzi et al., 2005). The hydroxyl group in the 2'-position of the ribose ring was found to be important; the 2'-deoxy derivatives of UDP (**4-02**) and (S)-methanocarba-UDP (**4-23**) were 5-fold less potent than their corresponding 2'-hydroxy-substituted analogues (**2-09**) and **4-22** (Maruoka et al., 2010; Meltzer et al., 2015). Similar to the P2Y₂R and the P2Y₄R, 2'-amino-2'-deoxy and 2'-azido-2'-deoxy substitutions at the ribose moiety caused a reduction in potency at the P2Y₆R (Besada et al., 2006; Jacobson et al., 2006).

4.1.9. Fluorescence-labelled P2Y₆ receptor agonists

The BODIPY® 630/650 conjugate MRS4162 (**3-40**, Fig. 3 and Table 3) that was mentioned in Section 2.1.5. can be used to characterize not only the P2Y₂ and the P2Y₄R but for the P2Y₆R as well. At the P2Y₆R, it displayed an EC₅₀ value of 23 nM (Jayasekara et al., 2014). MRS4129 (**3-42**, Fig. 3 and Table 3) is another tool compound, which is coupled to a different fluorescent dye: Alexa Fluor® 488. It showed an EC₅₀ value of 9 nM at the P2Y₆R and, in contrast to **3-40**, is highly selective versus the P2Y₂R (280-fold) and the P2Y₄R (>1100-fold) (Jayasekara et al., 2013). Both compounds are cytosine nucleotide derivatives with

alkyloxyimino substitutions in the N^4 -position. MRS4162 (**3-40**) contains three phosphate groups but MRS4129 (**3-42**) is a diphosphate (Jayasekara et al., 2013; Jayasekara et al., 2014).

4.2. P2Y₆ receptor antagonists

4.2.1. Therapeutic potential of P2Y₆ receptor antagonists

P2Y₆R activation was shown to prolong the survival of osteoclasts by activating NF- κ B, a key transcription factor regulating osteoclastogenesis. It was also found to stimulate the formation of osteoclasts from precursor cells (Korcok et al., 2005; Orriss et al., 2011). Antagonists of the P2Y₆R may therefore find use in the treatment of inflammatory bone diseases, such as rheumatoid arthritis and periodontitis (Table 1). P2Y₆R antagonism could also be beneficial in treating other excessive inflammatory reactions, including inflammatory bowel disease, gout, allergic airway inflammation, atherosclerosis, and CNS disorders (Cox et al., 2005; Garcia et al., 2014; Grbic et al., 2012; Grbic, Degagne, Langlois, Dupuis, & Gendron, 2008; Guns, Hendrickx, van Assche, Fransen, & Bult, 2010; Hao, Liang, Chow, Cheung, & Ko, 2014; Khine et al., 2006; Kim et al., 2011; Morioka et al., 2013; Müller et al., 2017; Nakano et al., 2017; Riegel et al., 2011; Shi et al., 2016; Sil et al., 2017; Somers, Hammet, Trute, Southey, & Venter, 1998; Stachon et al., 2014; Uratsuji et al., 2012; Vieira et al., 2011; Warny et al., 2001). Pro-nociceptive effects were attributed to the P2Y₆R in a rat model of inflammatory pain; P2Y₆R antagonists could thus be useful for pain therapy (Barragán-Iglesias et al., 2015). The P2Y₆R was also shown to mediate ATP release from the urothelium, leading to increased bladder sensation and detrusor overactivity in patients with benign prostatic hyperplasia. P2Y₆R antagonism was thus proposed as a novel therapeutic strategy to control persistent storage symptoms in patients with bladder outflow obstruction (Silva, Ferreirinha, Magalhães-Cardoso, Silva-Ramos, & Correia-de-Sá, 2015). Since the P2Y₆R was found to induce interleukin-8 production and epithelial barrier dysfunction following exposure to *Clostridium difficile* toxins, it may further become a target for the treatment of nosocomial diarrhea (Hansen et al., 2013). Moreover, P2Y₆R antagonists may be useful for treating obesity, as P2Y₆R activation in orexigenic agouti-related peptide-expressing neurons in the arcuate nucleus of the hypothalamus was shown to promote feeding in mice. Accordingly, P2Y₆R deficiency in these neurons led to a reduced food intake and improved systemic insulin sensitivity (Steculorum et al., 2015; Steculorum et al., 2017). P2Y₆R mRNA was upregulated in breast cancer cells and its activation facilitated breast cancer metastasis both *in vitro* and *in vivo*. P2Y₆R activation further prolonged survival of different cell lines (Azimi et al., 2016; Gendaszewska-Darmach & Szustak, 2016; Ma et al., 2016). P2Y₆R antagonists could, similar to P2Y₂R antagonists, be useful for cancer therapy. The P2Y₆R appears to be involved in cardiac development, as is the P2Y₄R, but with seemingly opposite effects. P2Y₆R-null mice showed macrocardia and amplified pathological cardiac hypertrophy (Clouet et al., 2016; Horckmans et al., 2012). It was thus postulated that P2Y₆R antagonists could potentially be used in the therapy of these pathological conditions in humans. Furthermore, blockade of the P2Y₆R was proposed as a strategy to prevent pressure overload-induced cardiac fibrosis (Nishida et al., 2008). P2Y₆R mRNA expression was found to be upregulated in response to the inflammatory mediator interleukin-1 β in the rat aorta, where it promotes vascular smooth muscle growth and differentiation (Hou et al., 2002). It is therefore possible that the P2Y₆R contributes to atherosclerosis and the formation of neointima. As the P2Y₆R is also involved in the contraction of human cerebral arteries, antagonists may be useful for preventing cerebral vasospasms (Malmjö, Hou, Pendergast, Erlinge, & Edvinsson, 2003). Moreover, the P2Y₆R was associated with oxidative stress in neurons and neural degeneration (Qian, Xu, Yang, & Xiao, 2017). P2Y₆R antagonism could thus be beneficial for the treatment of neurodegenerative disorders, including Parkinson's disease.

4.2.2. Diisothiocyanate derivatives

The first selective P2Y₆R antagonists described in the literature were from a series of symmetrical aryl diisothiocyanate derivatives (Mamedova, Joshi, Gao, von Kügelgen, & Jacobson, 2004). They were developed following the observation that 4,4'-diisothiocyanatostilbene-2,2'-disulfonic acid disodium salt and its dihydro derivative were capable of inhibiting UDP-induced rat P2Y₆R activation at micromolar concentrations. Substitution of one of the two isothiocyanate groups caused a loss in activity, indicating that both isothiocyanate functions were essential (Mamedova et al., 2004). The most potent and selective compounds of this series are MRS2578 (**10-08**, compound **08** in Table 10 and Fig. 10), MRS2567 (**10-09**), and MRS2575 (**10-10**). They showed IC₅₀ values of 37 nM, 126 nM, and 155 nM, respectively, at the human P2Y₆R. At the related P2Y₁R, P2Y₂R, P2Y₄R, and P2Y₁₁R, all three compounds were inactive at 10 μ M; the only exception was a slightly higher potency of **10-08** at the P2Y₁R. Compound **10-09** had similar effects on the human and the rat P2Y₆Rs, while **10-08** was slightly less potent at the rat ortholog (IC₅₀ of 98 nM) and **10-10** was inactive at the rat receptor (Mamedova et al., 2004). These aryl diisothiocyanate derivatives were described as insurmountable antagonists, as they were unable to induce a parallel shift of UDP concentration-response curves. Due to the absence of radioligands for the P2Y₆R, selected derivatives of this compound series that showed activity at the P2Y₁R were assessed in P2Y₁R radioligand binding assays. They were incapable of displacing the radioligand from its binding site on the P2Y₁R and thus were proposed to be non-competitive antagonists (Mamedova et al., 2004). Owing to their hydrophobic nature, the diisothiocyanate derivatives display low solubility in aqueous buffer, which limits their usefulness as pharmacological tools. Disadvantages include their reactivity towards nucleophiles and their instability in aqueous medium. They are likely to bind covalently, and thus irreversibly, to the P2Y₆R (Jacobson, Ivanov, de Castro, Harden, & Ko, 2009). Compound **10-08** is commercially available from several sources and currently the most widely used P2Y₆R antagonist. However, it may not be suitable for *in vivo* applications.

4.2.3. Other P2Y₆ receptor antagonists

In order to obtain alternatives to the diisothiocyanate derivatives, it was attempted to develop nucleotide-based P2Y₆R antagonists by reducing the efficacy of UDP through different substitutions at the nucleobase, the ribose moiety, and the phosphate chain (Meltzer et al., 2015). These efforts yielded comprehensive SAR data and an antagonist – uridylyl phosphosulfate (**10-13**, Fig. 10 and Table 10) – that displayed modest potency (P2Y₆R IC₅₀ of 112 μ M) but selectivity versus P2Y₂ and P2Y₄Rs.

The three non-specific P2 receptor antagonists suramin (**10-04**), RB-2 (**10-07**), and pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid (PPADS, **10-12**) are weakly active at the P2Y₆R (see Fig. 10 and Table 10). At 100 μ M, RB-2 achieved 87 % inhibition (IC₅₀ of 31 μ M), PPADS 69 %, and suramin only 27 % inhibition of the effect mediated by 2 μ M UDP (Robaye et al., 1997). RB-2 may behave as a competitive antagonist, as increasing concentrations caused a parallel shift of UDP concentration-response curves without affecting the upper plateau (Robaye et al., 1997).

5. The P2Y₁₄ receptor

The P2Y₁₄R was discovered in 1994 following a systematic approach to identify uncharacterized human genes through predicting their coding sequences in an immature human myeloid cell line (Nomura et al., 1994). It was initially referred to as KIAA0001 or GPR105 until its deorphanization in the year 2000. The endogenous agonist was found via the screening of a library of 700 known and putative GPCR ligands (Chambers et al., 2000). The P2Y₁₄R sequence contains several features that are commonly found in GPCRs. These include a DRY motif between transmembrane domain 3 and the second intracellular

loop as well as consensus sites for asparagine-linked glycosylation on extracellular sequences and for phosphorylation by the protein kinases A and C on the third intracellular loop (Chambers et al., 2000). Unlike the other uracil-activated P2YRs, the P2Y₁₄R signals mainly through G_{i/o}, thereby inhibiting cAMP production by adenylate cyclase. It was also found to activate extracellular signal-regulated kinase 1/2 (Carter et al., 2009; Fricks, Carter, Lazarowski, & Harden, 2009; Hamel et al., 2011; Scrivens & Dickenson, 2006). The protein sequence of the P2Y₁₄R is 18–45 % identical to that of the other P2YRs, and it shares greatest sequence homology with the P2Y₁₂R and the P2Y₁₃R (particularly in the transmembrane regions) (Abbracchio et al., 2006). The rat ortholog of the P2Y₁₄R, also known as VTR15–20, is 80 % and the mouse ortholog 83 % identical to the human receptor. The sequence identity shared between the rat and mouse receptors is 89 % (Freeman et al., 2001; Fricks et al., 2008). The P2Y₁₄R appears to be ubiquitously expressed and was detected at protein level in most tissues and organs (Table 1) (Moore et al., 2001; Uhlén et al., 2015). The physiological function of the P2Y₁₄R remains to be confirmed, but experimental observations and its prominent expression in immune cells suggest an involvement in inflammation and immune responses (Azroyan et al., 2015; Barrett et al., 2013; Charlton, Williams, Fogliano, Sweetnam, & Duman, 1997; Lazarowski & Harden, 2015; Li et al., 2016).

5.1. P2Y₁₄ receptor agonists

5.1.1. Therapeutic potential of P2Y₁₄ receptor agonists

The P2Y₁₄R, together with other P2YRs, was shown to be expressed on human plasmacytoid dendritic cells. Activation through extracellular nucleotides inhibited the release of interferon- α , possibly to avoid excessive tissue damage or the induction of autoimmunity. It was thus postulated that P2Y₁₄R agonists could be useful in treating autoimmune diseases (Shin et al., 2008). P2Y₁₄R knockout mice showed impaired insulin secretion, and agonists might be useful for treating diabetes (Meister et al., 2014).

5.1.2. Physiological agonists

The main function of UDP-glucose (8-01, compound 01 in Fig. 8 and Table 8) was commonly believed to serve as a glycosyl donor in the enzymatic biosynthesis of carbohydrates. Later, it was discovered to be released from cells as a signaling molecule and to function as an agonist at the P2Y₁₄R (Carter et al., 2009; Chambers et al., 2000; Hamel et al., 2011; Lazarowski & Harden, 2015; Lazarowski, Shea, Boucher, & Harden, 2003). Other sugar derivatives of UDP (Fig. 8 and Table 8), including UDP-galactose (8-02), UDP-glucuronic acid (8-03), and UDP-N-acetylglucosamine (8-04), can also activate the P2Y₁₄R, albeit less potently (Chambers et al., 2000; Hamel et al., 2011; Ko et al., 2009; Ko, Fricks, Ivanov, Harden, & Jacobson, 2007). Some of these nucleotide sugars have also been described as partial agonists, but reported discrepancies could be due to different receptor expression levels (Fujioka & Omori, 2012; Ko et al., 2007; Scrivens & Dickenson, 2005). The pharmacological profile of UDP-glucose, UDP-galactose, UDP-glucuronic acid, and UDP-N-acetylglucosamine is similar for the rat and mouse orthologs (Freeman et al., 2001). In contrast, the glucose conjugates of CDP, ADP, and guanosine 5'-diphosphate (GDP) were inactive (Chambers et al., 2000; Ko et al., 2007). The responsiveness to nucleotide sugars is a characteristic but not an exclusive feature of the P2Y₁₄R, as UDP-glucose was also shown to activate the P2Y₂R and P2Y₆R at mid-micromolar concentrations (Das et al., 2010; Ko et al., 2009).

UDP (2-09, Fig. 2 and Table 2) was initially found inactive at the P2Y₁₄R in a yeast transcription reporter assay, in an assay using transient expression with a promiscuous G α_{16} protein, and also in [³⁵S] GTP γ S binding assays. It was also previously described as a competitive antagonist or as a partial agonist at the human P2Y₁₄R (and an agonist at the rat ortholog) when using a chimeric G $\alpha_{q/i}$ protein that couples G_i-activating receptors to phospholipase C β and inositol lipid hydrolysis

(Ault & Broach, 2006; Chambers et al., 2000; Fricks et al., 2008). However, these test systems were later found unsuitable, and UDP was subsequently confirmed as a full P2Y₁₄R agonist in a more natural assay system: when measuring P2Y₁₄R-mediated adenylate cyclase inhibition in three stably transfected cell lines, UDP was consistently shown to be even slightly more potent than UDP-glucose and to exhibit similar efficacy (Carter et al., 2009; Hamel et al., 2011; Harden, Sesma, Fricks, & Lazarowski, 2010). Other mononucleotides and different dinucleotides (e.g. Ap_{3/4/5/6}A, UP₄U, Up₂U, and Cp₂C) were less potent or inactive at the P2Y₁₄R. This is a distinction from the other P2YRs and further emphasizes the unique pharmacological profile of the P2Y₁₄R (Carter et al., 2009; Chambers et al., 2000; Ko et al., 2007).

5.1.3. Synthetic nucleotide analogues

A thio substitution in the 2-position of the uracil moiety caused an improvement in potency at the P2Y₁₄R: 2-thio-UDP (3-02, Fig. 3 and Table 3) was 37- to 83-fold more potent than UDP (Carter et al., 2009; Das et al., 2010). However, this substitution is also well tolerated by the P2Y₆R. Although being 230-fold selective for the P2Y₁₄R over the P2Y₆R, 3-02 activates the P2Y₆R at nanomolar concentrations. It is thus not ideal for discerning these two receptors experimentally (Das et al., 2010; El-Tayeb et al., 2006; Jacobson et al., 2006). The benefits of a 2-thio substitution could be combined with other modifications at the phosphate groups of UDP to obtain compounds with (sub) nanomolar potency and significantly enhanced selectivity versus the P2Y₆R (Das et al., 2010). Among the best P2Y₁₄R agonists is α,β -methylene-2-thio-UDP (MRS2905, 6-08, Fig. 6 and Table 6). It showed an EC₅₀ value of 0.92 nM at the P2Y₁₄R, which makes this compound 170-fold more potent than UDP and 2000-fold selective versus the P2Y₆R. In contrast, an analogue without substitution in the 2-position, α,β -methylene-UDP (5-13, Fig. 5 and Table 5), was merely 15-fold more potent at the P2Y₁₄R than UDP and 31-fold selective versus the P2Y₆R. The derivative α,β -difluoromethylene-UDP (MRS2802, 5-15, Fig. 5 and Table 5) may also be used to pharmacologically distinguish the P2Y₁₄R (EC₅₀ of 63 nM) from the P2Y₆R (inactive at 10 μ M) (Das et al., 2010). Due to the replacement of the diphosphate by a phosphonate, it is expected to be metabolically stable and therefore useful for *in vivo* studies.

5.1.4. Synthetic nucleotide-sugar derivatives

A simple 2-thio substitution of UDP-glucose led to a 36-fold improvement in potency at the P2Y₁₄R. Unlike 2-thio-UDP (3-02, Fig. 3 and Table 3), 2-thio-UDP-glucose (MRS2690, 8-15, Fig. 8 and Table 8) was inactive at the related P2Y₆R (Das et al., 2010). The selectivity versus the P2Y₂R was also improved through this substitution: MRS2690 was inactive at the P2Y₂R at 10 μ M. In contrast, UDP-glucose (8-01, Fig. 8 and Table 8) showed half-maximal P2Y₂R activation at the same concentration (Ko et al., 2009). A thio substitution in the 4-position was either not tolerated at all (4-thio-Up₂-OMe was inactive), or it had no significant effect on the potency (UDP-glucose and 4-thio-UDP-glucose (MRS2670, 8-16, Fig. 8 and Table 8) were equipotent) (Das et al., 2010; Ko et al., 2007). Different substitutions in 5-position or complete replacement of the uracil moiety by adenine, guanine, and cytosine resulted in inactive compounds (Ko et al., 2007).

The glucose moiety of UDP-glucose was proposed to form H-bonding interactions with charged residues in the putative ligand binding pocket. Yet, this region appears to be the part at which substitutions are best tolerated by the P2Y₁₄R. The glucose moiety of UDP-glucose was replaced by different sugars in a systematic approach to investigate the tolerance of the P2Y₁₄R to modifications at this site (Ko et al., 2009). Most analogues were found active (e.g. compounds 8-05 to 8-13, Fig. 8 and Table 8), further indicating a large potential for substitutions at the glucose. However, their potencies were mostly lower or similar to that of UDP-glucose, and it has not yet been possible to exploit the steric freedom at the glucose moiety to enhance the potency (Ko et al., 2009).

UDP could further be substituted by smaller ester groups at the terminal phosphate. Potencies in the nanomolar and low micromolar range were achieved with a series of UDP- β -esters, which was further enhanced if combined with a 2-thio substitution. For example, Up₂-OMe (not shown) exhibited an EC₅₀ value of 2.73 μ M, while its 2-thio analogue **6-17** (Fig. 6 and Table 6; EC₅₀ of 0.056 μ M) was 50-fold more potent (Das et al., 2010). The ethoxy (MRS2906, **6-18**), propoxy (MRS2907, **6-19**), and *tert*-butyl (**6-20**) derivatives of 2-thio-Up₂ as well as **6-21** were similarly of mid-nanomolar potency (Fig. 6 and Table 6). They were either inactive at the P2Y₆R (**6-17**, **6-18**, **6-21**), or 64-fold (**6-20**) and 230-fold (**6-19**) selective for the P2Y₁₄R. Larger branched β -alkyl esters (e.g. cyclohexyl) or β -aryloxy esters were less potent (Carter et al., 2009; Das et al., 2010). The chemical and metabolic stability of these phosphoric acid ester derivatives has not been investigated but may be limited.

5.1.5. Modifications at the ribose moiety

Replacement of the flexible ribose moiety with a constrained methanocarba (bicyclo[3.1.0]hexane) ring led to inactivity at the P2Y₁₄R (Das et al., 2010). This is a disparity from the other uracil-activated P2YRs, where a ring constraint in the preferred position (North for the P2Y₂R and P2Y₄R, South for the P2Y₆R) was tolerated (Kim et al., 2002; Maruoka et al., 2010). In fact, the P2Y₁₄R appears to be highly restrictive with regard to modifications at the ribose. Several different substitutions in the 2'- and 3'-position, some of which enhanced the potency or selectivity for the P2Y₂R and P2Y₄R, caused a complete loss of activity at the P2Y₁₄R (Ko et al., 2007).

5.1.6. Nucleotide-dendrimer conjugates

Dendrimers are branched, polymeric macromolecules that often form a spherical three-dimensional shape and are typically symmetrical around the core. They can serve as nano-carriers for drug delivery. UDP-glucuronic (**8-03**, Fig. 8 and Table 8) acid as well as its ethylenediamine analogue (MRS2892, **8-21**, Fig. 8 and Table 8) were covalently coupled to polyamidoamine (PAMAM) dendrimers. These consist of several bifurcating layers of methyl acrylate and ethylenediamine (Das et al., 2009). A PAMAM conjugate containing four bound UDP-glucuronic acid moieties was 2-fold more potent (EC₅₀ of 159 nM) than UDP-glucuronic acid alone (EC₅₀ of 370 nM). If 20 UDP-glucuronic acid molecules were bound, the PAMAM conjugate was 150-fold more potent (EC₅₀ of 2.4 nM) than UDP-glucuronic acid alone, which corresponds to an 8-fold increase in potency per dendrimer-bound UDP-glucuronic acid moiety compared to monomeric UDP-glucuronic acid. The larger PAMAM conjugate **8-24** (Fig. 8 and Table 8) that contained 147 bound UDP-glucuronic acid moieties was 460-fold more potent (EC₅₀ of 0.8 nM) than UDP-glucuronic acid, which still corresponds to a 3-fold increase in potency per UDP-glucuronic acid moiety.

Molecular docking of a UDP-glucuronic acid dendrimer to a P2Y₁₄R homology model suggested that the dendrimer branches extend far beyond the dimensions of the receptor. They could possibly be available for multivalent binding to receptor aggregates. It was proposed that these multivalent conjugates span several P2Y₁₄R binding sites simultaneously in dimers and other higher-order receptor aggregates (Das et al., 2009).

In an attempt to target heteromeric receptor assemblies, a bifunctional hybrid dendrimer conjugate (MRS5259, **8-29**, Fig. 8 and Table 8) was developed and patented (Jacobson & Tosh, 2010; Tosh et al., 2010). MRS5259 features the selective A₃AR agonist MRS3558 coupled to amide-linked UDP-glucuronic acid *via* a PAMAM dendrimer structure. Both the P2Y₁₄R and the A₃AR are involved in immune functions and possibly expressed on the same cells. Thus, coactivation of these two receptors with one hybrid compound could be advantageous over separate receptor agonists for therapeutic use. Linking UDP-glucuronic acid to the A₃AR agonist in a PAMAM dendrimer did not alter the potency at the P2Y₁₄R in comparison to monofunctional

dendrimer analogues. However, the potency and selectivity at the A₃AR was reduced (Tosh et al., 2010).

Moreover, different prosthetic groups were coupled to dendrimer-nucleotide conjugates to create tools for *in vivo* detection and characterization of the P2Y₁₄R. These include biotin for avidin complexation (**8-25**), the fluorophore Alexa Fluor® 488 (**8-26**), and the metal-chelating group diethylenetriaminepentaacetic acid (DTPA; **8-27** and **8-28**) that might be useful for magnetic resonance imaging (Fig. 8 and Table 8) (Das et al., 2009).

5.1.7. Fluorescence-labelled agonists

SAR studies revealed that UDP-glucuronic acid (**8-03**, Fig. 8 and Table 8) could be linked to larger groups at the C6 carbon atom of the hexose ring without loss in potency (Kiselev et al., 2015). It was therefore used as a precursor for chain extensions to develop fluorescent probes. The diaminoalkyl-linked BODIPY® conjugate MRS4183 (**8-23**, Fig. 8 and Table 8) and an analogue with a shorter linker were first docked into a P2Y₁₄R homology model and subsequently synthesized and assessed (Kiselev et al., 2015; Trujillo, Paoletta, Kiselev, & Jacobson, 2015). According to the model, the pharmacophore binds inside the deep orthosteric binding pocket while the fluorophore is located at the extracellular surface. The pharmacophore and the fluorophore are connected by a linker consisting of 14 atoms in MRS4183. The 7-atom linker of the shorter analogue was insufficient to enable optimal binding of the fluorophore on the outside of the receptor (Kiselev et al., 2015). MRS4183 exhibited an EC₅₀ value of 0.96 nM and was thus 50-fold more potent than UDP-glucuronic acid, while the shorter analogue was 100-fold less active (EC₅₀ of 91 nM). MRS4183 showed high specific binding to the P2Y₁₄R in a flow cytometry assay with an apparent binding constant (K_{d app}) of 21.4 nM and a half-life of 23.9 min for the binding of 50 nM. It was successfully used to detect P2Y₁₄R-expressing Chinese hamster ovary cells using confocal microscopy (Kiselev et al., 2015).

5.1.8. Radioligands

A tritiated form of UDP-glucose, uridine diphospho-D-[6-³H]glucose, was used as a radioligand for the P2Y₁₄R. It showed an affinity of 8.1 nM (Chambers et al., 2000). However, the detected specific binding was essentially identical in non-transfected and P2Y₁₄R-transfected HEK293 cells, 1321N1 astrocytoma cells, and Chinese hamster ovary cells. The compound is thus unsuitable as a radioligand for P2Y₁₄R assays (Chambers et al., 2000; Hamel et al., 2011). In contrast, a radioligand binding assay with [³H]UDP could be established. It was used to characterize UDP-glucose (**8-01**, Fig. 8 and Table 8), UDP (**2-09**, Fig. 2 and Table 2), UMP (**2-10**, Fig. 2 and Table 2), and a non-nucleotide antagonist (**10-22**, Fig. 10 and Table 10) in competition binding assays using HEK cells transiently-transfected with the P2Y₁₄R (Hamel et al., 2011). A K_d value of 16.8 nM was determined for [³H]UDP in saturation binding experiments. [³H]UDP was found to be stable under the assay conditions of radioligand binding for at least one hour, as detected by C18 reverse-phase high-performance liquid chromatography. In contrast, [³H]UTP was rapidly dephosphorylated to UDP. However, in the presence of 25 % human plasma, [³H]UDP was also broken down, yielding uridine as the final product (Hamel et al., 2011).

5.2. P2Y₁₄ receptor antagonists

5.2.1. Therapeutic potential of P2Y₁₄ receptor antagonists

The P2Y₁₄R is prominently expressed in different cells of the immune system. Furthermore, P2Y₁₄R activation was shown to promote chemotaxis and recruitment of neutrophils and macrophages, as well as the release of proinflammatory cytokines, chemokines, and mast cell mediators (Amison et al., 2017; Arase et al., 2009; Barrett et al., 2013; Ferreira et al., 2017; Gao, Ding, & Jacobson, 2010; Gao, Wei, Jayasekara, & Jacobson, 2013; Gendaszewska-Darmach et al., 2016; Jokela et al., 2014; Li et al., 2016; Müller et al., 2005; Sesma et al.,

2012; Sesma et al., 2016; Xu et al., 2012). P2Y₁₄R expression was found to be upregulated in rat brain and spleen following challenge with lipopolysaccharide. It was also upregulated in spinal microglia following peripheral nerve injury, where it contributes to mechanical pain hypersensitivity (Kobayashi, Yamanaka, Yanamoto, Okubo, & Noguchi, 2012; Moore et al., 2003). Nucleotide-sugar conjugates are resistant to hydrolysis by classical ectonucleotidases (such as CD39). As a result, UDP-glucose is found in high concentrations in the extracellular tissue surrounding airway epithelial cells as well as in lung secretions of cystic fibrosis patients. All of these experimental observations indicated that UDP-glucose functions as an extracellular mediator of inflammation by exerting its actions via the P2Y₁₄R (Barrett et al., 2013; Sesma et al., 2016). Antagonists may thus find use in treating excessive inflammatory reactions, including asthma, as had also been postulated for other P2YRs. The concept of diagnosing renal inflammation through measuring the concentration of UDP-glucose in, for example, urine samples and treatment with a P2Y₁₄R antagonist has been patented (Breton, Brown, Azroyan, Cortez-Retamozo, & Pittet, 2014). A reduced resistance to insulin was observed in obese P2Y₁₄R knockout mice. Antagonists could thus be of interest for treating type II diabetes (Xu et al., 2012). Moreover, UDP-sugars were shown to stimulate osteoclastogenesis, and P2Y₁₄R downregulation by RNA interference inhibited osteoclast formation. Antagonists might thus also be useful for treating bone disorders, such as osteoporosis (Lee, Park, & Lee, 2013).

5.2.2. Naphthoic acid derivatives as competitive P2Y₁₄ receptor antagonists

High-throughput screening led to the discovery of 4,7-disubstituted 2-naphthoic acid **10-17** (compound **17** in Fig. 10 and Table 10) to be a weak competitive P2Y₁₄R antagonist (IC₅₀ of 3.5 μ M at both the mouse and chimpanzee P2Y₁₄Rs, K_i of 0.16 μ M at chimpanzee P2Y₁₄R) (Gauthier et al., 2011). In an attempt to optimize its potency and pharmacokinetic properties, different substitutions in the 3-, 4- and 7-position were explored. The most potent compound (**10-16**, Fig. 10 and Table 10) displayed IC₅₀ values of 8 nM and 1 nM at the mouse and chimpanzee P2Y₁₄Rs, respectively (Belley et al., 2008; Gauthier et al., 2011). It showed a bioavailability of 67 % following oral administration and a low intrinsic clearance of 1.6 ml/min/kg in mice (Gauthier et al., 2011). In the presence of 5 % human serum albumin, more than 99 % of the compound was bound, which may have a negative impact on *in vivo* activity (Robichaud et al., 2011). The K_i value at the chimpanzee P2Y₁₄R was reduced to 1.29 μ M in the presence of 2 % human serum albumin while it was 4 nM in the absence of human serum albumin, which corresponds to a 320-fold difference (Gauthier et al., 2011; Robichaud et al., 2011). Consequently, 4-(4-(piperidin-4-yl)-phenyl)-7-(4-(trifluoromethyl)phenyl)-2-naphthoic acid (PPTN, **10-14**, Fig. 10 and Table 10) was developed. It showed less protein binding and K_i values at the chimpanzee P2Y₁₄R of 1.9 nM and 35 nM in the absence and presence of 2 % human serum albumin (Belley et al., 2008; Robichaud et al., 2011). PPTN exhibited a K_B value calculated from Schild analysis of 0.434 nM and a K_i value of 0.3 nM at the human P2Y₁₄R. It was completely inactive at all other P2YR subtypes at a concentration of 1 μ M, and is thus a highly selective P2Y₁₄R antagonist (Barrett et al., 2013; Kiselev et al., 2014). A clear parallel rightward shift towards higher concentrations was observed for concentration-effect curves of UDP-glucose in the presence of increasing concentrations of PPTN, which suggests competitive antagonism. However, the zwitterionic PPTN displayed a very low oral bioavailability. To overcome this issue, an ester prodrug was developed with superior pharmacokinetic properties and efficient conversion to the active drug *in vivo* (Robichaud et al., 2011). The potency of PPTN could be preserved by chain extension at the piperidine moiety, as predicted by homology modeling and confirmed through the synthesis and assessment of a series of chain-elongated alkynyl and amino derivatives of PPTN (Kiselev et al., 2014). The most potent derivative, MRS4174 (**10-18**, Fig. 10 and Table 10), displayed a K_i value of 80 pM, while those of most other compounds in this series were in the low nanomolar range. MRS4174 is an analogue

of PPTN coupled to the fluorophore Alexa Fluor® 488. It is astonishing that the addition of the large fluorophore did not cause a reduction in potency, but in fact led to a 160-fold increase. In contrast, a BODIPY® conjugate was significantly less active (K_i > 100 nM). The potent fluorescent antagonist MRS4174 was employed as a tracer in flow cytometry binding assays, where it displayed high affinity and low non-specific binding (Kiselev et al., 2014). Its synthesis could subsequently be improved to yield sufficient quantity for use in routine assays (Junker et al., 2016). It will be a useful probe for detecting the P2Y₁₄R in different experimental settings and can be an alternative to the fluorescent P2Y₁₄R agonist MRS4183 (**8-23**, Fig. 8 and Table 8) (Kiselev et al., 2015). The fact that both fluorescent probes, MRS4174 (**10-18**) and MRS4183 (**8-23**), were designed using a P2Y₁₄R homology model based on the P2Y₁₂R X-ray structure emphasizes the value of computer modeling for medicinal chemistry, especially after the publication of several P2YR X-ray structures (Zhang et al., 2015; Zhang, Zhang, Gao, Paoletta, et al., 2014; Zhang, Zhang, Gao, Zhang, et al., 2014).

5.2.3. Analogues of PPTN

The hydrophobic naphthalene ring of PPTN (**10-14**, Fig. 10 and Table 10) is responsible for the low solubility of the compound. It was thus attempted to replace the naphthalene structure with a bioisostere to improve its physicochemical properties (Junker et al., 2016). Docking studies and molecular dynamics simulations on a P2Y₁₄R homology model led to the design of alkyne and triazole derivatives. The alkyne derivative **10-20** (Fig. 10 and Table 10; IC₅₀ value of 5.69 μ M) was 950-fold less potent than PPTN. The corresponding triazole derivative **10-19** (Fig. 10 and Table 10) was more potent (IC₅₀ of 32 nM, 5-fold less potent than PPTN) and therefore used as a lead structure for further optimization. However, an improvement in potency has not been achieved; the IC₅₀ values were in between 72 nM and 481 nM (Junker et al., 2016). Despite the 5-fold lower potency of **10-19**, it may nevertheless be preferably used in biological experiments if its physicochemical properties are indeed superior. No data with regard to this have been published so far, but computer calculations predicted a higher solubility and lower plasma protein binding (Junker et al., 2016).

PPTN and **10-19** were assessed in the Psychoactive Drug Screening Program for off-target activities (Besnard et al., 2012). PPTN displayed micromolar K_i values at the dopamine D₃ and the δ -opioid receptors, and **10-19** showed affinity for the α_{2A} and α_{2C} adrenergic receptors. A nanomolar K_i value (170 nM) was observed for **10-19** only at the H₁ histamine receptor. PPTN and **10-19** were inactive at all other receptors, ion channels, and transporters assessed in the standard diverse screen of the Psychoactive Drug Screening Program (Besnard et al., 2012; Junker et al., 2016). Thus, they showed only few off-target interactions (Junker et al., 2016).

5.2.4. Dihydropyridopyrimidines – non-competitive P2Y₁₄ receptor antagonists

High-throughput screening at Merck Frosst (Canada) led to the discovery of another non-nucleotide P2Y₁₄R antagonist with a dihydropyridopyrimidine core. It showed micromolar potency at the mouse P2Y₁₄R (Guay et al., 2011). This hit compound formed the basis for comprehensive SAR studies that resulted in several derivatives with potencies in the nano- to micromolar range at the mouse and chimpanzee receptors (Guay et al., 2008). The 3,4-methylenedioxyphenyl derivative **10-21** (Fig. 10 and Table 10) was among the most promising antagonists, exhibiting IC₅₀ values of 10 nM and 81 nM on the mouse and chimpanzee P2Y₁₄Rs, respectively (Guay et al., 2008; Guay et al., 2011). It was found to be readily bioavailable following oral administration and provided sufficient exposure in mice to be a suitable tool for *in vivo* studies. The compound did not displace [³H]UDP in filtration binding assays and was thus proposed to be a non-competitive antagonist. Although several analogues showed slightly higher potencies, they were exempted from further

considerations due to binding to the human Ether-à-go-go-Related Gene (hERG) channel that may cause serious side-effects (Guay et al., 2011).

5.2.5. Phosphonates

One of the first non-nucleotide P2Y₁₄R antagonists, **10-22** (Fig. 10 and Table 10), was discovered through the screening of a compound library that consisted of 608 phosphonates (Hamel et al., 2011). Compound **10-22** exhibited an IC₅₀ value of 2.3 μM in calcium mobilization assays and a K_i value of 1.28 μM in competition binding assays with [³H]UDP. It was suggested that the activity of **10-22** could be improved by altering its structure in a manner to more closely resemble a diphosphate compound, but no such attempts have been reported so far (Hamel et al., 2011).

6. Conclusions

6.1. Agonists

The P2YR family of GPCRs is of considerable therapeutic interest. Consequently, extensive research efforts have been directed towards the design of ligands suitable for target validation studies and drug development. A relatively large number of agonists has been developed in the past years as a result, and the most useful ones with respect to potency, selectivity, and physicochemical properties are collected in Fig. 11.

Among the most potent and selective P2Y₂R agonists is 2'-amino-2'-deoxy-2-thio-UTP (MRS2698, **6-01** compound **01** in Fig. 6 and Table 6). It shows an EC₅₀ value in the low nanomolar range, is 6-fold more potent than the endogenous agonist UTP, and is at least 300-fold selective versus the P2Y₄R and P2Y₆R. Highly selective (>6600-fold) P2Y₂R agonists were also developed by replacing the nucleobase of nucleotide

triphosphates with 'unnatural' bicyclic aromatic residues (e.g. **3-35**). Owing to its enhanced ectonucleotidase stability, 4-thio-β,γ-difluoromethylene-UTP (PSB-1114, **6-05**) is of great interest as well, despite its 8-fold lower potency than UTP. Dinucleotides are also more resistant to enzymatic hydrolysis as compared to mononucleotides, but are generally less potent (except at the P2Y₆R), and lack selectivity in many cases. Eye drops (Diquas®) containing Up₄U (3 % solution of the tetrasodium salt, diquafosol, INS365, **7-03**) was approved in Japan and other Asian countries as a treatment for dry eye syndrome. All of these compounds except **3-35** can be obtained from commercial distributors. Very recently, a 4(1*H*)-quinoline derivative (**9-01**) was described as a non-nucleotide ago-allosteric agonist with micromolar potency at the P2Y₂R and no detectable response at most other P2YRs. It may serve as a lead structure for the development of more potent non-nucleotide P2Y₂R agonists.

For the P2Y₄R, fewer ligands have been described so far. Among the most potent and selective agonists is the commercially available N⁴-(phenylpropoxy)-CTP (MRS4062, **3-15**), which is twice as potent as the endogenous agonist UTP and 27-fold selective.

The development of agonists for the P2Y₆R has been far more successful and yielded several potent and highly selective compounds. The R_p isomer of 5-MeO-uridine 5'-O-(α-boranodiphosphate) (**6-12**) is one of the most potent (EC₅₀ of 8 nM) and selective (>12,000-fold) P2Y₆R agonists known to date. It was reported to be relatively stable towards degradation by ectonucleotidases. Very recently, prostaglandin E₂ glyceryl ester (**9-02**) was described as an extraordinarily potent (EC₅₀ in the low picomolar range) physiological P2Y₆R agonist. Should this be confirmed by other laboratories, it will enable new possibilities for compound/drug design, which, however, will be challenging due to its instability under physiological conditions and its lipid character. Other useful P2Y₆R agonists include 3-phenacyl-UDP (PSB-0474, **3-11**; 570-fold selective), N⁴-(benzyloxy)-CDP (MRS2964, **3-17**; 44-fold




Fig. 11. The most potent and selective agonists for the human (A) P2Y₂R, (B) P2Y₄R, (C) P2Y₆R, and (D) P2Y₁₄R. No data is available in cases where no bar chart is indicated.

selective), 5-MeO-UDP (**3-31**), 5-iodo-UDP (MRS2693, **3-26**), (S)-methanocarba-UDP (MRS2795, **4-22**), as well as the dinucleotides monobenzylacetal-Up₃U (INS48823, **7-11**) and N⁴-MeO-Cp₃U (MRS2957, **7-08**). The selective partial agonist TMPS (**5-08**) may also be of value.

The pharmacological profile of the P2Y₁₄R is more distinct. Unlike the other three uracil nucleotide-activated receptors, it is potently activated by UDP-sugars. UDP (**2-09**) is another physiological agonist of the P2Y₁₄R, overlapping with the P2Y₆R. Thus, a challenge was to confer selectivity for the P2Y₁₄R over the P2Y₆R. A 2-thio substitution of the uracil base was highly favorable for the P2Y₁₄R. As a result, α,β-methylene-2-thio-UDP (MRS2905, **6-08**), 2-thio-UDP-glucose (MRS2690, **8-15**), and the 2-thio-UDP-β-esters **6-17**, **6-18**, and **6-21** are highly potent and selective P2Y₁₄R agonists. They were inactive at the P2Y₆R. In addition, 2-thio-UDP (**3-02**) was found to be 230-fold selective for the P2Y₁₄R but could activate the P2Y₆R at high nanomolar concentrations. Alternatively, α,β-difluoromethylene-UDP (MRS2802, **5-15**) showed >160-fold selectivity for the P2Y₁₄R over the P2Y₆R. PAMAM dendrimers add further diversity to the selection of P2Y₁₄R tools. They were connected to the P2Y₁₄R agonist UDP-glucuronic acid (**8-03**) in addition to different prosthetic groups, such as an A₃AR agonist, a fluorophore, biotin for avidin complexation, and the metal chelator DTPA for use in magnetic resonance imaging. The fluorescent BODIPY® conjugate MRS4183 (**8-23**) and the radioligand [³H]UDP enable further characterization of this receptor. Despite being the most recently described P2YR subtype, ligand development efforts for the P2Y₁₄R have been most successful.

In addition to the compounds described above, the fluorescent probes MRS4129 (**3-42**; selective for the P2Y₆R) and MRS4162 (**3-40**; potent agonist at the P2Y₂R, P2Y₄R, and P2Y₆R) serve as practical tools for detecting and quantifying these receptors in living cells.

6.2. Antagonists

The availability of antagonists for the four uracil nucleotide-activated P2YR subtypes is more limited. The thiouracil derivative AR-C118925 (**10-01**, compound **01** in Fig. 10 and Table 10; 50-fold selectivity) for the P2Y₂R, and the anthraquinone derivative PSB-16133 (**10-06**) as 37-fold selective P2Y₄R antagonist, are superior to the non-selective, moderately-potent standard antagonists suramin (**10-04**), RB-2 (**10-07**), and PPADS (**10-12**) for these two receptors. Among the most frequently used P2Y₆R antagonists is the non-competitive, insurmountable diisothiocyanate derivative MRS2578 (**10-08**) with mid-nanomolar potency and at least 270-fold selectivity. However, the hydrophobic structure of MRS2578 and its reactivity towards nucleophiles and its instability in aqueous media constitute severe limitations. MRS2578 binds covalently, and thus irreversibly, to

the P2Y₆R. The 2-naphthoic acid derivative PPTN (**10-14**) is a competitive P2Y₁₄R antagonist with (sub-)nanomolar potency and 2300-fold selectivity. However, it showed low aqueous solubility and poor oral bioavailability. An ester prodrug (**10-15**) was subsequently developed for *in vivo* studies. Several potent derivatives were also synthesized, including **10-16**, **10-19**, and the fluorescent P2Y₁₄R antagonist MRS4174 (**10-18**) that exhibited a sub-nanomolar *K_i* value at the P2Y₁₄R. Moreover, dihydropyridopyrimidines were reported to function as non-competitive P2Y₁₄R antagonists, with **10-21** showing nanomolar potency and high oral bioavailability. A summary of the most potent and selective antagonists is shown in Fig. 12.

6.3. Outlook

Several agonists showing potencies in the low nanomolar range and good selectivity profiles could be developed for the P2Y₂R, P2Y₆R, and P2Y₁₄R, while agonists for the P2Y₄R remain scarce. With very few exceptions, the described agonists are all nucleotide derivatives. These have generally poor oral bioavailability due to the negative charges at physiological pH, lack brain penetration, and may be subject to degradation by ectonucleotidases, although the stability could be improved for some synthetic analogues. Only few antagonists have been reported. Although most of them are not derived from nucleotides, they still have disadvantages in many cases, such as a low water solubility, high plasma protein binding, or insufficient selectivity. Compounds with higher potency in the low nanomolar range are required for the development of radioligands that would advance further receptor characterization. In conclusion, future efforts are necessary towards the development of (non-nucleotide) agonists and improved antagonists for the uracil nucleotide-activated P2YRs, and the P2Y₄R in particular.

Conflict of Interest

The authors declare that there are no conflicts of interest.

References

- Abbracchio, M. P., & Burnstock, G. (1994). Purinoceptors: Are there families of P2X and P2Y purinoceptors? *Pharmacology & Therapeutics* 64, 445–475.
- Abbracchio, M. P., Burnstock, G., Boeynaems, J. -M., Barnard, E. A., Boyer, J. L., Kennedy, C., ... Weisman, G. A. (2006). International Union of Pharmacology LVIII: Update on the P2Y G protein-coupled nucleotide receptors: From molecular mechanisms and pathophysiology to therapy. *Pharmacological Reviews* 58, 281–341.
- Accurso, F. J., Moss, R. B., Wilmott, R. W., Anbar, R. D., Schaberg, A. E., Durham, T. A., & Ramsey, B. W. (2011). Denufisol tetrasodium in patients with cystic fibrosis and normal to mildly impaired lung function. *American Journal of Respiratory and Critical Care Medicine* 183, 627–634.
- Agca, Y., Qian, S., Agca, C., & Seye, C. I. (2016). Direct evidence for P2Y₂ receptor involvement in vascular response to injury. *Journal of Vascular Research* 53, 163–171.
- Ajit, D., Woods, L. T., Camden, J. M., Thebeau, C. N., El-Sayed, F. G., Greeson, G. W., ... Weisman, G. A. (2014). Loss of P2Y(2) nucleotide receptors enhances early pathology in the TgCRND8 mouse model of Alzheimer's disease. *Molecular Neurobiology* 49, 1031–1042.
- Akbar, G. K., Dasari, V. R., Webb, T. E., Ayyanathan, K., Pillarisetti, K., Sandhu, A. K., ... Kunapuli, S. P. (1996). Molecular cloning of a novel P2 purinoceptor from human erythroleukemia cells. *The Journal of Biological Chemistry* 271, 18363–18367.
- Amison, R. T., Arnold, S., O'Shaughnessy, B. G., Cleary, S. J., Ofoedu, J., Idzko, M., ... Pitchford, S. C. (2017). Lipopolysaccharide (LPS) induced pulmonary neutrophil recruitment and platelet activation is mediated via the P2Y₁ and P2Y₁₄ receptors in mice. *Pulmonary Pharmacology & Therapeutics* 45, 62–68.
- Arase, T., Uchida, H., Kajitani, T., Ono, M., Tamaki, K., Oda, H., ... Maruyama, T. (2009). The UDP-glucose receptor P2RY₁₄ triggers innate mucosal immunity in the female reproductive tract by inducing IL-8. *Journal of Immunology* 182, 7074–7084.
- Aschrafi, A., Sadler, S., Niculescu, C., Rettinger, J., & Schmalzing, G. (2004). Trimeric architecture of homomeric P2X2 and heteromeric P2X1 + 2 receptor subtypes. *Journal of Molecular Biology* 342, 333–343.
- Ault, A. D., & Broach, J. R. (2006). Creation of GPCR-based chemical sensors by directed evolution in yeast. *Protein Engineering, Design & Selection* 19, 1–8.
- Azimi, I., Beilby, H., Davis, F. M., Marcial, D. L., Kenny, P. A., Thompson, E. W., ... Monteith, G. R. (2016). Altered purinergic receptor-Ca²⁺ signaling associated with hypoxia-induced epithelial-mesenchymal transition in breast cancer cells. *Molecular Oncology* 10, 166–178.




Fig. 12. The most potent and selective P2YR antagonists. No data is available in cases where no bar chart is indicated.

- Azroyan, A., Cortez-Retamozo, V., Bouley, R., Liberman, R., Ruan, Y. C., Kiselev, E., ... Breton, S. (2015). Renal intercalated cells sense and mediate inflammation via the P2Y₁₄ receptor. *PLoS One* 10, e0121419.
- Bagchi, S., Liao, Z., Gonzalez, F. A., Chorna, N. E., Seye, C. I., Weisman, G. A., & Erb, L. (2005). The P2Y₂ nucleotide receptor interacts with α 5 integrins to activate G α and induce cell migration. *The Journal of Biological Chemistry* 280, 39050–39057.
- Balasubramanian, R., Ruiz de Azua, I., Wess, J., & Jacobson, K. A. (2010). Activation of distinct P2Y receptor subtypes stimulates insulin secretion in MIN6 mouse pancreatic beta cells. *Biochemical Pharmacology* 79, 1317–1326.
- Baltensperger, K., & Porzig, H. (1997). The P2U purinoceptor obligatorily engages the heterotrimeric G protein G16 to mobilize intracellular Ca²⁺ in human erythroleukemia cells. *The Journal of Biological Chemistry* 272, 10151–10159.
- Baqi, Y., Atzler, K., Köse, M., Glänzel, M., & Müller, C. E. (2009). High-affinity, non-nucleotide-derived competitive antagonists of platelet P2Y₁₂ receptors. *Journal of Medicinal Chemistry* 52, 3784–3793.
- Baqi, Y., Hausmann, R., Rosefort, C., Rettinger, J., Schmalzing, G., & Müller, C. E. (2011). Discovery of potent competitive antagonists and positive modulators of the P2X₂ receptor. *Journal of Medicinal Chemistry* 54, 817–830.
- Baqi, Y., Lee, S. -Y., Iqbal, J., Ripphausen, P., Lehr, A., Scheiff, A. B., ... Müller, C. E. (2010). Development of potent and selective inhibitors of ecto-5'-nucleotidase based on an anthraquinone scaffold. *Journal of Medicinal Chemistry* 53, 2076–2086.
- Bar, I., Guns, P. -J., Metallo, J., Cammarata, D., Wilkin, F., Boeynaems, J. -M., ... Robaye, B. (2008). Knockout mice reveal a role for P2Y₆ receptor in macrophages, endothelial cells, and vascular smooth muscle cells. *Molecular Pharmacology* 74, 777–784.
- Barragán-Iglesias, P., Mendoza-Garcés, L., Pineda-Farias, J. B., Solano-Olivares, V., Rodríguez-Silverio, J., Flores-Murrieta, F. J., ... Rocha-González, H. I. (2015). Participation of peripheral P2Y₁, P2Y₆ and P2Y₁₁ receptors in formalin-induced inflammatory pain in rats. *Pharmacology, Biochemistry, and Behavior* 128, 23–32.
- Barrett, M. O., Sesma, J. I., Ball, C. B., Jayasekara, P. S., Jacobson, K. A., Lazarowski, E. R., & Harden, T. K. (2013). A selective high-affinity antagonist of the P2Y₁₄ receptor inhibits UDP-glucose-stimulated chemotaxis of human neutrophils. *Molecular Pharmacology* 84, 41–49.
- Belley, M., Deschenes, D., Fortin, R., Fournier, J.-F., Gagné, S., Gareau, Y., ... Wang, Z. (2008). Merck Frosst Canada Ltd. WO2009070873A1.
- Bender, E., Buist, A., Jurzak, M., Langlois, X., Baggerman, G., Verhasselt, P., ... Luyten, W. (2002). Characterization of an orphan G protein-coupled receptor localized in the dorsal root ganglia reveals adenine as a signaling molecule. *Proceedings of the National Academy of Sciences of the United States of America* 99, 8573–8578.
- Besada, P., Shin, D. H., Costanzi, S., Ko, H., Mathe, C., Gagneron, J., ... Jacobson, K. A. (2006). Structure-activity relationships of uridine 5'-diphosphate analogues at the human P2Y₆ receptor. *Journal of Medicinal Chemistry* 49, 5532–5543.
- Besnard, J., Ruda, G. F., Setola, V., Abecassis, K., Rodriguiz, R. M., Huang, X. -P., ... Hopkins, A. L. (2012). Automated design of ligands to polypharmacological profiles. *Nature* 492, 215–220.
- Bogdanov, Y. D., Dale, L., King, B. F., Whittlock, N., & Burnstock, G. (1997). Early expression of a novel nucleotide receptor in the neural plate of *Xenopus* embryos. *The Journal of Biological Chemistry* 272, 12583–12590.
- Bogdanov, Y. D., Wildman, S. S., Clements, M. P., King, B. F., & Burnstock, G. (1998). Molecular cloning and characterization of rat P2Y₄ nucleotide receptor. *British Journal of Pharmacology* 124, 428–430.
- Boyer, J. L., Shaver, S. R., Douglass, J. G. III, & Redick, C. C. (2006). Inspire Pharmaceuticals Inc. WO002007002945A2.
- Breton, S., Brown, D., Azroyan, A., Cortez-Retamozo, V., & Pittet, M. J. (2014). General Hospital Corp. WO002015070001A2.
- Brookings, D., Davenport, R. J., Davis, J., Galvin, F. C. A., Lloyd, S., Mack, S. R., ... Wynn, J. (2007). Novel nucleotide triphosphates as potent P2Y₂ agonists. *Bioorganic & Medicinal Chemistry Letters* 17, 562–565.
- Brown, C., Tanna, B., & Boarder, M. R. (1995). PPADS: An antagonist at endothelial P2Y-purinoceptors but not P2U-purinoceptors. *British Journal of Pharmacology* 116, 2413–2416.
- Brunschweiler, A., & Müller, C. E. (2006). P2 receptors activated by uracil nucleotides—An update. *Current Medicinal Chemistry* 13, 289–312.
- Brüser, A., Zimmermann, A., Crews, B. C., Sliwoski, G., Meiler, J., König, G. M., ... Schöneberg, T. (2017). Prostaglandin E₂ glyceryl ester is an endogenous agonist of the nucleotide receptor P2Y₆. *Scientific Reports* 7, 2380.
- Burnstock, G. (1976). Do some nerve cells release more than one transmitter? *Neuroscience* 1, 239–248.
- Burnstock, G., & Knight, G. E. (2017). Cell culture: Complications due to mechanical release of ATP and activation of purinoceptors. *Cell and Tissue Research* 370, 1–11.
- Camden, J. M., Schrader, A. M., Camden, R. E., Gonzalez, F. A., Erb, L., Seye, C. I., & Weisman, G. A. (2005). P2Y₂ nucleotide receptors enhance alpha-secretase-dependent amyloid precursor protein processing. *The Journal of Biological Chemistry* 280, 18696–18702.
- Carter, R. L., Fricks, I. P., Barrett, M. O., Burianek, L. E., Zhou, Y., Ko, H., ... Harden, T. K. (2009). Quantification of Gi-mediated inhibition of adenylyl cyclase activity reveals that UDP is a potent agonist of the human P2Y₁₄ receptor. *Molecular Pharmacology* 76, 1341–1348.
- Cavaliere, F., Amadio, S., Angelini, D. F., Sanesario, G., Bernardi, G., & Volonté, C. (2004). Role of the metabotropic P2Y(4) receptor during hypoglycemia: Cross talk with the ionotropic NMDAR1 receptor. *Experimental Cell Research* 300, 149–158.
- Chambers, J. K., Macdonald, L. E., Sarau, H. M., Ames, R. S., Freeman, K., Foley, J. J., ... Livi, G. P. (2000). A G protein-coupled receptor for UDP-glucose. *The Journal of Biological Chemistry* 275, 10767–10771.
- Chang, K., Hanaoka, K., Kumada, M., & Takuwa, Y. (1995). Molecular cloning and functional analysis of a novel P2 nucleotide receptor. *The Journal of Biological Chemistry* 270, 26152–26158.
- Charlton, M. E., Williams, A. S., Fogliano, M., Sweetnam, P. M., & Duman, R. S. (1997). The isolation and characterization of a novel G protein-coupled receptor regulated by immunologic challenge. *Brain Research* 764, 141–148.
- Charlton, S. J., Brown, C. A., Weisman, G. A., Turner, J. T., Erb, L., & Boarder, M. R. (1996a). PPADS and suramin as antagonists at cloned P2Y- and P2U-purinoceptors. *British Journal of Pharmacology* 118, 704–710.
- Charlton, S. J., Brown, C. A., Weisman, G. A., Turner, J. T., Erb, L., & Boarder, M. R. (1996b). Cloned and transfected P2Y4 receptors: Characterization of a suramin and PPADS-insensitive response to UTP. *British Journal of Pharmacology* 119, 1301–1303.
- Chen, X., Qian, S., Hoggatt, A., Tang, H., Hacker, T. A., Obukhov, A. G., ... Seye, C. I. (2017). Endothelial cell-specific deletion of P2Y2 receptor promotes plaque stability in atherosclerosis-susceptible ApoE-null mice. *Arteriosclerosis, Thrombosis, and Vascular Biology* 37, 75–83.
- Chen, Y., Corriden, R., Inoue, Y., Yip, L., Hashiguchi, N., Zinkernagel, A., ... Junger, W. G. (2006). ATP release guides neutrophil chemotaxis via P2Y₂ and A₃ receptors. *Science* 314, 1792–1795.
- Chen, Z. P., Krull, N., Xu, S., Levy, A., & Lightman, S. L. (1996). Molecular cloning and functional characterization of a rat pituitary G protein-coupled adenosine triphosphate (ATP) receptor. *Endocrinology* 137, 1833–1840.
- Christopoulos, A., Changeux, J. -P., Catterall, W. A., Fabbro, D., Burris, T. P., Cidlowski, J. A., ... Langmead, C. J. (2014). International union of basic and clinical pharmacology. XC. multisite pharmacology: Recommendations for the nomenclature of receptor allosterism and allosteric ligands. *Pharmacological Reviews* 66, 918–947.
- Ciruela, F., Fernández-Duenas, V., & Jacobson, K. A. (2015). Lighting up G protein-coupled purinergic receptors with engineered fluorescent ligands. *Neuropharmacology* 98, 58–67.
- Clouet, S., Di Pietrantonio, L., Daskalopoulos, E. -P., Esfahani, H., Horckmans, M., Vanorle, M., ... Communi, D. (2016). Loss of mouse P2Y₆ nucleotide receptor is associated with physiological macrocardia and amplified pathological cardiac hypertrophy. *The Journal of Biological Chemistry* 291, 15841–15852.
- Coddou, C., Yan, Z., Obsil, T., Huidobro-Toro, J. P., & Stojilkovic, S. S. (2011). Activation and regulation of purinergic P2X receptor channels. *Pharmacological Reviews* 63, 641–683.
- Cohen, R., Shainberg, A., Hochhauser, E., Cheporko, Y., Tobar, A., Birk, E., ... Porat, E. (2011). UTP reduces infarct size and improves late heart function after myocardial infarct via P2Y₂ receptor. *Biochemical Pharmacology* 82, 1126–1133.
- Communi, D., Motte, S., Boeynaems, J. M., & Pirotton, S. (1996). Pharmacological characterization of the human P2Y₄ receptor. *European Journal of Pharmacology* 317, 383–389.
- Communi, D., Parmentier, M., & Boeynaems, J. M. (1996). Cloning, functional expression and tissue distribution of the human P2Y₆ receptor. *Biochemical and Biophysical Research Communications* 222, 303–308.
- Communi, D., Pirotton, S., Parmentier, M., & Boeynaems, J. M. (1995). Cloning and functional expression of a human uridine nucleotide receptor. *The Journal of Biological Chemistry* 270, 30849–30852.
- Conroy, S., Kindon, N., Kellam, B., & Stocks, M. J. (2016). Drug-like antagonists of P2Y receptors from lead identification to drug development. *Journal of Medicinal Chemistry* 59, 9981–10005.
- Cosentino, S., Banfi, C., Burbiel, J. C., Luo, H., Tremoli, E., & Abbraccio, M. P. (2012). Cardiac myocyte death induced by ischaemic/hypoxic stress is differentially affected by distinct purinergic P2 receptors. *Journal of Cellular and Molecular Medicine* 16, 1074–1084.
- Costanzi, S., Joshi, B. V., Maddileti, S., Mamedova, L., Gonzalez-Moa, M. J., Marquez, V. E., ... Jacobson, K. A. (2005). Human P2Y(6) receptor: Molecular modeling leads to the rational design of a novel agonist based on a unique conformational preference. *Journal of Medicinal Chemistry* 48, 8108–8111.
- Costanzi, S., Mamedova, L., Gao, Z. -G., & Jacobson, K. A. (2004). Architecture of P2Y nucleotide receptors: Structural comparison based on sequence analysis, mutagenesis, and homology modeling. *Journal of Medicinal Chemistry* 47, 5393–5404.
- Cosyn, L., van Calenbergh, S., Joshi, B. V., Ko, H., Carter, R. L., Kendall Harden, T., & Jacobson, K. A. (2009). Synthesis and P2Y receptor activity of nucleoside 5'-phosphonate derivatives. *Bioorganic & Medicinal Chemistry Letters* 19, 3002–3005.
- Cowlen, M., Yexxa, B. R., Jones, A. C., & Brown, E. G. (2002). Inspire Pharmaceuticals Inc. WO002003000056A1.
- Cox, M. A., Gomes, B., Palmer, K., Du, K., Wiekowski, M., Wilburn, B., ... Jenh, C. -H. (2005). The pyrimidinergic P2Y₆ receptor mediates a novel release of proinflammatory cytokines and chemokines in monocytic cells stimulated with UDP. *Biochemical and Biophysical Research Communications* 330, 467–473.
- Cressman, V. L., Lazarowski, E., Homolya, L., Boucher, R. C., Koller, B. H., & Grubb, B. R. (1999). Effect of loss of P2Y(2) receptor gene expression on nucleotide regulation of murine epithelial Cl(-) transport. *The Journal of Biological Chemistry* 274, 26461–26468.
- Dale, C. L., Hill, S. J., & Kellam, B. (2012). New potent, short-linker BODIPY-630/650™ labelled fluorescent adenosine receptor agonists. *Medicinal Chemistry Communications* 3, 333–338.
- Das, A., Ko, H., Burianek, L. E., Barrett, M. O., Harden, T. K., & Jacobson, K. A. (2010). Human P2Y(14) receptor agonists: Truncation of the hexose moiety of uridine 5'-diphosphoglucose and its replacement with alkyl and aryl groups. *Journal of Medicinal Chemistry* 53, 471–480.
- Das, A., Zhou, Y., Ivanov, A. A., Carter, R. L., Harden, T. K., & Jacobson, K. A. (2009). Enhanced potency of nucleotide-dendrimer conjugates as agonists of the P2Y₁₄ receptor: Multivalent effect in G protein-coupled receptor recognition. *Bioconjugate Chemistry* 20, 1650–1659.
- Davenport, R. J., Diaz, P., Galvin, F. C. A., Lloyd, S., Mack, S. R., Owens, R., ... Wynn, J. (2007). Novel nucleotide triphosphates as potent P2Y₂ agonists with enhanced stability over UTP. *Bioorganic & Medicinal Chemistry Letters* 17, 558–561.
- Davis, J. M., Mack, S. R., Sabin, V. M., & Davenport, R. J. (2002). Cell-tech R&D Ltd. WO002002062816A1.

- Delbro, D. S., Nylund, G., & Nordgren, S. (2005). Demonstration of P2Y₄ purinergic receptors in the HT-29 human colon cancer cell line. *Autonomic & Autacoid Pharmacology* 25, 163–166.
- Deterding, R. R., Lavange, L. M., Engels, J. M., Mathews, D. W., Coquillotte, S. J., Brody, A. S., ... Ramsey, B. W. (2007). Phase 2 randomized safety and efficacy trial of nebulized denufosol tetrasodium in cystic fibrosis. *American Journal of Respiratory and Critical Care Medicine* 176, 362–369.
- Diego-García, L. d., Ramírez-Escudero, M., Sebastián-Serrano, Á., Díaz-Hernández, J. I., Pintor, J., Lucas, J. J., & Díaz-Hernández, M. (2017). Regulation of proteasome activity by P2Y₂ receptor underlies the neuroprotective effects of extracellular nucleotides. *Biochimica et Biophysica Acta* 1863, 43–51.
- Dissmore, T., Seye, C. I., Medeiros, D. M., Weisman, G. A., Bardford, B., & Mamedova, L. (2016). The P2Y₂ receptor mediates uptake of matrix-retained and aggregated low density lipoprotein in primary vascular smooth muscle cells. *Atherosclerosis* 252, 128–135.
- Dixon, C. J., Bowler, W. B., Littlewood-Evans, A., Dillon, J. P., Bilbe, G., Sharpe, G. R., & Gallagher, J. A. (1999). Regulation of epidermal homeostasis through P2Y₂ receptors. *British Journal of Pharmacology* 127, 1680–1686.
- Drury, A. N., & Szent-Györgyi, A. (1929). The physiological activity of adenine compounds with especial reference to their action upon the mammalian heart. *The Journal of Physiology* 68, 213–237.
- Drutz, D. J., Rideout, J. L., & Jacobus, K. M. (1997). Inspire Pharmaceuticals Inc. DE000069716255T2.
- DuBose, D. R., Wolff, S. C., Qi, A., -D., Naruszewicz, I., & Nicholas, R. A. (2013). Apical targeting of the P2Y₄ receptor is directed by hydrophobic and basic residues in the cytoplasmic tail. *American Journal of Physiology. Cell Physiology* 304, C228–39.
- Dulong, S., Bernard, K., & Ehrenfeld, J. (2007). Enhancement of P2Y₆-induced Cl⁻ secretion by IL-13 and modulation of SK4 channels activity in human bronchial cells. *Cellular Physiology and Biochemistry: International Journal of Experimental Cellular Physiology, Biochemistry, and Pharmacology* 20, 483–494.
- Ecke, D., Hanck, T., Tulapurkar, M. E., Schafer, R., Kassack, M., Stricker, R., & Reiser, G. (2008). Hetero-oligomerization of the P2Y₁₁ receptor with the P2Y₁ receptor controls the internalization and ligand selectivity of the P2Y₁₁ receptor. *The Biochemical Journal* 409, 107–116.
- Eliahu, S. E., Camden, J., Lecka, J., Weisman, G. A., Sévigny, J., Gélinas, S., & Fischer, B. (2009). Identification of hydrolytically stable and selective P2Y₁ receptor agonists. *European Journal of Medicinal Chemistry* 54, 1525–1536.
- Elliott, M. R., Chekeni, F. B., Trampont, P. C., Lazarowski, E. R., Kadl, A., Walk, S. F., ... Ravichandran, K. S. (2009). Nucleotides released by apoptotic cells act as a find-me signal to promote phagocytic clearance. *Nature* 461, 282–286.
- El-Tayeb, A., Qi, A., & Müller, C. E. (2006). Synthesis and structure-activity relationships of uracil nucleotide derivatives and analogues as agonists at human P2Y₂, P2Y₄, and P2Y₆ receptors. *Journal of Medicinal Chemistry* 49, 7076–7087.
- El-Tayeb, A., Qi, A., Nicholas, R. A., & Müller, C. E. (2011). Structural modifications of UMP, UDP, and UTP leading to subtype-selective agonists for P2Y₂, P2Y₄, and P2Y₆ receptors. *Journal of Medicinal Chemistry* 54, 2878–2890.
- Erb, L., Lustig, K. D., Sullivan, D. M., Turner, J. T., & Weisman, G. A. (1993). Functional expression and photoaffinity labeling of a cloned P_{2U} purinergic receptor. *Proceedings of the National Academy of Sciences of the United States of America* 90, 10449–10453.
- Ferreira, M. A. R., Jansen, R., Willemssen, G., Penninx, B., Bain, L. M., Vicente, C. T., ... Phipps, S. (2017). Gene-based analysis of regulatory variants identifies 4 putative novel asthma risk genes related to nucleotide synthesis and signaling. *The Journal of Allergy and Clinical Immunology* 139, 1148–1157.
- Filipov, A. K., Webb, T. E., Barnard, E. A., & Brown, D. A. (1997). Inhibition by heterologously-expressed P2Y₂ nucleotide receptors of N-type calcium currents in rat sympathetic neurones. *British Journal of Pharmacology* 121, 849–851.
- Filipov, A. K., Webb, T. E., Barnard, E. A., & Brown, D. A. (1998). P2Y₂ nucleotide receptors expressed heterologously in sympathetic neurons inhibit both N-type Ca²⁺ and M-type K⁺ currents. *The Journal of Neuroscience* 18, 5170–5179.
- Filipov, A. K., Webb, T. E., Barnard, E. A., & Brown, D. A. (1999). Dual coupling of heterologously-expressed rat P2Y₆ nucleotide receptors to N-type Ca²⁺ and M-type K⁺ currents in rat sympathetic neurones. *British Journal of Pharmacology* 126, 1009–1017.
- Fischer, B., & Elyahu, S. (2008). Bar-Ilan University WO002009066298A1.
- Fischer, B., & Nahum, V. (2006). Bar-Ilan University US020060287271A1.
- Fischer, B., Pintor, J. J., Elyahu, S., & Ginsburg-Shmuel, T. (2011). Bar-Ilan University; Universidad Complutense de Madrid WO002012073237A1.
- Fischer, B., Sevigny, J., Elyahu, S., & Lecka, J. (2011). Bar-Ilan University WO002012032513A1.
- Freeman, K., Tsui, P., Moore, D., Emson, P. C., Vawter, L., Naheed, S., ... Chambers, J. K. (2001). Cloning, pharmacology, and tissue distribution of G-protein-coupled receptor GPR105 (KIAA0001) rodent orthologs. *Genomics* 78, 124–128.
- Fricks, I. P., Carter, R. L., Lazarowski, E. R., & Harden, T. K. (2009). G_i-dependent cell signaling responses of the human P2Y₁₄ receptor in model cell systems. *The Journal of Pharmacology and Experimental Therapeutics* 330, 162–168.
- Fricks, I. P., Maddileti, S., Carter, R. L., Lazarowski, E. R., Nicholas, R. A., Jacobson, K. A., & Harden, T. K. (2008). UDP is a competitive antagonist at the human P2Y₁₄ receptor. *The Journal of Pharmacology and Experimental Therapeutics* 325, 588–594.
- Fujioka, M., & Omori, N. (2012). Subtleties in GPCR drug discovery: A medicinal chemistry perspective. *Drug Discovery Today* 17, 1133–1138.
- Gao, Z., -G., Ding, Y., & Jacobson, K. A. (2010). UDP-glucose acting at P2Y₁₄ receptors is a mediator of mast cell degranulation. *Biochemical Pharmacology* 79, 873–879.
- Gao, Z., -G., Wei, Q., Jayasekara, M. P. S., & Jacobson, K. A. (2013). The role of P2Y₁₄ and other P2Y receptors in degranulation of human LAD2 mast cells. *Purinergic Signalling* 9, 31–40.
- García, R. A., Yan, M., Search, D., Zhang, R., Carson, N. L., Ryan, C. S., ... Gargalovic, P. S. (2014). P2Y₆ receptor potentiates pro-inflammatory responses in macrophages and exhibits differential roles in atherosclerotic lesion development. *PLoS One* 9, e111385.
- Gauthier, J. Y., Belley, M., Deschenes, D., Fournier, J. -F., Gagné, S., Gareau, Y., ... Black, W. C. (2011). The identification of 4,7-disubstituted naphthoic acid derivatives as UDP-competitive antagonists of P2Y₁₄. *Bioorganic & Medicinal Chemistry Letters* 21, 2836–2839.
- Gendaszewska-Darmach, E., & Szustak, M. (2016). Thymidine 5'-O-monophosphorothioate induces HeLa cell migration by activation of the P2Y₆ receptor. *Purinergic Signalling* 12, 199–209.
- Gendaszewska-Darmach, E., Weglowska, E., Walczak-Drzewiecka, A., & Karas, K. (2016). Nucleoside 5'-O-monophosphorothioates as modulators of the P2Y₁₄ receptor and mast cell degranulation. *Oncotarget* 7, 69358–69370.
- Ghanem, E., Robaye, B., Leal, T., Leipziger, J., van Driessche, W., Beauwens, R., & Boeynaems, J. -M. (2005). The role of epithelial P2Y₂ and P2Y₄ receptors in the regulation of intestinal chloride secretion. *British Journal of Pharmacology* 146, 364–369.
- Ginsburg-Shmuel, T., Haas, M., Grbic, D., Arguin, G., Nadel, Y., Gendron, F. -P., ... Fischer, B. (2012). UDP made a highly promising stable, potent, and selective P2Y₆-receptor agonist upon introduction of a boranophosphate moiety. *Bioorganic & Medicinal Chemistry* 20, 5483–5495.
- Ginsburg-Shmuel, T., Haas, M., Schumann, M., Reiser, G., Kalid, O., Stern, N., & Fischer, B. (2010). 5-OME-UDP is a potent and selective P2Y₆-receptor agonist. *Journal of Medicinal Chemistry* 53, 1673–1685.
- Glänzel, M., Bültmann, R., Starke, K., & Frahm, A. W. (2003). Constitutional isomers of reactive blue 2-selective P2Y-receptor antagonists? *European Journal of Medicinal Chemistry* 38, 303–312.
- Grbic, D. M., Degagne, E., Langlois, C., Dupuis, A. -A., & Gendron, F. -P. (2008). Intestinal inflammation increases the expression of the P2Y₆ receptor on epithelial cells and the release of CXC chemokine ligand 8 by UDP. *Journal of Immunology* 180, 2659–2668.
- Grbic, D. M., Degagne, E., Larivee, J. -F., Bildeau, M. S., Vinette, V., Arguin, G., ... Gendron, F. -P. (2012). P2Y₆ receptor contributes to neutrophil recruitment to inflamed intestinal mucosa by increasing CXC chemokine ligand 8 expression in an AP-1-dependent manner in epithelial cells. *Inflammatory Bowel Diseases* 18, 1456–1469.
- Greig, A. V. H., Linge, C., Cambrey, A., & Burnstock, G. (2003). Purinergic receptors are part of a signaling system for keratinocyte proliferation, differentiation, and apoptosis in human fetal epidermis. *The Journal of Investigative Dermatology* 121, 1145–1149.
- Guay, D., Beaulieu, C., Belley, M., Crane, S. N., DeLuca, J., Fortin, R., ... Wang, Z. (2008). Merck Frost Canada Ltd. WO00200900087A1.
- Guay, D., Beaulieu, C., Belley, M., Crane, S. N., DeLuca, J., Gareau, Y., ... Black, W. C. (2011). Synthesis and SAR of pyrimidine-based, non-nucleotide P2Y₁₄ receptor antagonists. *Bioorganic & Medicinal Chemistry Letters* 21, 2832–2835.
- Guile, S. D., Ince, F., Ingall, A. H., Kindon, N. D., Meghani, P., & Mortimore, M. P. (2001). The medicinal chemistry of the P2 receptor family. *Progress in Medicinal Chemistry* 38, 115–187.
- Guns, P. -J. D. F., Hendrickx, J., van Assche, T., Fransen, P., & Bult, H. (2010). P2Y receptors and atherosclerosis in apolipoprotein E-deficient mice. *British Journal of Pharmacology* 159, 326–336.
- Haas, M., Ginsburg-Shmuel, T., Fischer, B., & Reiser, G. (2014). 5-OME-uridine-5'-O-(alpha-boranodiphosphate), a novel nucleotide derivative highly active at the human P2Y₆ receptor protects against death-receptor mediated glial apoptosis. *Neuroscience Letters* 578, 80–84.
- Hamel, M., Henault, M., Hyjazie, H., Morin, N., Bayly, C., Skorey, K., ... Kargman, S. (2011). Discovery of novel P2Y₁₄ agonist and antagonist using conventional and nonconventional methods. *Journal of Biomolecular Screening* 16, 1098–1105.
- Hansen, A., Alston, L., Tulk, S. E., Schenck, L. P., Grassie, M. E., Alhassan, B. F., ... Hirota, S. A. (2013). The P2Y₆ receptor mediates clostridium difficile toxin-induced CXCL8/IL-8 production and intestinal epithelial barrier dysfunction. *PLoS One* 8, e81491.
- Hao, Y., Liang, J. F., Chow, A. W., Cheung, W. -T., & Ko, W. -H. (2014). P2Y₆ receptor-mediated proinflammatory signaling in human bronchial epithelia. *PLoS One* 9, e106235.
- Harden, T. K., Lazarowski, E. R., & Boucher, R. C. (1997). Release, metabolism and interconversion of adenine and uridine nucleotides: Implications for G protein-coupled P2 receptor agonist selectivity. *Trends in Pharmacological Sciences* 18, 43–46.
- Harden, T. K., Sesma, J. I., Fricks, I. P., & Lazarowski, E. R. (2010). Signalling and pharmacological properties of the P2Y receptor. *Acta Physiologica* 199, 149–160.
- Herold, C. L., Li, Q., Schachter, J. B., Harden, T. K., & Nicholas, R. A. (1997). Lack of nucleotide-promoted second messenger signaling responses in 1321N1 cells expressing the proposed P2Y receptor, p2y7. *Biochemical and Biophysical Research Communications* 235, 717–721.
- Herold, C. L., Qi, A. -D., Harden, T. K., & Nicholas, R. A. (2004). Agonist versus antagonist action of ATP at the P2Y₄ receptor is determined by the second extracellular loop. *The Journal of Biological Chemistry* 279, 11456–11464.
- Hillmann, P., Ko, G. -Y., Spinrath, A., Raulf, A., von Kügelgen, I., Wolff, S. C., ... Müller, C. E. (2009). Key determinants of nucleotide-activated G protein-coupled P2Y₂ receptor function revealed by chemical and pharmacological experiments, mutagenesis and homology modeling. *Journal of Medicinal Chemistry* 52, 2762–2775.
- Hochhauser, E., Cohen, R., Waldman, M., Maksin, A., Isak, A., Aravot, D., ... Shainberg, A. (2013). P2Y₂ receptor agonist with enhanced stability protects the heart from ischemic damage in vitro and in vivo. *Purinergic Signalling* 9, 633–642.
- Hoebertz, A., Mahendran, S., Burnstock, G., & Arnett, T. R. (2002). ATP and UTP at low concentrations strongly inhibit bone formation by osteoblasts: A novel role for the P2Y₂ receptor in bone remodeling. *Journal of Cellular Biochemistry* 86, 413–419.
- Hoffmann, K., Soltysiak, K., West, P. L., & Jacobson, K. A. (2004). Shift in purine/pyrimidine base recognition upon exchanging extracellular domains in P2Y_{1/6} chimeric receptors. *Biochemical Pharmacology* 68, 2075–2086.
- Hoffmann, K., Baqi, Y., Morena, M. S., Glänzel, M., Müller, C. E., & von Kügelgen, I. (2009). Interaction of new, very potent non-nucleotide antagonists with Arg256 of the human platelet P2Y₁₂ receptor. *The Journal of Pharmacology and Experimental Therapeutics* 331, 648–655.

- Horckmans, M., Esfahani, H., Beauloye, C., Clouet, S., Di Pietrantonio, L., Robaye, B., ... Communi, D. (2015). Loss of mouse P2Y₄ nucleotide receptor protects against myocardial infarction through endothelin-1 downregulation. *Journal of Immunology* 194, 1874–1881.
- Horckmans, M., Robaye, B., Leon-Gomicronmez, E., Lantz, N., Unger, P., Dol-Gleizes, F., ... Communi, D. (2012). P2Y₄ nucleotide receptor: A novel actor in post-natal cardiac development. *Angiogenesis* 15, 349–360.
- Hou, M., Harden, T. K., Kuhn, C. M., Baldetorp, B., Lazarowski, E., Pendergast, W., ... Erlinge, D. (2002). UDP acts as a growth factor for vascular smooth muscle cells by activation of P2Y₆ receptors. *American Journal of Physiology. Heart and Circulatory Physiology* 282, H784–92.
- Ide, S., Nishimaki, N., Tsukimoto, M., & Kojima, S. (2014). Purine receptor P2Y₆ mediates cellular response to gamma-ray-induced DNA damage. *The Journal of Toxicological Sciences* 39, 15–23.
- Idzko, M.,ammad, H., van Nimwegen, M., Kool, M., Willart, M. A. M., Muskens, F., ... Lambrecht, B. N. (2007). Extracellular ATP triggers and maintains asthmatic airway inflammation by activating dendritic cells. *Nature Medicine* 13, 913–919.
- Inoue, K., Koizumi, S., Kataoka, A., Tozaki-Saitoh, H., & Tsuda, M. (2009). P2Y₆-evoked microglial phagocytosis. *International Review of Neurobiology* 85, 159–163.
- Ito, M., Egashira, S., I., Yoshida, K., Mineno, T., Kumagai, K., Kojima, H., ... Matsuoka, I. (2017). Identification of novel selective P2Y₆ receptor antagonists by high-throughput screening assay. *Life Sciences* 180, 137–142.
- Ivanov, A. A., Fricks, I., Kendall Harden, T., & Jacobson, K. A. (2007). Molecular dynamics simulation of the P2Y₁₄ receptor. Ligand docking and identification of a putative binding site of the distal hexose moiety. *Bioorganic & Medicinal Chemistry Letters* 17, 761–766.
- Ivanov, A. A., Ko, H., Cosyn, L., Maddileti, S., Besada, P., Fricks, I., ... Jacobson, K. A. (2007). Molecular modeling of the human P2Y₂ receptor and design of a selective agonist, 2'-amino-2'-deoxy-2'-thiouridine 5'-triphosphate. *Journal of Medicinal Chemistry* 50, 1166–1176.
- Jacobson, K. A., & Civan, M. M. (2016). Ocular purine receptors as drug targets in the eye. *Journal of Ocular Pharmacology and Therapeutics* 32, 534–547.
- Jacobson, K. A., Costanzi, S., Ivanov, A. A., Tchilibon, S., Besada, P., Gao, Z., ... Harden, T. K. (2006). Structure activity and molecular modeling analyses of ribose- and base-modified uridine 5'-triphosphate analogues at the human P2Y₂ and P2Y₄ receptors. *Biochemical Pharmacology* 71, 540–549.
- Jacobson, K. A., Fischer, B., & Maillard, M. (1995). USA Department of Health and Human Services US000005620676A.
- Jacobson, K. A., Ivanov, A. A., de Castro, S., Harden, T. K., & Ko, H. (2009). Development of selective agonists and antagonists of P2Y receptors. *Purinergic Signalling* 5, 75–89.
- Jacobson, K. A., & Marquez, V. E. (2001). Jacobson, Kenneth A.; Marquez, Victor E. WO002001051490A1.
- Jacobson, K. A., & Tosh, D. K. (2010). USA Department of Health and Human Services WO002011068978A1.
- Jacobus, K. M., & Leighton, J. H. (1997). Inspire Pharmaceuticals Inc. DE000069725160T2.
- Janssens, R., Paindavoinne, P., Parmentier, M., & Boeynaems, J. M. (1999). Human P2Y₂ receptor polymorphism: identification and pharmacological characterization of two allelic variants. *British Journal of Pharmacology* 127, 709–716.
- Jayasekara, P. S., Barrett, M. O., Ball, C. B., Brown, K. A., Hammes, E., Balasubramanian, R., ... Jacobson, K. A. (2014). 4-Alkylxoyimino derivatives of uridine-5'-triphosphate: distal modification of potent agonists as a strategy for molecular probes of P2Y₂, P2Y₄, and P2Y₆ receptors. *Journal of Medicinal Chemistry* 57, 3874–3883.
- Jayasekara, P. S., Barrett, M. O., Ball, C. B., Brown, K. A., Kozma, E., Costanzi, S., ... Jacobson, K. A. (2013). 4-Alkylxoyimino-cytosine nucleotides: tethering approaches to molecular probes for the P2Y₆ receptor. *Medicinal Chemistry Communications* 4, 1156–1165.
- Johnson, F. L., Donohue, J. F., & Shaffer, C. L. (2002). Improved sputum expectoration following a single dose of INS316 in patients with chronic bronchitis. *Chest* 122, 2021–2029.
- Jokela, T. A., Karna, R., Makkonen, K. M., Laitinen, J. T., Tammi, R. H., & Tammi, M. I. (2014). Extracellular UDP-glucose activates P2Y₁₄ receptor and induces signal transducer and activator of transcription 3 (STAT3) Tyr705 phosphorylation and binding to hyaluronan synthase 2 (HAS2) promoter, stimulating hyaluronan synthesis of keratinocytes. *The Journal of Biological Chemistry* 289, 18569–18581.
- Junker, A., Balasubramanian, R., Ciancetta, A., Uliassi, E., Kiselev, E., Martiriggiano, C., ... Jacobson, K. A. (2016). Structure-based design of 3-(4-Aryl-1H-1,2,3-triazol-1-yl)-bi-phenyl derivatives as P2Y₁₄ receptor antagonists. *Journal of Medicinal Chemistry* 59, 6149–6168.
- Kaczmarek-Hájek, K., Lorinczi, E., Hausmann, R., & Nicke, A. (2012). Molecular and functional properties of P2X receptors—recent progress and persisting challenges. *Purinergic Signalling* 8, 375–417.
- Kargman, S., Hamel, M., Mancini, J. A., & Bayly, C. (2008). Merck US020090148850A1.
- Kaulich, M., Streicher, F., Mayer, R., Müller, I., & Müller, C. E. (2003). Flavonoids - novel lead compounds for the development of P2Y₂ receptor antagonists. *Drug Development Research* 59, 72–81.
- Kawashita, E., Tsuji, D., Kanno, Y., Tsuchida, K., & Itoh, K. (2016). Enhancement by uridine diphosphate of macrophage inflammatory protein-1 alpha production in microglia derived from sandhoff disease model mice. *JIMD Reports* 28, 85–93.
- Keating, G. M. (2015). Diquafosol ophthalmic solution 3 %: A review of its use in dry eye. *Drugs* 75, 911–922.
- Kellerman, D., Rossi Mospan, A., Engels, J., Schaberg, A., Gorden, J., & Smiley, L. (2008). Denufosol: A review of studies with inhaled P2Y₂ agonists that led to phase 3. *Pulmonary Pharmacology & Therapeutics* 21, 600–607.
- Kemp, P. A., Sugar, R. A., & Jackson, A. D. (2004). Nucleotide-mediated mucin secretion from differentiated human bronchial epithelial cells. *American Journal of Respiratory Cell and Molecular Biology* 31, 446–455.
- Kennedy, C., Qi, A. D., Herold, C. L., Harden, T. K., & Nicholas, R. A. (2000). ATP, an agonist at the rat P2Y₄ receptor, is an antagonist at the human P2Y₄ receptor. *Molecular Pharmacology* 57, 926–931.
- Kerr, D. I. B., & Krantis, A. (1979). A new class of ATP antagonist. *Proceedings of the Australian Physiological and Pharmacological Society* 10, 156.
- Khelashvili, G., Dorff, K., Shan, J., Camacho-Artacho, M., Skrabanek, L., Vroiling, B., ... Filizola, M. (2010). GPCR-OKB: The G protein coupled receptor oligomer knowledge base. *Bioinformatics* 26, 1804–1805.
- Khine, A. A., Del Sorbo, L., Vaschetto, R., Voglis, S., Tullis, E., Slutsky, A. S., ... Zhang, H. (2006). Human neutrophil peptides induce interleukin-8 production through the P2Y₆ signaling pathway. *Blood* 107, 2936–2942.
- Kikuta, Y., Ohiwa, E., Okada, K., Watanabe, A., & Haruki, S. (1999). Clinical application of diadenosine tetraphosphate (Ap4A:F-1500) for controlled hypotension. *Acta Anaesthesiologica Scandinavica* 43, 82–86.
- Kim, B., Jeong, H., Kim, J., Lee, S. Y., Jou, I., & Joe, E. (2011). Uridine 5'-diphosphate induces chemokine expression in microglia and astrocytes through activation of the P2Y₆ receptor. *Journal of Immunology* 186, 3701–3709.
- Kim, C., Kim, H., Lee, H. S., Chang, S. O., Oh, S., Lee, J. H. (2010). P2Y₄-mediated regulation of Na⁺ absorption in the Reissner's membrane of the cochlea. *The Journal of Neuroscience* 30, 3762–3769.
- Kim, H. J., Ajit, D., Peterson, T. S., Wang, Y., Camden, J. M., Gibson Wood, W., ... Weisman, G. A. (2012). Nucleotides released from Abeta(1)(-)(4)(2) -treated microglial cells increase cell migration and Abeta(1)(-)(4)(2) uptake through P2Y(2) receptor activation. *Journal of Neurochemistry* 121, 228–238.
- Kim, H. S., Ravi, R. G., Marquez, V. E., Maddileti, S., Wihlborg, A., Erlinge, D., ... Jacobson, K. A. (2002). Methanocarpa modification of uracil and adenine nucleotides: high potency of Northern ring conformation at P2Y₁, P2Y₂, P2Y₄, and P2Y₁₁, but not P2Y₆ receptors. *Journal of Medicinal Chemistry* 45, 208–218.
- Kim, S. G., Gao, Z., G., Soltysiak, K. A., Chang, T. -S., Brodie, C., & Jacobson, K. A. (2003). P2Y₆ nucleotide receptor activates PKC to protect 1321N1 astrocytoma cells against tumor necrosis factor-induced apoptosis. *Cellular and Molecular Neurobiology* 23, 401–418.
- Kim, S. G., Soltysiak, K. A., Gao, Z., G., Chang, T. -S., Chung, E., & Jacobson, K. A. (2003). Tumor necrosis factor alpha-induced apoptosis in astrocytes is prevented by the activation of P2Y₆, but not P2Y₄ nucleotide receptors. *Biochemical Pharmacology* 65, 923–931.
- Kindon, N., Davis, A., Dougall, I., Dixon, J., Johnson, T., Walters, I., ... Stocks, M. J. (2017). From UTP to AR-C118925, the discovery of a potent non nucleotide antagonist of the P2Y₂ receptor. *Bioorganic & Medicinal Chemistry Letters* 27, 4849–4853.
- Kindon, N., Meghani, P., & Thom, S. (1998). Astra Pharmaceuticals Ltd. WO001999002501A1.
- Kiselev, E., Balasubramanian, R., Uliassi, E., Brown, K. A., Trujillo, K., Katritch, V., ... Jacobson, K. A. (2015). Design, synthesis, pharmacological characterization of a fluorescent agonist of the P2Y(1)(4) receptor. *Bioorganic & Medicinal Chemistry Letters* 25, 4733–4739.
- Kiselev, E., Barrett, M. O., Katritch, V., Paoletta, S., Weitzer, C. D., Brown, K. A., ... Jacobson, K. A. (2014). Exploring a 2-naphthoic acid template for the structure-based design of P2Y₁₄ receptor antagonist molecular probes. *ACS Chemical Biology* 9, 2833–2842.
- Kishore, B. K., Carlson, N. G., Ecelbarger, C. M., Kohan, D. E., Müller, C. E., Nelson, R. D., ... Zhang, Y. (2015). Targeting renal purinergic signalling for the treatment of lithium-induced nephrogenic diabetes insipidus. *Acta Physiologica* 214, 176–188.
- Kishore, B. K., Carlson, N. G., & Zhang, Y. (2012). University of Utah Research Foundation WO002013033178A1.
- Knoblauch, B. H. A., Müller, C. E., Järlbark, L., Lawoko, G., Kottke, T., Wikström, M. A., & Heilbronn, E. (1999). 5-Substituted UTP derivatives as P2Y₂ receptor agonists. *European Journal of Medicinal Chemistry*, 809–824.
- Ko, H., Carter, R. L., Cosyn, L., Petrelli, R., de Castro, S., Besada, P., ... Jacobson, K. A. (2008). Synthesis and potency of novel uracil nucleotides and derivatives as P2Y₂ and P2Y₆ receptor agonists. *Bioorganic & Medicinal Chemistry* 16, 6319–6332.
- Ko, H., Das, A., Carter, R. L., Fricks, I. P., Zhou, Y., Ivanov, A. A., ... Jacobson, K. A. (2009). Molecular recognition in the P2Y(14) receptor: Probing the structurally permissive terminal sugar moiety of uridine-5'-diphosphoglucose. *Bioorganic & Medicinal Chemistry* 17, 5298–5311.
- Ko, H., Fricks, I., Ivanov, A. A., Harden, T. K., & Jacobson, K. A. (2007). Structure-activity relationship of uridine 5'-diphosphoglucose analogues as agonists of the human P2Y14 receptor. *Journal of Medicinal Chemistry* 50, 2030–2039.
- Kobayashi, K., Yamanaka, H., Yanamoto, F., Okubo, M., & Noguchi, K. (2012). Multiple P2Y subtypes in spinal microglia are involved in neuropathic pain after peripheral nerve injury. *Glia* 60, 1529–1539.
- Koizumi, S., Shigemoto-Mogami, Y., Nasu-Tada, K., Shinozaki, Y., Ohsawa, K., Tsuda, M., ... Inoue, K. (2007). UDP acting at P2Y₆ receptors is a mediator of microglial phagocytosis. *Nature* 446, 1091–1095.
- Korcok, J., Raimundo, L. N., Du, X., Sims, S. M., & Dixon, S. J. (2005). P2Y₆ nucleotide receptors activate NF-kappaB and increase survival of osteoclasts. *The Journal of Biological Chemistry* 280, 16909–16915.
- Köttgen, M., Löffler, T., Jacobi, C., Nitschke, R., Pavenstädt, H., Schreiber, R., ... Leipziger, J. (2003). P2Y₆ receptor mediates colonic NaCl secretion via differential activation of cAMP-mediated transport. *The Journal of Clinical Investigation* 111, 371–379.
- Kozak, K. R., Crews, B. C., Ray, J. L., Tai, H. H., Morrow, J. D., & Marnett, L. J. (2001). Metabolism of prostaglandin glycerol esters and prostaglandin ethanolamides in vitro and in vivo. *The Journal of Biological Chemistry* 276, 36993–36998.
- Kozma, E., Kumar, T. S., Federico, S., Phan, K., Balasubramanian, R., Gao, Z., ... Jacobson, K. A. (2012). Novel fluorescent antagonist as a molecular probe in A(3) adenosine receptor binding assays using flow cytometry. *Biochemical Pharmacology* 83, 1552–1561.
- Kunzelmann, K., & Mall, M. (2003). Pharmacotherapy of the ion transport defect in cystic fibrosis: Role of purinergic receptor agonists and other potential therapeutics. *American Journal of Respiratory Medicine: Drugs, Devices, and other Interventions* 2, 299–309.

- Lai, M. K. P., Tan, M. G. K., Kirvell, S., Hobbs, C., Lee, J., Esiri, M. M., ... Francis, P. T. (2008). Selective loss of P2Y₂ nucleotide receptor immunoreactivity is associated with Alzheimer's disease neuropathology. *Journal of Neural Transmission* 115, 1165–1172.
- Lambrecht, G. (1996). Design and pharmacology of selective P2-purinoceptor antagonists. *Journal of Autonomic Pharmacology* 16, 341–344.
- Lau, O. C. F., Samarawickrama, C., & Skalkicky, S. E. (2014). P2Y₂ receptor agonists for the treatment of dry eye disease: A review. *Clinical Ophthalmology* 8, 327–334.
- Lazarowski, E. R., Boucher, R. C., & Harden, T. K. (2000). Constitutive release of ATP and evidence for major contribution of ecto-nucleotide pyrophosphatase and nucleoside diphosphokinase to extracellular nucleotide concentrations. *The Journal of Biological Chemistry* 275, 31061–31068.
- Lazarowski, E. R., Boucher, R. C., & Harden, T. K. (2003). Mechanisms of release of nucleotides and integration of their action as P2X- and P2Y-receptor activating molecules. *Molecular Pharmacology* 64, 785–795.
- Lazarowski, E. R., & Harden, T. K. (1994). Identification of a uridine nucleotide-selective G-protein-linked receptor that activates phospholipase C. *The Journal of Biological Chemistry* 269, 11830–11836.
- Lazarowski, E. R., & Harden, T. K. (2015). UDP-sugars as extracellular signaling molecules: Cellular and physiologic consequences of P2Y₁₄ receptor activation. *Molecular Pharmacology* 88, 151–160.
- Lazarowski, E. R., Rochelle, L. G., O'Neal, W. K., Ribeiro, C. M., Grubb, B. R., Zhang, V., ... Boucher, R. C. (2001). Cloning and functional characterization of two murine uridine nucleotide receptors reveal a potential target for correcting ion transport deficiency in cystic fibrosis gallbladder. *The Journal of Pharmacology and Experimental Therapeutics* 297, 43–49.
- Lazarowski, E. R., Shea, D. A., Boucher, R. C., & Harden, T. K. (2003). Release of cellular UDP-glucose as a potential extracellular signaling molecule. *Molecular Pharmacology* 63, 1190–1197.
- Lazarowski, E. R., Watt, W. C., Stutts, M. J., Boucher, R. C., & Harden, T. K. (1995). Pharmacological selectivity of the cloned human P_{2U}-purinoceptor: Potent activation by diadenosine tetraphosphate. *British Journal of Pharmacology* 116, 1619–1627.
- Lazarowski, E. R., Watt, W. C., Stutts, M. J., Brown, H. A., Boucher, R. C., & Harden, T. K. (1996). Enzymatic synthesis of UTPγS, a potent hydrolysis resistant agonist of P_{2U}-purinoceptors. *British Journal of Pharmacology* 117, 203–209.
- Lee, S. A., Park, J. H., & Lee, S. Y. (2013). Selective induction of P2Y₁₄ receptor by RANKL promotes osteoclast formation. *Molecules and Cells* 36, 273–277.
- Li, F., Li, W., Li, X., Li, F., Zhang, L., Wang, B., ... Ma, J. (2016). Geniposide attenuates inflammatory response by suppressing P2Y₁₄ receptor and downstream ERK1/2 signaling pathway in oxygen and glucose deprivation-induced brain microvascular endothelial cells. *Journal of Ethnopharmacology* 185, 77–86.
- Li, H.-q., Chen, C., Dou, Y., Wu, H.-j., Liu, Y.-j., Lou, H.-F., ... Duan, S. (2013). P2Y₄ receptor-mediated pinocytosis contributes to amyloid beta-induced self-uptake by microglia. *Molecular and Cellular Biology* 33, 4282–4293.
- Li, Q., Olesky, M., Palmer, R. K., Harden, T. K., & Nicholas, R. A. (1998). Evidence that the p2y3 receptor is the avian homologue of the mammalian P2Y₆ receptor. *Molecular Pharmacology* 54, 541–546.
- Li, R., Tan, B., Yan, Y., Ma, X., Zhang, N., Zhang, Z., ... Du, B. (2014). Extracellular UDP and P2Y₆ function as a danger signal to protect mice from vesicular stomatitis virus infection through an increase in IFN-beta production. *Journal of Immunology* 193, 4515–4526.
- Liao, Z., Seye, C. I., Weisman, G. A., & Erb, L. (2007). The P2Y₂ nucleotide receptor requires interaction with alpha v integrins to access and activate G₁₂. *Journal of Cell Science* 120, 1654–1662.
- Lin, T. A., Lustig, K. D., Sportiello, M. G., Weisman, G. A., & Sun, G. Y. (1993). Signal transduction pathways coupled to a P_{2U} receptor in neuroblastoma x glioma (NG108-15) cells. *Journal of Neurochemistry* 60, 1115–1125.
- Lustig, K. D., Shiau, A. K., Brake, A. J., & Julius, D. (1993). Expression cloning of an ATP receptor from mouse neuroblastoma cells. *Proceedings of the National Academy of Sciences of the United States of America* 90, 5113–5117.
- Ma, X., Pan, X., Wei, Y., Tan, B., Yang, L., Ren, H., ... Du, B. (2016). Chemotherapy-induced uridine diphosphate release promotes breast cancer metastasis through P2Y₆ activation. *Oncotarget* 7, 29036–29050.
- Mack, S. R., Sabin, V. M., & Galvin, F. C. A. (2002). Cell-tech R&D Ltd. WO002003011885A1.
- Magni, G., Merli, D., Verderio, C., Abbraccio, M. P., & Ceruti, S. (2015). P2Y₂ receptor antagonists as anti-allodynic agents in acute and sub-chronic trigeminal sensitization: role of satellite glial cells. *Glia* 63, 1256–1269.
- Malam-Souley, R., Seye, C., Gadeau, A. P., Loirand, G., Pillois, X., Campan, M., ... Desgranges, C. (1996). Nucleotide receptor P_{2U} partially mediates ATP-induced cell cycle progression of aortic smooth muscle cells. *Journal of Cellular Physiology* 166, 57–65.
- Malik, E. M., & Müller, C. E. (2016). Anthraquinones as pharmacological tools and drugs. *Medicinal Research Reviews* 36, 705–748.
- Malin, S. A., Davis, B. M., Koerber, H. R., Reynolds, I. J., Albers, K. M., & Molliver, D. C. (2008). Thermal nociception and TRPV1 function are attenuated in mice lacking the nucleotide receptor P2Y₂. *Pain* 138, 484–496.
- Malmström, M., Adner, M., Harden, T. K., Pendergast, W., Edvinsson, L., & Erlinge, D. (2000). The stable pyrimidines UDPbetasA and UTPgammaSa discriminate between the P2 receptors that mediate vascular contraction and relaxation of the rat mesenteric artery. *British Journal of Pharmacology* 131, 51–56.
- Malmström, M., Hou, M., Pendergast, W., Erlinge, D., & Edvinsson, L. (2003). Potent P2Y₆ receptor mediated contractions in human cerebral arteries. *BMC Pharmacology* 3, 4.
- Mamedova, L. K., Joshi, B. V., Gao, Z.-G., von Kügelgen, I., & Jacobson, K. A. (2004). Diisothiocyanate derivatives as potent, insurmountable antagonists of P2Y₆ nucleotide receptors. *Biochemical Pharmacology* 67, 1763–1770.
- Mamedova, L. K., Wang, R., Besada, P., Liang, B. T., & Jacobson, K. A. (2008). Attenuation of apoptosis *in vitro* and ischemia/reperfusion injury *in vivo* in mouse skeletal muscle by P2Y₆ receptor activation. *Pharmacological Research* 58, 232–239.
- Marcus, D. C., Liu, J., Lee, J. H., Scherer, E. Q., Scofield, M. A., & Wangemann, P. (2005). Apical membrane P2Y₄ purinergic receptor controls K⁺ secretion by strial marginal cell epithelium. *Cell Communication and Signaling: CCS* 3, 13.
- Markovskaya, A., Crooke, A., Guzman-Aranguez, A. I., Peral, A., Ziganshin, A. U., & Pintor, J. (2008). Hypotensive effect of UDP on intraocular pressure in rabbits. *European Journal of Pharmacology* 579, 93–97.
- Martínez-Ramírez, A. S., Garay, E., García-Carrancá, A., & Vázquez-Cuevas, F. G. (2016). The P2RY2 receptor induces carcinoma cell migration and EMT through cross-talk with epidermal growth factor receptor. *Journal of Cellular Biochemistry* 117, 1016–1026.
- Maruoka, H., Barrett, M. O., Ko, H., Tosh, D. K., Melman, A., Burianek, L. E., ... Jacobson, K. A. (2010). Pyrimidine ribonucleotides with enhanced selectivity as P2Y(6) receptor agonists: Novel 4-alkyloxymino, (S)-methanocarpa, and 5'-triphosphate gamma-ester modifications. *Journal of Medicinal Chemistry* 53, 4488–4501.
- Maruoka, H., Jayasekara, M. P. S., Barrett, M. O., Franklin, D. A., de Castro, S., Kim, N., ... Jacobson, K. A. (2011). Pyrimidine nucleotides with 4-alkyloxymino and terminal tetraphosphate delta-ester modifications as selective agonists of the P2Y(4) receptor. *Journal of Medicinal Chemistry* 54, 4018–4033.
- Matos, J. E., Robaye, B., Boeynaems, J. M., Beauwens, R., & Leipziger, J. (2005). K⁺ secretion activated by luminal P2Y₂ and P2Y₄ receptors in mouse colon. *The Journal of Physiology* 564, 269–279.
- Meister, J., Le Duc, D., Ricken, A., Burkhardt, R., Thiery, J., Pfannkuche, H., ... Schulz, A. (2014). The G protein-coupled receptor P2Y₁₄ influences insulin release and smooth muscle function in mice. *The Journal of Biological Chemistry* 289, 23353–23366.
- Mellor, E. A., Frank, N., Soler, D., Hodge, M. R., Lora, J. M., Austen, K. F., & Boyce, J. A. (2003). Expression of the type 2 receptor for cysteinyl leukotrienes (CysLT2R) by human mast cells: Functional distinction from CysLT1R. *Proceedings of the National Academy of Sciences of the United States of America* 100, 11589–11593.
- Mellor, E. A., Maekawa, A., Austen, K. F., & Boyce, J. A. (2001). Cysteinyl leukotriene receptor 1 is also a pyrimidinergic receptor and is expressed by human mast cells. *Proceedings of the National Academy of Sciences of the United States of America* 98, 7964–7969.
- Meltzer, D., Ethan, O., Arguin, G., Nadel, Y., Danino, O., Lecka, J., ... Fischer, B. (2015). Synthesis and structure-activity relationship of uracil nucleotide derivatives towards the identification of human P2Y₆ receptor antagonists. *Bioorganic & Medicinal Chemistry* 23, 5764–5773.
- Miyagi, Y., Kobayashi, S., Ahmed, A., Nishimura, J., Fukui, M., & Kanaide, H. (1996). P_{2U} purinergic activation leads to the cell cycle progression from the G₁ to the S and M phases but not from the G₀ to G₁ phase in vascular smooth muscle cells in primary culture. *Biochemical and Biophysical Research Communications* 222, 652–658.
- Moore, D. J., Chambers, J. K., Wahlin, J. P., Tan, K. B., Moore, G. B., Jenkins, O., ... Murdock, P. R. (2001). Expression pattern of human P2Y₂ receptor subtypes: A quantitative reverse transcription-polymerase chain reaction study. *Biochimica et Biophysica Acta* 1521, 107–119.
- Moore, D. J., Murdock, P. R., Watson, J. M., Faull, R. L. M., Waldvogel, H. J., Szekeres, P. G., ... Emson, P. C. (2003). GPR105, a novel G_{i/o}-coupled UDP-glucose receptor expressed on brain glia and peripheral immune cells, is regulated by immunologic challenge: Possible role in neuroimmune function. *Brain Research. Molecular Brain Research* 118, 10–23.
- Morioka, N., Tokuhara, M., Harano, S., Nakamura, Y., Hisaoka-Nakashima, K., & Nakata, Y. (2013). The activation of P2Y₆ receptor in cultured spinal microglia induces the production of CCL2 through the MAP kinases-NF-kappaB pathway. *Neuropharmacology* 75, 116–125.
- Moss, R. B. (2013). Pitfalls of drug development: Lessons learned from trials of denofosol in cystic fibrosis. *The Journal of Pediatrics* 162, 676–680.
- Müller, C. (2002). P2-Pyrimidinergic Receptors and Their Ligands. *Current Pharmaceutical Design* 8, 2353–2369.
- Müller, C. E., Schiedel, A. C., & Baqi, Y. (2012). Allosteric modulators of rhodopsin-like G protein-coupled receptors: Opportunities in drug development. *Pharmacology & Therapeutics* 135, 292–315.
- Müller, T., Bayer, H., Myrtek, D., Ferrari, D., Sorichter, S., Ziegenhagen, M. W., ... Idzko, M. (2005). The P2Y₁₄ receptor of airway epithelial cells: Coupling to intracellular Ca²⁺ and IL-8 secretion. *American Journal of Respiratory Cell and Molecular Biology* 33, 601–609.
- Müller, T., Fay, S., Vieira, R. P., Karmouty-Quintana, H., Cicko, S., Ayata, C. K., ... Idzko, M. (2017). P2Y₆ receptor activation promotes inflammation and tissue remodeling in pulmonary fibrosis. *Frontiers in Immunology* 8, 1028.
- Mundas, M. V., Novack, G. D., Allgood, V. E., Evans, R. M., Gorden, J. C., & Yexxa, B. R. (2001). Ocular safety of INS365 ophthalmic solution: a P2Y(2) agonist in healthy subjects. *Journal of Ocular Pharmacology and Therapeutics* 17, 173–179.
- Muscella, A., Elia, M. G., Greco, S., Storelli, C., & Marsigliante, S. (2003). Activation of P2Y₂ receptor induces c-FOS protein through a pathway involving mitogen-activated protein kinases and phosphoinositide 3-kinases in HeLa cells. *Journal of Cellular Physiology* 195, 234–240.
- Nakano, M., Ito, K., Yuno, T., Soma, N., Aburakawa, S., Kasai, K., ... Takami, H. (2017). UDP/P2Y₆ receptor signaling regulates IgE-dependent degranulation in human basophils. *Allergy International* 66, 574–580.
- Nguyen, T., Erb, L., Weisman, G. A., Marchese, A., Heng, H. H., Garrad, R. C., ... O'Dowd, B. F. (1995). Cloning, expression, and chromosomal localization of the human uridine nucleotide receptor gene. *The Journal of Biological Chemistry* 270, 30845–30848.
- Nicholas, R. A., Watt, W. C., Lazarowski, E. R., Li, Q., & Harden, K. (1996). Uridine nucleotide selectivity of three phospholipase C-activating P2 receptors: Identification of a UDP-selective, a UTP-selective, and an ATP- and UTP-specific receptor. *Molecular Pharmacology* 50, 224–229.
- Nichols, K. K., Yexxa, B., & Kellerman, D. J. (2004). Diquafosol tetrasodium: A novel dry eye therapy. *Expert Opinion on Investigational Drugs* 13, 47–54.

- Nishida, M., Sato, Y., Uemura, A., Narita, Y., Tozaki-Saitoh, H., Nakaya, M., ... Kurose, H. (2008). P2Y₆ receptor-Galpha12/13 signalling in cardiomyocytes triggers pressure overload-induced cardiac fibrosis. *The EMBO Journal* 27, 3104–3115.
- Noguchi, K., Ishii, S., & Shimizu, T. (2003). Identification of p2y9/GPR23 as a novel G protein-coupled receptor for lysophosphatidic acid, structurally distant from the Edg family. *The Journal of Biological Chemistry* 278, 25600–25606.
- Nomura, N., Miyajima, N., Sazuka, T., Tanaka, A., Kawarabayashi, Y., Sato, S., ... Tabata, S. (1994). Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1. *DNA Research: an International Journal for Rapid Publication of Reports on Genes and Genomes* 1, 27–35.
- North, R. A. (1996). P2X receptors: A third major class of ligand-gated ion channels. *Ciba foundation symposium* 198, (pp. 91–105) (discussion 105–9).
- Nylund, G., Hultman, L., Nordgren, S., & Delbro, D. S. (2007). P2Y₂- and P2Y₄ purinergic receptors are over-expressed in human colon cancer. *Autonomic & Autacoid Pharmacology* 27, 79–84.
- Ohno, M., Costanzi, S., Kim, H. S., Kempeneers, V., Vastmans, K., Herdewijn, P., ... Jacobson, K. A. (2004). Nucleotide analogues containing 2-oxa-bicyclo[2.2.1]heptane and 1-alpha-thiofuranosyl ring systems: interactions with P2Y receptors. *Bioorganic & Medicinal Chemistry* 12, 5619–5630.
- Ohtani, M., Suzuki, J., -I., Jacobson, K. A., & Oka, T. (2008). Evidence for the possible involvement of the P2Y(6) receptor in Ca(2+) mobilization and insulin secretion in mouse pancreatic islets. *Purinergic Signalling* 4, 365–375.
- Önnheim, K., Christenson, K., Gabl, M., Burbiel, J. C., Müller, C. E., Oprea, T. I., ... Forsman, H. (2014). A novel receptor cross-talk between the ATP receptor P2Y₂ and formyl peptide receptors reactivates desensitized neutrophils to produce superoxide. *Experimental Cell Research* 323, 209–217.
- Orriss, I. R., Guneri, D., Hajjawi, M. O. R., Shaw, K., Patel, J. J., & Arnett, T. R. (2017). Activation of the P2Y₂ receptor regulates bone cell function by enhancing ATP release. *The Journal of Endocrinology* 233, 341–356.
- Orriss, I. R., Wang, N., Burnstock, G., Arnett, T. R., Gartland, A., Robaye, B., & Boeynaems, J. -M. (2011). The P2Y(6) receptor stimulates bone resorption by osteoclasts. *Endocrinology* 152, 3706–3716.
- Parandeh, F., Abaraviciene, S. M., Amisten, S., Erlinge, D., & Salehi, A. (2008). Uridine diphosphate (UDP) stimulates insulin secretion by activation of P2Y₆ receptors. *Biochemical and Biophysical Research Communications* 370, 499–503.
- Parr, C. E., Sullivan, D. M., Paradiso, A. M., Lazarowski, E. R., Burch, L. H., Olsen, J. C., ... Turner, J. T. (1994). Cloning and expression of a human P_{2U} nucleotide receptor, a target for cystic fibrosis pharmacotherapy. *Proceedings of the National Academy of Sciences of the United States of America* 91, 3275–3279.
- Patel, K., Barnes, A., Camacho, J., Paterson, C., Boughtflower, R., Cousins, D., & Marshall, F. (2001). Activity of diadenosine polyphosphates at P2Y receptors stably expressed in 1321N1 cells. *European Journal of Pharmacology* 430, 203–210.
- Pendergast, W., Rideout, J. L., Siddiqi, S. M., & Yerxa, B. R. (1998). Inspire Pharmaceuticals Inc. WO001998034942A2.
- Pendergast, W., Shaver, S. R., Drutz, D. J., & Rideout, J. L. (1999). Inspire Pharmaceuticals Inc. WO002000030629A2.
- Pendergast, W., Yerxa, B. R., Douglass, J. G., III, Shaver, S. R., Dougherty, R. W., Redick, C. C., ... Rideout, J. L. (2001). Synthesis and P2Y receptor activity of a series of uridine dinucleoside 5'-polyphosphates. *Bioorganic & Medicinal Chemistry Letters* 11, 157–160.
- Peterson, W. (2001). Inspire Pharmaceuticals Inc. WO002001087913A2.
- Peterson, W., & Yerxa, B. R. (2002). Inspire Pharmaceuticals Inc. WO002002060454A2.
- Pintor, J., Carracedo, G., Alonso, M. C., Bautista, A., & Peral, A. (2002). Presence of diadenosine polyphosphates in human tears. *Pflügers Archiv - European Journal of Physiology* 443, 432–436.
- Pintor, J., Peral, A., Peláez, T., Carracedo, G., Bautista, A., & Hoyle, C. H. V. (2003). Nucleotides and dinucleotides in ocular physiology: New possibilities of nucleotides as therapeutic agents in the eye. *Drug Development Research* 59, 136–145.
- Qian, Y., Xu, S., Yang, X., & Xiao, Q. (2017). Purinergic receptor P2Y₆ contributes to 1-methyl-4-phenylpyridinium-induced oxidative stress and cell death in neuronal SH-SY5Y cells. *Journal of Neuroscience Research* 96, 253–264.
- Qiu, Y., Liu, Y., Li, W., -H., Zhang, H. -Q., Tian, X. -X., & Fang, W. -G. (2018). P2Y₂ receptor promotes the migration and invasion of breast cancer cells via EMT-related genes Snail and E-cadherin. *Oncology Reports* 39, 138–150.
- Rafehi, M., Burbiel, J. C., Attah, I. Y., Abdelrahman, A., & Müller, C. E. (2017). Synthesis, characterization, and *in vitro* evaluation of the selective P2Y₂ receptor antagonist AR-C118925. *Purinergic Signalling* 13, 89–103.
- Rafehi, M., Malik, E. M., Neumann, A., Abdelrahman, A., Hanck, T., Namasivayam, V., ... Baqi, Y. (2017). Development of potent and selective antagonists for the UTP-activated P2Y₄ receptor. *Journal of Medicinal Chemistry* 60, 3020–3038.
- Rafehi, M., Neumann, A., Baqi, Y., Malik, E. M., Wiese, M., Namasivayam, V., & Müller, C. E. (2017). Molecular recognition of agonists and antagonists by the nucleotide-activated P2Y₂ receptor. *Journal of Medicinal Chemistry* 60, 8425–8440.
- Ratjen, F., Durham, T., Navratil, T., Schaberg, A., Accurso, F. J., Wainwright, C., ... Moss, R. B. (2012). Long term effects of denufisol tetrasodium in patients with cystic fibrosis. *Journal of Cystic Fibrosis* 11, 539–549.
- Ribeiro-Filho, A. C., Buri, M. V., Barros, C. C., Dreyfuss, J. L., Nader, H. B., Justo, G. Z., ... Paredes-Gamero, E. J. (2016). Functional and molecular evidence for heteromeric association of P2Y1 receptor with P2Y₂ and P2Y₄ receptors in mouse granulocytes. *BMC Pharmacology and Toxicology* 17, 29.
- Rideout, J. L., Yerxa, B. R., Shaver, S. R., & Douglass, J. G. III. (2003). Inspire Pharmaceuticals Inc. US000006867199B2.
- Riegel, A. -K., Faigle, M., Zug, S., Rosenberger, P., Robaye, B., Boeynaems, J. -M., ... Eltzschig, H. K. (2011). Selective induction of endothelial P2Y₆ nucleotide receptor promotes vascular inflammation. *Blood* 117, 2548–2555.
- Robaye, B., Boeynaems, J. M., & Communi, D. (1997). Slow desensitization of the human P2Y₆ receptor. *European Journal of Pharmacology* 329, 231–236.
- Robaye, B., Ghanem, E., Wilkin, F., Fokan, D., van Driessche, W., Schurmans, S., ... Beauwens, R. (2003). Loss of nucleotide regulation of epithelial chloride transport in the jejunum of P2Y₄-null mice. *Molecular Pharmacology* 63, 777–783.
- Robichaud, J., Fournier, J. -F., Gagné, S., Gauthier, J. Y., Hamel, M., Han, Y., ... Black, W. C. (2011). Applying the pro-drug approach to afford highly bioavailable antagonists of P2Y(14). *Bioorganic & Medicinal Chemistry Letters* 21, 4366–4368.
- Robles-Martinez, L., Garay, E., Martel-Gallegos, M. G., Cisneros-Mejorado, A., Perez-Montiel, D., Lara, A., & Arellano, R. O. (2017). Kca3.1 activation via P2Y₂ purinergic receptors promotes human ovarian cancer cell (Skov-3) migration. *Scientific Reports* 7, 4340.
- Sakuma, K., Nakagawa, H., Oikawa, T., Noda, M., & Ikeda, S. (2017). Effects of 4(1H)-quinolinone derivative, a novel non-nucleotide allosteric purinergic P2Y₂ agonist, on cardiomyocytes in neonatal rats. *Scientific Reports* 7, 6050.
- Sauer, R., El-Tayeb, A., Kaulich, M., & Müller, C. E. (2009). Synthesis of uracil nucleotide analogs with a modified, acyclic ribose moiety as P2Y(2) receptor antagonists. *Bioorganic & Medicinal Chemistry* 17, 5071–5079.
- Sauzeau, V., LeJeune, H., Cario-Toumaniantz, C., Vaillant, N., Gadeau, A. P., Desgranges, C., ... Loirand, G. (2000). P2Y(1), P2Y(2), P2Y(4), and P2Y(6) receptors are coupled to Rho and Rho kinase activation in vascular myocytes. *American Journal of Physiology. Heart and Circulatory Physiology* 278, H1751–61.
- Schafer, R., Sedehizade, F., Welte, T., & Reiser, G. (2003). ATP- and UTP-activated P2Y receptors differently regulate proliferation of human lung epithelial tumor cells. *American Journal of Physiology. Lung Cellular and Molecular Physiology* 285, L376–85.
- Schreiber, R., & Kunzelmann, K. (2005). Purinergic P2Y₆ receptors induce Ca²⁺ and CFTR dependent Cl⁻ secretion in mouse trachea. *Cellular Physiology and Biochemistry: International Journal of Experimental Cellular Physiology, Biochemistry, and Pharmacology* 16, 99–108.
- Schumacher, D., Strilic, B., Sivaraj, K. K., Wetschurack, N., & Offermanns, S. (2013). Platelet-derived nucleotides promote tumor-cell transendothelial migration and metastasis via P2Y₂ receptor. *Cancer Cell* 24, 130–137.
- Scrivens, M., & Dickinson, J. M. (2005). Functional expression of the P2Y₁₄ receptor in murine T-lymphocytes. *British Journal of Pharmacology* 146, 435–444.
- Scrivens, M., & Dickinson, J. M. (2006). Functional expression of the P2Y₁₄ receptor in human neutrophils. *European Journal of Pharmacology* 543, 166–173.
- Seifert, R., & Schultz, G. (1989). Involvement of pyrimidinoreceptors in the regulation of cell functions by uridine and by uracil nucleotides. *Trends in Pharmacological Sciences* 10, 365–369.
- Sesma, J. I., Kreda, S. M., Steinckwich-Besancon, N., Dang, H., Garcia-Mata, R., Harden, T. K., & Lazarowski, E. R. (2012). The UDP-sugar-sensing P2Y(14) receptor promotes Rho-mediated signaling and chemotaxis in human neutrophils. *American Journal of Physiology. Cell Physiology* 303, C490–8.
- Sesma, J. I., Weitzer, C. D., Livraghi-Butrico, A., Dang, H., Donaldson, S., Alexis, N. E., ... Lazarowski, E. R. (2016). UDP-glucose promotes neutrophil recruitment in the lung. *Purinergic Signalling* 12, 627–635.
- Seye, C. I., Kong, Q., Erb, L., Garrad, R. C., Krugh, B., Wang, M., ... Weisman, G. A. (2002). Functional P2Y₂ nucleotide receptors mediate uridine 5'-triphosphate-induced intimal hyperplasia in collared rabbit carotid arteries. *Circulation* 106, 2720–2726.
- Shaffer, C. L., Boucher, R. C., Rideout, J. L., & Jacobus, K. M. (1997). Inspire Pharmaceuticals Inc. WO001998019685A1.
- Shaver, S. R., Pendergast, W., Siddiqi, S. M., Yerxa, B. R., Croom, D. K., Dougherty, R. W., ... Rideout, J. L. (1997). 4-Substituted uridine 5'-triphosphates as agonists of the P2Y₂ purinergic receptor. *Nucleosides & Nucleotides*, 1099.
- Shaver, S. R., Rideout, J. L., Pendergast, W., Douglass, J. G., Brown, E. G., Boyer, J. L., ... Yerxa, B. R. (2005). Structure-activity relationships of dinucleotides: Potent and selective agonists of P2Y receptors. *Purinergic Signalling* 1, 183–191.
- Shen, J., Seye, C. I., Wang, M., Weisman, G. A., Wilden, P. A., & Sturek, M. (2004). Cloning, up-regulation, and mitogenic role of porcine P2Y₂ receptor in coronary artery smooth muscle cells. *Molecular Pharmacology* 66, 1265–1274.
- Shi, J. -P., Wang, S. -Y., Chen, L. -L., Zhang, X. -Y., Zhao, Y. -H., Du, B., ... Ren, H. (2016). P2Y₆ contributes to ovalbumin-induced allergic asthma by enhancing mast cell function in mice. *Oncotarget* 7, 60906–60918.
- Shi, Y., Qin, W., Nie, F., Wen, H., Lu, K., & Cui, J. (2017). Ulinastatin attenuates neuropathic pain via the ATP/P2Y₂ receptor pathway in rat models. *Gene* 627, 263–270.
- Shin, A., Toy, T., Rothenfusser, S., Robson, N., Vorac, J., Dauer, M., ... Schnurr, M. (2008). P2Y receptor signaling regulates phenotype and IFN-alpha secretion of human plasmacytoid dendritic cells. *Blood* 111, 3062–3069.
- Shinozaki, Y., Kashiwagi, K., Namekata, K., Takeda, A., Ohno, N., Robaye, B., ... Koizumi, S. (2017). Purinergic dysregulation causes hypertensive glaucoma-like optic neuropathy. *JCI Insight* 2, 93456.
- Sil, P., Hayes, C. P., Reaves, B. J., Breen, P., Quinn, S., Sokolove, J., & Rada, B. (2017). P2Y6 receptor antagonist MRS2578 inhibits neutrophil activation and aggregated neutrophil extracellular trap formation induced by gout-associated monosodium urate crystals. *Journal of Immunology* 198, 428–442.
- Silva, I., Ferreirinha, F., Magalhães-Cardoso, M. T., Silva-Ramos, M., & Correia-de-Sá, P. (2015). Activation of P2Y₆ receptors facilitates nonneuronal adenosine triphosphate and acetylcholine release from urothelium with the lamina propria of men with bladder outlet obstruction. *The Journal of Urology* 194, 1146–1154.
- Singh Grewal, A., Pandita, D., Bhardwaj, S., & Lather, V. (2016). Recent updates on development of drug molecules for human african trypanosomiasis. *Current Topics in Medicinal Chemistry* 16, 2245–2265.
- Skrabaneck, L., Murcia, M., Bouvier, M., Devi, L., George, S. R., Lohse, M. J., ... Filizola, M. (2007). Requirements and ontology for a G protein-coupled receptor oligomerization knowledge base. *BMC Bioinformatics* 8, 177.

- Somers, G. R., Hammet, F. M., Trute, L., Southey, M. C., & Venter, D. J. (1998). Expression of the P2Y₆ purinergic receptor in human T cells infiltrating inflammatory bowel disease. *Laboratory Investigation* 78, 1375–1383.
- Song, L., Risseuw, M. D. P., Karalic, I., Barrett, M. O., Brown, K. A., Harden, T. K., & van Calenbergh, S. (2014). Synthesis of extended uridine phosphonates derived from an allosteric P2Y₂ receptor ligand. *Molecules* 19, 4313–4325.
- Stachon, P., Peikert, A., Michel, N. A., Hergeth, S., Marchini, T., Wolf, D., ... Zirlik, A. (2014). P2Y₆ deficiency limits vascular inflammation and atherosclerosis in mice. *Arteriosclerosis, Thrombosis, and Vascular Biology* 34, 2237–2245.
- Steculorum, S. M., Paeger, L., Bremser, S., Evers, N., Hinze, Y., Idzko, M., ... Bruning, J. C. (2015). Hypothalamic UDP increases in obesity and promotes feeding via P2Y₆-dependent activation of AgRP neurons. *Cell* 162, 1404–1417.
- Steculorum, S. M., Timper, K., Engstrom Ruud, L., Evers, N., Paeger, L., Bremser, S., ... Bruning, J. C. (2017). Inhibition of P2Y₆ signaling in AgRP neurons reduces food intake and improves systemic insulin sensitivity in obesity. *Cell Reports* 18, 1587–1597.
- Suarez-Huerta, N., Pouillon, V., Boeynaems, J., & Robaye, B. (2001). Molecular cloning and characterization of the mouse P2Y₄ nucleotide receptor. *European Journal of Pharmacology* 416, 197–202.
- Suzuki, T., Namba, K., Tsuga, H., & Nakata, H. (2006). Regulation of pharmacology by hetero-oligomerization between A₁ adenosine receptor and P2Y₂ receptor. *Biochemical and Biophysical Research Communications* 351, 559–565.
- Tauber, J., Davitt, W. F., Bokosky, J. E., Nichols, K. K., Yerxa, B. R., Schaberg, A. E., ... Kellerman, D. J. (2004). Double-masked, placebo-controlled safety and efficacy trial of difafofosol tetrasodium (INS365) ophthalmic solution for the treatment of dry eye. *Cornea* 23, 784–792.
- Tosh, D. K., & Jacobson, K. A. (2013). Methanocarba ring as a ribose modification in ligands of G protein-coupled purine and pyrimidine receptors: Synthetic approaches. *MedChemComm* 2013, 619–630.
- Tosh, D. K., Yoo, L. S., Chinn, M., Hong, K., Kilbey, S. M., II, Barrett, M. O., ... Jacobson, K. A. (2010). Polyamidamine (PAMAM) dendrimer conjugates of “clickable” agonists of the A₃ adenosine receptor and coactivation of the P2Y₁₄ receptor by a tethered nucleotide. *Bioconjugate Chemistry* 21, 372–384.
- Tran, M. D. (2011). P2 receptor stimulation induces amyloid precursor protein production and secretion in rat cortical astrocytes. *Neuroscience Letters* 492, 155–159.
- Trujillo, K., Paoletta, S., Kiselev, E., & Jacobson, K. A. (2015). Molecular modeling of the human P2Y₁₄ receptor: A template for structure-based design of selective agonist ligands. *Bioorganic & Medicinal Chemistry* 23, 4056–4064.
- Tu, M. T., Luo, S. F., Wang, C. C., Chien, C. S., Chiu, C. T., Lin, C. C., & Yang, C. M. (2000). P2Y₂ (2) receptor-mediated proliferation of C(6) glioma cells via activation of Ras/Raf/MEK/MAPK pathway. *British Journal of Pharmacology* 129, 1481–1489.
- Tulapurkar, M. E., Laubinger, W., Nahum, V., Fischer, B., & Reiser, G. (2004). Subtype specific internalization of P2Y₁ and P2Y₂ receptors induced by novel adenosine 5'-O-(1-boranotriphosphate) derivatives. *British Journal of Pharmacology* 142, 869–878.
- Uhlén, M., Fagerberg, L., Hallström, B. M., Lindskog, C., Oksvold, P., Mardinoglu, A., ... Pontén, F. (2015). Proteomics. Tissue-based map of the human proteome. *Science* 347, 1260419.
- UniProt Consortium (2015). UniProt: A hub for protein information. *Nucleic Acids Research* 43, D204–12.
- Uratsujii, H., Tada, Y., Kawashima, T., Kamata, M., Hau, C. S., Asano, Y., ... Tamaki, K. (2012). P2Y₆ receptor signaling pathway mediates inflammatory responses induced by monosodium urate crystals. *Journal of Immunology* 188, 436–444.
- van Poecke, S., Barrett, M. O., Santhosh Kumar, T., Sinnaeve, D., Martins, J. C., Jacobson, K. A., ... van Calenbergh, S. (2012). Synthesis and P2Y₂(2) receptor agonist activities of uridine 5'-phosphonate analogues. *Bioorganic & Medicinal Chemistry* 20, 2304–2315.
- Vieira, R. P., Müller, T., Grimm, M., von Gernler, V., Vetter, B., Durk, T., ... Idzko, M. (2011). Purinergic receptor type 6 contributes to airway inflammation and remodeling in experimental allergic airway inflammation. *American Journal of Respiratory and Critical Care Medicine* 184, 215–223.
- Vigne, P., Pacaud, P., Urbach, V., Feolde, E., Breitmayer, J. P., & Frelin, C. (1996). The effect of PPADS as an antagonist of inositol (1,4,5)trisphosphate induced intracellular calcium mobilization. *British Journal of Pharmacology* 119, 360–364.
- Voogd, T. E., Vansterkenburg, E. L., Wilting, J., & Janssen, L. H. (1993). Recent research on the biological activity of suramin. *Pharmacological Reviews* 45, 177–203.
- Wan, H., Xie, R., Xu, J., He, J., Tang, B., Liu, Q., ... Dong, H. (2017). Anti-proliferative effects of nucleotides on gastric cancer via a novel P2Y₆/SOCE/Ca²⁺/beta-catenin pathway. *Scientific Reports* 7, 2459.
- Wang, S., Iring, A., Strilic, B., Albarrán Juárez, J., Kaur, H., Troidl, K., ... Offermanns, S. (2015). P2Y₂(2) and Gq/G(11) control blood pressure by mediating endothelial mechanotransduction. *The Journal of Clinical Investigation* 125, 3077–3086.
- Warny, M., Aboudola, S., Robson, S. C., Sevigny, J., Communi, D., Soltoff, S. P., & Kelly, C. P. (2001). P2Y₆ nucleotide receptor mediates monocyte interleukin-8 production in response to UDP or lipopolysaccharide. *The Journal of Biological Chemistry* 276, 26051–26056.
- Webb, T. E., Henderson, D., King, B. F., Wang, S., Simon, J., Bateson, A. N., ... Barnard, E. A. (1996). A novel G protein-coupled P2 purinoceptor (P2Y₃) activated preferentially by nucleoside diphosphates. *Molecular Pharmacology* 50, 258–265.
- Webb, T. E., Henderson, D. J., Roberts, J. A., & Barnard, E. A. (1998). Molecular cloning and characterization of the rat P2Y₄ receptor. *Journal of Neurochemistry* 71, 1348–1357.
- Welch, B. D., Carlson, N. G., Shi, H., Myatt, L., & Kishore, B. K. (2003). P2Y₂ receptor-stimulated release of prostaglandin E₂ by rat inner medullary collecting duct preparations. *American Journal of Physiology. Renal Physiology* 285, F711–711.
- Weyler, S., Baqi, Y., Hillmann, P., Kaulich, M., Hunder, A. M., Müller, I. A., & Müller, C. E. (2008). Combinatorial synthesis of anilinoanthraquinone derivatives and evaluation as non-nucleotide-derived P2Y₂ receptor antagonists. *Bioorganic & Medicinal Chemistry Letters* 18, 223–227.
- Wilden, P. A., Agazie, Y. M., Kaufman, R., & Halenda, S. P. (1998). ATP-stimulated smooth muscle cell proliferation requires independent ERK and PI3K signaling pathways. *The American Journal of Physiology* 275, H1209–15.
- Wildman, S. S., Unwin, R. J., & King, B. F. (2003). Extended pharmacological profiles of rat P2Y₂ and rat P2Y₄ receptors and their sensitivity to extracellular H⁺ and Zn²⁺ ions. *British Journal of Pharmacology* 140, 1177–1186.
- Wu, L., Oshima, T., Fukui, H., Watari, J., & Miwa, H. (2016). Adenosine triphosphate induces P2Y₂ activation and IL-8 release in human esophageal epithelial cells. *Journal of Gastroenterology and Hepatology* 32, 1341–1347.
- Xing, M., Post, S., Ostrom, R. S., Samardzija, M., & Insel, P. A. (1999). Inhibition of phospholipase A₂-mediated arachidonic acid release by cyclic AMP defines a negative feedback loop for P2Y receptor activation in Madin-Darby canine kidney D1 cells. *The Journal of Biological Chemistry* 274, 10035–10038.
- Xu, J., Chalimoniuk, M., Shu, Y., Simonyi, A., Sun, A. Y., Gonzalez, F. A., ... Sun, G. Y. (2003). Prostaglandin E₂ production in astrocytes: Regulation by cytokines, extracellular ATP, and oxidative agents. *Prostaglandins, Leukotrienes, and Essential Fatty Acids* 69, 437–448.
- Xu, J., Morinaga, H., Oh, D., Li, P., Chen, A., Talukdar, S., ... Kim, J. J. (2012). GPR105 ablation prevents inflammation and improves insulin sensitivity in mice with diet-induced obesity. *Journal of Immunology* 189, 1992–1999.
- Xu, J., Weng, Y. -I., Simonyi, A., Krugh, B. W., Liao, Z., Weisman, G. A., & Sun, G. Y. (2002). Role of PKC and MAPK in cytosolic PLA₂ phosphorylation and arachidonic acid release in primary murine astrocytes. *Journal of Neurochemistry* 83, 259–270.
- Yamane, M., Ogawa, Y., Fukui, M., Kamoi, M., Saijo-Ban, Y., Yaguchi, S., ... Tsubota, K. (2015). Long-term rebamipide and difafofosol in two cases of immune-mediated dry eye. *Optometry and Vision Science* 92, S25–32.
- Yerxa, B. R., Pendergast, W., Rideout, J. L., Picher, M., Boucher, R. C., & Stutts, M. J. (1999). Inspire Pharmaceuticals Inc. WO001999061012A2.
- Yerxa, B. R., Peterson, W., Rideout, J. L., & Pendergast, W. (2009). Inspire Pharmaceuticals Inc. US000008008274B2.
- Yerxa, B. R., Rideout, J. L., Pendergast, W., Shaver, S. R., Zhang, Z., Peterson, W., & Cowlen, M. (2000). Inspire Pharmaceuticals Inc. WO002001045691A2.
- Yerxa, B. R., Sabater, J. R., Davis, C. W., Stutts, M. J., Lang-Furr, M., Picher, M., ... Boucher, R. C. (2002). Pharmacology of INS37217 P(1)-(uridine 5')-P(4)-(2'-deoxycytidine 5') tetraphosphate, tetrasodium salt, a next-generation P2Y₂ receptor agonist for the treatment of cystic fibrosis. *The Journal of Pharmacology and Experimental Therapeutics* 302, 871–880.
- Yoshioka, K., Hosoda, R., Kuroda, Y., & Nakata, H. (2002). Hetero-oligomerization of adenosine A₁ receptors with P2Y₁ receptors in rat brains. *FEBS Letters* 531, 299–303.
- Yoshioka, K., Saitoh, O., & Nakata, H. (2001). Heteromeric association creates a P2Y₂-like adenosine receptor. *Proceedings of the National Academy of Sciences of the United States of America* 98, 7617–7622.
- Zamboni, A. C., Hughes, R. J., Meszaros, J. G., Wu, J. J., Torres, B., Brunton, L. L., & Insel, P. A. (2000). P2Y₂ receptor of MDCK cells: Cloning, expression, and cell-specific signaling. *American Journal of Physiology. Renal Physiology* 279, F1045–52.
- Zhang, D., Gao, Z. -G., Zhang, K., Kiselev, E., Crane, S., Wang, J., ... Wu, B. (2015). Two disparate ligand-binding sites in the human P2Y₁ receptor. *Nature* 520, 317–321.
- Zhang, J., Zhang, K., Gao, Z. -G., Paoletta, S., Zhang, D., Han, G. W., ... Zhao, Q. (2014). Agonist-bound structure of the human P2Y₁₂ receptor. *Nature* 509, 119–122.
- Zhang, J. -L., Liu, Y., Yang, H., Zhang, H. -Q., Tian, X. -X., & Fang, W. -G. (2017). ATP-P2Y₂-beta-catenin axis promotes cell invasion in breast cancer cells. *Cancer Science* 108, 1318–1327.
- Zhang, K., Zhang, J., Gao, Z. -G., Zhang, D., Zhu, L., Han, G. W., ... Zhao, Q. (2014). Structure of the human P2Y₁₂ receptor in complex with an antithrombotic drug. *Nature* 509, 115–118.
- Zhang, Z., Wang, Z., Ren, H., Yue, M., Huang, K., Gu, H., ... Qian, M. (2011). P2Y₆ agonist uridine 5'-diphosphate promotes host defense against bacterial infection via monocyte chemoattractant protein-1-mediated monocytes/macrophages recruitment. *Journal of Immunology* 186, 5376–5387.
- Zimmermann, H. (2016). Extracellular ATP and other nucleotides-ubiquitous triggers of intercellular messenger release. *Purinergic Signalling* 12, 25–57.
- Zimmermann, H., Zebisch, M., & Sträter, N. (2012). Cellular function and molecular structure of ecto-nucleotidases. *Purinergic Signalling* 8, 437–502.
- Zizzo, M. G., Mastropalo, M., Grahlert, J., Mule, F., & Serio, R. (2012). Pharmacological characterization of uracil nucleotide-prefering P2Y receptors modulating intestinal motility: a study on mouse ileum. *Purinergic Signalling* 8, 275–285.