

0327 Distinct patterns of EEG-EMG-coherence in various stages of disease severity in patients with sleep-disordered breathing [Abstract]

K. Bahr, H. Gouveris, T. Huppertz, E. Martin, C. Matthias, Muthuraman
Muthuraman

Angaben zur Veröffentlichung / Publication details:

Bahr, K., H. Gouveris, T. Huppertz, E. Martin, C. Matthias, and Muthuraman Muthuraman. 2018. "0327 Distinct patterns of EEG-EMG-coherence in various stages of disease severity in patients with sleep-disordered breathing [Abstract]." *Sleep* 41 (Suppl. 1): A125.
<https://doi.org/10.1093/sleep/zsy061.326>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



0327

DISTINCT PATTERNS OF EEG-EMG-COHERENCE IN VARIOUS STAGES OF DISEASE SEVERITY IN PATIENTS WITH SLEEP-DISORDERED BREATHING

Bahr K¹, Gouveris H¹, Huppertz T¹, Martin E¹, Matthias C¹, Muthuraman M¹

¹University Medical Center Mainz, Mainz, GERMANY, ²University Medical Center Mainz, Mainz, GERMANY, ³University Medical Center Mainz, Mainz, GERMANY

Introduction: We investigated whether using EEG-EMG-coherence (EEC) as a feature fed to a support vector machine (SVM) algorithm may allow staging of disease severity among sleep-disordered-breathing (SDB) patients.

Methods: EEG-EMG-coherence data resulted by applying a multi-taper processing for estimating the power spectrums separately and calculating the coherence on raw C3-/C4-EEG- and EMG- chin data of polysomnographic (PSG) recordings of 102 SDB patients (33 female; age: 53, \pm 12,4 yrs) acquired on two consecutive nights in each patient. Four epochs (30 seconds each, classified manually by AASM 2012-criteria) of each sleep stage were marked (in total 1632 epochs / night) and were included in the analysis. After multitaper processing, EEC values were fed to a SVM algorithm to classify SDB disease severity based on respiratory disturbance index (RDI). Twenty patients had a mild ($RDI \geq 10 / h$ and $<15 / h$), 30 patients had a moderate ($RDI \geq 15 / h$ and $<30 / h$) and 27 patients had a severe OSA ($RDI \geq 30 / h$). Twenty five patients had a $RDI <10 / h$. The AUC (area under the curve) value was calculated for each receiver operator characteristic (ROC) curve.

Results: EEG-EMG coherence values could distinguish between SDB-patients without OSA and OSA patients of the above three severity groups using an SVM algorithm. Using PSG data of the first night the AUC for mild OSA was 0.602 ($p = 0.032$), in moderate OSA the AUC was 0.6319 ($p = 0.021$), and in severe OSA the AUC was 0.823 ($p < 0.001$). On the second night, in mild OSA the AUC was 0.616 ($p = 0.024$), in moderate OSA the AUC was 0.659 ($p = 0.003$), and in severe OSA the AUC was 0.823 ($p < 0.001$).

Conclusion: Grading disease severity in SDB patients can be performed using PSG-based multitaper-processed EEC values processed with a SVM algorithm. No clinically relevant first night- effect was shown in EEC values. EEG-EMG-coherence appears to be a robust marker for SDB severity classification.