

The derivative map for diffeomorphism of disks: an example

Diarmuid Crowley, Thomas Schick, Wolfgang Steimle

Angaben zur Veröffentlichung / Publication details:

Crowley, Diarmuid, Thomas Schick, and Wolfgang Steimle. 2023. "The derivative map for diffeomorphism of disks: an example." *Geometry & Topology* 27 (9): 3699–3713.
<https://doi.org/10.2140/gt.2023.27.3699>.



Geometry & Topology

Volume 27 (2023)

The derivative map for diffeomorphism of disks: an example

DIARMUID CROWLEY

THOMAS SCHICK

WOLFGANG STEIMLE



The derivative map for diffeomorphism of disks: an example

DIARMUID CROWLEY

THOMAS SCHICK

WOLFGANG STEIMLE

We prove that the derivative map $d: \text{Diff}_\partial(D^k) \rightarrow \Omega^k \text{SO}_k$, defined by taking the derivative of a diffeomorphism, can induce a nontrivial map on homotopy groups. Specifically, for $k = 11$ we prove that the following homomorphism is nonzero:

$$d_*: \pi_5 \text{Diff}_\partial(D^{11}) \rightarrow \pi_5 \Omega^{11} \text{SO}_{11} \cong \pi_{16} \text{SO}_{11}.$$

As a consequence we give a counterexample to a conjecture of Burghlelea and Lashof by giving an example of a nontrivial vector bundle E over a sphere which is trivial as a topological \mathbb{R}^k -bundle (the rank of E is $k = 11$ and the base sphere is S^{17}).

The proof relies on a recent result of Burklund and Senger which determines the homotopy 17-spheres bounding 8-connected manifolds, the plumbing approach to the Gromoll filtration due to Antonelli, Burghlelea and Kahn, and an explicit construction of low-codimension embeddings of certain homotopy spheres.

57R50, 57S05; 57R60

1 Introduction

The derivative map

$$d: \text{Diff}_\partial(D^k) \rightarrow \text{Map}((D^k, \partial D^k), (\text{SO}_k, \text{Id})) \simeq \Omega^k \text{SO}_k, \quad f \mapsto (x \mapsto D_x f),$$

is a basic invariant of the diffeomorphism group of the k -disk; in fact the first-order approximation in the embedding calculus approach to the diffeomorphism group. While $d_{\mathbb{Q}}: \text{Diff}_\partial(D^k)_{\mathbb{Q}} \rightarrow (\Omega^k \text{SO}_k)_{\mathbb{Q}}$, the rationalisation of d , is nullhomotopic, as we explain in Section 3, much less is known about the derivative map d integrally. For example, to the best of our knowledge, it was not yet known whether the map induced by d on homotopy groups,

$$d_*: \pi_i \text{Diff}_\partial(D^k) \rightarrow \pi_{i+k} \text{SO}_k,$$

was ever nontrivial. Burghelea and Lashof showed that d_* vanishes for $i = 0, 1$. At odd primes p , they also showed that $d_* = 0$ provided $i < k - 3$ and they made a conjecture equivalent to the claim that this holds for $p = 2$ as well [6, Conjecture, page 40]. Burghelea and Lashof also report A'Campo informing them about a proof that $d_* = 0$ for $i = 2$ (however, a written proof has not appeared).

Using smoothing theory, or an explicit geometric construction we introduce here, the map d_* admits an interpretation as describing the normal bundle of certain homotopy spheres embedded in euclidean space. Combining this interpretation with recent results of Burklund and Senger and the refined plumbing construction of Antonelli, Burghelea and Kahn, we obtain a counterexample to the conjecture of Burghelea and Lashof.

In more detail, in [3; 4] Antonelli, Burghelea and Kahn constructed families of diffeomorphisms of the disk using a pairing

$$\sigma: \pi_p \mathrm{SO}_{q-a} \otimes \pi_q \mathrm{SO}_{p-b} \rightarrow \pi_{a+b+1} \mathrm{Diff}_{\partial}(D^{p+q-a-b-1})$$

for $0 \leq a \leq q$ and $0 \leq b \leq p$, refining Milnor's plumbing pairing; see below. Now $\pi_8 \mathrm{SO}_6 \cong \mathbb{Z}/24$ (see [12, page 162]) and we have:

Theorem 1.1 *Let $\xi \in \pi_8 \mathrm{SO}_6 \cong \mathbb{Z}/24$ be a generator. The image of $\sigma(\xi, \xi)$ under the derivative map*

$$d_*: \pi_5 \mathrm{Diff}_{\partial}(D^{11}) \rightarrow \pi_{16} \mathrm{SO}_{11}$$

is nonzero.

Using Morlet's smoothing theory isomorphism, the derivative map d_* on π_k is identified with the boundary map

$$\partial: \pi_{n+k+1} \mathrm{PL}_k / O_k \rightarrow \pi_{n+k} \mathrm{SO}_k$$

of the fibration sequence $\mathrm{SO}_k \rightarrow \mathrm{SPL}_k \rightarrow \mathrm{PL}_k / O_k$ (and this allows for our interpretation of [6, Conjecture, page 40] in terms of the derivative map). We conclude that the map $\mathrm{SO}_{11} \rightarrow \mathrm{SPL}_{11}$ is not injective on π_{16} . More specifically, if $\tau_{11}: S^{11} \rightarrow B\mathrm{SO}_{11}$ represents the tangent bundle of the 11-sphere and $f: S^{17} \rightarrow S^{11}$ represents the unique nontrivial homotopy class (see [21, Proposition 5.11]), we have:

Corollary 1.2 *The pullback $f^* \tau_{11}$ is a nontrivial vector bundle which becomes trivial as an \mathbb{R}^{11} -bundle, even when considered as a bundle with structure group SPL_{11} .*

To the best of our knowledge, this is the first example of a nontrivial vector bundle over a sphere which is known to be trivial as a topological \mathbb{R}^n -bundle. Milnor famously gave examples of nontrivial vector bundles over Moore spaces, for example the Moore space $M(\mathbb{Z}/7, 7) = S^7 \cup_7 D^8$, which are trivial as \mathbb{R}^n -bundles [17, Lemma 9.1]. These examples are stable bundles over 4-connected spaces and so the vector bundles are trivial as piecewise linear bundles too.

Acknowledgements

We would like to thank Alexander Kupers for helpful comments on an earlier draft, as well as two referees for helpful comments.

2 Proofs

In this section we give the proofs of Theorem 1.1 and Corollary 1.2. We first recall Gromoll's map $A = C \circ \lambda$ from [9],

$$A: \pi_{n-k} \text{Diff}_\partial(D^k) \xrightarrow{\lambda} \pi_0 \text{Diff}_\partial(D^n) \xrightarrow{C} \Theta_{n+1},$$

where Θ_{n+1} is the group of homotopy $(n+1)$ -spheres. The first map λ includes fibrewise diffeomorphisms of $D^{n-k} \times D^k$ into all diffeomorphisms, and the second C uses a diffeomorphism of $D^n \subset S^n$ as a datum to clutch two $(n+1)$ -disks and make a homotopy sphere.¹

Lemma 2.1 *For any $[\psi] \in \pi_{n-k} \text{Diff}_\partial(D^k)$, the homotopy sphere $A([\psi]) \in \Theta_{n+1}$ admits an embedding into \mathbb{R}^{n+k+1} whose normal bundle is classified (up to possible sign) by $d_*([\psi]) \in \pi_n \text{SO}_k \cong \pi_{n+1} \text{BSO}_k$.*

We will offer two proofs of this result; one by an explicit geometric construction and a more abstract one by the classification of smoothings through Rourke and Sanderson's theory of block bundles.

Next we recall that in [16, Section 1], Milnor constructed exotic spheres by plumbing linear disk bundles and taking the boundary sphere; this construction gives rise to a pairing

$$\sigma_M: \pi_p \text{SO}_q \otimes \pi_q \text{SO}_p \rightarrow \Theta_{p+q+1}.$$

¹The map C is denoted by Σ in [8].

By [3, Proposition 3.1], the pairing of Antonelli, Burghlea and Kahn refines this pairing in the sense that we have a commutative diagram

$$(1) \quad \begin{array}{ccc} \pi_p \mathrm{SO}_{q-a} \otimes \pi_q \mathrm{SO}_{p-b} & \xrightarrow{\sigma} & \pi_{a+b+1} \mathrm{Diff}_\partial(D^{p+q-a-b-1}) \\ \downarrow & & \downarrow A \\ \pi_p \mathrm{SO}_q \otimes \pi_q \mathrm{SO}_p & \xrightarrow{\sigma_M} & \Theta_{p+q+1} \end{array}$$

where the map on the left is the tensor product of the canonical stabilisations. We now consider the homotopy 17–sphere

$$\Sigma_{\xi, \xi} := A(\sigma(\xi, \xi)),$$

recalling that $\xi \in \pi_8 \mathrm{SO}_6 \cong \mathbb{Z}/24$ denotes a generator. By the commutativity of (1) and the definition of σ_M , $\Sigma_{\xi, \xi}$ is the boundary of an 8–connected compact 18–manifold and so by a recent result of Burklund and Senger [7, Theorem 1.4] its image under the normal invariant map $\Theta_{17} \rightarrow \mathrm{coker}(J_{17})$ must be either 0 or $[\eta\eta_4]$. We will show:

Lemma 2.2 *The homotopy sphere $\Sigma_{\xi, \xi}$ represents $[\eta\eta_4] \in \mathrm{coker}(J_{17})$.*

We deduce from this that every embedding $\Sigma_{\xi, \xi} \hookrightarrow S^{28}$ has a nontrivial normal bundle. Indeed, recall that the map $\Theta_{17} \rightarrow \mathrm{coker}(J_{17})$ is obtained by embedding a homotopy 17–sphere into some euclidean space with trivial normal bundle, and performing the Pontryagin–Thom collapse as to obtain an element in π_{17}^s which is well-defined modulo the image of J . Now, assuming by contradiction that $\Sigma_{\xi, \xi}$ embeds into S^{28} with trivial normal bundle, then $[\eta\eta_4]$ would have a representative in $\pi_{28} S^{11}$. However, this contradicts the computations of Toda [21, Theorem 12.17 and Proposition 12.20] on the stabilisation map $\pi_{28} S^{11} \rightarrow \pi_{17}^s$, which we display below:

$$\begin{array}{ccc} \pi_{28} S^{11} = \mathbb{Z}/2((\eta^2 \rho)_{11}) \oplus \mathbb{Z}/2((\mu_{17})_{11}) \oplus \mathbb{Z}/2((\nu\kappa)_{11}) & & \\ \downarrow & \swarrow & \swarrow \\ \pi_{17}^s = \mathbb{Z}/2(\eta^2 \rho) \oplus \mathbb{Z}/2(\mu_{17}) \oplus \mathbb{Z}/2(\nu\kappa) \oplus \mathbb{Z}/2(\eta\eta_4) & & \end{array}$$

Here the notation is such that an element $(\delta)_{11}$ stabilises to δ and the stable class $\eta^2 \rho$ generates $\mathrm{im}(J_{17}: \pi_{17}(\mathrm{SO}) \rightarrow \pi_{17}^s)$. Lemma 2.1 then implies that $d_*(\sigma(\xi, \xi)) \neq 0$, which concludes the proof of Theorem 1.1, modulo Lemmas 2.1 and 2.2.

To prove Corollary 1.2 we note that by Lemma 2.1 the normal bundle $\nu(\Sigma_{\xi, \xi} \subset S^{28})$ has clutching function $\pm d_*(\sigma(\xi, \xi)) \in \pi_{16} \mathrm{SO}_{11}$ and $d_*(\sigma(\xi, \xi)) \neq 0$ by Theorem 1.1.

Moreover, $d_*(\sigma(\xi, \xi))$ maps to $0 \in \pi_{16}\text{SPL}_{11}$; this is explained following Theorem 1.1 using the exact sequence $\pi_{*+1}(\text{PL}_{11}/O_{11}) \xrightarrow{d_*} \pi_*(\text{SO}_{11}) \longrightarrow \pi_*(\text{SPL}_{11})$. Now by Antonelli [2], the normal bundle of every homotopy 17–sphere embedded in euclidean space in codimension 12 is zero. Hence

$$\nu(\Sigma_{\xi, \xi} \subset S^{28}) \in \ker(\pi_{17}BSO_{11} \rightarrow \pi_{17}BSO_{12}) = \text{im}(\pi_{17}S^{11} \rightarrow \pi_{17}BSO_{11}),$$

where the last map is induced by the classifying map of the tangent bundle of the 11–sphere.

It remains to prove Lemmas 2.1 and 2.2.

Proof of Lemma 2.1 Choose a smooth map $\psi: D^{n-k} \times D^k \rightarrow D^k$ representing the class $[\psi] \in \pi_{n-k}(\text{Diff}_{\partial}(D^k))$, ie for $x \in D^{n-k}$ we have that $\psi_x := \psi(x, -) \in \text{Diff}_{\partial}(D^k)$, and $\psi_x = \text{Id}_{D^k}$ for $x \in \partial D^{n-k}$. Then $\lambda([\psi])$ is represented by

$$\Psi: D^k \times D^{n-k} \rightarrow D^k \times D^{n-k}, \quad (x, y) \mapsto (x, \psi(x, y)),$$

and $A([\psi])$ is represented by the homotopy sphere Σ_{Ψ}^{n+1} obtained by gluing two copies of D^{n+1} along the boundary using the diffeomorphism Ψ . Note also that the image of $[\psi]$ under the derivative map is represented by $d\psi: D^{n-k} \times D^k \rightarrow \text{Gl}_k(\mathbb{R})$ with $d\psi(x, y) = D_y\psi_x$.

For technical reasons, we actually assume without loss of generality that the maps are the identity maps in a neighbourhood of the boundaries.

We construct an explicit embedding $\iota_{\psi}: \Sigma_{\Psi}^{n+1} \hookrightarrow S^{n+k+1}$ of Σ_{Ψ}^{n+1} , compute the normal bundle of this embedding and show explicitly that it is obtained by clutching with $d\psi$.

As might be expected, given that all our data is on disks (and trivial near the boundary of the disks), we actually produce an interesting embedding ι_{ψ} of $D^{n+1} = D^{n-k} \times D^k \times [0, 1]$ into $D^{n+k+1} = D^{n-k} \times D^k \times D^k \times [0, 1]$, which has standard form near the boundary, and then obtain an embedding of Σ_{Ψ}^{n+1} by gluing with a standard embedding of D^{n+1} into D^{n+k+1} in the appropriate way.

The desired embedding ι_{ψ} is explicitly given by

$$\begin{aligned} \iota_{\psi}: D^{n-k} \times D^k \times [0, 1] &\rightarrow D^{n-k} \times D^k \times D^k \times [0, 1], \\ (x, y, t) &\mapsto (x, \alpha(t)y, \beta(t)\psi_x(y), t). \end{aligned}$$

Here, $\alpha, \beta: [0, 1] \rightarrow [0, 1]$ are smooth maps such that $\alpha(t) = 1$ for $t < 0.6$ and $\alpha(t) = 0$ for $t > 0.9$, and $\beta(t) = \alpha(1 - t)$.

This is evidently a smooth embedding whose image we denote by S_ψ , and we let $\partial S_\psi = \iota_\psi(\partial D^{n+1})$. Then $\partial S_\psi \subset \partial D^{n+k+1}$: to see this, observe that if either the x - or the t -coordinate is in the boundary, then the first or fourth coordinate of the image point is so, too. For each $t \in [0, 1]$, then $\alpha(t) = 1$ or $\beta(t) = 1$. If $y \in \partial D^k$, then therefore either the second or the third component of the image point is in the boundary (or both). As $\partial(D^{n-k} \times D^k \times [0, 1])$ is the union of those points with at least one component in the boundary, this proves the claim.

We also note that the subset $\partial S_\psi \subset \partial D^{n+k+1}$ is in fact independent of ψ (as ψ is fixed to be the identity map near the boundary). Let us identify this image set ∂S_ψ with $S^n = \partial(D^{n-k} \times D^k \times [0, 1])$ via the restriction of ι_{Id} to $\partial(D^{n-k} \times D^k \times [0, 1])$.

Then $\iota_\psi|: \partial(D^{n-k} \times D^k \times [0, 1]) \rightarrow \partial(D^{n-k} \times D^k \times D^k \times [0, 1])$ is supported on the disk $D^{n-k} \times D^k \times \{1\}$, where it is given by Ψ . Therefore, we can glue two copies of $D^{n-k} \times D^k \times D^k \times [0, 1]$ along the boundary by the identity map to obtain S^{n+k+1} , and the embeddings ι_ψ in one copy and ι_{Id} in the other glue together to form the desired embedding of Σ_Ψ^{n+1} into S^{n+k+1} .

Strictly speaking, one has to round the corners off to get an actual smooth embedding. This can easily be achieved, as ψ is the identity in a neighbourhood of the boundaries. We omit spelling out the somewhat cumbersome details.

It remains to compute the normal bundle of the embedding. To do this, we first compute the differential

$$D\iota_\psi: (D^{n-k} \times D^k \times [0, 1]) \times (\mathbb{R}^{n-k} \oplus \mathbb{R}^k \oplus \mathbb{R}) \rightarrow S_\psi \times (\mathbb{R}^{n-k} \oplus \mathbb{R}^k \oplus \mathbb{R}^k \oplus \mathbb{R})$$

to be given in each fibre by

$$D_{(x,y,t)}\iota_\psi = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \alpha(t) & \alpha'(t)y \\ \beta(t)\partial_x\psi & \beta(t)d\psi & \beta'(t)\psi_x(y) \\ 0 & 0 & 1 \end{pmatrix}.$$

We obtain an explicit trivialisation of the normal bundle of this embedding via the fibrewise linear map covering ι_ψ ,

$$\begin{aligned} \nu: D^{n-k} \times D^k \times [0, 1] \times \mathbb{R}^k &\rightarrow S_\psi \times (\mathbb{R}^{n-k} \oplus \mathbb{R}^k \oplus \mathbb{R}^k \oplus \mathbb{R}), \\ \nu_{(x,y,t)} &= \begin{pmatrix} 0 \\ -\beta(t)(d\psi)^{-1} \\ \alpha(t) \\ 0 \end{pmatrix}. \end{aligned}$$

To observe that this really describes the normal bundle, for dimension reasons we just have to check that the image of v intersects the tangent bundle of S_ψ , ie the image of $D\iota_\psi$, trivially. It is clear that $v(v)$ can only be equal to a tangent vector of the form $(0, \alpha(t)w, \beta(t) d\psi(w), 0)$ for $w \in \mathbb{R}^k$. This implies $\alpha(t)v = \beta(t) d\psi(w)$ and $-\beta(t)v = \alpha(t) d\psi(w)$; the two equations imply $\alpha(t)^2 v = -\beta(t)^2 v$ and finally (as $\alpha(t)^2 + \beta(t)^2 > 0$) then $v = 0$ and then also $w = 0$. It follows that the image of v represents the normal bundle of S_ψ in D^{n+k+1} .

For the other half-disk which produces the embedding of Σ_ψ^{n+1} into S^{n+k+1} , we obtain a trivialisation of the normal bundle by the same recipe, replacing ψ by Id . We observe then that we obtain the global normal bundle by gluing these two explicitly chosen normal subbundles of TD^{n+k+1} along the boundary, where they coincide. The trivialisations differ precisely on the half-disk $\iota(D^{n-k} \times D^k \times \{1\})$, and there they differ by the derivative map $d\psi$. On the other half-disk, the two trivialisations coincide.

Consequently, the normal bundle of the embedding ι_ψ is obtained by clutching with $d\psi$, precisely as claimed, and the lemma is proved. \square

Remark 2.3 It is tempting to hope that the explicit geometric construction of d_* as the normal bundle of the embedding ι_ψ can be used to get some new information about d_* . On the other hand, the information obtained by the formulas in the proof given above seems rather limited. At least in the case where ψ lies in the image of σ , we present Conjecture 3.1 below on $d_* \circ \sigma$.

Proof of Lemma 2.2 Let A_8^{18} denote the group of bordism classes relative boundary of 8-connected 18-manifolds with boundary a homotopy sphere, which are defined in [23, Section 17]. Specifically, elements of A_8^{18} are represented by compact oriented 8-connected 18-manifolds W with boundary a homotopy sphere, and W_1 is bordant to W_2 if there is an h -cobordism Y between their boundaries such that the closed manifold $W_1 \cup Y \cup -W_2$ bounds an 8-connected 19-manifold. According to [23, Section 17] and [22, Theorem 2(5)], we have an isomorphism

$$(2) \quad A_8^{18} \rightarrow \mathbb{Z}/2 \oplus \mathbb{Z}/2, \quad [W] \mapsto (\Phi(\varphi_W), \varphi_W(\hat{\chi}_W)).$$

Here $\varphi_W: H_9(W; \mathbb{Z}) \rightarrow \mathbb{Z}/2$ is a quadratic refinement of the mod 2 intersection form defined as follows. By [22, Lemma 2], representing an integral homology class by an embedded sphere and taking its normal bundle gives rise to a quadratic map

$$\alpha_W: H_9(W; \mathbb{Z}) \rightarrow \pi_8 \text{SO}_9.$$

Since the stabilisation map $S: \pi_8 \mathrm{SO}_9 \rightarrow \pi_8 \mathrm{SO} = \mathbb{Z}/2$ is split surjective with kernel $\mathbb{Z}/2$, from α_W we obtain a quadratic map $\varphi_W: H_9(W; \mathbb{Z}) \rightarrow \mathbb{Z}/2$ with values in $\mathbb{Z}/2 = \ker(S)$ by fixing a splitting of $\pi_8 \mathrm{SO}_9$. The first component of (2) is the Arf invariant of φ_W and we next define the second component. Let $S\alpha_W: H_9(W; \mathbb{Z}) \rightarrow \mathbb{Z}/2 = \pi_8(\mathrm{SO})$ be the composition of α_W with the stabilisation map S above. Using [22, Lemma 2] again, we see that $S\alpha_W$ is a homomorphism. Define $\chi_W \in H_9(W; \mathbb{Z}/2)$ to be the Poincaré dual of

$$S\alpha_W \in \mathrm{Hom}(H_9(W; \mathbb{Z}), \mathbb{Z}/2) = H^9(W; \mathbb{Z}/2) \cong H^9(W, \partial W; \mathbb{Z}/2).$$

The second component of (2) is given by evaluating φ_W on any integral lift $\hat{\chi}_W$ of χ_W .

Let $S^3(\xi) \in \pi_8 \mathrm{SO}_9$ be the image of $\xi \in \pi_8 \mathrm{SO}_6$ under the inclusion $\mathrm{SO}_6 \rightarrow \mathrm{SO}_9$. By the commutativity of (1), $\Sigma_{\xi, \xi}$ is the boundary of the Milnor plumbing W of $S^3(\xi) \in \pi_8 \mathrm{SO}_9$ with itself, and we compute $\varphi_W(\hat{\chi}_W)$ as follows: with $H_9(W; \mathbb{Z}) = \mathbb{Z}(x) \oplus \mathbb{Z}(y)$ the normal bundles obtained from representing x and y by embeddings are both given by $S^3(\xi)$. We conclude that $\varphi_W(x) = \varphi_W(y)$. Moreover, we may use that in this basis the intersection form λ_W of W has matrix

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Furthermore, $S^3(\xi)$ stabilises to a generator of $\pi_8 \mathrm{SO}$, by Lemma 2.4 below. Thus, $S\alpha_W$ maps both x and y to a generator and so we may take $\hat{\chi}_W = x + y$. We now compute that

$$\varphi_W(\hat{\chi}_W) = \varphi_W(x + y) = \underbrace{\varphi_W(x) + \varphi_W(y)}_{=0} + \rho_2(\lambda_W(x, y)) = 1,$$

where ρ_2 denotes reduction mod 2.

Now, taking the homotopy sphere on the boundary defines a homomorphism

$$(3) \quad \partial: A_8^{18} \rightarrow \Theta_{17}.$$

From the short exact sequence

$$0 \rightarrow bP_{18}(= \mathbb{Z}/2) \rightarrow \Theta_{17} \rightarrow \mathrm{coker}(J_{17}) \rightarrow 0$$

and [7, Theorem 1.4], we see that the image of the map ∂ from (3) consists of precisely 4 different elements, so the map ∂ is injective. Each of the bP -spheres is the boundary of a manifold P which satisfies $S\alpha_P = 0$ and therefore $\varphi_P(\hat{\chi}_P) = 0$ and it follows that the element W from above must map under ∂ to a non- bP -sphere, which then represents $[\eta\eta_4]$ in view of [7, Theorem 1.4]. \square

Lemma 2.4 *The map $\mathbb{Z}/24 \cong \pi_8 \mathrm{SO}_6 \rightarrow \pi_8 \mathrm{SO} \cong \mathbb{Z}/2$ is surjective.*

Proof By [14, Theorem 1.4], $(\mathbb{Z}/2)^3 \cong \pi_8 \mathrm{SO}_8 \rightarrow \pi_8 \mathrm{SO} \cong \mathbb{Z}/2$ is onto and therefore has a kernel of 4 elements. (We refer to [12] for the computation of the relevant homotopy groups.) On the other hand, $(\mathbb{Z}/2)^2 \cong \pi_8 \mathrm{SO}_7 \rightarrow \pi_8 \mathrm{SO}_8$ is injective (its cokernel injects into $\pi_8(S^7) \cong \mathbb{Z}/2$) and so has an image of 4 elements. These two subgroups do not coincide: Since the maximal number of pointwise linearly independent vector fields on S^9 is 1 [1, Theorem 1.1], the tangent bundle of S^9 defines an element in $\pi_8 \mathrm{SO}_8$ that is not in the image of $\pi_8 \mathrm{SO}_7$ but maps to $0 \in \pi_8 \mathrm{SO}$.

Therefore, $(\mathbb{Z}/2)^2 \cong \pi_8 \mathrm{SO}_7 \rightarrow \pi_8 \mathrm{SO}$ is surjective and has a kernel of precisely two elements; similarly the image of $\mathbb{Z}/24 \cong \pi_8 \mathrm{SO}_6 \rightarrow \pi_8 \mathrm{SO}_7 \cong (\mathbb{Z}/2)^2$ consists of precisely two elements (its cokernel injects into $\pi_8 S^6 \cong \mathbb{Z}/2$), and we are left to show that these two subgroups do not agree. To see this, we consider the element $a := (2\gamma)_7 \eta_7$ where $(2\gamma)_7$ is a generator of $\mathbb{Z} \cong \pi_7 \mathrm{SO}_7$ and $\eta_7: S^8 \rightarrow S^7$ is the nontrivial class: By [14, Theorem 1.4], $(2\gamma)_7$ stabilises to an element divisible by 2 and so a is in the kernel of the stabilisation; and it does not lift to $\pi_8 \mathrm{SO}_6$ by the commutativity of the following diagram with exact rows:

$$\begin{array}{ccccc} \pi_7 \mathrm{SO}_7 & \longrightarrow & \pi_7 S^6 & \longrightarrow & \pi_6 \mathrm{SO}_6 (= 0) \\ \downarrow \eta_7 & & \cong \downarrow \eta_7 & & \\ \pi_8 \mathrm{SO}_6 & \longrightarrow & \pi_8 \mathrm{SO}_7 & \longrightarrow & \pi_8 S^6 \end{array} \quad \square$$

We conclude this section by giving the promised second proof of Lemma 2.1. To this end we recall from [18, Section 6] that a smoothing of S^{n+1} in S^{n+k+1} consists of a smooth manifold W and a PL homeomorphism $H: W \rightarrow S^{n+k+1}$, such that $\Sigma := H^{-1}(S^{n+1}) \subset W$ is a smooth submanifold, and such that H is concordant to the identity smoothing of S^{n+k+1} ; and recall the group \mathfrak{d}_{n+1}^k of concordance classes of such smoothings.² We note that up to diffeomorphism, W is a standard sphere mapping to S^{n+k+1} by a PL homeomorphism concordant to the identity, so that elements of \mathfrak{d}_{n+1}^k are represented by PL homeomorphisms $H: S^{n+k+1} \rightarrow S^{n+k+1}$ which are concordant to the identity (ie orientation-preserving). Note also that Σ is a homotopy $(n+1)$ -sphere, oriented through the PL homeomorphism $h := H|_{\Sigma}$, which is smoothly embedded into S^{n+k+1} .

²The group \mathfrak{d}_{n+1}^k is denoted by Γ_{n+1}^k in [18]. We have used different notation, to avoid confusion with the notation Γ_k^{n+1} for the subgroups of the Gromoll filtration.

There are two obvious homomorphisms out of \mathfrak{d}_{n+1}^k ,

$$\Theta_{n+1} \xleftarrow{F} \mathfrak{d}_{n+1}^k \xrightarrow{\nu} \pi_{n+1} BSO_k,$$

the left one mapping the class of H to the diffeomorphism class of Σ , and the right one to the classifying map of the normal bundle of $\Sigma \subset S^{n+k+1}$ (where, as usual, we identify a homotopy sphere up to homotopy equivalence with a standard sphere using the given orientation). Then, Lemma 2.1 is clearly implied by the following result:

Lemma 2.5 *There exists a group homomorphism $B: \pi_{n-k} \text{Diff}_{\partial}(D^k) \rightarrow \mathfrak{d}_{n+1}^k$ such that the following diagram commutes up to possible signs:*

$$\begin{array}{ccccc} \pi_{n-k} \text{Diff}_{\partial}(D^k) & \xrightarrow{d_*} & \pi_n \text{SO}_k & & \\ \swarrow A & \downarrow B & \downarrow \cong & & \\ \Theta_{n+1} & \xleftarrow{F} \mathfrak{d}_{n+1}^k & \xrightarrow{\nu} & \pi_{n+1} BSO_k & \end{array}$$

Proof We recall the homotopy equivalence

$$M_k: \text{Diff}_{\partial}(D^k) \rightarrow \Omega^{k+1} \text{PL}_k / \text{SO}_k$$

of Morlet (see [5, Theorem 4.4]) and consider the diagram

$$(4) \quad \begin{array}{ccccccc} \pi_0 \text{Diff}_{\partial}(D^n) & \xleftarrow{\lambda} & \pi_{n-k} \text{Diff}_{\partial}(D^k) & \xrightarrow{d_*} & \pi_n \text{SO}_k & & \\ & & \downarrow (M_k)_* & & \parallel & & \\ & & \pi_{n+1} \text{PL}_k / O_k & \xrightarrow{\partial} & \pi_n \text{SO}_k & & \\ & \swarrow S & \downarrow \tilde{i} & & \parallel & & \\ \pi_{n+1} \text{PL} / O & \xleftarrow{\tilde{S}} & \pi_{n+1} \widetilde{\text{PL}}_k / O_k & \xrightarrow{\partial} & \pi_n \text{SO}_k & & \\ \uparrow \cong \uparrow \Psi & & \uparrow \cong & & \uparrow \cong & & \\ \Theta_{n+1} & \xleftarrow{F} & \mathfrak{d}_{n+1}^k & \xrightarrow{\nu} & \pi_{n+1} BSO_k & & \end{array}$$

Here Ψ is the map which sends a homotopy sphere Σ to the element represented by the tangent PL microbundle of the mapping cylinder $\text{cyl}(h: \Sigma \rightarrow S^{n+1})$ of an orientation-preserving PL homeomorphism h , along with its linear structure induced by the smooth structure of Σ on the Σ end of the cylinder and its canonical trivialisation at the S^{n+1} end. The map Ψ is an isomorphism by surgery theory; see eg [15, Theorem 6.48]. The map $\mathfrak{d}_{n+1}^k \rightarrow \pi_{n+1} \widetilde{\text{PL}}_k / O_k$ is an isomorphism by [18, Corollary 6.7]: it is defined

by sending the class of $(H, h): (S^{n+k+1}, \Sigma^{n+1}) \rightarrow (S^{n+k+1}, S^{n+1})$ to the normal block bundle ν_{cyl} of $\text{cyl}(h)$ inside $\text{cyl}(H)$ along with its linear reduction at the Σ^{n+1} end of the cylinder and its canonical trivialisation at the other end. Finally, the map \tilde{S} is obtained from the fact that the inclusion $\text{PL} \rightarrow \widetilde{\text{PL}}$ is an equivalence [19, Corollary 5.5(ii)]; that is, there is no essential difference between stable PL (micro)bundles and stable block bundles.

We claim that the lower left square of (4) is commutative up to sign. To see this, we may assume, increasing k if necessary, that the normal block bundle ν_{cyl} is given by a PL microbundle. Then, the sum of the two composites, applied to $[(H, h)]$, is represented by the direct sum microbundle $T\text{cyl}(h) \oplus \nu_{\text{cyl}}$ over $\text{cyl}(h)$ along with its linear reduction at the front end and its canonical trivialisation at the other end. But now, we have an isomorphism $T\text{cyl}(h) \oplus \nu_{\text{cyl}} \cong T\text{cyl}(H)|_{\text{cyl}(h)}$ of microbundles which extends isomorphisms $T\Sigma \oplus \nu_{\Sigma \subset S^{n+k+1}} \cong TS^{n+k+1}|_{\Sigma}$ and $TS^{n+1} \oplus \nu_{S^{n+1} \subset S^{n+k+1}} \cong TS^{n+k+1}|_{S^{n+1}}$ of vector bundles.

Since H is PL isotopic to the identity (being an orientation-preserving PL homeomorphism of the sphere), we conclude that the sum of the two composite maps, applied to $[(H, h)]$, represents the zero element.

All other parts of this diagram commute up to possible signs: the commutativity of the squares on the right and of the triangle in the middle follows from the definitions. That $M_{n*} \circ \lambda = S \circ M_{k*}$ follows from [8, Lemma 2.5], and that $M_{n*} = \Psi \circ C$ is proven in [8, Lemma 2.7]. The lemma now follows by a diagram chase. \square

3 Concluding remarks

In this section we discuss some of the background to our results and state a conjecture about the map $d_* \circ \sigma$.

(1) The homotopy fibre of $d: \text{Diff}_{\partial}(D^k) \rightarrow \Omega^k \text{SO}_k$ is the H -space $\text{Diff}_{\partial}^{\text{fr}}(D^k)$ of framing-preserving diffeomorphisms. It is the loop space of the classifying space $B\text{Diff}_{\partial}^{\text{fr}}(D^k)$, which features in the recent work of Kupers and Randal-Williams [13] on the rational homotopy groups of $\text{Diff}_{\partial}(D^k)$. We see that d is rationally trivial because the Alexander trick implies that d becomes nullhomotopic after composition with the natural map $\Omega^k \text{SO}_k \rightarrow \Omega^k \text{SPL}_k$. It is well known that $(\text{SO}_k)_{\mathbb{Q}}$ is Eilenberg–Mac Lane, detected by the suspensions of the rational Pontryagin classes and rational Euler class. Since these classes are defined on $(\text{SPL}_k)_{\mathbb{Q}}$, it follows that $(\text{SO}_k)_{\mathbb{Q}}$ is a homotopy

retract of $(\mathrm{SPL}_k)_{\mathbb{Q}}$. If $\Omega_0^k X$ denotes the connected component of the constant map, then it follows that $(\Omega_0^k \mathrm{SO}_k)_{\mathbb{Q}} \simeq \Omega_0^k (\mathrm{SO}_k)_{\mathbb{Q}}$ is a homotopy retract of $(\Omega_0^k \mathrm{SPL}_k)_{\mathbb{Q}}$, showing that the map $d: \mathrm{Diff}_{\partial}(D^k) \rightarrow \Omega_0^k \mathrm{SO}_k$ is rationally nullhomotopic.

(2) The proof of Theorem 1.1 relies on the fact that the normal bundle of any embedding $\Sigma_{\xi, \xi} \hookrightarrow S^{28}$ is nontrivial. Despite the elementary argument we give for this in Section 2, computing the normal bundle of an embedding of a homotopy sphere $g: \Sigma^{n+1} \hookrightarrow S^{n+k+1}$ is a subtle problem. Provided one is in the metastable range $n < 2k-4$, Haefliger [10] proved that the isotopy class of g depends only on the diffeomorphism type of Σ , so that, in particular, the normal bundle is independent of the choice of embedding. Hsiang, Levine and Szczarba [11] proved that the latter statement holds even for $n < 2k-2$, defined the homomorphism

$$\phi_{n+1}^k: \Theta_{n+1} \rightarrow \pi_{n+1} B\mathrm{SO}_k, \quad \Sigma \mapsto \nu(\Sigma \subset S^{n+k+1}), \quad \text{where } n < 2k-2,$$

and proved that $\phi_{16}^{13} \neq 0$; ie the exotic 16–sphere embeds into S^{29} with nontrivial normal bundle. Then Antonelli [2] made a systematic study of normal bundles of homotopy spheres in the metastable range, which includes the statement that $\phi_{17}^{11} \neq 0$.

(3) Concerning A’Campo’s claim that d_* vanishes for $i = 2$, we note that, since $\phi_{16}^{13} \neq 0$, Lemma 2.1 entails that if A’Campo’s claim holds, then the exotic 16–sphere does not lie in the image of the map $A: \pi_2 \mathrm{Diff}_{\partial}(D^{13}) \rightarrow \Theta_{16} = \mathbb{Z}/2$. This is consistent with computations we have made for the refined plumbing pairing

$$\sigma: \pi_8 \mathrm{SO}_6 \otimes \pi_7 \mathrm{SO}_8 \rightarrow \pi_2 \mathrm{Diff}_{\partial}(D^{13}),$$

which show that $A \circ \sigma = 0$, even though $\sigma_M: \pi_8 \mathrm{SO}_7 \otimes \pi_7 \mathrm{SO}_8 \rightarrow \Theta_{16}$ is nontrivial, a statement which can be deduced from [20, Satz 12.1].

(4) Finally, we present a conjectural description of the homomorphism

$$d_* \circ \sigma: \pi_p \mathrm{SO}_{q-a} \otimes \pi_q \mathrm{SO}_{p-b} \rightarrow \pi_{p+q} \mathrm{SO}_{p+q-a-b-1}$$

in purely homotopy-theoretic terms.

Let $h: \pi_i \mathrm{SO}_j \rightarrow \pi_i(S^{j-1})$ be the map induced by the canonical projection $\mathrm{SO}_j \rightarrow S^{j-1}$. For maps $f: W \rightarrow X$ and $g: Y \rightarrow Z$ let $f * g: W * Y \rightarrow X * Z$ be their *join*. Let $\partial: \pi_{m+1}(S^k) \rightarrow \pi_m \mathrm{SO}_k$ denote the boundary map in the homotopy long exact sequence of the fibration $\mathrm{SO}_k \rightarrow \mathrm{SO}_{k+1} \rightarrow S^k$. For compactness, we use the notation

$$p' := p - b \quad \text{and} \quad q' := q - a$$

and let $\xi_1 \in \pi_p \mathrm{SO}_{q'}$ and $\xi_2 \in \pi_q \mathrm{SO}_{p'}$. Then we have

$$h(\xi_1) \in \pi_p(S^{q'-1}), \quad h(\xi_2) \in \pi_q(S^{p'-1}) \quad \text{and} \quad h(\xi_1) * h(\xi_2) \in \pi_{p+q+1}(S^{p'+q'-1}),$$

so that $\partial(h(\xi_1) * h(\xi_2)) \in \pi_{p+q} \mathrm{SO}_{p'+q'-1}$.

In addition, we have the J -homomorphisms

$$J_{p,q'}: \pi_p \mathrm{SO}_{q'} \rightarrow \pi_{p+q'} S^{q'} \quad \text{and} \quad J_{q,p'}: \pi_q \mathrm{SO}_{p'} \rightarrow \pi_{q+p'} S^{p'},$$

and we can suspend in the target of each of these to get the homomorphisms

$$\Sigma^a \circ J_{p,q'}: \pi_p \mathrm{SO}_{q'} \rightarrow \pi_{p+q} S^q \quad \text{and} \quad \Sigma^b \circ J_{q,p'}: \pi_q \mathrm{SO}_{p'} \rightarrow \pi_{p+q} S^p.$$

We then take compositions with the maps induced by ξ_i for $i = 1, 2$ and the inclusions $i_{p'}: \mathrm{SO}_{p'} \rightarrow \mathrm{SO}_{p'+q'-1}$ and $i_{q'}: \mathrm{SO}_{q'} \rightarrow \mathrm{SO}_{p'+q'-1}$. Hence we have homomorphisms

$$\begin{aligned} \bar{\xi}_{2*}: \pi_p \mathrm{SO}_{q'} &\xrightarrow{\Sigma^a \circ J_{p,q'}} \pi_{p+q} S^q \xrightarrow{\xi_{2*}} \pi_{p+q} \mathrm{SO}_{p'} \xrightarrow{i_{p'*}} \pi_{p+q} \mathrm{SO}_{p'+q'-1}, \\ \bar{\xi}_{1*}: \pi_q \mathrm{SO}_{p'} &\xrightarrow{\Sigma^b \circ J_{q,p'}} \pi_{p+q} S^p \xrightarrow{\xi_{1*}} \pi_{p+q} \mathrm{SO}_{q'} \xrightarrow{i_{q'*}} \pi_{p+q} \mathrm{SO}_{p'+q'-1}. \end{aligned}$$

Conjecture 3.1 *Up to sign, the homomorphism*

$$d_* \circ \sigma: \pi_p \mathrm{SO}_{q'} \otimes \pi_q \mathrm{SO}_{p'} \rightarrow \pi_{p+q} \mathrm{SO}_{p'+q'-1}$$

is given by

$$d_*(\sigma(\xi_1, \xi_2)) = \partial(h(\xi_1) * h(\xi_2)) + \bar{\xi}_{1*}(\xi_2) + \bar{\xi}_{2*}(\xi_1).$$

We briefly discuss Conjecture 3.1 in light of Theorem 1.1 and Corollary 1.2. For $\xi \in \pi_8 \mathrm{SO}_6$ a generator, $h(\xi) \in \pi_8 S^5 \cong \pi_3^S$ is again a generator and we choose ξ so that $h(\xi) = \nu_5$. Hence Conjecture 3.1 gives $d_*(\sigma(\xi, \xi)) = \partial(\nu_5 * \nu_5) + 2\bar{\xi}_*(\xi)$. Now $\pi_{16} S^8 \cong (\mathbb{Z}/2)^4$, which entails that $2\bar{\xi}_*(\xi) = 0$ and the proof of Corollary 1.2 shows that $d_*(\sigma(\xi, \xi)) = \partial(\nu_{11}^2)$. Since $\nu_5 * \nu_5 = \nu_{11}^2$, Conjecture 3.1 is consistent with Theorem 1.1 and Corollary 1.2, with both giving the same nonzero expression for $d_* \circ \sigma: \pi_8 \mathrm{SO}_6 \otimes \pi_8 \mathrm{SO}_6 \rightarrow \pi_{16} \mathrm{SO}_{11}$.

References

- [1] **J F Adams**, *Vector fields on spheres*, Ann. of Math. 75 (1962) 603–632 MR Zbl
- [2] **P L Antonelli**, *On stable diffeomorphism of exotic spheres in the metastable range*, Canadian J. Math. 23 (1971) 579–587 MR Zbl

- [3] **P Antonelli, D Burghlelea, P J Kahn**, *Gromoll groups, $\text{Diff } S^n$ and bilinear constructions of exotic spheres*, Bull. Amer. Math. Soc. 76 (1970) 772–777 MR Zbl
- [4] **P L Antonelli, D Burghlelea, P J Kahn**, *The nonfinite homotopy type of some diffeomorphism groups*, Topology 11 (1972) 1–49 MR Zbl
- [5] **D Burghlelea, R Lashof**, *The homotopy type of the space of diffeomorphisms, I*, Trans. Amer. Math. Soc. 196 (1974) 1–36 MR Zbl
- [6] **D Burghlelea, R Lashof**, *The homotopy type of the space of diffeomorphisms, II*, Trans. Amer. Math. Soc. 196 (1974) 37–50 MR Zbl
- [7] **R Burklund, A Senger**, *On the high-dimensional geography problem*, preprint (2020) arXiv 2007.05127
- [8] **D Crowley, T Schick**, *The Gromoll filtration, KO –characteristic classes and metrics of positive scalar curvature*, Geom. Topol. 17 (2013) 1773–1789 MR Zbl
- [9] **D Gromoll**, *Differenzierbare Strukturen und Metriken positiver Krümmung auf Sphären*, Math. Ann. 164 (1966) 353–371 MR Zbl
- [10] **A Haefliger**, *Plongements différentiables de variétés dans variétés*, Comment. Math. Helv. 36 (1961) 47–82 MR Zbl
- [11] **W C Hsiang, J Levine, R H Szczarba**, *On the normal bundle of a homotopy sphere embedded in Euclidean space*, Topology 3 (1965) 173–181 MR Zbl
- [12] **M A Kervaire**, *Some nonstable homotopy groups of Lie groups*, Illinois J. Math. 4 (1960) 161–169 MR Zbl
- [13] **A Kupers, O Randal-Williams**, *On diffeomorphisms of even-dimensional discs*, J. Amer. Math. Soc. (online ahead of print October 2023)
- [14] **J P Levine**, *Lectures on groups of homotopy spheres*, from “Algebraic and geometric topology” (A Ranicki, N Levitt, F Quinn, editors), Lecture Notes in Math. 1126, Springer (1985) 62–95 MR Zbl
- [15] **W Lück**, *A basic introduction to surgery theory*, from “Topology of high-dimensional manifolds, I, II” (F T Farrell, L Göttsche, W Lück, editors), ICTP Lect. Notes 9, Abdus Salam Int. Cent. Theoret. Phys., Trieste (2002) 1–224 MR Zbl
- [16] **J Milnor**, *Differentiable structures on spheres*, Amer. J. Math. 81 (1959) 962–972 MR Zbl
- [17] **J Milnor**, *Microbundles, I*, Topology 3 (1964) 53–80 MR Zbl
- [18] **C P Rourke, B J Sanderson**, *Block bundles, I*, Ann. of Math. 87 (1968) 1–28 MR Zbl
- [19] **C P Rourke, B J Sanderson**, *Block bundles, III: Homotopy theory*, Ann. of Math. 87 (1968) 431–483 MR Zbl
- [20] **S Stolz**, *Hochzusammenhängende Mannigfaltigkeiten und ihre Ränder*, Lecture Notes in Math. 1116, Springer (1985) MR Zbl

- [21] **H Toda**, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies 49, Princeton Univ. Press (1962) MR Zbl
- [22] **C T C Wall**, *Classification of $(n-1)$ -connected $2n$ -manifolds*, Ann. of Math. 75 (1962) 163–189 MR Zbl
- [23] **C T C Wall**, *Classification problems in differential topology, VI: Classification of $(s-1)$ -connected $(2s+1)$ -manifolds*, Topology 6 (1967) 273–296 MR Zbl

*School of Mathematics and Statistics, University of Melbourne
Parkville, Victoria, Australia*

*Mathematisches Institut, Universität Göttingen
Göttingen, Germany*

*Institut für Mathematik, Universität Augsburg
Augsburg, Germany*

`dcrowley@unimelb.edu.au`, `thomas.schick@math.uni-goettingen.de`,
`wolfgang.steimle@math.uni-augsburg.de`

`https://www.dcrowley.net`,
`https://www.uni-math.gwdg.de/schick`, `https://www.uni-augsburg.de/de/fakultaet/mntf/math/prof/diff/team/wolfgang-steimle`

Proposed: Stefan Schwede
Seconded: Ulrike Tillmann, Nathalie Wahl

Received: 6 May 2021
Revised: 12 January 2022

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITOR

András I Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

Dan Abramovich	Brown University dan_abramovich@brown.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Frances Kirwan	University of Oxford frances.kirwan@balliol.oxford.ac.uk
Mark Behrens	University of Notre Dame mbehren1@nd.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Mladen Bestvina	University of Utah bestvina@math.utah.edu	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Martin R Bridson	University of Oxford bridson@maths.ox.ac.uk	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	John Lott	University of California, Berkeley lott@math.berkeley.edu
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Tobias H Colding	Massachusetts Institute of Technology colding@math.mit.edu	Haynes Miller	Massachusetts Institute of Technology hmm@math.mit.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Tomasz Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Walter Neumann	Columbia University neumann@math.columbia.edu
Benson Farb	University of Chicago farb@math.uchicago.edu	Jean-Pierre Otal	Université Paul Sabatier, Toulouse otal@math.univ-toulouse.fr
Steve Ferry	Rutgers University sferry@math.rutgers.edu	Peter Ozsváth	Princeton University petero@math.princeton.edu
David M Fisher	Rice University davidfisher@rice.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
Mike Freedman	Microsoft Research michaelf@microsoft.com	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
David Gabai	Princeton University gabai@princeton.edu	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Peter Teichner	Max Planck Institut für Mathematik teichner@mac.com
Cameron Gordon	University of Texas gordon@math.utexas.edu	Richard P Thomas	Imperial College, London richard.thomas@imperial.ac.uk
Lothar Götsche	Abdus Salam Int. Centre for Th. Physics gotsche@ictp.trieste.it	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Mark Gross	University of Cambridge mgross@dpmms.cam.ac.uk	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de

See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2023 is US \$740/year for the electronic version, and \$1030/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2023 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 27 Issue 9 (pages 3387–3831) 2023

Hyperbolic groups acting improperly	3387
DANIEL GROVES and JASON FOX MANNING	
Cyclic homology, S^1 -equivariant Floer cohomology and Calabi–Yau structures	3461
SHEEL GANATRA	
Congruences on K-theoretic Gromov–Witten invariants	3585
JÉRÉMY GUÉRE	
Moduli of spherical tori with one conical point	3619
ALEXANDRE EREMENKO, GABRIELE MONDELLO and DMITRI PANOV	
The derivative map for diffeomorphism of disks: an example	3699
DIARMUID CROWLEY, THOMAS SCHICK and WOLFGANG STEIMLE	
On self-shrinkers of medium entropy in \mathbb{R}^4	3715
ALEXANDER MRAMOR	
The Gromov–Hausdorff distance between spheres	3733
SUNHYUK LIM, FACUNDO MÉMOLI and ZANE SMITH	
Contact three-manifolds with exactly two simple Reeb orbits	3801
DANIEL CRISTOFARO-GARDINER, UMBERTO HRYNIEWICZ, MICHAEL HUTCHINGS and HUI LIU	