

Salivary gland ultrasound elastography requires interpretation of “normal” [Letter]

Henry T. Hoffman, Piper Wenzel, Johannes Zenk, Antonio Marcelino, Harry Quon

Angaben zur Veröffentlichung / Publication details:

Hoffman, Henry T., Piper Wenzel, Johannes Zenk, Antonio Marcelino, and Harry Quon. 2024. “Salivary gland ultrasound elastography requires interpretation of ‘normal’ [Letter].” *Laryngoscope Investigative Otolaryngology* 9 (5): e70016. <https://doi.org/10.1002/liv.2.70016>.

LETTER TO THE EDITOR

Salivary gland ultrasound elastography requires interpretation of “normal”

Strong support has evolved for elastography as a supplement to salivary gland ultrasound assessment.¹ We advocate for the clinical value of this tool but offer caution in considering “normal” shear wave velocity (also reported as Young's modulus) to consistently reflect normal gland function.

Standard ultrasound assessment of parotid tumors has been reported to identify malignancy with a 91% accuracy.² Jering et al. reported that additional evaluation with elastography improved the diagnostic accuracy by identifying malignant tumors to be associated with faster shear wave velocities and larger areas of stiff tissue than benign tumors.³

Assessments of non-neoplastic salivary disorders with ultrasound elastography have reported the capacity to discriminate between normal salivary glands with slower shear wave velocity from those in patients with Sjogren's syndrome with faster shear wave velocity.⁴ Dai et al. through a meta-analysis of 15 articles addressing primary Sjogren's syndrome (pSS) concluded that ultrasound elastography “demonstrates high accuracy in differentiating between pSS and healthy/disease control groups”.⁵

Chang and Wang identified their experience with ultrasound shear wave elastography to characterize glands affected with sialolithiasis.⁶ Sequential assessments identified changes to gland stiffness following treatment of obstructive sialadenitis to further support the clinical utility of shear wave elastography as was similarly reported by a group in Munich, Germany.^{7,8}

We concur with Chang and Wang's contention that the significant decrease in shear wave velocity they identified following stone removal does “imply the diseased gland became softer after removal of sialolithiasis” but offer caution in their interpretation that softening and slower speed reflects “recovery of salivary gland function.”⁶ Although others have similarly identified the value of elastography to “indirectly reflect organ function,”⁵ it is important to acknowledge that fatty replacement of diseased glands may dominate the exam to provide a normal shear wave analysis of a poorly functioning or nonfunctioning gland.

Takagi et al. employed MR analysis to identify fatty degeneration occurring within the salivary glands of patients with long-standing Sjogren's syndrome.⁹ Study of patients with post-irradiation xerostomia employing MRI and advanced CT imaging has also correlated poor salivary gland function associated with fat infiltration.¹⁰ An elegant

study of surgically resected human breast tissue identified a broad range of viscoelastic properties within the tissue subject to elastography analysis.¹¹ These investigators reported the modulus (stiffness) of fat tissue in the specimens to be significantly lower than that of glandular and fibrous tissue.

Our experience in examining patients following long-term injury from chronic salivary duct obstruction and from previous irradiation often identifies a slow (“normal”) speed to elastography analysis in a way that correlates with fat replacement identified on CT imaging. This finding is highlighted by our report of a patient who had undergone parotid duct ligation with long-term follow-up identifying an elevated shear wave velocity in an infarcted tail of one parotid with normal velocity seen in the other parotid regions.¹² The normal shear wave velocity measurements correlated with CT assessment identifying parotid gland atrophy with fat replacement.



Advances in salivary ultrasound are occurring at a rapid rate and will benefit from efforts to standardize evaluation and reporting.¹³ The addition of novel approaches to elastography coupled with advances in complementary approaches such as intravenous and intracavitary contrast enhanced ultrasonography (CEUS) will provide greater utility to salivary ultrasound evaluation.^{14,15} Multidimensional ultrasound assessment coupled with deep learning offers promise to further enhance diagnostic capabilities.¹⁶

FUNDING INFORMATION

This study was supported by Otolaryngology Department, University of Iowa.

CONFLICT OF INTEREST STATEMENT

Henry T. Hoffman: MeiraGtx research study participant; RiboX scientific advisor board; and UpToDate author (no conflicts related to this publication). Harry Quon: MeiraGtx research study participant (no conflicts related to this publication).

Henry T. Hoffman MD¹ 
Piper Wenzel BS¹ 
Johannes Zenk MD²
Antonio Marcelino MD³
Harry Quon MD⁴

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.
© 2024 The Author(s). *Laryngoscope Investigative Otolaryngology* published by Wiley Periodicals LLC on behalf of The Triological Society.

¹Department of Otolaryngology, University of Iowa, Iowa City, Iowa, USA

²Department of ENT, University Hospital Augsburg, Augsburg, Germany

³Department of Radiology, University of Iowa, Iowa City, Iowa, USA

⁴Department of Radiation Oncology and Molecular Radiation Sciences, Johns Hopkins University, Baltimore, Maryland, USA

Correspondence

Henry T. Hoffman, Department of Otolaryngology, University of Iowa, 200 Hawkins Drive, Department of Otolaryngology, University of Iowa, Iowa City, IA 52242, USA.

Email: henry-hoffman@uiowa.edu

ORCID

Henry T. Hoffman  <https://orcid.org/0000-0003-4486-855X>

Piper Wenzel  <https://orcid.org/0009-0001-3040-0908>

REFERENCES

- Yu Q, Ding A, Chen Q, Zuo J, Cao J, Xiong P. Diagnostic value of shear wave elastography quantification combined with conventional ultrasound in salivary gland tumors. *Quant Imaging Med Surg.* 2023; 13(9):5863-5876. doi:10.21037/qims-23-103
- Mansour N, Bas M, Stock KF, Strassen U, Hofauer B, Knopf A. Multimodal ultrasonographic pathway of parotid gland lesions. *Ultraschall Med.* 2017;38(2):166-173. doi:10.1055/s-0035-1553267
- Jering M, Zenk J, Thölken R, Rüger H, Psychogios G. Can ultrasound in combination with virtual touch imaging quantification predict the dignity of a parotid tumor? *Ultrasound Med Biol.* 2021;47(5):1192-1203. doi:10.1016/j.ultrasmedbio.2020.12.027
- Ma H, Liu L, Sun X, Liu M, Liu Z. The diagnostic value of shear-wave elastography in the salivary glands of patients with primary Sjögren syndrome. *Int J Rheum Dis.* 2023;26(7):1314-1320. doi:10.1111/1756-185X.14734
- Dai X, Sui X, Chen S, Zhao B, Liu Z, Wang X. The diagnostic performance of salivary gland ultrasound elastography in Sjögren's syndrome and sicca symptoms: a systematic review and meta-analysis. *Eur Radiol.* 2024;34(3):1545-1555. doi:10.1007/s00330-023-10166-5
- Chang CF, Wang HK. Ultrasound shear wave elastography for patients with sialolithiasis undergoing interventional sialendoscopy. *Laryngoscope Investig Otolaryngol.* 2023;8(1):76-81. doi:10.1002/lio2.1007
- Zengel P, Reichel CA, Vincek T, Clevert DA. Ultrasound elastography in diagnosis and follow-up for patients with chronic recurrent parotitis. *Clin Hemorheol Microcirc.* 2017;67(3-4):389-397. doi:10.3233/CH-179219
- Reichel CA, Vincek T, Gellrich D, Schrötzmair F, Clevert D, Zengel P. Ultrasound elastography in diagnosis and follow-up for patients with sialolithiasis. *Dentomaxillofac Radiol.* 2018;47(7):20170424. doi:10.1259/dmfr.20170424
- Takagi Y, Sasaki M, Eida S, et al. Comparison of salivary gland MRI and ultrasonography findings among patients with Sjögren's syndrome over a wide age range. *Rheumatology.* 2022;61(5):1986-1996. doi:10.1093/rheumatology/keab560
- Sheikh K, Lee SH, Cheng Z, et al. Predicting acute radiation induced xerostomia in head and neck cancer using MR and CT radiomics of parotid and submandibular glands. *Radiat Oncol.* 2019;14(1):131. doi:10.1186/s13014-019-1339-4
- Krouskop TA, Wheeler TM, Kallel F, Garra BS, Hall T. Elastic moduli of breast and prostate tissues under compression. *Ultrason Imaging.* 1998;20(4):260-274. doi:10.1177/016173469802000403
- Wenzel P, Maley J, Marcelino AZ, Hoffman H. Clinical and radiographic follow-up after the Wilkie procedure at 28 years: A case report. *Radiol Case Rep.* 2023;19(1):1-6. doi:10.1016/j.radcr.2023.09.071
- Hoffman HT, Koch M, Witt RL, et al. Proposal for standardized ultrasound analysis of the salivary glands: Part 1 submandibular gland. *Laryngoscope Investig Otolaryngol.* 2024;9(1):e1224. doi:10.1002/lio2.1224
- Jia Y, Yang J, Zhu Y, et al. Ultrasound-based radiomics: current status, challenges and future opportunities. *Med Ultrason.* 2022;24(4):451-460. doi:10.11152/mu-3248
- Vollert K, Clevert DA, Kleffel T. Endocavitary contrast-enhanced ultrasound. *Rofo.* 2022;194(11):1204-1215. doi:10.1055/a-1826-0325
- Liu XH, Miao YY, Qian L, et al. Deep learning based ultrasound analysis facilitates precise distinction between parotid pleomorphic adenoma and Warthin tumor. *Front Oncol.* 2024;27(14):1337631. doi:10.3389/fonc.2024.1337631