

Derived category of the spinor 15-fold

Vladimiro Benedetti, Daniele Faenzi, Maxim N. Smirnov

Angaben zur Veröffentlichung / Publication details:

Benedetti, Vladimiro, Daniele Faenzi, and Maxim N. Smirnov. 2025. "Derived category of the spinor 15-fold." *Journal of Algebra* 677: 460-96.
<https://doi.org/10.1016/j.jalgebra.2025.04.019>.



Contents lists available at ScienceDirect

Journal of Algebra

journal homepage: www.elsevier.com/locate/jalgebra

Research Paper

Derived category of the spinor 15-fold

Vladimiro Benedetti^{a,*}, Daniele Faenzi^b, Maxim N. Smirnov^c^a *Université Côte d'Azur, CNRS – Laboratoire J.-A. Dieudonné, Parc Valrose, F-06108 Nice CEDEX 2, France*^b *Université Bourgogne Europe, CNRS, IMB UMR 5584, F-21000 Dijon, France*^c *Institut für Mathematik, Fakultäten Mathematisch-Naturwissenschaftlich-Technische Fakultät, Universitätsstrasse 14, 86159 Augsburg, Germany*

ARTICLE INFO

ABSTRACT

Article history:

Received 26 February 2024

Available online 22 April 2025

Communicated by Hannah Markwig

MSC:

14F08

Keywords:

Full exceptional Lefschetz collections

Derived categories

Spinor varieties

Isotropic Grassmannians

Freudenthal's magic square

We construct a full exceptional Lefschetz collection on the spinor 15-fold consisting of a connected component of the space of orthogonal 6-dimensional subspaces of a 12-dimensional complex vector space, isotropic with respect to a fixed non-degenerate quadratic form. The collection is made of 2 twists of a 4-item block and 8 twists of a 3-item block, confirming a conjecture of Kuznetsov and Smirnov. We speculate that a similar collection might work for the Freudenthal E_7 -variety.

© 2025 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

It is widely expected that, for any parabolic subgroup P of a reductive complex algebraic group G , the associated rational homogeneous variety $X = G/P$ admits a full exceptional collection (E_1, \dots, E_r) . This means that there is a sequence (E_1, \dots, E_r) of

* Corresponding author.

E-mail addresses: vladimiro.benedetti@univ-cotedazur.fr (V. Benedetti), daniele.faezi@ube.fr (D. Faenzi), maxim.smirnov@math.uni-augsburg.de, maxim.n.smirnov@gmail.com (M.N. Smirnov).

objects of the derived category of coherent sheaves $\mathbf{D}^b(X)$, which is *exceptional*, namely $\text{Ext}_X^p(E_i, E_j) = 0$ for all $1 \leq i, j \leq r$ for all $p \in \mathbb{Z}$ when $i > j$ or when $i = j$ and $p \neq 0$, and which is *full*, i.e. the smallest triangulated subcategory $\langle E_1, \dots, E_r \rangle$ of $\mathbf{D}^b(X)$ containing E_1, \dots, E_r is the whole $\mathbf{D}^b(X)$. This implies that $\mathbf{D}^b(X)$ admits a tilting bundle. Moreover (E_1, \dots, E_r) should consist of G -equivariant vector bundles and the collection should admit a natural partial order induced by the Bruhat-Chevalley order, see for instance [5] for an account. While full exceptional collections were given for flags of type A_n and quadrics in [1,14], in the remaining classical types exceptional collections of maximal length were constructed much later, see [15]. Some more cases admitting full exceptional collections were studied, notably for isotropic Grassmannians in the symplectic case, we refer for instance to [27,25,12,21]. Full exceptional collections on some homogeneous varieties of exceptional type were studied in [23,10,29,3,19]. However, the questions of existence of a full exceptional collection, which is moreover G -equivariant and compatible with the Bruhat-Chevalley order (conjecturally consisting of vector bundles), has been settled in full generality for any rational homogeneous variety according to the authors of the very recent preprint [30].

A slightly different point of view on the structure of the derived category and on exceptional collections stems from homological projective duality, as in [20]. In this context, the emphasis is on Lefschetz properties with respect to a given ample line bundle $\mathcal{O}_X(1)$, so that a full exceptional collection should be obtained from an initial set of objects by twisting them with $\mathcal{O}_X(t)$, for $t = 0, \dots, \ell - 1$ and occasionally removing some objects. Here ℓ is some integer which is often the Fano index of X , see below. Full exceptional Lefschetz collections were given in some classical and exceptional types in [21,11,10,3]. The question of when one should remove objects along the construction of a Lefschetz collection is a very interesting point giving rise to the study of residual categories, conjecturally related to the structure of the quantum cohomology of X , according to a refinement of Dubrovin’s conjecture, see [16,17,28].

In this paper we focus on two specific varieties, one of classical type, namely the *spinor 15-fold* $OG_+(6, 12)$, which is one connected component of the variety of maximal isotropic subspaces for a non-degenerate quadratic form in 12 variables, and the other of exceptional type, namely Freudenthal’s 27-dimensional variety E_7/P_7 . We construct a full exceptional Lefschetz collection on the first one and providing numerical evidence on the second one, based on the ansatz that they should share some common features as they sit on the same row of Freudenthal’s magic square related to real division algebras, cf. [22]. We write X_m for the varieties sitting in the third row of the Freudenthal’s magic square, where the index m refers to the dimension of the corresponding real division algebra \mathbb{A}_m . These varieties are homogeneous for the action of a group G listed below. They are Fano varieties whose Picard group is generated by a very ample line bundle $\mathcal{O}_{X_m}(1)$, hence $\omega_{X_m} \simeq \mathcal{O}_{X_m}(-\iota_{X_m})$ for some integer ι_{X_m} called the Fano index of X_m . We have $\dim(X_m) = 3(m + 1)$ and $\iota_{X_m} = 2(m + 1)$. Explicitly, we have the following table:

m	1	2	4	8
\mathbb{A}_m	\mathbb{R}	\mathbb{C}	\mathbb{H}	\mathbb{O}
G	Sp_3	GL_6	$Spin_{12}$	E_7
X_m	$LG(3, 6)$	$G(3, 6)$	$OG_+(6, 12)$	E_7/P_7
$\dim(X_m)$	6	9	15	27
ι_{X_m}	4	6	10	18
$\text{rk}(K_0(X_m))$	8	20	32	56

Excluding $LG(3, 6)$, that does not quite fit into this picture, we have $\text{rk}(K_0(X_m)) = 6m + 8$. We would expect that for $m = 2, 4, 8$ the derived category of X_m has a full exceptional Lefschetz collection of the following form:

$$(\mathcal{A}, \mathcal{A}(1), \mathcal{B}(2), \dots, \mathcal{B}(2m + 1))$$

with:

$$\begin{aligned} \mathcal{A} &= (\mathcal{O}_X, O, P, Q) \\ \mathcal{B} &= (\mathcal{O}_X, O, P) \end{aligned} \tag{1.1}$$

Here, denoting by \mathcal{U}_ω the irreducible G -homogeneous bundle of maximal weight ω , the bundles O, P and Q should be, respectively, $\mathcal{U}_{\omega_1}, \wedge^2 \mathcal{U}_{\omega_1}$ and $S^{2,1} \mathcal{U}_{\omega_1}$ with the caveat that, in case such bundles are not exceptional, we should replace them by some equivariant extension with homogeneous bundles of lower maximal weight (for precise definitions see the next section) or projections on the semiorthogonal summand we are interested in. For $m = 2$, i.e. for $G(3, 6)$, no extension is necessary. The resulting full exceptional collection was studied in [8] in the attempt to verify Homological Projective Duality for $G(3, 6)$. On the other hand, this gets more tricky for $m = 4$ and $m = 8$.

The goal of this paper is to prove the statement for $m = 4$ and provide a partial proof of a closely related statement for $m = 8$. For $m = 4$ we prove:

Theorem 1. *Let $X = OG_+(6, 12)$ and set $O = \mathcal{U}_{\omega_1}$. Then, there are unique $Spin_{12}$ -homogeneous exceptional bundles P and Q fitting into:*

$$0 \rightarrow \mathcal{O}_X \rightarrow P \rightarrow \mathcal{U}_{\omega_2} \rightarrow 0, \quad 0 \rightarrow \mathcal{U}_{\omega_1} \rightarrow Q \rightarrow \mathcal{U}_{\omega_1 + \omega_2} \rightarrow 0,$$

such that, defining \mathcal{A} and \mathcal{B} as in (1.1), we get a full Lefschetz exceptional collection:

$$\mathbf{D}^b(X) = \langle \mathcal{A}, \mathcal{A}(1), \mathcal{B}(2), \dots, \mathcal{B}(9) \rangle.$$

Moreover, $Q' = L_{(\mathcal{B})}(Q)$ is a homogeneous exceptional bundle and Q and $Q'(1)$ are completely orthogonal.

This proves [17, Conjecture 1.3] and [16, Conjecture 1.2] for the spinor 15-fold, including the statement about the complete orthogonality of the generators of the residual category with respect to the rectangular part of the Lefschetz collection.

The two conjectures above are related to Dubrovin’s conjecture and quantum cohomology. Indeed, the structure of the small quantum cohomology of $X = OG_+(6, 12)$, which can be deduced from [6,4], should govern the structure of our exceptional collection. The fact that Q and $Q'(1)$ are completely orthogonal corresponds to the fact that the operator of small quantum multiplication by $-K_X$ has zero as an eigenvalue of multiplicity two (see [2] for a nice picture of the spectrum). We refer to [16,17] for more details on these conjectures.

Let us notice that the nontrivial extensions that appear in our collections can be realised by the method of Kuznetsov and Polishchuk in [15]. For $m = 8$ and $X := X_8 = E_7/P_7$, we prove a weaker result. Let us define O as the unique non-trivial E_7 -equivariant extension fitting into

$$0 \rightarrow \mathcal{O}_X \rightarrow O \rightarrow \mathcal{U}_{\omega_1} \rightarrow 0.$$

Let us define P as the projection of \mathcal{U}_{ω_3} to the left orthogonal of $\langle \mathcal{O}_X(1), O(1), \dots, \mathcal{O}_X(18), O(18) \rangle$, and Q as the projection (see Remark 7.3) of $\mathcal{U}_{\omega_1+\omega_3}$ to the left orthogonal of $\langle \mathcal{O}_X(1), O(1), P(1), \dots, \mathcal{O}_X(18), O(18), P(18) \rangle$.

Theorem 2. *On $X = E_7/P_7$ the collection $(\mathcal{O}_X, O, \dots, \mathcal{O}_X(17), O(17))$ is exceptional. Moreover, defining \mathcal{A} and \mathcal{B} as in (1.1), we get a numerically exceptional collection of maximal length:*

$$(\mathcal{A}, \mathcal{A}(1), \mathcal{B}(2), \dots, \mathcal{B}(17)), \quad \text{with} \quad K_0(X) = K_0(\langle \mathcal{A}, \mathcal{A}(1), \mathcal{B}(2), \dots, \mathcal{B}(17) \rangle).$$

Here, by numerically exceptional collection we mean a collection E_1, \dots, E_r whose numerical properties reproduce those of an exceptional collection: $\chi(E_i, E_j) = 0$ if $i > j$ and $\chi(E_i, E_i) = 1$ for all i . Of course having a numerically exceptional collection is a priori a much weaker condition than having an exceptional collection (not to mention having a full exceptional collection). However, due to the analogy with the other cases of the Freudenthal magic square, we believe that this collection is indeed a full exceptional collection. As a further element of comparison with the very interesting paper [30], let us note that the collections constructed in [30], besides being far from explicit, are not Lefschetz collections and that, a priori, the point of whether they consist of vector bundles remains conjectural. Our Lefschetz collection for $OG_+(6, 12)$ does consist of equivariant vector bundles, while for E_7/P_7 this is not clear.

The paper is organised as follows. In Section 2 we define our Lefschetz collection. The main tools are the theorem of Borel-Bott-Weil and a result about the non-degeneracy of a cup-product owing to Dimitrov and Roth. In Section 3 we outline our strategy to prove fullness and use it to reprove fullness of a natural Lefschetz collection on $OG_+(5, 10)$. Here we use a complex constructed in Section 4, where we also construct an analogous complex for $OG_+(6, 12)$ which in turn we use in Sections 5 and 6. In Section 5 we show that certain homogeneous bundles belong to the subcategory \mathcal{D} generated by our exceptional

collection. We use this in Section 6 to prove fullness of our collection on $OG_+(6, 12)$. In Section 7 we provide some remarks on our numerically exceptional collection on the Freudenthal variety E_7/P_7 .

Acknowledgements. V.B. and D.F. partially supported by FanoHK ANR-20-CE40-0023, SupToPhAG/EIPHI ANR-17-EURE-0002, Région Bourgogne-Franche-Comté, Feder Bourgogne and Bridges ANR-21-CE40-0017. D.F. partially supported by “CAPES-COFECUB” programme (project number: Ma 1017/24, Géométrie des équations différentielles et des variétés algébriques), funded by the French Ministry for Europe and Foreign Affairs, the French Ministry for Higher Education and Research, and CAPES. M.S. was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), Projektnummer 448537907. We warmly thank Sasha Kuznetsov for useful discussions.

2. A Lefschetz exceptional collection on the spinor 15-fold

Here we begin by sketching the exceptional collection we want to work with. We first introduce the setting about spinor varieties and homogeneous bundles on them, then define the bundles appearing in the desired Lefschetz collection and finally show that this is indeed an exceptional Lefschetz collection.

By convention, we italicize varieties such as the Grassmannian $G(k, n)$ of k -dimensional vector subspaces of \mathbb{C}^n while groups are written in roman letters, with the exception of the Levi factor L , see below.

Concerning derived categories, here is the notation we use besides the elements defined in the introduction. We write $L_{\mathcal{E}}(\mathcal{F})$ the left mutation of an object \mathcal{F} along an object \mathcal{E} of $\mathbf{D}^b(X)$. If \mathcal{B} generated by an exceptional sequence (E_1, \dots, E_r) then, for an object \mathcal{E} of $\mathbf{D}^b(X)$, we write $L_{\mathcal{B}}(\mathcal{E}) = L_{E_1} \cdots L_{E_r}(\mathcal{E})$. In the same setting, we write \mathcal{B}^\perp as the full subcategory of $\mathbf{D}^b(X)$ whose objects \mathcal{E} satisfy $\text{Ext}^p(E_i, \mathcal{E}) = 0$ for all $p \in \mathbb{Z}$ and $1 \leq i \leq r$.

2.1. A lemma on non-degeneracy of cup product maps

We will use the following special case of [9, Theorem I]. Let G be a semisimple complex algebraic group, fix a Borel subgroup B of G , a parabolic subgroup $P \supset B$ of G and let λ, μ be P -dominant weights of G . Write $X = G/P$. Set ρ for the sum of fundamental weights of G .

Lemma 2.1. *Assume μ is G -dominant and $H^1(X, \mathcal{U}_\lambda) \neq 0 \neq H^1(X, \mathcal{U}_{\lambda+\mu})$. Assume that $\lambda + \rho$ and $\lambda + \mu + \rho$ are sent to a dominant weight by the same reflection w in the Weyl group, i.e. $w(\lambda + \rho)$ and $w(\lambda + \mu + \rho)$ are both dominant. Then the cup product map $H^1(X, \mathcal{U}_\lambda) \otimes H^0(X, \mathcal{U}_\mu) \rightarrow H^1(X, \mathcal{U}_{\lambda+\mu})$ is a surjection.*

Proof. Put $Y = G/B$. For a B -dominant weight ν , we write \mathcal{L}_ν for the associated line bundle on $Y = G/B$. The inclusion $B \subset P$ induces a G -equivariant projection $\pi : Y \rightarrow X$. The weights λ, μ and $\lambda + \mu$ are B -dominant and we have natural isomorphisms $\pi_*(\mathcal{L}_\lambda) \simeq \mathcal{U}_\lambda$, $\pi_*(\mathcal{L}_\mu) \simeq \mathcal{U}_\mu$ and $\pi_*(\mathcal{L}_{\lambda+\mu}) \simeq \mathcal{U}_{\lambda+\mu}$. Under these isomorphism, the cup product map under consideration is identified with the cup product:

$$H^1(Y, \mathcal{L}_\lambda) \otimes H^0(Y, \mathcal{L}_\mu) \rightarrow H^1(Y, \mathcal{L}_{\lambda+\mu}).$$

Note that all these cohomology groups have natural structure of irreducible G -representation and that the cup product map under consideration is G -equivariant.

The inversion sets of λ and $\lambda + \mu$ (as defined in [9]) both consist of the fundamental root α defining the reflection w , as $w(\lambda + \rho)$ and $w(\lambda + \mu + \rho)$ are dominant and $H^1(Y, \mathcal{L}_\lambda)$ and $H^1(Y, \mathcal{L}_{\lambda+\mu})$ are non-zero. Therefore, by [9, Theorem I] the cup-product map under consideration is surjective. \square

2.2. Homogeneous bundles on spinor varieties

We consider the group Spin_{2n} , namely the universal cover of the group of linear automorphisms of \mathbb{C}^{2n} preserving a non-degenerate quadratic form q . Let P_n be the parabolic subgroup of Spin_{2n} defining the spinor Grassmannian $X = \text{Spin}_{2n}/P_n := OG_+(n, 2n)$, one of the two isomorphic connected components parametrizing n -dimensional isotropic subspaces of a $2n$ -dimensional subspace endowed with a non-degenerate symmetric form. We denote by $L(P_n)$ its Levi factor.

We will denote by \mathcal{U}_ω the homogeneous bundle on Spin_{2n}/P_n associated to the $L(P_n)$ -weight ω . We write $\mathcal{O}_X(1) := \mathcal{U}_{\omega_n}$ and $\mathcal{U} := \mathcal{U}_{\omega_1}^\vee$. These correspond to the ample generator of $\text{Pic}(X)$, providing the equivariant embedding of X into $\mathbb{P}(V^{\omega_n})$, and to the tautological sub-bundle on $G(n, 2n)$, restricted to X . Here we denoted by V^λ the Spin_{2n} -representation of highest weight λ .

Unless specified otherwise, we will set $n = 6$ from now on and work on $X = \text{Spin}_{12}/P_6$. This is the spinor 15-fold that we are interested in. It is a Fano variety of Picard number one and index 10. The rank of its K_0 group is 32. We note that

$$\mathcal{U} = \mathcal{U}_{\omega_1}^\vee \simeq \mathcal{U}_{\omega_5}(-1), \quad \wedge^2 \mathcal{U} \simeq \mathcal{U}_{\omega_2}^\vee \simeq \mathcal{U}_{\omega_4}(-2), \quad \Sigma^{2,1} \mathcal{U} \simeq \mathcal{U}_{\omega_1+\omega_2}^\vee \simeq \mathcal{U}_{\omega_4+\omega_5}(-3). \tag{2.1}$$

2.3. The bundles of the exceptional collection

Let us introduce the homogeneous vector bundles appearing in our exceptional collection.

Lemma 2.2. *On X we have a canonical Spin_{2n} -equivariant exceptional bundle P fitting into:*

$$0 \rightarrow \mathcal{O}_X \rightarrow P \rightarrow \mathcal{U}_{\omega_2} \rightarrow 0. \tag{2.2}$$

Moreover, $P^\vee(2)$ is the normal bundle of X inside $\mathbb{P}(V^{\omega_6})$, while \mathcal{U}_{ω_2} is the tangent bundle of X .

Proof. The spinor 15-fold X is a cominuscule variety, which means that the tangent bundle of X is an irreducible homogeneous Spin_{12} -bundle (we refer for instance to [26]). More precisely, the tangent bundle of X is isomorphic $\wedge^2 \mathcal{U}_{\omega_1} \simeq \mathcal{U}_{\omega_2}$. On the other hand, the tangent bundle of $\mathbb{P}(V^{\omega_6})$ restricted to X is the quotient $V^{\omega_6} \otimes \mathcal{O}_X(1)/\mathcal{O}_X$. Since the irreducible factors of $V^{\omega_6} \otimes \mathcal{O}_X(1)$ are \mathcal{O}_X , \mathcal{U}_{ω_2} , $\mathcal{U}_{\omega_2}^\vee(2)$ and $\mathcal{O}_X(2)$, we obtain that the normal bundle \mathcal{N} of X inside $\mathbb{P}(V^{\omega_6})$ is a Spin_{12} -equivariant extension $\gamma \in \text{Ext}_X^1(\mathcal{U}_{\omega_2}, \mathcal{O}_X)$ giving:

$$0 \rightarrow \mathcal{U}_{\omega_2}^\vee(2) \rightarrow \mathcal{N} \rightarrow \mathcal{O}_X(2) \rightarrow 0$$

Let us check that this extension is not trivial, which is to say, $\gamma \neq 0$. By the Bott-Borel-Weil (BBW) Theorem, we have $\text{Ext}_X^1(\mathcal{O}_X(2), \mathcal{U}_{\omega_2}^\vee(2)) = H^1(X, \mathcal{U}_{\omega_2}^\vee) \simeq \mathbb{C}$. Hence the sheaf fitting as middle term of a non-trivial extension as above is unique up to isomorphism. Since $\mathcal{N}(-1)$ is a quotient of $V^{\omega_6} \otimes \mathcal{O}_X$ of half its rank, by autoduality of V^{ω_6} we get an exact sequence

$$0 \rightarrow \mathcal{N}^\vee(1) \rightarrow V^{\omega_6} \otimes \mathcal{O}_X \rightarrow \mathcal{N}(-1) \rightarrow 0.$$

Since $\mathcal{O}_X(-2)$ and $\mathcal{U}_{\omega_2}(-2)$ have no cohomology, \mathcal{N}^\vee has no cohomology as well. From the short exact sequence above we deduce that \mathcal{N} is a non-trivial extension, and thus $\mathcal{N} = P^\vee(2)$; indeed, if it were not the case, one would deduce that $\mathbb{C} \simeq H^0(\mathcal{O}_X) \simeq H^0(\mathcal{N}(-2)) \simeq V^{\omega_6} \otimes H^0(\mathcal{O}_X(-1))$, which is false.

Since $\mathcal{U}_{\omega_2}(-2)$ has no cohomology and $\mathcal{U}_{\omega_2} \otimes \mathcal{U}_{\omega_2}(-2)$ has no cohomology except for $H^1(\mathcal{U}_{\omega_2} \otimes \mathcal{U}_{\omega_2}(-2)) = \mathbb{C}$, we get that $\mathcal{U}_{\omega_2} \otimes P(-2)$ has no cohomology except for $H^1(\mathcal{U}_{\omega_2} \otimes P(-2)) = \mathbb{C}$. By twisting the exact sequence defining P by $P(-2)$ we deduce that $P \otimes P(-2)$ has no cohomology except for $H^1(P \otimes P(-2)) = \mathbb{C}$. Now let us consider the exact sequence

$$0 \rightarrow P \otimes P(-2) \rightarrow V^{\omega_6} \otimes P(-1) \rightarrow P^\vee \otimes P \rightarrow 0.$$

Since $\mathcal{O}_X(-1)$ and $\mathcal{U}_{\omega_2}(-1)$ have no cohomology, the same is true for $P(-1)$ and $V^{\omega_6} \otimes P(-1)$. We deduce that $H^0(P^\vee \otimes P) = \mathbb{C}$ and all other cohomologies of $P^\vee \otimes P$ vanish. So the bundle P is exceptional. \square

Lemma 2.3. *On X , we have a Spin_{12} -homogeneous exceptional bundle Q fitting into a canonical equivariant extension:*

$$0 \rightarrow \mathcal{U}_{\omega_1} \rightarrow Q \rightarrow \mathcal{U}_{\omega_1+\omega_2} \rightarrow 0 \tag{2.3}$$

Moreover, we have $\text{Ext}_X^\bullet(Q, Q(-1)) = 0$.

Proof. We recall (2.1) and, using [31] to compute tensor products of representations (and homogeneous vector bundles), we obtain:

$$\mathcal{U}_{\omega_4+\omega_5} \otimes \mathcal{U}_{\omega_1} \simeq \mathcal{U}_{2\omega_5}(1) \oplus \mathcal{U}_{\omega_4}(1) \oplus \mathcal{U}_{\omega_1+\omega_4+\omega_5}. \tag{2.4}$$

We compute $H^\bullet(\mathcal{U}_{2\omega_5}(-2)) = H^\bullet(\mathcal{U}_{\omega_1+\omega_4+\omega_5}(-3)) = 0$, hence:

$$\begin{aligned} \text{Ext}_X^\bullet(\mathcal{U}_{\omega_1+\omega_2}, \mathcal{U}_{\omega_1}) &= \text{Ext}_X^1(\mathcal{U}_{\omega_1+\omega_2}, \mathcal{U}_{\omega_1}) \simeq H^1(\mathcal{U}_{\omega_4+\omega_5} \otimes \mathcal{U}_{\omega_1}(-3)) \\ &\simeq H^1(\mathcal{U}_{\omega_4}(-2)) = \mathbb{C}. \end{aligned} \tag{2.5}$$

Choosing a nonzero element ζ of $\text{Ext}_X^1(\mathcal{U}_{\omega_1+\omega_2}, \mathcal{U}_{\omega_1}) \simeq \mathbb{C}$ defines the desired equivariant vector bundle Q .

To compute $\text{Ext}_X^\bullet(Q, Q)$, we consider:

$$\begin{aligned} \mathcal{U}_{\omega_1+\omega_2}^\vee \otimes \mathcal{U}_{\omega_1} &\simeq \mathcal{U}_{\omega_1} \otimes \mathcal{U}_{\omega_4+\omega_5}(-3), \\ \mathcal{U}_{\omega_1}^\vee \otimes \mathcal{U}_{\omega_1+\omega_2} &\simeq \mathcal{U}_{\omega_1+\omega_2} \otimes \mathcal{U}_{\omega_5}(-1), \end{aligned} \tag{2.6}$$

$$\mathcal{U}_{\omega_1+\omega_2}^\vee \otimes \mathcal{U}_{\omega_1+\omega_2} \simeq \mathcal{U}_{\omega_1+\omega_2} \otimes \mathcal{U}_{\omega_4+\omega_5}(-3). \tag{2.7}$$

We computed the first item and its cohomology in (2.4) and (2.5). Using this, the fact that \mathcal{U}_{ω_1} is exceptional and that Q is defined by the non-zero extension ζ , applying $\text{Ext}_X^\bullet(-, \mathcal{U}_{\omega_1})$ to the sequence (2.3) defining Q we get

$$\text{Ext}_X^\bullet(Q, \mathcal{U}_{\omega_1}) = 0. \tag{2.8}$$

Therefore:

$$\text{Ext}_X^\bullet(Q, Q) \simeq H^\bullet(Q^\vee \otimes \mathcal{U}_{\omega_1+\omega_2}).$$

To compute the term on the right-hand-side, we need to compute the cohomology of (2.6) and (2.7). For (2.6) we get:

$$\mathcal{U}_{\omega_1+\omega_2} \otimes \mathcal{U}_{\omega_5} \simeq \mathcal{U}_{\omega_2}(1) \qquad H^\bullet(\mathcal{U}_{\omega_2}) \simeq H^0(\mathcal{U}_{\omega_2}) \simeq V^{\omega_2}, \tag{2.9}$$

$$\oplus \mathcal{U}_{2\omega_1}(1) \qquad H^\bullet(\mathcal{U}_{2\omega_1}) \simeq H^0(\mathcal{U}_{2\omega_1}) \simeq V^{2\omega_1}, \tag{2.10}$$

$$\oplus \mathcal{U}_{\omega_1+\omega_2+\omega_5} \qquad H^\bullet(\mathcal{U}_{\omega_1+\omega_2+\omega_5}(-1)) = 0. \tag{2.11}$$

Next, we compute the cohomology of $\mathcal{U}_{\omega_1+\omega_2} \otimes \mathcal{U}_{\omega_1+\omega_2}^\vee$. We use the duality isomorphisms mentioned above and get:

$$\begin{aligned}
 \mathcal{U}_{\omega_1+\omega_2} \otimes \mathcal{U}_{\omega_4+\omega_5} &\simeq \mathcal{O}_X(3) & H^\bullet(\mathcal{O}_X) &= H^0(\mathcal{O}_X) \simeq \mathbb{C}, \\
 \oplus \mathcal{U}_{\omega_2+2\omega_5}(1) & & H^\bullet(\mathcal{U}_{\omega_2+2\omega_5}(-2)) &= 0, \\
 \oplus \mathcal{U}_{\omega_2+\omega_4}(1) & & H^\bullet(\mathcal{U}_{\omega_2+\omega_4}(-2)) &= H^1(\mathcal{U}_{\omega_2+\omega_4}(-2)) \simeq V^{\omega_2}, \\
 \oplus \mathcal{U}_{\omega_1+\omega_5}(2)^{\oplus 2} & & H^\bullet(\mathcal{U}_{\omega_1+\omega_5}(-1)) &= 0, \\
 \oplus \mathcal{U}_{\omega_1+\omega_2+\omega_4+\omega_5} & & H^\bullet(\mathcal{U}_{\omega_1+\omega_2+\omega_4+\omega_5}(-3)) &= 0, \\
 \oplus \mathcal{U}_{2\omega_1+2\omega_5}(1) & & H^\bullet(\mathcal{U}_{2\omega_1+2\omega_5}(-2)) &= 0, \\
 \oplus \mathcal{U}_{2\omega_1+\omega_4}(1) & & H^\bullet(\mathcal{U}_{2\omega_1+\omega_4}(-2)) &= H^1(\mathcal{U}_{2\omega_1+\omega_4}(-2)) \simeq V^{2\omega_1}.
 \end{aligned}$$

Having computed this, we get that $H^i(Q^\vee \otimes \mathcal{U}_{\omega_1+\omega_2}) = 0$ for all $i > 0$ if and only if the boundary map induced by ζ :

$$V^{\omega_2} \oplus V^{2\omega_1} \simeq H^0(\mathcal{U}_{\omega_1}^\vee \otimes \mathcal{U}_{\omega_1+\omega_2}) \rightarrow H^1(\mathcal{U}_{\omega_1+\omega_2}^\vee \otimes \mathcal{U}_{\omega_1+\omega_2}) \simeq V^{\omega_2} \oplus V^{2\omega_1}$$

is an isomorphism, and in this case $H^0(Q^\vee \otimes \mathcal{U}_{\omega_1+\omega_2}) \simeq \mathbb{C}$. In other words, Q is exceptional if and only if the following Yoneda map is an isomorphism:

$$\text{Ext}_X^1(\mathcal{U}_{\omega_1+\omega_2}, \mathcal{U}_{\omega_1}) \otimes \text{Hom}_X(\mathcal{U}_{\omega_1}, \mathcal{U}_{\omega_1+\omega_2}) \rightarrow \text{Ext}_X^1(\mathcal{U}_{\omega_1+\omega_2}, \mathcal{U}_{\omega_1+\omega_2}).$$

In view of the cohomologies of $\mathcal{U}_{\omega_4+\omega_5} \otimes \mathcal{U}_{\omega_1}(-3)$, $\mathcal{U}_{\omega_1+\omega_2} \otimes \mathcal{U}_{\omega_5}(-1)$, $\mathcal{U}_{\omega_1+\omega_2} \otimes \mathcal{U}_{\omega_4+\omega_5}(-3)$ computed above, this happens if and only if the cup-product maps below are isomorphisms:

$$\begin{aligned}
 H^1(\mathcal{U}_{\omega_4-2\omega_6}) \otimes H^0(\mathcal{U}_{\omega_2}) &\rightarrow H^1(\mathcal{U}_{\omega_2+\omega_4-2\omega_6}), \\
 H^1(\mathcal{U}_{\omega_4-2\omega_6}) \otimes H^0(\mathcal{U}_{2\omega_1}) &\rightarrow H^1(\mathcal{U}_{2\omega_1+\omega_4-2\omega_6}).
 \end{aligned}$$

However, this follows at once from Lemma 2.1, the required reflection being about the root α_6 .

Concerning $\text{Ext}_X^1(Q, Q(-1))$, it is sufficient to check that all irreducible bundles in $\mathcal{U}_{\omega_1+\omega_2}^\vee \otimes \mathcal{U}_{\omega_1}(-1)$, $\mathcal{U}_{\omega_1+\omega_2}^\vee \otimes \mathcal{U}_{\omega_1+\omega_2}(-1)$, $\mathcal{U}_{\omega_1}^\vee \otimes \mathcal{U}_{\omega_1}(-1)$, $\mathcal{U}_{\omega_1}^\vee \otimes \mathcal{U}_{\omega_1+\omega_2}(-1)$ have no non-vanishing cohomology (by BBW). \square

2.4. The exceptional Lefschetz collection

Let us define the following collections of Spin_{12} -homogeneous vector bundles

$$\begin{aligned}
 \mathcal{A} &= (\mathcal{O}_X, \mathcal{U}_{\omega_1}, P, Q), \\
 \mathcal{B} &= (\mathcal{O}_X, \mathcal{U}_{\omega_1}, P).
 \end{aligned}$$

Lemma 2.4. *The following is an exceptional collection in $\mathbf{D}^b(X)$:*

$$(\mathcal{B}, \mathcal{B}(1), \dots, \mathcal{B}(9)).$$

Proof. Recall that \mathcal{O}_X and P are exceptional. We compute:

$$\mathcal{U}_{\omega_5} \otimes \mathcal{U}_{\omega_1} \simeq \mathcal{O}_X(1) \oplus \mathcal{U}_{\omega_1+\omega_5} \tag{2.12}$$

We compute the largest intervals of integers where the twists of the bundles appearing in the right-hand-side have vanishing cohomology by BBW. This gives:

$$\begin{aligned} H^\bullet(\mathcal{O}_X(-t)) &= 0, & \text{for } t \in \{1, \dots, 9\}, \\ H^\bullet(\mathcal{U}_{\omega_1+\omega_5}(-t)) &= 0, & \text{for } t \in \{1, \dots, 11\}. \end{aligned}$$

Then, using (2.1), we get that \mathcal{U}_{ω_1} is exceptional. We also get the required vanishing of twisted endomorphisms of \mathcal{O}_X and \mathcal{U}_{ω_1} . Also, we have the vanishing of $\text{Ext}_X^\bullet(\mathcal{O}_X(i), \mathcal{U}_{\omega_1}(j))$ for $0 \leq j < i \leq 9$ and of $\text{Ext}_X^\bullet(\mathcal{U}_{\omega_1}(i), \mathcal{O}_X(j))$ for $0 \leq j \leq i \leq 9$.

It remains to deal with P . Looking at the extension defining P and using BBW, we get $\text{Ext}_X^\bullet(\mathcal{O}_X(i), P(j)) = 0$ for $0 \leq j < i \leq 10$, so Serre duality ensures also $\text{Ext}_X^\bullet(P(i), \mathcal{O}_X(j)) = 0$ for $0 \leq j \leq i \leq 9$.

Next we show $\text{Ext}_X^\bullet(\mathcal{U}_{\omega_1}(i), P(j)) = 0$ for $0 \leq j < i \leq 9$ and note that the vanishing holds true even for $i = 10$. We recall (2.1) and use $\mathcal{U}_{\omega_5} \otimes \mathcal{U}_{\omega_2} \simeq \mathcal{U}_{\omega_1}(1) \oplus \mathcal{U}_{\omega_2+\omega_5}$. Then, tensoring the sequence (2.2) defining P with $\mathcal{U}_{\omega_5}(-1-t)$, for $1 \leq t \leq 9$, we get the desired vanishing by using:

$$H^\bullet(\mathcal{U}_{\omega_5}(-1-t)) = H^\bullet(\mathcal{U}_{\omega_1}(-t)) = H^\bullet(\mathcal{U}_{\omega_2+\omega_5}(-1-t)) = 0.$$

Now Serre duality gives $\text{Ext}_X^\bullet(P(i), \mathcal{U}_{\omega_1}(j)) = 0$ for $0 \leq j \leq i \leq 9$.

Finally we check $\text{Ext}_X^\bullet(P(i), P(j)) = 0$ for $0 \leq j < i \leq 9$. We compute:

$$\mathcal{U}_{\omega_2} \otimes \mathcal{U}_{\omega_4} \simeq \mathcal{O}_X(2) \oplus \mathcal{U}_{\omega_1+\omega_5}(1) \oplus \mathcal{U}_{\omega_2+\omega_4}.$$

Tensoring the sequence (2.2) defining P with its dual and using (2.1), we deduce the desired vanishing results from the following ones, which in turn are given by BBW for $1 \leq t \leq 9$:

$$\begin{aligned} H^\bullet(\mathcal{O}_X(-t)) &= H^\bullet(\mathcal{U}_{\omega_2}(-t)) = H^\bullet(\mathcal{U}_{\omega_4}(-2-t)) \\ &= H^\bullet(\mathcal{U}_{\omega_2+\omega_4}(-t)) = H^\bullet(\mathcal{U}_{\omega_1+\omega_5}(-1-t)) = 0. \quad \square \end{aligned}$$

Lemma 2.5. *The following is an exceptional collection in $\mathbf{D}^b(X)$:*

$$(\mathcal{A}, \mathcal{A}(1), \mathcal{B}(2), \dots, \mathcal{B}(9)). \tag{2.13}$$

Proof. By the previous lemma and thanks to Serre duality, we will be done once we prove $\text{Ext}_X^\bullet(Q, Q(-1)) = 0$ (which we did in Lemma 2.3) and:

$$\text{Ext}_X^\bullet(\mathcal{O}_X, Q(-t)) = \text{Ext}_X^\bullet(\mathcal{U}_{\omega_1}, Q(-t)) = \text{Ext}_X^\bullet(P, Q(-t)) = 0,$$

for $1 \leq t \leq 10$. Looking at the sequence (2.3) defining Q , we see that BBW directly implies $H^\bullet(Q(-t)) = 0$ for $1 \leq t \leq 10$. As for $\text{Ext}_X^\bullet(\mathcal{U}_{\omega_1}, Q(-t)) = 0$, note that the case $t = 10$ is (2.8) by Serre duality. On the other hand, for $1 \leq t \leq 9$, this follows from (2.1) and from the vanishing

$$H^\bullet(\mathcal{U}_{\omega_5} \otimes \mathcal{U}_{\omega_1}(-1-t)) = 0, \quad H^\bullet(\mathcal{U}_{\omega_5} \otimes \mathcal{U}_{\omega_1+\omega_2}(-1-t)) = 0,$$

for $1 \leq t \leq 9$, which in turn is a consequence of (2.9), (2.10), (2.11) and (2.12).

Finally, let us show that $\text{Ext}_X^\bullet(P, Q(-t)) = 0$. For $t \neq 10$, this follows from BBW, (2.1) and from the isomorphisms:

$$\begin{aligned} \mathcal{U}_{\omega_4} \otimes \mathcal{U}_{\omega_1} &\simeq \mathcal{U}_{\omega_5}(1) \oplus \mathcal{U}_{\omega_1+\omega_4}, \\ \mathcal{U}_{\omega_4} \otimes \mathcal{U}_{\omega_1+\omega_2} &\simeq \mathcal{U}_{\omega_1}(2) \oplus \mathcal{U}_{\omega_2+\omega_5}(1) \oplus \mathcal{U}_{\omega_1+\omega_2+\omega_4} \oplus \mathcal{U}_{2\omega_1+\omega_5}. \end{aligned}$$

For $t = 10$, the statement is equivalent to $\text{Ext}_X^\bullet(Q, P) = 0$. To check this last vanishing, using the isomorphisms of the previous display, we are reduced to show $\text{Ext}_X^\bullet(Q, \mathcal{U}_{\omega_2}) = 0$ and in turn to show that cupping with $\zeta \in \text{Ext}_X^1(\mathcal{U}_{\omega_1+\omega_2}, \mathcal{U}_{\omega_1}) \simeq H^1(\mathcal{U}_{\omega_4}(-2)) \simeq \mathbb{C}$ induces an isomorphism:

$$V^{\omega_1} \simeq H^0(\mathcal{U}_{\omega_1}) \simeq \text{Hom}_X(\mathcal{U}_{\omega_1}, \mathcal{U}_{\omega_2}) \rightarrow \text{Ext}_X^1(\mathcal{U}_{\omega_1+\omega_2}, \mathcal{U}_{\omega_2}) \simeq H^1(\mathcal{U}_{\omega_1+\omega_4}(-2)) \simeq V^{\omega_1}$$

Then we have to show that the cup product map below is an isomorphism

$$H^1(\mathcal{U}_{\omega_4-2\omega_6}) \otimes H^0(\mathcal{U}_{\omega_1}) \rightarrow H^1(\mathcal{U}_{\omega_1+\omega_4-2\omega_6}).$$

This follows from Lemma 2.1, taking w to be the reflection about the root α_6 . \square

3. Warmup for fullness

Let \mathcal{D} be the full triangulated subcategory of $\mathbf{D}^b(X)$ generated by our exceptional collection, i.e. define

$$\mathcal{D} = \langle \mathcal{A}, \mathcal{A}(1), \mathcal{B}(2), \dots, \mathcal{B}(9) \rangle \subset \mathbf{D}^b(X).$$

Thus, we have a semiorthogonal decomposition

$$\mathbf{D}^b(X) = \langle \mathcal{D}^\perp, \mathcal{D} \rangle.$$

Our aim is to prove $\mathcal{D}^\perp = 0$. To achieve this, we will restrict to a covering family of smaller spinor varieties whose derived category is well-known and prove that any object orthogonal to \mathcal{D} restricts to zero over such varieties by showing that the structure sheaf of these subvarieties is resolved by objects in \mathcal{D} . This technique has already been used in the literature, see for instance [27,21,10,13]. After describing such a covering family,

we will show fullness in the easier and well known case of spinor 10-folds as a warmup. In doing so, we will make use of an exact complex appearing in Section 4, pointing out that the existence of such complex is fundamental for our proof of fullness.

3.1. A covering family of spinor varieties

Let us come back to the general case of a vector space V of dimension $2n$. Let

$$q: V \times V \rightarrow \mathbb{C}$$

be the symmetric bilinear form defining $X := \text{Spin}_{2n}/\mathbb{P}_n$, i.e. we have

$$X = OG_+(n, V).$$

Recall that $\mathcal{U} = \mathcal{U}_{\omega_1}$ we have

$$H^0(X, \mathcal{U}^\vee) = V^\vee \xrightarrow{\simeq} V.$$

Since q is non-degenerate, there is a bijection between elements $w \in V$ and sections $s_w \in H^0(X, \mathcal{U}^\vee) = V^\vee$ that sends w to $s_w = q(w, \cdot)$. It is easy to see that we have

$$q|_W \text{ is non-degenerate} \iff q(w, w) \neq 0, \tag{3.1}$$

where $W = \ker s_w$. If s_w satisfies (3.1), then we can define two things:

1. A morphism of algebraic varieties

$$\begin{aligned} \varphi_w: OG_+(n, V) &\rightarrow OG(n-1, W) \\ U &\mapsto U \cap W. \end{aligned} \tag{3.2}$$

2. The natural morphism $s_w: \mathcal{O}_X \rightarrow \mathcal{U}^\vee$ does not vanish anywhere (since there are no n -dimensional isotropic subspaces in W) and, therefore, defines a short exact sequence of vector bundles

$$0 \rightarrow \mathcal{O}_X \xrightarrow{s_w} \mathcal{U}^\vee \rightarrow \mathcal{E}^\vee \rightarrow 0, \tag{3.3}$$

where \mathcal{E}^\vee is a vector bundle of rank $n-1$ with $H^0(X, \mathcal{E}^\vee) = W^\vee$.

Let us fix a section s_w satisfying (3.1). First recall the very classical identification of $OG_+(n, 2n)$ and $OG(n-1, 2n-1)$. We give a proof of it in the next lemma for the reader's convenience.

Lemma 3.1. *If s_w satisfies (3.1), the morphism φ_w is an isomorphism.*

Proof. Since both varieties are smooth and we are in characteristic zero, it suffices to show that the morphism is one to one. Let $[T] \in OG(n - 1, W)$; the preimage of $[T]$ via φ_w is given by those $[U] \in OG_+(n, V)$ such that $U \cap W = T$. This is equivalent to $T \subset U$ since isotropic subspaces of W have dimension at most equal to $n - 1$. Thus the set of isotropic subspaces U in V containing T are parametrized by the zero-dimensional quadric inside $\mathbb{P}(T^\perp/T) \cong \mathbb{P}^1$, i.e. two points U_+ and U_- . These spaces intersect exactly in T , i.e. in codimension one, so they must belong to different components of the set of maximal isotropic spaces in V ; we can assume that $[U_+] \in OG_+(n, V)$ and $[U_-] \in OG_-(n, V)$. Thus the preimage of $[T]$ via φ_w consists of the only point $[U_+]$ for every $T \in OG(n - 1, W)$, and the morphism is one to one. \square

Lemma 3.2. *Let $s \in H^0(X, \mathcal{E}^\vee) = W^\vee$ and consider the zero-locus Y_s of s .*

i) *If s is general enough, then $Y_s \simeq OG(n - 1, 2n - 2)$. Let us denote the inclusion by*

$$i_s: Y_s \rightarrow X.$$

ii) *For any section s as in i), we have*

$$i_s^* \mathcal{U}^\vee = \mathcal{U}_{n-1}^\vee \oplus \mathcal{O}_{Y_s}$$

iii) *Varying $s \in H^0(X, \mathcal{E}^\vee)$ as above we can cover X by copies of $OG(n - 1, 2n - 2)$.*

iv) *If for any object $F \in \mathbf{D}^b(X)$ the restrictions $i_s^* F$ vanish for all $s \in H^0(X, \mathcal{E}^\vee)$ as above, then $F = 0$.*

Proof. For s to be general enough, it is enough to satisfy the analogue of (3.1). i.e. the restriction of q to $L := \ker(s) \subset W$ should be non-degenerate.

- i) Under (3.2) the vector bundle \mathcal{E}^\vee corresponds to the dual of the tautological subbundle on $OG(n - 1, W)$. Hence, we get the claim.
- ii) Under (3.2) the sequence (3.3) shows that there is a non-trivial extension between the dual of the tautological subbundle and the structure sheaf on $OG(n - 1, W)$. However, by BBW on $OG(n - 1, L) = OG(n - 1, 2n - 2)$ such extensions vanish and the sequence splits.
- iii) Indeed, for any $(n - 1)$ -dimensional isotropic subspace $U_{n-1} \subset W$ we can consider U_{n-1}^\perp , take any element $u \in U_{n-1}^\perp \setminus U_{n-1}$ and take $L = u^\perp$. Clearly we have $U_{n-1} \subset L$ and $q|_L$ is non-degenerate by (3.1).
- iv) This is [21, Lemma 4.5]. \square

3.2. Full exceptional collection on the spinor 10-fold

The orthogonal Grassmannian $Y = OG(5, 10)$ has two connected components that we denote by

$$Y_- = OG_-(5, 10) \quad \text{and} \quad Y_+ = OG_+(5, 10).$$

These components are isomorphic to each other, we call them spinor 10-folds.

As usual, on $Y = OG(5, 10)$ we can consider the tautological subbundle \mathcal{U}_5 of rank 5. We denote as $\mathcal{U}_{5,\pm} := \mathcal{U}_5|_{Y_{\pm}}$ its restrictions to Y_{\pm} .

We prove the following result as a useful warm-up to the case of $X = \text{Spin}_{12}/\mathbb{P}_6 = OG_+(6, 12)$.

Theorem 3.3. *We have*

$$\mathbf{D}^b(Y_{\pm}) = \left\langle \mathcal{O}, \mathcal{U}_{5,\pm}^{\vee}, \mathcal{O}(1), \mathcal{U}_{5,\pm}^{\vee}(1), \dots, \mathcal{O}(7), \mathcal{U}_{5,\pm}^{\vee}(7) \right\rangle$$

Proof. Let us fix the + sign and let us define $\mathcal{D}_5 := \langle \mathcal{O}, \mathcal{U}_5^{\vee}, \mathcal{O}(1), \mathcal{U}_5^{\vee}(1), \dots, \mathcal{O}(7), \mathcal{U}_5^{\vee}(7) \rangle$ on $Y := Y_+$. Let us take an object $F \in \mathcal{D}_5^{\perp}$, i.e. we have

$$\text{Ext}_X^{\bullet}(A, F) = 0 \quad \text{for any} \quad A \in \mathcal{D}_5.$$

Let $s \in H^0(Y, \mathcal{E}^{\vee})$ be a general section and $i_s: Z_s \rightarrow Y$ the embedding of its zero locus, as in Lemma 3.2(1).

Let us consider the set of vector bundles on Y defined by

$$\Upsilon_5 := \{ \mathcal{O}_Y(t) \mid t \in [2, 7] \} \cup \{ \mathcal{U}_5^{\vee}(2) \}.$$

Let us denote by \mathcal{E}_5^{\vee} the vector bundle defined in (3.3). We claim that for any $E \in \Upsilon_5$ and any j the bundle $E \otimes \wedge^j \mathcal{E}_5^{\vee}$ lies in \mathcal{D}_5 . Let us for the moment assume that the claim is true. Then we have

$$\text{Ext}_X^{\bullet}(E \otimes \wedge^j \mathcal{E}_5^{\vee}, F) = H^{\bullet}(Y, \wedge^j \mathcal{E}_5 \otimes E^{\vee} \otimes F) = 0 \quad \text{for all } j,$$

and making use of the Koszul complex

$$0 \rightarrow \wedge^4 \mathcal{E}_5 \rightarrow \dots \rightarrow \mathcal{E}_5 \rightarrow \mathcal{O}_Y \rightarrow i_{s*} \mathcal{O}_{Z_s} \rightarrow 0,$$

we obtain

$$H^{\bullet}(Y, (E^{\vee} \otimes F) \otimes i_{s*} \mathcal{O}_Z) = 0.$$

Now, by the projection formula we rewrite

$$H^{\bullet}(Y, (E^{\vee} \otimes F) \otimes i_{s*} \mathcal{O}_Z) = H^{\bullet}(Z_s, i_s^*(E^{\vee} \otimes F)) = \text{Ext}_Z^{\bullet}(i_s^* E, i_s^* F) = 0.$$

Recall that $Z_s \simeq OG(4, 8)$ has two connected components Z_{s+} and Z_{s-} which are two six-dimensional quadrics. We denote the compositions $Z_{s\pm} \subset Z_s \xrightarrow{i_s} Y$ by $i_{s\pm}$. Using this notation we have

$$\text{Ext}_{Z_s}^\bullet(i_s^*E, i_s^*F) = \text{Ext}_{Z_{s+}}^\bullet(i_{s+}^*E, i_{s+}^*F) \oplus \text{Ext}_{Z_{s-}}^\bullet(i_{s-}^*E, i_{s-}^*F).$$

Hence, we have

$$\text{Ext}_{Z_{s+}}^\bullet(i_{s+}^*E, i_{s+}^*F) = 0 \quad \text{and} \quad \text{Ext}_{Z_{s-}}^\bullet(i_{s-}^*E, i_{s-}^*F) = 0.$$

Applying Lemma 3.2(2) and the fact that the six-dimensional quadrics $Z_{s\pm}$ admit the following full exceptional collection (see [14])

$$\mathbf{D}^b(Z_{s\pm}) = \langle \mathcal{O}(2), \mathcal{U}_{4,\pm}^\vee(2), \mathcal{O}(3), \dots, \mathcal{O}(7) \rangle,$$

we obtain

$$i_{s+}^*F = 0 \quad \text{and} \quad i_{s-}^*F = 0.$$

Hence, we conclude $i_s^*F = 0$. Finally, since the above argument works for any general $s \in H^0(Y, \mathcal{E}_5^\vee)$, by Lemma 3.2(iii,iv) we obtain $F = 0$.

Now, let us prove the claim. We need to prove that $\wedge^j \mathcal{E}_5^\vee(t) \in \mathcal{D}_5$ for $t \in [2, 7]$ and $\mathcal{U}_5^\vee \otimes \wedge^j \mathcal{E}_5^\vee(2) \in \mathcal{D}_5$ for all possible j 's. From the exact sequence

$$0 \rightarrow \mathcal{O}_Y \rightarrow \mathcal{U}_5^\vee \rightarrow \mathcal{E}_5^\vee \rightarrow 0$$

we deduce that our claim is implied by the fact that $\wedge^j \mathcal{U}_5^\vee(t) \in \mathcal{D}_5$ for $t \in [2, 7]$ and $\mathcal{U}_5^\vee \otimes \wedge^j \mathcal{U}_5^\vee(2) \in \mathcal{D}_5$ for all possible j 's.

The bundles $\mathcal{O}(t)$ and $\mathcal{U}_5^\vee(t)$ for $t \in [0, 7]$ generate \mathcal{D}_5 . From the exact sequence $0 \rightarrow \mathcal{U}_5 \rightarrow V_{10} \otimes \mathcal{O}_Y \rightarrow \mathcal{U}_5^\vee \rightarrow 0$, where V_{10} is a ten-dimensional vector space, we deduce that $\mathcal{U}_5(t) \in \mathcal{D}_5$ for $t \in [0, 7]$. Thus $\mathcal{O}_Y(t), \mathcal{U}_5(t), \wedge^4 \mathcal{U}_5(t) = \mathcal{U}^\vee(t-2), \wedge^5 \mathcal{U}_5(t) = \mathcal{O}_Y(t-2)$ all belong to \mathcal{D}_5 for $t \in [2, 7]$.

Denote by S^+ (respectively S^-) the even (resp. odd) Spin representation for the Spin group - see also Section 4.1.1. Recall that $\mathbf{ss}(S^- \otimes \mathcal{O}_Y) = \mathcal{U}_5(-1) \oplus \wedge^3 \mathcal{U}_5^\vee(-1) \oplus \mathcal{O}_Y(1)$, where \mathbf{ss} denotes the semisimplification of the bundle, so we deduce that $\wedge^3 \mathcal{U}_5^\vee(t-2) = \wedge^2 \mathcal{U}(t) \in \mathcal{D}_5$ for $t \in [2, 7]$. Similarly the fact that $\mathbf{ss}(S^+ \otimes \mathcal{O}_Y) = \mathcal{O}_Y(-1) \oplus \wedge^2 \mathcal{U}_5^\vee(-1) \oplus \mathcal{U}_5(1)$ implies that $\wedge^2 \mathcal{U}_5^\vee(t-2) = \wedge^3 \mathcal{U}(t) \in \mathcal{D}_5$ for $t \in [2, 7]$.

Now we need to deal with $\mathcal{U}_5^\vee \otimes \wedge^j \mathcal{U}_5(2)$. When $j = 0$ and $j = 5$, $\mathcal{U}_5^\vee \otimes \wedge^j \mathcal{U}_5(2) \in \mathcal{D}_5$. For $j = 1$ use the decomposition $\mathbf{ss}(\wedge^2 V_{10} \otimes \mathcal{O}_Y) = \wedge^2 \mathcal{U}_5 \oplus \mathcal{U}_5 \otimes \mathcal{U}_5^\vee \oplus \wedge^2 \mathcal{U}_5^\vee$ to deduce that $\mathcal{U}_5 \otimes \mathcal{U}_5^\vee(t) \in \mathcal{D}_5$ for $t \in [2, 5]$. For $j = 3$ use the decomposition $\mathbf{ss}(S^+ \otimes \mathcal{U}_5^\vee) = \mathcal{U}_5^\vee(-1) \oplus \wedge^3 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(1) \oplus \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(1)$ to deduce that $\wedge^3 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(t) \in \mathcal{D}_5$ for $t \in [2, 5]$. The cases $j = 2$ and $j = 4$ can be dealt with in parallel. Indeed one can use the decomposition $\mathbf{ss}(S^- \otimes \mathcal{U}_5^\vee) = \mathcal{U}_5^\vee(1) \oplus \wedge^2 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(1) \oplus \wedge^4 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(1)$ to deduce that, if $t \in [1, 6]$ then: $\wedge^2 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(t) \in \mathcal{D}_5$ if and only if $\wedge^4 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(t) \in \mathcal{D}_5$.

Finally, we want to prove for instance that $\wedge^2 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(2) \in \mathcal{D}_5$. For this the exact sequence appearing in Proposition 4.2 is crucial (Section 4.2 is independent of this proof, so we can use the results therein), and in particular the extension R_5 whose existence is

ensured by the same proposition. Indeed from that complex one deduces that $R_5(t) \in \mathcal{D}_5$ for $t \in [0, 5]$, which in turn implies that $\mathcal{U}_5^{\omega_1+\omega_2}(t) \in \mathcal{D}_5$ for $t \in [0, 3]$. Then, using the decomposition $\text{ss}(\wedge^2 \mathcal{U}_5^\vee \otimes V_{10}) = \mathcal{U}_5^{\omega_1+\omega_2} \oplus \wedge^3 \mathcal{U}_5^\vee \oplus \mathcal{U}_5^\vee \oplus \mathcal{U}_5^{\omega_2+\omega_4}(-2)$, we obtain that $\mathcal{U}_5^{\omega_2+\omega_4}(t) \in \mathcal{D}_5$ for $t \in [-2, 1]$. Notice also that previously we showed that $\mathcal{U}_5 \otimes \mathcal{U}_5^\vee(t) = \mathcal{O}_Y(t) \oplus \mathcal{U}_5^{\omega_1+\omega_4}(t-2) \in \mathcal{D}_5$ for $t \in [2, 5]$, so $\mathcal{U}_5^{\omega_1+\omega_4}(t) \in \mathcal{D}_5$ for $t \in [0, 3]$. These two facts imply that $\mathcal{U}_5^{\omega_1+\omega_3}(t) \in \mathcal{D}_5$ for $t = 0, 1$ because of the decomposition $\text{ss}(\mathcal{U}_5^{\omega_1+\omega_4}(-2) \otimes V_{10}) = \mathcal{U}_5^{\omega_1+\omega_3}(-2) \oplus \mathcal{U}_5^{\omega_2+\omega_4}(-2) \oplus \mathcal{U}_5 \oplus \mathcal{U}_5^\vee$. Then from the decomposition $\text{ss}(\wedge^2 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee) = \mathcal{U}_5^{\omega_1+\omega_3}(-2) \oplus \mathcal{U}_5$ we deduce that $\wedge^2 \mathcal{U}_5 \otimes \mathcal{U}_5^\vee(t) \in \mathcal{D}_5$ for $t = 2, 3$. \square

Remark 3.4. Theorem 3.3 was already known from [18, Section 6.2]. The proof given here is more direct and corresponds better to our approach.

4. Dissecting Spin bundles

In this section, we look more closely to the vector bundles on $X = OG_+(6, 12)$ induced by the spinor representations. The main goal is to prove Proposition 4.2 and 4.3, which in turn will be used in Section 5 in view of showing fullness of our collection. We often abbreviate \mathcal{O}_X to \mathcal{O} .

4.1. The Spin representations

In the following we will recall a selection of generalities about the Clifford algebra and Spin representations that can be found, for instance, in [24]. Let us begin with an even-dimensional vector space V endowed with a non-degenerate symmetric form q . The Clifford algebra is defined as the quotient of the tensor algebra V^{\otimes} by all relations of the form $v \otimes v - q(v, v)$ for $v \in V$. Notice that both V and $\wedge^2 V \simeq \mathfrak{so}_V$ embed inside the Clifford algebra.

4.1.1. The Spin representation and exterior powers

Let us fix a maximal isotropic subspace U of V . Any other maximal isotropic subspace intersecting U transversally can be identified through q with U^\vee ; we thus get a decomposition of $V = U \oplus U^\vee$. The Spin representations can be identified, as vector spaces, as follows:

$$S^+ := \wedge^+ U^\vee = \bigoplus_i \wedge^{2i} U^\vee,$$

$$S^- := \wedge^- U^\vee = \bigoplus_i \wedge^{2i+1} U^\vee.$$

There is a natural action

$$\eta_\pm := V \otimes S^\pm \rightarrow S^\mp$$

defined as follows: $\eta_{\pm}(v \otimes \omega) = v \wedge \omega$ if $v \in U^{\vee}$ and $\eta_{\pm}(v \otimes \omega) = v \lrcorner \omega$ if $v \in U$, where \lrcorner is the contraction. This induces an action of the Clifford algebra, and hence of \mathfrak{so}_V , on S^{\pm} , which endows this vector space with a structure of Spin-representation; S^{\pm} are the so-called Spin representations. It turns out that, if the dimension of V is $2n$, then $(S^{\pm})^{\vee} = S^{(-1)^n \pm}$ as representations. Moreover if n is odd then $S^+ = V^{\omega_{n-1}}$ and $S^- = V^{\omega_n}$ while if n is even then $S^+ = V^{\omega_n}$ and $S^- = V^{\omega_{n-1}}$. Notice moreover that the action η_{\pm} naturally induces a Spin-equivariant morphism $\eta_{\pm}^{\otimes i} : V^{\otimes i} \otimes S^{\pm} \rightarrow S^{(-1)^i \pm}$, and thus also a Spin-equivariant morphism

$$\wedge^i \eta_{\pm} : \wedge^i V \otimes S^{\pm} \rightarrow S^{(-1)^i \pm}.$$

In the following we want to use the morphism $\wedge^i \eta_{\pm}$ to construct some exact complexes on $\text{Spin}_{2n}/\mathbb{P}_n$ for $n = 5, 6$. Before doing so, we will recall basic linear algebra facts in order to explain how to rewrite $\wedge^i \eta_{\pm}$ as a morphism $\xi : S^{\pm} \otimes (S^{(-1)^i \pm})^{\vee} \rightarrow \wedge^i V^{\vee} \simeq \wedge^i V$.

4.1.2. Linear algebra digression

Let us begin with a linear morphism $u : A \otimes B \rightarrow C$ for three vector spaces A, B, C . This means that $u \in A \otimes B \otimes C^{\vee} = \text{Hom}(A \otimes B, C^{\vee})$, so it defines another morphism $t : A \otimes C^{\vee} \rightarrow B^{\vee}$. Clearly one can recover u from t as well.

Lemma 4.1. *$\text{Im}(u)^{\perp}$ is identified with the subspace $\{x \in C^{\vee} \mid t(a, x) = 0 \ \forall a \in A\} \subset C^{\vee}$.*

Proof. Let us denote by D the above subspace. By definition of t , for any $x \in C^{\vee}$, $a \in A$ and $b \in B$, $x(u(a \otimes b)) = t(a \otimes x)(b)$. It is straightforward to deduce that $x \in \text{Im}(u)^{\perp}$ if and only if $x \in D$. \square

4.2. Spinor bundles

Let us consider the variety $\text{Spin}_{2n}/\mathbb{P}_n = \text{OG}_+(n, 2n)$ which is one of the two isomorphic connected components of the variety parametrizing maximal isotropic subspaces of V . Let us denote by $\epsilon := (n \bmod 2)$. The line bundle $\mathcal{O}(1) = \mathcal{U}_{\omega_n}$ embeds $\text{Spin}_{2n}/\mathbb{P}_n$ inside $\mathbb{P}(V^{\omega_n}) = \mathbb{P}(S^{(-1)^{\epsilon}})$. Thus $\mathcal{O}(1)$ is a G -equivariant quotient of $S^{(-1)^{\epsilon}} \otimes \mathcal{O}$. In fact, one can construct a filtration of G -equivariant vector bundles

$$0 =: \mathcal{F}_0 \subset \mathcal{F}_1 \subset \dots \subset \mathcal{F}_{[n]+\epsilon} := S^{(-1)^{\epsilon}}$$

such that

$$\mathcal{F}_{i+1}/\mathcal{F}_i = (\wedge^{2i+\epsilon} \mathcal{U}^{\vee})(-1).$$

This is the relative version of the filtration of $S^{(-1)^{\epsilon}} = \wedge^+ U^{\vee}$ given by the subspaces $F_{i+1} := \sum_{j \leq i} \wedge^{2j+\epsilon} U^{\vee}$. A similar filtration exists for $S^{(-1)^{\epsilon}} \otimes \mathcal{O}$. For instance, we get that $\mathcal{F}_1 = \mathcal{U}^{\vee}(-1)$ is a subbundle of $S^- \otimes \mathcal{O}$. This filtration was described in [21, Proposition 6.3].

4.2.1. An exact complex in low dimension

We will now construct an exact complex of vector spaces using the morphisms $\wedge^i \eta_{\pm}$ when $n = 5$ and $n = 6$. We believe that this type of complexes can be generalized for higher n and will be crucial in proving fullness of exceptional collections on $\text{Spin}_{2n}/\mathbb{P}_n$ for higher n . From now on we fix $\eta := \eta_+$

4.2.2. The case $n = 5$

In this case we have the following decomposition of representations: $S^+ \otimes S^- = \mathbb{C} \oplus \wedge^2 V \oplus V^{\omega_4 + \omega_5}$. This implies that there exists a unique G -equivariant morphism $S^+ \otimes S^- \rightarrow \wedge^2 V$, which must then be equal to $\wedge^2 \eta$ (notice that $(S^-)^\vee = S^+$ since n is odd). As a consequence of BBW $H^0(\mathcal{U}(1)) = S^-$ and thus there exists a unique G -equivariant morphism $\mathcal{U}^\vee(-1) \otimes S^+ \rightarrow \wedge^2 V$. This morphism must then be the composition $\wedge^2 \eta \circ (i \otimes \text{id})$ where i is the inclusion $i : \mathcal{U}^\vee(-1) \rightarrow S^- \otimes \mathcal{O}$. All in all we get a G -equivariant morphism $\wedge^2 \eta \circ (i \otimes \text{id}) : \mathcal{U}^\vee(-1) \otimes S^+ \rightarrow \wedge^2 V$. The aim of this section is to prove the following:

Proposition 4.2. *There exists a G -equivariant extension*

$$0 \rightarrow \mathcal{U}^\vee(-2) \rightarrow R_5 \rightarrow \mathcal{U}_{\omega_1 + \omega_2}(-2) \rightarrow 0$$

and a G -equivariant exact complex of vector bundles

$$0 \rightarrow R_5 \rightarrow \mathcal{U}^\vee(-1) \otimes S^+ \rightarrow \wedge^2 V \otimes \mathcal{O} \rightarrow \wedge^2 \mathcal{U}^\vee \rightarrow 0,$$

where the central map is $\wedge^2 \eta \circ (i \otimes \text{id})$.

Proof. The morphism $\wedge^2 V \otimes \mathcal{O} \rightarrow \wedge^2 \mathcal{U}^\vee$ above is the natural projection induced by the exact sequence

$$0 \rightarrow \mathcal{U} \rightarrow V \otimes \mathcal{O} \rightarrow \mathcal{U}^\vee \rightarrow 0.$$

Since this map, as well as $\wedge^2 \eta \circ (i \otimes \text{id})$, is a G -equivariant morphism of G -homogeneous vector bundles, it is sufficient to restrict to any fiber of $\text{Spin}_{10}/\mathbb{P}_5$ to prove exactness. More precisely we will show that, if $[U] \in \text{Spin}_{10}/\mathbb{P}_5$, the induced complex of vector spaces

$$(\mathcal{U}^\vee(-1) \otimes S^+) |_{[U]} \rightarrow (\wedge^2 V \otimes \mathcal{O}) |_{[U]} \rightarrow (\wedge^2 \mathcal{U}^\vee) |_{[U]} \rightarrow 0 \tag{4.1}$$

is exact. From this it will follow that the complex

$$\mathcal{U}^\vee(-1) \otimes S^+ \rightarrow \wedge^2 V \otimes \mathcal{O} \rightarrow \wedge^2 \mathcal{U}^\vee \rightarrow 0$$

is exact. The result will then follow by noticing that, since $\text{ss}(\mathcal{U}^\vee(-1) \otimes S^+) = \mathcal{U}^\vee(-2) \oplus \mathcal{U}_{\omega_1 + \omega_2}(-2) \oplus \wedge^3 \mathcal{U}^\vee(-2) \oplus \mathfrak{sl}(\mathcal{U}) \oplus \mathcal{O}$ and $\text{ss}(\wedge^2 V) = \wedge^2 \mathcal{U} \oplus \mathcal{O} \oplus \mathfrak{sl}(\mathcal{U}) \oplus \wedge^2 \mathcal{U}^\vee$, the

semisimple reduction of the kernel of $\wedge^2 \eta \circ (i \otimes \text{id})$ is necessarily equal to $\mathcal{U}^\vee(-2) \oplus \mathcal{U}_{\omega_1 + \omega_2}(-2)$.

Let $[U] \in \text{Spin}_{10}/P_5$ be any point. Then $(\mathcal{U}^\vee(-1))|_{[U]} \simeq U^\vee$ is a subspace of $(S^- \otimes \mathcal{O})|_{[U]} = S^- = \bigoplus_i \wedge^{2i+1} U^\vee$ - here the last equality only holds as an equality of $L(P_5)$ -representations. Following the linear algebra digression, the morphism $\xi \circ (i \otimes \text{id})$ corresponds to the morphism $\wedge^i \eta \circ (i \otimes \text{id})$. Moreover, letting

$$t := (\wedge^2 \eta \circ (i \otimes \text{id}))|_{[U]} : U^\vee \otimes \wedge^2 V^\vee \rightarrow S^- = \bigoplus_i \wedge^{2i+1} U^\vee$$

and

$$u := (\xi \circ (i \otimes \text{id}))|_{[U]} : U^\vee \otimes S^+ \rightarrow \wedge^2 V$$

and applying Lemma 4.1, we deduce the following:

$$\text{Im}(u)^\perp = \{v \in \wedge^2 V^\vee \mid \forall f \in U^\vee \subset S^- = \bigoplus_i \wedge^{2i+1} U^\vee, t(f \otimes v) = 0\} \subset \wedge^2 V^\vee.$$

We bothered going through all of this because we have a very explicit description of the map t , which is the one induced by η ; let us see how to use it. First notice that t is a P_5 -equivariant morphism, so in particular let us treat it as a $L(P_5)$ -equivariant morphism. Thus we can decompose $\wedge^2 V = \wedge^2 U \oplus (U \otimes U^\vee) \oplus \wedge^2 U^\vee = \wedge^2 U \oplus \mathbb{C} \oplus \mathfrak{sl}(U) \oplus \wedge^2 U^\vee$. By $L(P_5)$ -equivariance, each of these factors is either completely contained in $\text{Im}(u)^\perp$ or intersects $\text{Im}(u)^\perp$ trivially. In order to distinguish the two cases it is thus sufficient to decide whether a non-zero vector in a given factor belongs to $\text{Im}(u)^\perp$ or not. We thus have four cases to deal with. We will denote by u_1, \dots, u_5 a basis of U and by w_1, \dots, w_5 the dual basis. We will denote by $u_{ij} = u_i \wedge u_j$ and by $w_{ij} = w_i \wedge w_j$; $\delta_{i,j}$ will denote Kronecker's delta.

$\wedge^2 U$: Let $0 \neq u_{ij} \in \wedge^2 U$ and $w_k \in U^\vee$. Then $t(w_k \otimes u_{ij}) = u_{i \lrcorner} (u_j \lrcorner w_k) - u_j \lrcorner (u_i \lrcorner w_k) = 0$, for any $k = 1, \dots, 5$, so $\wedge^2 U \subset \text{Im}(u)^\perp$.

\mathbb{C} : Let $0 \neq \sum_i u_i \wedge w_i \in \mathbb{C} \subset \wedge^2 V$ and $w_k \in U^\vee$. Then $t(w_k \otimes (\sum_i u_i \wedge w_i)) = \sum_i (u_i \lrcorner (w_{ik}) - \delta_{i,k} w_k) = \sum_i ((1 - \delta_{i,k}) w_k - \delta_{i,k} w_k) = \sum_i (1 - 2\delta_{i,k}) w_k = 2w_k \neq 0$, so $\mathbb{C} \cap \text{Im}(u)^\perp = 0$.

$\mathfrak{sl}(U)$: Let $0 \neq u_i \wedge w_j \in \mathfrak{sl}(U)$ for $i \neq j$, and $w_k \in U^\vee$. Then $t(w_k \otimes u_i \wedge w_j) = u_i \lrcorner (w_{jk}) - (u_i \lrcorner w_k) w_j = -2\delta_{i,k} w_k \neq 0$, so $\mathfrak{sl}(U) \cap \text{Im}(u)^\perp = 0$.

$\wedge^2 U^\vee$: Let $0 \neq w_{ij} \in \wedge^2 U^\vee$ and $w_k \in U^\vee$. Then $t(w_k \otimes w_{ij}) = w_{ijk} \neq 0$, so $\wedge^2 U^\vee \cap \text{Im}(u)^\perp = 0$.

The previous computations imply that $\text{Im}(u)^\perp = \wedge^2 U \subset \wedge^2 V \simeq \wedge^2 V^\vee$. This is equivalent to the fact that $\text{Im}(u)$ is the kernel of $\wedge^2 V \rightarrow \wedge^2 U^\vee$. Moreover the latter morphism is clearly surjective, so we deduce that the complex in (4.1) is exact. The statement of the proposition follows. \square

4.2.3. The case $n = 6$

In this case we have the following decomposition of representations: $S^+ \otimes S^- = V \oplus \wedge^3 V \oplus V^{\omega_5 + \omega_6}$. This implies that there exists a unique G -equivariant morphism $S^+ \otimes S^- \rightarrow \wedge^3 V$, which must then be equal to $\wedge^3 \eta$ (notice that $(S^+)^{\vee} = S^+$ since n is even). As a consequence of the BBW Theorem $H^0(\mathcal{U}(1)) = S^-$ and thus there exists a unique G -equivariant morphism $\mathcal{U}^{\vee}(-1) \otimes S^+ \rightarrow \wedge^3 V$. This morphism must then be the composition $\wedge^3 \eta \circ (i \otimes \text{id})$ where i is the inclusion $i : \mathcal{U}^{\vee}(-1) \rightarrow S^- \otimes \mathcal{O}$. All in all we get a G -equivariant morphism $\wedge^3 \eta \circ (i \otimes \text{id}) : \mathcal{U}^{\vee}(-1) \otimes S^+ \rightarrow \wedge^3 V$. The aim of this section is to prove an analogue of Proposition 4.2. In order to do so, let us begin by defining the vector bundle \mathcal{C} as the cokernel of the unique G -equivariant inclusion $\mathcal{U}^{\vee} \rightarrow V \otimes \wedge^2 \mathcal{U}^{\vee}$; we thus have an exact sequence

$$0 \rightarrow \mathcal{U}^{\vee} \rightarrow V \otimes \wedge^2 \mathcal{U}^{\vee} \rightarrow \mathcal{C} \rightarrow 0.$$

Proposition 4.3. *There exists a G -equivariant extension R_6 whose semisimple reduction is*

$$\text{ss}(R_6) = \mathcal{U}^{\vee}(-2) \oplus \mathcal{U}_{\omega_1 + \omega_2}(-2)$$

and a G -equivariant exact complex of vector bundles

$$0 \rightarrow R_6 \rightarrow \mathcal{U}^{\vee}(-1) \otimes S^+ \rightarrow \wedge^3 V \otimes \mathcal{O} \rightarrow \mathcal{C} \rightarrow \mathcal{U}_{\omega_1 + \omega_2} \rightarrow 0,$$

where the map $\mathcal{U}^{\vee}(-1) \otimes S^+ \rightarrow \wedge^3 V \otimes \mathcal{O}$ is $\wedge^3 \eta \circ (i \otimes \text{id})$.

Proof. The morphism $\wedge^3 V \otimes \mathcal{O} \rightarrow \mathcal{C}$ is the unique G -equivariant morphism and it is the one induced on the quotient from the natural one $\wedge^3 V \otimes \mathcal{O} \rightarrow V \otimes \wedge^2 \mathcal{U}^{\vee}$. Since the cokernel of the latter is $\mathcal{U}_{\omega_1 + \omega_2}$, this is also the cokernel of the former. Since the morphism $\wedge^2 \eta \circ (i \otimes \text{id})$ is a G -equivariant morphism of G -homogeneous vector bundles, it is sufficient to restrict to any fiber of X to prove exactness, as we did in the proof of Proposition 4.2. More precisely we will show that, if $[U] \in X$, the induced complex of vector spaces

$$(\mathcal{U}^{\vee}(-1) \otimes S^+)_{|[U]} \rightarrow (\wedge^3 V \otimes \mathcal{O})_{|[U]} \rightarrow \mathcal{C}_{|[U]}$$

is exact. From this it will follow that the complex

$$\mathcal{U}^{\vee}(-1) \otimes S^+ \rightarrow \wedge^3 V \otimes \mathcal{O} \rightarrow \mathcal{C} \rightarrow \mathcal{U}_{\omega_1 + \omega_2} \rightarrow 0$$

is exact. The result will then follow by noticing that, since $\text{ss}(\mathcal{U}^{\vee}(-1) \otimes S^+) = \mathcal{U}^{\vee}(-2) \oplus \mathcal{U}_{\omega_1 + \omega_2}(-2) \oplus \wedge^3 \mathcal{U}^{\vee}(-2) \oplus \mathcal{U}_{\omega_1 + \omega_4}(-2) \oplus \mathcal{U} \oplus \mathcal{U}^{\vee}$ and $\text{ss}(\wedge^3 V) = \wedge^3 \mathcal{U} \oplus \mathcal{U}_{\omega_1 + \omega_4}(-2) \oplus \mathcal{U} \oplus \mathcal{U}_{\omega_2 + \omega_5}(-2) \oplus \mathcal{U}^{\vee} \oplus \wedge^3 \mathcal{U}^{\vee}$, the semisimple reduction of the kernel of $\wedge^3 \eta \circ (i \otimes \text{id})$ is necessarily equal to $\mathcal{U}^{\vee}(-2) \oplus \mathcal{U}_{\omega_1 + \omega_2}(-2)$.

Let $[U] \in X$ be any point. Then $(U^\vee(-1))|_{[U]} \simeq U^\vee$ is a subspace of $(S^- \otimes \mathcal{O})|_{[U]} = S^- = \bigoplus_i \wedge^{2i+1} U^\vee$ as $L(\mathbb{P}_6)$ -representations. Letting

$$t := (\wedge^3 \eta \circ (i \otimes \text{id}))|_{[U]} : U^\vee \otimes \wedge^3 V^\vee \rightarrow S^+ = \bigoplus_i \wedge^{2i} U^\vee$$

and

$$u := (\xi \circ (i \otimes \text{id}))|_{[U]} : U^\vee \otimes S^+ \rightarrow \wedge^3 V$$

and applying Lemma 4.1, we deduce the following:

$$\text{Im}(u)^\perp = \{v \in \wedge^3 V^\vee \mid \forall f \in U^\vee \subset S^- = \bigoplus_i \wedge^{2i+1} U^\vee, t(f \otimes v) = 0\} \subset \wedge^3 V^\vee.$$

Since we can treat everything as $L(\mathbb{P}_6)$ -equivariant/homogeneous, we can decompose $\wedge^3 V = \wedge^3 U \oplus U^{\omega_1+\omega_4} \oplus U \oplus U^{\omega_2+\omega_5} \oplus U^\vee \oplus \wedge^3 U^\vee$ (here, by abuse of notation, we denoted by U^ω the $\text{SL}(U)$ -representation with highest weight ω). By $L(\mathbb{P}_6)$ -equivariance, each of these factors is either completely contained in $\text{Im}(u)^\perp$ or it intersects $\text{Im}(u)^\perp$ trivially. In order to distinguish the two cases it is thus sufficient to decide whether a non-zero vector in a given factor belongs to $\text{Im}(u)^\perp$ or not. We thus have six cases to deal with. We will denote by u_1, \dots, u_6 a basis of U and by w_1, \dots, w_6 the dual basis. We will denote by $u_{ijk} = u_i \wedge u_j \wedge u_k$, $u_{ij} = u_i \wedge u_j$, $w_{ij} = w_i \wedge w_j$ and $w_{ijk} = w_i \wedge w_j \wedge w_k$; $\delta_{i,j}$ will denote Kronecker's delta.

$\wedge^3 U$: Let $0 \neq u_{ijk} \in \wedge^3 U$ and $w_h \in U^\vee$. Then $t(w_h \otimes u_{ijk}) = u_{ij} \lrcorner (u_k \lrcorner w_k) - u_{ik} \lrcorner (u_j \lrcorner w_h) + u_{ij} \lrcorner (u_k \lrcorner w_h) = 0$, for any $k = 1, \dots, 6$, so $\wedge^3 U \subset \text{Im}(u)^\perp$.

$U^{\omega_1+\omega_4}$: Let $0 \neq u_{ij} \wedge w_k \in U^{\omega_1+\omega_4}$ for $i \neq k$ and $j \neq k$, and $w_h \in U^\vee$. Then $t(w_h \otimes u_{ij} \wedge w_k) = u_{ij} \lrcorner (w_{kh}) - (u_i \lrcorner w_k)(u_j \lrcorner w_h) + (u_j \lrcorner w_k)(u_i \lrcorner w_h) = 0$, so $U^{\omega_1+\omega_4} \subset \text{Im}(u)^\perp$.

U : Let $0 \neq \sum_i u_{ij} \wedge w_j \in U \subset \wedge^3 V$ and $w_h \in U^\vee$. Then $t(w_h \otimes (\sum_i u_{ij} \wedge w_j)) = \sum_i (u_{ij} \lrcorner w_{jh} - (u_i \lrcorner w_j)(u_j \lrcorner w_h) + (u_j \lrcorner w_j)(u_i \lrcorner w_h)) = \sum_i (-2\delta_{i,h} + \delta_{i,h}) \neq 0$, so $U \cap \text{Im}(u)^\perp = 0$.

U^\vee : Let $0 \neq \sum_i u_i \wedge w_{ij} \in U^\vee \subset \wedge^3 V$ and $w_h \in U^\vee$. Then $t(w_h \otimes (\sum_i u_i \wedge w_{ij})) = \sum_i (u_i \lrcorner w_{ijh} - w_i \wedge (u_i \lrcorner w_{jh}) + w_j \wedge (u_i \lrcorner w_{ih})) = \sum_i (2w_{jh} + \delta_{i,h} w_{ij}) = \sum_i w_{jh} \neq 0$, so $U^\vee \cap \text{Im}(u)^\perp = 0$.

$U^{\omega_2+\omega_5}$: Let $0 \neq u_i \wedge w_{jk} \in U^{\omega_2+\omega_5}$ for $i \neq j$ and $i \neq k$, and $w_h \in U^\vee$. Then $t(w_h \otimes u_i \wedge w_{jk}) = u_i \lrcorner (w_{jkh}) - w_j \wedge (u_i \lrcorner w_{kh}) + w_{jk} \lrcorner (u_i \lrcorner w_h) = 3\delta_{i,h} w_{jk} \neq 0$, so $U^{\omega_2+\omega_5} \cap \text{Im}(u)^\perp = 0$.

$\wedge^3 U^\vee$: Let $0 \neq w_{ijk} \in \wedge^3 U^\vee$ and $w_h \in U^\vee$. Then $t(w_h \otimes w_{ijk}) = w_{ijkh} \neq 0$, so $\wedge^3 U^\vee \cap \text{Im}(u)^\perp = 0$.

The previous computations imply that $\text{Im}(u)^\perp = \wedge^3 U \oplus U^{\omega_4+\omega_1} \subset \wedge^3 V \simeq \wedge^3 V^\vee$. This is equivalent to the fact that $\text{Im}(u)$ is the kernel of $\wedge^3 V \rightarrow \mathcal{C}|_{[U]}$. The statement of the proposition follows. \square

4.2.4. Complete orthogonality

Here we show the following result. We define Q' using a left mutation (see soon before §2.2), as follows:

$$Q' = L_{\langle \mathcal{B} \rangle}(Q).$$

Proposition 4.4. *The exceptional bundles Q and $Q'(1)$ are completely orthogonal.*

Proof. We know that $Q'(1)$ is an exceptional object that $\text{Ext}_X^\bullet(Q'(1), Q) = 0$, so we have to check that Q' is concentrated in degree 0 and that $\text{Ext}_X^\bullet(Q, Q'(1)) = 0$. First we check that:

$$\text{Ext}_X^\bullet(P, Q) = \text{Hom}_X(P, Q) = V^{\omega_1}. \tag{4.2}$$

To see this, recall (2.3) and use that $(\mathcal{U}_{\omega_1}, P)$ is exceptional to get, for all $p \geq 0$:

$$\text{Ext}_X^p(P, Q) \simeq \text{Ext}_X^p(P, \mathcal{U}_{\omega_1+\omega_2}).$$

Next, apply $\text{Hom}_X(-, \mathcal{U}_{\omega_1+\omega_2})$ to the sequence (2.2) defining P and work as in Lemma 2.3 to show that:

$$\begin{aligned} \text{Ext}_X^\bullet(\mathcal{O}_X, \mathcal{U}_{\omega_1+\omega_2}) &= H^0(\mathcal{U}_{\omega_1+\omega_2}) \simeq V^{\omega_1+\omega_2}, \\ \text{Ext}_X^{>0}(\mathcal{U}_{\omega_2}, \mathcal{U}_{\omega_1+\omega_2}) &= \text{Ext}_X^1(\mathcal{U}_{\omega_2}, \mathcal{U}_{\omega_1+\omega_2}) \simeq H^1(\mathcal{U}_{\omega_1+\omega_2+\omega_4}(-2)) \simeq V^{\omega_1+\omega_2}, \\ \text{Hom}_X(\mathcal{U}_{\omega_2}, \mathcal{U}_{\omega_1+\omega_2}) &= H^0(\mathcal{U}_{\omega_1}) \simeq V^{\omega_1}. \end{aligned}$$

Hence (4.2) holds if and only if the cup-product map below is non-degenerate:

$$\text{Ext}_X^1(\mathcal{U}_{\omega_2}, \mathcal{O}_X) \otimes H^0\mathcal{U}_{\omega_1+\omega_2} \rightarrow \text{Ext}_X^1(\mathcal{U}_{\omega_2}, \mathcal{U}_{\omega_1+\omega_2})$$

However, by the above analysis, using the notation of the proof of Lemma 2.3, this map is the cup-product

$$H^1(\mathcal{U}_{\omega_4-2\omega_6}) \otimes H^0(\mathcal{U}_{\omega_1+\omega_2}) \rightarrow H^1(\mathcal{U}_{\omega_1+\omega_2+\omega_4-2\omega_6})$$

and therefore it is non-degenerate by Lemma 2.1, once again using the reflection about the root α_6 . So (4.2) is proved. The resulting evaluation map $V^{\omega_1} \otimes P \rightarrow Q$ is surjective, as it results by tensoring (2.2) by V^{ω_1} and considering the evaluation map to (2.3); more precisely $L_P(Q)$ is an exceptional homogeneous bundle fitting into:

$$0 \rightarrow \mathcal{U}_{\omega_1}^\vee \rightarrow L_P(Q) \rightarrow N \rightarrow 0, \quad \text{with} \quad 0 \rightarrow \mathcal{U}_{\omega_1} \xrightarrow{\alpha} N \rightarrow K \rightarrow 0, \tag{4.3}$$

where K is the kernel of the map $\mathcal{C} \rightarrow \mathcal{U}_{\omega_1+\omega_2}$ of Proposition 4.3. Note that N fits into:

$$0 \rightarrow N \rightarrow V^{\omega_1} \otimes \mathcal{U}_{\omega_2} \rightarrow \mathcal{U}_{\omega_1+\omega_2} \rightarrow 0. \tag{4.4}$$

Next, we show that \mathcal{U}_{ω_1} is completely orthogonal to $L_P(Q)$, so we need to prove

$$\text{Ext}_X^\bullet(\mathcal{U}_{\omega_1}, L_P(Q)) = 0. \tag{4.5}$$

Using (4.3) and (4.4), one checks that (4.5) is proved once we show that α induces a non-zero map:

$$\text{Ext}_X^1(N, \mathcal{U}_{\omega_1}^\vee) \rightarrow \text{Ext}_X^1(\mathcal{U}_{\omega_1}, \mathcal{U}_{\omega_1}^\vee) \tag{4.6}$$

To achieve this, first note that, as a consequence of the definition of K in terms of \mathcal{C} and the definition of \mathcal{C} , K is an extension

$$0 \rightarrow \mathcal{U}_{\omega_2+\omega_5}(-1) \rightarrow K \rightarrow \mathcal{U}_{\omega_3} \rightarrow 0.$$

Next, we work as in the proof of Lemma 2.2 to check:

$$\text{Ext}_X^\bullet(\mathcal{U}_{\omega_2+\omega_5}(-1), \mathcal{U}_{\omega_1}^\vee) = \text{Ext}_X^1(\mathcal{U}_{\omega_2+\omega_5}(-1), \mathcal{U}_{\omega_1}^\vee) \simeq H^1(\mathcal{U}_{\omega_4}(-2)) \simeq H^1(\Omega_X) \simeq \mathbb{C}, \tag{4.7}$$

$$\text{Ext}_X^\bullet(\mathcal{U}_{\omega_3}, \mathcal{U}_{\omega_1}^\vee) = \text{Ext}_X^2(\mathcal{U}_{\omega_3}, \mathcal{U}_{\omega_1}^\vee) \simeq H^2(\mathcal{U}_{\omega_3+\omega_5}(-3)) \simeq H^2(\Omega_X^2) \simeq \mathbb{C}, \tag{4.8}$$

$$\text{Ext}_X^1(\mathcal{U}_{\omega_3}, \mathcal{U}_{\omega_2+\omega_5}(-1)) \simeq H^1(\mathcal{U}_{\omega_4}(-2)) \simeq H^1(\Omega_X) \simeq \mathbb{C}. \tag{4.9}$$

Then, $\text{Ext}_X^\bullet(K, \mathcal{U}_{\omega_1}^\vee)$ vanishes if and only if the following cup-product map is non-degenerate:

$$\text{Ext}_X^1(\mathcal{U}_{\omega_3}, \mathcal{U}_{\omega_2+\omega_5}(-1)) \otimes \text{Ext}_X^1(\mathcal{U}_{\omega_2+\omega_5}(-1), \mathcal{U}_{\omega_1}^\vee) \rightarrow \text{Ext}_X^2(\mathcal{U}_{\omega_3}, \mathcal{U}_{\omega_1}^\vee)$$

But from the identifications (4.7), (4.8) and (4.9), this map is the cup-product in cohomology:

$$H^1(\Omega_X) \otimes H^1(\Omega_X) \rightarrow H^2(\Omega_X^2),$$

and therefore it is non-degenerate. This proves $\text{Ext}_X^\bullet(K, \mathcal{U}_{\omega_1}^\vee) = 0$.

Now we can check that (4.6) is non-zero. Indeed, assume it was. Then α induces an exact sequence:

$$0 \rightarrow \mathcal{U}_{\omega_1} \oplus \mathcal{U}_{\omega_1}^\vee \rightarrow L_P(Q) \rightarrow K \rightarrow 0.$$

But then, since $\text{Ext}_X^1(K, \mathcal{U}_{\omega_1}^\vee) = 0$, $\mathcal{U}_{\omega_1}^\vee$ is a direct summand of $L_P(Q)$, which cannot happen since $L_P(Q)$ is exceptional.

We have now proved (4.5). Moreover, we get that α induces a diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{U}_{\omega_1}^\vee & \xlongequal{\quad} & \mathcal{U}_{\omega_1}^\vee & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & V^{\omega_1} \otimes \mathcal{O}_X & \longrightarrow & L_P(Q) & \longrightarrow & K \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 & \longrightarrow & \mathcal{U}_{\omega_1} & \xrightarrow{\quad \alpha \quad} & N & \longrightarrow & K \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

The leftmost column is the tautological sequence (5.5) because the cup-product above is non-degenerate and thus (4.5) is proved. Moreover we get, from the previous diagram:

$$Q' = L_{\langle \mathcal{B} \rangle}(Q) \simeq L_{\mathcal{O}_X}(L_P(Q)) \simeq L_{\mathcal{O}_X}(K).$$

Finally using the definition of K we check $H^\bullet(K) = H^0(K) = V^{\omega_3}$ hence, by Proposition 4.3, Q' is concentrated in degree 0 and we obtain:

$$Q'(1) \in \langle \mathcal{U}_{\omega_1}(-1), \mathcal{U}_{\omega_1+\omega_2}(-1), \mathcal{U}_{\omega_1} \rangle.$$

Therefore, using Lemma 2.5 and (2.3) we get $\text{Ext}_X^\bullet(Q, Q'(1)) = 0$. \square

Remark 4.5. One can actually prove that the bundle R_6 appearing in Proposition 4.3 satisfies:

$$R_6 \simeq Q(-2) \simeq L_{\mathcal{U}_{\omega_1}(-1)}(L_{\langle \mathcal{B} \rangle}(Q)).$$

Indeed, we checked that $K \simeq L_{\langle \mathcal{B} \rangle}(Q)$ and one can prove $\text{Ext}_X^\bullet(\mathcal{U}_{\omega_1}(-1), K) \simeq V_{\omega_5}$, so $R_6 \simeq L_{\mathcal{U}_{\omega_1}(-1)}(K)$ is exceptional, hence indecomposable, so by Proposition 4.3 it must be isomorphic to $Q(-2)$.

5. Generating more objects

We come back to $X = \text{Spin}_{12}/P_6$. The goal of this section is to show that the exceptional full triangulated subcategory \mathcal{D} of $\mathbf{D}^b(X)$ generated by the exceptional Lefschetz collection of Section 3 contains a bunch of vector bundles, which will be needed in the proof that $\mathcal{D}^\perp = 0$. From (2.13) we immediately have

$$\mathcal{O}(t) \in \mathcal{D} \quad \text{for } t \in [0, 9], \tag{5.1}$$

$$\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 9], \tag{5.2}$$

$$\wedge^2 \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 9], \tag{5.3}$$

$$\Sigma^{2,1} \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 1]. \tag{5.4}$$

Often we are going to use the tautological exact sequence

$$0 \rightarrow \mathcal{U} \rightarrow V \otimes \mathcal{O} \rightarrow \mathcal{U}^\vee \rightarrow 0. \tag{5.5}$$

Twisting (5.5) by $\mathcal{O}(t)$ with $t \in [0, 9]$ and using (5.1), (5.2) we immediately obtain

$$\mathcal{U}(t) \in \mathcal{D} \quad \text{for } t \in [0, 9]. \tag{5.6}$$

We also note that for $j \in [0, 6]$ we have isomorphisms

$$\wedge^j \mathcal{U}^\vee \simeq \wedge^{6-j} \mathcal{U}(2) \quad \text{and} \quad \wedge^j \mathcal{U} \simeq \wedge^{6-j} \mathcal{U}^\vee(-2).$$

Lemma 5.1. *Considering the spinor representations V^{ω_5} and V^{ω_6} as vector bundles on X , we have:*

i) *the vector bundle $V^{\omega_5} \otimes \mathcal{O}$ has an increasing filtration, whose factors are of the form*

$$\wedge^{2i+1} \mathcal{U}^\vee(-1) \quad \text{for } t \in [0, 2]$$

ii) *the vector bundle $V^{\omega_6} \otimes \mathcal{O}$ has an increasing filtration, whose factors are of the form*

$$\wedge^{2i} \mathcal{U}^\vee(-1) \quad \text{for } t \in [0, 3]$$

Proof. This follows from [21, Proposition 6.3] (and is the same filtration \mathcal{F}_\bullet described in Section 4.2). \square

As a corollary we obtain the following.

Corollary 5.2. *We have*

$$\wedge^j \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for} \quad \begin{cases} t \in [0, 9] & \text{if } j \in [0, 2], \\ t \in [0, 7] & \text{if } j \in [3, 4], \\ t \in [-2, 7] & \text{if } j = 5, \\ t \in [-2, 7] & \text{if } j = 6. \end{cases} \tag{5.7}$$

Proof. The cases with $j \in [0, 2]$ we have already considered. We treat each $j \in [3, 6]$ separately.

- (1) *Case* $j = 3$. Twisting $V^{\omega_5} \otimes \mathcal{O}$ by $\mathcal{O}(t)$ with $t \in [1, 8]$, using Lemma 5.1, the isomorphism $\mathcal{U}(1) \simeq \wedge^5 \mathcal{U}^\vee(-1)$, (5.1), (5.2), (5.6), we obtain the claim.
- (2) *Case* $j = 4$. Twisting $V^{\omega_6} \otimes \mathcal{O}$ by $\mathcal{O}(t)$ with $t \in [1, 8]$, using Lemma 5.1, (5.1), (5.3), we obtain the claim.
- (3) *Case* $j = 5$. Twisting the isomorphism $\mathcal{U}(2) \simeq \wedge^5 \mathcal{U}^\vee$ by $\mathcal{O}(t)$ with $t \in [-2, 7]$ and using (5.6), we obtain the claim.
- (4) *Case* $j = 6$. Since $\wedge^6 \mathcal{U}^\vee \simeq \det(\mathcal{U}^\vee) \simeq \mathcal{O}(2)$, the claim follows from (5.1). \square

Lemma 5.3. *We have*

$$S^j \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for} \quad \begin{cases} t \in [0, 9] & \text{if } j \in [0, 1], \\ t \in [2, 9] & \text{if } j \geq 2. \end{cases} \tag{5.8}$$

Proof. For $j \in [0, 1]$ the statement are known by (5.1) and (5.2).

Case $j = 2$. From (5.5) we obtain the exact sequence

$$0 \rightarrow \wedge^2 \mathcal{U} \rightarrow \wedge^2 V \otimes \mathcal{O} \rightarrow V \otimes \mathcal{U}^\vee \rightarrow S^2 \mathcal{U}^\vee \rightarrow 0.$$

Twisting this sequence by $\mathcal{O}(t)$ with $t \in [2, 9]$, using (5.1), (5.2), the isomorphism $\wedge^2 \mathcal{U} \simeq \wedge^4 \mathcal{U}^\vee(-2)$, (5.7), we see that all the terms of the sequence except for $S^2 \mathcal{U}^\vee(t)$ are contained in \mathcal{D} . Hence, the same holds for $S^2 \mathcal{U}^\vee(t)$.

Cases $j \geq 3$. We argue by induction. For each $j \geq 3$ we consider the exact sequence

$$0 \rightarrow \wedge^j \mathcal{U} \rightarrow \wedge^j V \otimes \mathcal{O} \rightarrow \wedge^{j-1} V \otimes \mathcal{U}^\vee \rightarrow \wedge^{j-2} V \otimes S^2 \mathcal{U}^\vee \rightarrow \dots \rightarrow V \otimes S^{j-1} \mathcal{U}^\vee \rightarrow S^j \mathcal{U}^\vee.$$

All the middle terms twisted by $\mathcal{O}(t)$ with $t \in [2, 9]$ are contained in \mathcal{D} by the induction assumption. For $j \in [3, 6]$ the term $\wedge^j \mathcal{U}(t) = \wedge^{6-j} \mathcal{U}^\vee(t-2)$ is also in \mathcal{D} for $t \in [2, 9]$ by (5.7). For $j \geq 7$ this term vanishes. Hence, the claim follows. \square

Lemma 5.4. *We have*

$$S^2 \mathcal{U}(t) \in \mathcal{D} \quad \text{for } t \in [0, 9], \tag{5.9}$$

$$\mathcal{U} \otimes \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 9], \tag{5.10}$$

$$\mathcal{U}^\vee \otimes \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 9]. \tag{5.11}$$

Proof. From (5.5) we get an exact sequence

$$0 \rightarrow S^2 \mathcal{U} \rightarrow S^2 V \otimes \mathcal{O} \rightarrow V \otimes \mathcal{U}^\vee \rightarrow \wedge^2 \mathcal{U}^\vee \rightarrow 0.$$

Twisting this sequence by $\mathcal{O}(t)$ with $t \in [0, 9]$ and using (5.1), (5.2), (5.7) we obtain (5.9).

One can reformulate (5.5) by saying that the bundle $V \otimes \mathcal{O}$ has a filtration with factors \mathcal{U} and \mathcal{U}^\vee . Then, taking the symmetric square, we obtain on $S^2 V \otimes \mathcal{O}$ a filtration with

factors $S^2\mathcal{U}, \mathcal{U} \otimes \mathcal{U}^\vee, S^2\mathcal{U}^\vee$. Twisting it by $\mathcal{O}(t)$ with $t \in [2, 9]$, using (5.1), (5.9), and (5.8), we get (5.10).

Finally, tensoring (5.5) with \mathcal{U}^\vee we get $0 \rightarrow \mathcal{U} \otimes \mathcal{U}^\vee \rightarrow V \otimes \mathcal{U}^\vee \rightarrow \mathcal{U}^\vee \otimes \mathcal{U}^\vee \rightarrow 0$. Together with (5.2) and (5.10) it implies (5.11). \square

Recall that from Proposition 4.3 and Remark 4.5 we have the exact sequence

$$0 \rightarrow Q(-2) \rightarrow \mathcal{U}^\vee(-1) \otimes S^+ \rightarrow \wedge^3 V \otimes \mathcal{O} \rightarrow \mathcal{C} \rightarrow \Sigma^{2,1}\mathcal{U}^\vee \rightarrow 0,$$

with \mathcal{C} defined by $0 \rightarrow \mathcal{U}^\vee \rightarrow V \otimes \wedge^2 \mathcal{U}^\vee \rightarrow \mathcal{C} \rightarrow 0$. Twisting this sequence by $\mathcal{O}(2)$ and using (5.1)-(5.4), we obtain $\Sigma^{2,1}\mathcal{U}^\vee(2) \in \mathcal{D}$. Iterating this process one shows

$$\Sigma^{2,1}\mathcal{U}^\vee \in \mathcal{D} \quad \text{for } t \in [0, 9]. \tag{5.12}$$

Lemma 5.5. *We have*

$$\wedge^2 \mathcal{U} \otimes \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 9], \tag{5.13}$$

$$\wedge^2 \mathcal{U} \otimes \mathcal{U}(t) \in \mathcal{D} \quad \text{for } t \in [2, 9], \tag{5.14}$$

$$\mathcal{U}^\vee \otimes \wedge^2 \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 7], \tag{5.15}$$

$$\mathcal{U} \otimes \wedge^2 \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 7]. \tag{5.16}$$

Proof. To show (5.13) we consider the exact sequence

$$0 \rightarrow \wedge^2 \mathcal{U} \rightarrow \wedge^2 V \otimes \mathcal{O} \rightarrow V \otimes \mathcal{U}^\vee \rightarrow S^2 \mathcal{U}^\vee \rightarrow 0$$

obtained from (5.5). Tensoring it by \mathcal{U}^\vee we obtain the exact sequence

$$0 \rightarrow \wedge^2 \mathcal{U} \otimes \mathcal{U}^\vee \rightarrow \wedge^2 V \otimes \mathcal{U}^\vee \rightarrow V \otimes \mathcal{U}^\vee \otimes \mathcal{U}^\vee \rightarrow S^2 \mathcal{U}^\vee \otimes \mathcal{U}^\vee \rightarrow 0.$$

Note that we have

$$\begin{aligned} \mathcal{U}^\vee \otimes \mathcal{U}^\vee &\simeq \wedge^2 \mathcal{U}^\vee \oplus S^2 \mathcal{U}^\vee, \\ S^2 \mathcal{U}^\vee \otimes \mathcal{U}^\vee &\simeq S^3 \mathcal{U}^\vee \oplus \Sigma^{2,1} \mathcal{U}^\vee. \end{aligned}$$

Twisting by $\mathcal{O}(t)$ with $t \in [2, 9]$ and using (5.7), (5.8), (5.12) we obtain the claim.

To show (5.14) we tensor the exact sequence (5.5) by $\wedge^2 \mathcal{U}$ and get the exact sequence

$$0 \rightarrow \mathcal{U} \otimes \wedge^2 \mathcal{U} \rightarrow V \otimes \wedge^2 \mathcal{U} \rightarrow \mathcal{U}^\vee \otimes \wedge^2 \mathcal{U} \rightarrow 0.$$

Twisting by $\mathcal{O}(t)$ with $t \in [2, 9]$, using the isomorphism $\wedge^2 \mathcal{U} \simeq \wedge^4 \mathcal{U}^\vee(-2)$, (5.7), (5.13) we get the claim.

To show (5.15) we note

$$\mathcal{U}^\vee \otimes \wedge^2 \mathcal{U}^\vee \simeq \wedge^3 \mathcal{U}^\vee \oplus \Sigma^{2,1} \mathcal{U}^\vee.$$

Twisting by $\mathcal{O}(t)$ with $t \in [0, 7]$, using (5.12) and (5.7) we obtain the claim.

To show (5.16) we tensor the exact sequence (5.5) by $\wedge^2 \mathcal{U}^\vee$ to get

$$0 \rightarrow \mathcal{U} \otimes \wedge^2 \mathcal{U}^\vee \rightarrow V \otimes \wedge^2 \mathcal{U}^\vee \rightarrow \mathcal{U}^\vee \otimes \wedge^2 \mathcal{U}^\vee \rightarrow 0.$$

Twisting by $\mathcal{O}(t)$ with $t \in [0, 7]$, using (5.15) and (5.7) we obtain the claim. \square

At this point we have proved the following.

Corollary 5.6. *We have:*

- i) $\mathcal{U}^\vee \otimes \wedge^0 \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [0, 9]$,
- ii) $\mathcal{U}^\vee \otimes \wedge^1 \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [2, 9]$,
- iii) $\mathcal{U}^\vee \otimes \wedge^2 \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [0, 7]$,
- iv) $\mathcal{U}^\vee \otimes \wedge^4 \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [0, 7]$,
- v) $\mathcal{U}^\vee \otimes \wedge^5 \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [0, 7]$,
- vi) $\mathcal{U}^\vee \otimes \wedge^6 \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [-2, 7]$.

Proof. We already proved these statements. Indeed, i) is (5.2), ii) is (5.11), iii) is (5.15), iv) follows from $\mathcal{U}^\vee \otimes \wedge^4 \mathcal{U}^\vee(t) \simeq \mathcal{U}^\vee \otimes \wedge^2 \mathcal{U}(t + 2)$ and (5.13), v) follows from $\mathcal{U}^\vee \otimes \wedge^5 \mathcal{U}^\vee(t) \simeq \mathcal{U}^\vee \otimes \mathcal{U}(t + 2)$ and (5.10) and vi) follows from $\wedge^6 \mathcal{U}^\vee \simeq \mathcal{O}(2)$. \square

Thus, we are still missing the objects $\mathcal{U}^\vee \otimes \wedge^3 \mathcal{U}^\vee(t)$, and the range of t for $\mathcal{U}^\vee \otimes \mathcal{U}^\vee(t)$ needs to be extended. This is our next goal.

Lemma 5.7. *We have*

$$\mathcal{U}^\vee \otimes \wedge^3 \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 7], \tag{5.17}$$

$$\mathcal{U} \otimes \wedge^3 \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 7], \tag{5.18}$$

$$\mathcal{U}^\vee \otimes \wedge^3 \mathcal{U}(t) \in \mathcal{D} \quad \text{for } t \in [4, 9]. \tag{5.19}$$

Proof. To show (5.17) we proceed as follows. By Lemma 5.1 we have

$$\text{ss}(V^{\omega_5} \otimes \mathcal{O}) = \mathcal{U}^\vee(-1) \oplus \wedge^3 \mathcal{U}^\vee(-1) \oplus \mathcal{U}(1);$$

Tensoring this by \mathcal{U}^\vee we get

$$\text{ss}(V^{\omega_5} \otimes \mathcal{U}^\vee) = \mathcal{U}^\vee \otimes \mathcal{U}^\vee(-1) \oplus \wedge^3 \mathcal{U}^\vee \otimes \mathcal{U}^\vee(-1) \oplus \mathcal{U} \otimes \mathcal{U}^\vee(1).$$

Twisting this by $\mathcal{O}(t)$ with $t \in [3, 8]$ and using (5.2), (5.10), (5.11) we obtain the claim.

To show (5.18) one can tensor the exact sequence (5.5) by $\wedge^3\mathcal{U}^\vee$ to get

$$0 \rightarrow \mathcal{U} \otimes \wedge^3\mathcal{U}^\vee \rightarrow V \otimes \wedge^3\mathcal{U}^\vee \rightarrow \mathcal{U}^\vee \otimes \wedge^3\mathcal{U}^\vee \rightarrow 0.$$

Now we twist by $\mathcal{O}(t)$ with $t \in [2, 7]$ and use (5.17) and (5.7).

To show (5.19) we use the inclusion (5.17) and the isomorphism $\wedge^3\mathcal{U}^\vee \simeq \wedge^3\mathcal{U}(2)$. \square

Lemma 5.8. *We have*

$$\wedge^2\mathcal{U} \otimes \wedge^2\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [4, 7], \tag{5.20}$$

$$\wedge^2\mathcal{U}^\vee \otimes \wedge^2\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 5]. \tag{5.21}$$

Proof. To show (5.20) we consider the decomposition

$$\text{ss}(\wedge^4V \otimes \mathcal{O}) = \wedge^2\mathcal{U}^\vee(-2) \oplus \wedge^3\mathcal{U} \otimes \mathcal{U}^\vee \oplus \wedge^2\mathcal{U} \otimes \wedge^2\mathcal{U}^\vee \oplus \mathcal{U} \otimes \wedge^3\mathcal{U}^\vee \oplus \wedge^4\mathcal{U}^\vee.$$

Twisting by $\mathcal{O}(t)$ with $t \in [4, 7]$ and using (5.1), (5.7), (5.18), (5.19) we get the claim.

To show (5.21) we consider decomposition Lemma 5.1

$$\text{ss}(V^{\omega_6} \otimes \mathcal{O}) = \mathcal{O}(-1) \oplus \wedge^2\mathcal{U}^\vee(-1) \oplus \wedge^2\mathcal{U}(1) \oplus \mathcal{O}(1),$$

where we have used that $\wedge^4\mathcal{U}^\vee(-1) \simeq \wedge^2\mathcal{U}(1)$ and $\wedge^6\mathcal{U}^\vee(-1) \simeq \mathcal{O}(1)$. Tensoring it by $\wedge^2\mathcal{U}^\vee$ we get

$$\text{ss}(V^{\omega_6} \otimes \wedge^2\mathcal{U}^\vee) = \wedge^2\mathcal{U}^\vee(-1) \oplus \wedge^2\mathcal{U}^\vee \otimes \wedge^2\mathcal{U}^\vee(-1) \oplus \wedge^2\mathcal{U} \otimes \wedge^2\mathcal{U}^\vee(1) \oplus \wedge^2\mathcal{U}^\vee(1).$$

Twisting it by $\mathcal{O}(t)$ with $t \in [3, 6]$ and using (5.7), (5.20) we obtain the claim. \square

Lemma 5.9. *We have*

$$\Sigma^{3,1}\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 9], \tag{5.22}$$

$$\Sigma^{2,1,1}\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 7], \tag{5.23}$$

$$\Sigma^{2,2}\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 7]. \tag{5.24}$$

Proof. Our first step is to note that by the Littlewood-Richardson rule we have

$$\begin{aligned} \mathcal{U}^\vee \otimes \wedge^3\mathcal{U}^\vee &\simeq \Sigma^{2,1,1}\mathcal{U}^\vee \oplus \wedge^4\mathcal{U}^\vee, \\ \wedge^2\mathcal{U}^\vee \otimes \wedge^2\mathcal{U}^\vee &\simeq \Sigma^{2,1,1}\mathcal{U}^\vee \oplus \Sigma^{2,2}\mathcal{U}^\vee \oplus \wedge^4\mathcal{U}^\vee. \end{aligned}$$

Hence, by (5.7) the inclusions (5.17) and (5.21) immediately imply

$$\Sigma^{2,1,1}\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 7], \tag{5.25}$$

$$\Sigma^{2,2}\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [2, 5]. \tag{5.26}$$

This proves (5.23), but it is not quite enough to prove (5.24).

Our second step is to deal with $\Sigma^{3,1}\mathcal{U}^\vee$. Let us consider the exact sequence

$$0 \rightarrow \wedge^3\mathcal{U} \rightarrow \wedge^3V \otimes \mathcal{O} \rightarrow \wedge^2V \otimes \mathcal{U}^\vee \rightarrow V \otimes S^2\mathcal{U}^\vee \rightarrow S^3\mathcal{U}^\vee \rightarrow 0$$

obtained from (5.5). After twisting by $\mathcal{O}(2)$ and using the isomorphism $\wedge^3\mathcal{U} \simeq \wedge^3\mathcal{U}^\vee(-2)$ we rewrite the above sequence as

$$0 \rightarrow \wedge^3\mathcal{U}^\vee \rightarrow \wedge^3V \otimes \mathcal{O}(2) \rightarrow \wedge^2V \otimes \mathcal{U}^\vee(2) \rightarrow V \otimes S^2\mathcal{U}^\vee(2) \rightarrow S^3\mathcal{U}^\vee(2) \rightarrow 0.$$

Tensoring it by \mathcal{U}^\vee we obtain the exact sequence

$$0 \rightarrow \wedge^3\mathcal{U}^\vee \otimes \mathcal{U}^\vee \rightarrow \wedge^3V \otimes \mathcal{U}^\vee(2) \rightarrow \wedge^2V \otimes \mathcal{U}^\vee \otimes \mathcal{U}^\vee(2) \rightarrow V \otimes S^2\mathcal{U}^\vee \otimes \mathcal{U}^\vee(2) \rightarrow S^3\mathcal{U}^\vee \otimes \mathcal{U}^\vee(2) \rightarrow 0.$$

Now we note that by the Littlewood-Richardson rule we have

$$S^2\mathcal{U}^\vee \otimes \mathcal{U}^\vee \simeq S^3\mathcal{U}^\vee \oplus \Sigma^{2,1}\mathcal{U}^\vee \quad \text{and} \quad S^3\mathcal{U}^\vee \otimes \mathcal{U}^\vee \simeq S^4\mathcal{U}^\vee \oplus \Sigma^{3,1}\mathcal{U}^\vee$$

Therefore, tensoring the above sequence by $\mathcal{O}(t)$ with $t \in [2, 7]$ and using (5.17), (5.2), (5.11), (5.8), (5.12), we conclude

$$\Sigma^{3,1}\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [4, 9]. \tag{5.27}$$

It is not quite enough for (5.22), but we are going to fix this soon.

For the third step we proceed as follows. Recall again the exact sequence from Proposition 4.3 and Remark 4.5

$$0 \rightarrow Q(-2) \rightarrow \mathcal{U}^\vee(-1) \otimes S^+ \rightarrow \wedge^3V \otimes \mathcal{O} \rightarrow \mathcal{C} \rightarrow \Sigma^{2,1}\mathcal{U}^\vee \rightarrow 0. \tag{5.28}$$

Using the Littlewood-Richardson rule and the definitions of P and Q we have

$$\text{ss}(P \otimes \mathcal{U}^\vee) = \mathcal{U}^\vee \oplus \wedge^3\mathcal{U}^\vee \oplus \Sigma^{2,1}\mathcal{U}^\vee$$

and

$$\text{ss}(Q \otimes \mathcal{U}^\vee) = \Sigma^{3,1}\mathcal{U}^\vee \oplus \Sigma^{2,2}\mathcal{U}^\vee \oplus \Sigma^{2,1,1}\mathcal{U}^\vee \oplus S^2\mathcal{U}^\vee \oplus \wedge^2\mathcal{U}^\vee.$$

Hence, for $P \otimes \mathcal{U}^\vee$ we have

$$P \otimes \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 7], \tag{5.29}$$

as each individual factor is contained in \mathcal{D} by (5.12) and (5.7).

In the same way we have

$$Q \otimes \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [4, 5], \tag{5.30}$$

as each individual factor is contained in \mathcal{D} by (5.7), (5.8), (5.25), (5.26), (5.27).

Tensoring (5.28) by $\mathcal{U}^\vee(t)$ with $t \in [6, 7]$ and using (5.29), (5.30), (5.2), (5.11), (5.15) we conclude that $Q \otimes \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [6, 7]$. Similarly, tensoring (5.28) by $\mathcal{U}^\vee(t)$ with $t \in [4, 5]$, we conclude $Q \otimes \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [2, 3]$. Finally, tensoring (5.28) by $\mathcal{U}^\vee(3)$, we obtain $Q \otimes \mathcal{U}^\vee(1) \in \mathcal{D}$. Thus, we have shown $Q \otimes \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [1, 7]$. This inclusion, together with (5.8), (5.7), (5.25), (5.26), (5.27) allows to conclude first that $\Sigma^{2,2}\mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [6, 7]$, as all the other factors are already contained in \mathcal{D} with these twists. Then, similarly, we conclude that $\Sigma^{3,1}\mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [2, 3]$. \square

Corollary 5.10. *We have*

$$\wedge^3 \mathcal{U}^\vee \otimes \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 7], \tag{5.31}$$

$$\Sigma^{2,1,1}\mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 7], \tag{5.32}$$

$$\mathcal{U}^\vee \otimes \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 9]. \tag{5.33}$$

Proof. Let us consider again the exact sequence

$$0 \rightarrow \wedge^3 \mathcal{U}^\vee \otimes \mathcal{U}^\vee \rightarrow \wedge^3 V \otimes \mathcal{U}^\vee(2) \rightarrow \wedge^2 V \otimes \mathcal{U}^\vee \otimes \mathcal{U}^\vee(2) \rightarrow V \otimes S^2 \mathcal{U}^\vee \otimes \mathcal{U}^\vee(2) \rightarrow S^3 \mathcal{U}^\vee \otimes \mathcal{U}^\vee(2) \rightarrow 0,$$

as in the proof of the previous lemma. From the previous lemma, (5.7), (5.8), (5.11), (5.12), we know that all its terms except for $\wedge^3 \mathcal{U}^\vee \otimes \mathcal{U}^\vee$ are contained in \mathcal{D} with twists in $[0, 7]$. Hence, the same holds for $\wedge^3 \mathcal{U}^\vee \otimes \mathcal{U}^\vee$.

The inclusion (5.32) follows from

$$\mathcal{U}^\vee \otimes \wedge^3 \mathcal{U}^\vee = \Sigma^{2,1,1}\mathcal{U}^\vee \oplus \wedge^4 \mathcal{U}^\vee,$$

combined with (5.31) and (5.7).

Finally, we show (5.33). Let us consider the decomposition

$$\text{ss}(V^{\omega_5} \otimes \mathcal{U}^\vee) \simeq \mathcal{U}^\vee \otimes \mathcal{U}^\vee(-1) \oplus \wedge^3 \mathcal{U}^\vee \otimes \mathcal{U}^\vee(-1) \oplus \mathcal{U} \otimes \mathcal{U}^\vee(1)$$

provided by Lemma 5.1. Twisting it by $\mathcal{O}(t)$ with $t \in [1, 2]$ and using (5.31), (5.10), (5.2) we obtain $\mathcal{U}^\vee \otimes \mathcal{U}^\vee(t) \in \mathcal{D}$ for $t \in [0, 1]$. Combining this with (5.11) we get the claim. \square

Lemma 5.11. *For any $j \in [0, 5]$ we have*

$$\mathcal{U}^\vee \otimes \wedge^j \mathcal{E}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 7]. \tag{5.34}$$

Proof. Let us consider the exact sequence

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{U}^\vee \rightarrow \mathcal{E}^\vee \rightarrow 0.$$

It implies that $\wedge^j \mathcal{U}^\vee$ has a filtration with factors $\mathcal{O}_X, \mathcal{E}^\vee, \wedge^2 \mathcal{E}^\vee, \dots, \wedge^j \mathcal{E}^\vee$. Therefore, arguing inductively with respect to j , if we know the inclusions

$$\mathcal{U}^\vee \otimes \wedge^j \mathcal{U}^\vee(t) \in \mathcal{D} \quad \text{for } t \in [0, 7], \tag{5.35}$$

for all $j \in [0, 5]$, then we know (5.34). Now we note that (5.35) holds by Corollary 5.6, (5.31) and (5.33). \square

6. Proof of fullness

We are now in position to prove fullness of our Lefschetz exceptional collection. Recall that this is defined in (2.13). We will use that X is covered by zero-loci of a general global sections of \mathcal{E} , where \mathcal{E}^\vee is the cokernel bundle of a general global section of $\mathcal{U}^\vee = \mathcal{U}_{\omega_1}$, which is nowhere vanishing. A section s of \mathcal{E} vanishes along a spinor 10-fold $Y_s \simeq OG(5, 10)_+$, as we recalled in §3.1.

Theorem 6.1. *The semiorthogonal exceptional collection appearing in (2.13) is full.*

Proof. Let us take an object $F \in \mathcal{D}^\perp$, i.e. we have

$$\text{Ext}_X^\bullet(A, F) = 0 \quad \text{for any } A \in \mathcal{D}.$$

Let $s \in H^0(X, \mathcal{E}^\vee)$ be a general section and $i_s: Y_s \rightarrow X$ the embedding of its zero locus, as in Lemma 3.2(1).

Let us consider the set of vector bundles on X defined by

$$\Upsilon := \{\mathcal{U}^\vee(t) \mid t \in [0, 7]\}.$$

By Lemma 5.11 for any $E \in \Upsilon$ and any j the bundle $E \otimes \wedge^j \mathcal{E}^\vee$ lies in \mathcal{D} . Hence, we have

$$\text{Ext}_X^\bullet(E \otimes \wedge^j \mathcal{E}^\vee, F) = H^\bullet(X, \wedge^j \mathcal{E} \otimes E^\vee \otimes F) = 0 \quad \text{for all } j,$$

and making use of the Koszul complex

$$0 \rightarrow \wedge^5 \mathcal{E} \rightarrow \dots \rightarrow \mathcal{E} \rightarrow \mathcal{O}_X \rightarrow i_{s*} \mathcal{O}_{Y_s} \rightarrow 0,$$

we obtain

$$H^\bullet(X, (E^\vee \otimes F) \otimes i_{s*} \mathcal{O}_Y) = 0.$$

Now, by the projection formula we rewrite

$$H^\bullet(X, (E^\vee \otimes F) \otimes i_{s*} \mathcal{O}_Y) = H^\bullet(Y_s, i_s^*(E^\vee \otimes F)) = \text{Ext}_{Y_s}^\bullet(i_s^* E, i_s^* F) = 0.$$

Recall that $Y_s \simeq OG(5, 10)$ has two connected components Y_{s+} and Y_{s-} . We denote the compositions $Y_{s\pm} \subset Y_s \xrightarrow{i_s} X$ by $i_{s\pm}$. Using this notation we have

$$\text{Ext}_{Y_s}^\bullet(i_s^* E, i_s^* F) = \text{Ext}_{Y_{s+}}^\bullet(i_{s+}^* E, i_{s+}^* F) \oplus \text{Ext}_{Y_{s-}}^\bullet(i_{s-}^* E, i_{s-}^* F).$$

Hence, we have

$$\text{Ext}_{Y_{s+}}^\bullet(i_{s+}^* E, i_{s+}^* F) = 0 \quad \text{and} \quad \text{Ext}_{Y_{s-}}^\bullet(i_{s-}^* E, i_{s-}^* F) = 0.$$

Applying Lemma 3.2(2) and Theorem 3.3 we obtain $i_{s+}^* F = 0$ and $i_{s-}^* F = 0$. Hence, we conclude $i_s^* F = 0$. Finally, since the above argument works for any general $s \in H^0(X, \mathcal{E}^\vee)$, by Lemma 3.2(3,4) we obtain $F = 0$. \square

7. A collection on the Freudenthal variety

Recall from the introduction that, in the third row of Freudenthal magic square, the homogeneous varieties $\mathbb{P}^2 \times \mathbb{P}^2$, $G(3, 6)$, $X = \text{Spin}_{12}/P_6$ and E_7/P_7 appear, so by the general philosophy of [22] these varieties should share a similar geometric behaviour. Our first observation here is that Lemma 2.2 can be transposed to E_7/P_7 .

Lemma 7.1. *On E_7/P_7 we have a canonical E_7 -equivariant extension*

$$0 \rightarrow \mathcal{O}_{E_7/P_7} \rightarrow O \rightarrow \mathcal{U}_{\omega_1} \rightarrow 0$$

which is an exceptional object, with $O^\vee(2)$ being the normal bundle of E_7/P_7 inside $\mathbb{P}(V^{\omega_7})$.

Proof. In the proof of Lemma 2.2 substitute: Spin_{12} with E_7 , P_6 with P_7 , ω_2 with ω_1 , ω_6 with ω_7 and P with O ; the modified proof still holds. \square

Lemma 7.2. *The collection $\langle \mathcal{O}_{E_7/P_7}, O, \mathcal{O}_{E_7/P_7}(1), O(1), \dots, \mathcal{O}_{E_7/P_7}(17), O(17) \rangle$ is exceptional.*

Proof. By an application of the BBW Theorem, we get, for $1 \leq i \leq 17$:

$$\text{Ext}_X^\bullet(\mathcal{O}_{E_7/P_7}(i), \mathcal{O}_{E_7/P_7}) = \text{Ext}_X^\bullet(O(i), \mathcal{O}_{E_7/P_7}) = \text{Ext}_X^\bullet(O(i), O) = 0.$$

Since O is a non-trivial extension of \mathcal{O}_{E_7/P_7} and \mathcal{U}_{ω_1} , we also get $\text{Ext}_X^\bullet(O, \mathcal{O}_{E_7/P_7}) = 0$. \square

One can also define a G -equivariant extension

$$0 \rightarrow O \rightarrow P' \rightarrow \mathcal{U}_{2\omega_1} \rightarrow 0$$

Let us define a *numerical exceptional collection* in the derived category $\mathbf{D}^b(X)$ of any smooth projective variety X as a collection of objects E_1, \dots, E_r such that $\chi(E_i, E_j) = 0$ if $i > j$ and $\chi(E_i, E_i) = 1$ for all i . Let us denote by

$$\mathcal{B}' := (\mathcal{O}_{E_7/P_7}, O, P').$$

Moreover we will denote by Q' the projection of $\mathcal{U}_{\omega_1+\omega_3}(-5)$ to the left orthogonal of

$$\langle \mathcal{B}', \dots, \mathcal{B}'(17) \rangle.$$

Remark 7.3. Here and later on by “projection” we mean that Q' is obtained as an extension of $\mathcal{U}_{\omega_1+\omega_3}(-5)$ with elements in the collection $(\mathcal{B}', \dots, \mathcal{B}'(17))$ so that, for any element $E \in (\mathcal{B}', \dots, \mathcal{B}'(17))$, $\chi(E, Q') = 0$; if we knew that $(\mathcal{B}', \dots, \mathcal{B}'(17))$ were an exceptional collection, then it would be admissible and the “projection” to its left orthogonal would be well defined. Notice however that Q' is uniquely defined in the Grothendieck group.

We consider the collection:

$$\mathcal{A}' := (Q', \mathcal{O}_{E_7/P_7}, O, P').$$

Proposition 7.4. *The collection $(\mathcal{A}', \mathcal{A}'(1), \mathcal{B}'(2), \dots, \mathcal{B}'(17))$ is a numerically exceptional collection of maximal length, i.e. of length equal to $\sum_p h^{p,p}(E_7/P_7) = 56$.*

Proof. The projection Q' can be computed numerically, i.e. in the Grothendieck group of E_7/P_7 . Let us explain the strategy. Let us denote by $R_0 := \mathcal{U}_{\omega_1+\omega_3}(-5)$ and let us define $R_1, R_2, \dots, R_{54} = Q'$ inductively. Write the collection $(\mathcal{B}', \dots, \mathcal{B}'(17))$ as (E_1, \dots, E_{54}) . The object R_{i+1} will be an extension of R_i by $\chi(E_{i+1}, R_i)E_{i+1}(-18)$ in the Grothendieck group. When this process finishes, by Serre duality one obtains an object Q' which is by definition left orthogonal to $\langle E_1, \dots, E_{54} \rangle$, and one checks that $\chi(Q', Q') = -\chi(Q'(2), Q') = 1$ and $\chi(Q'(1), Q') = 0$. Another computation with BBW Theorem yields the numerical exceptionality of the collection. \square

Some observations are in order. Let us write the element in the Grothendieck group corresponding to Q' :

$$\begin{aligned} & \mathcal{U}_{\omega_1+\omega_3}(-5) - P'(-7) + O(-6) + 56P'(-6) - 1673\mathcal{O}_{E_7/P_7}(-5) - 3137O(-5) + P'(-5) + \\ & - 94656\mathcal{O}_{E_7/P_7}(-4) - 56P'(-4) - 54342\mathcal{O}_{E_7/P_7}(-3) + 3271O(-3) - P'(-3) - 58576\mathcal{O}_{E_7/P_7}(-2) + \\ & - 968O(-2) + 56P'(-2) + 54342\mathcal{O}_{E_7/P_7}(-1) - 3137O(-1). \end{aligned}$$

Notice that by general properties of mutations we also obtain another numerically exceptional collection:

$$(Q', L_{\mathcal{B}}Q'(1), \mathcal{B}', \mathcal{B}'(1), \dots, \mathcal{B}'(17)).$$

The peculiar fact about this collection is that its residual collection $(Q', L_{\mathcal{B}'}Q'(1))$ is numerically completely orthogonal, meaning that $\chi(Q', L_{\mathcal{B}'}Q'(1)) = \chi(L_{\mathcal{B}'}Q'(1), Q') = 0$. Therefore, such a residual collection numerically satisfies Dubrovin’s refined conjecture, see [17, Conjecture 1.3] and [7, Corollary 1.2]. We believe that the collection above is an exceptional collection in $\mathbf{D}^b(E_7/P_7)$ (of maximal length), but we could not prove our claim due to the big number of cohomologies between $\mathcal{H}om$ ’s of the irreducible factors of the extensions in play. We even suspect that the collection is full.

If the above collection has the advantage of respecting Dubrovin’s conjecture’s expectation, we will briefly describe another numerically exceptional collection on E_7/P_7 which is closer to the collection of (2.13) on Spin_{12}/P_6 . Let us begin with the usual collection $(\mathcal{O}_{E_7/P_7}, O, \dots, \mathcal{O}_{E_7/P_7}(17), O(17))$. Consider the projection P of \mathcal{U}_{ω_3} to the left orthogonal of $\langle \mathcal{O}_{E_7/P_7}(1), O(1), \dots, \mathcal{O}_{E_7/P_7}(18), O(18) \rangle$. Moreover consider the projection (as in Remark 7.3) Q of $\mathcal{U}_{\omega_1+\omega_3}$ to the left orthogonal of $\langle \mathcal{O}_{E_7/P_7}(1), O(1), P(1), \dots, \mathcal{O}_{E_7/P_7}(18), O(18), P(18) \rangle$. Let us write

$$\mathcal{B} := \langle \mathcal{O}_{E_7/P_7}, O, P \rangle, \quad \mathcal{A} := \langle \mathcal{O}_{E_7/P_7}, O, P, Q \rangle.$$

By a repeated application of the BBW Theorem done with a Python script using [31] as in the proof of Proposition 7.4 one obtains the following result.

Proposition 7.5. *The homogeneous bundles P and Q are numerically exceptional and the collection*

$$\mathcal{D}' := (\mathcal{A}, \mathcal{A}(1), \mathcal{B}(2), \dots, \mathcal{B}(17))$$

is numerically exceptional of maximal length. Moreover Q and $L_{\mathcal{B}(1)}Q(1)$ are numerically completely orthogonal.

The computation is enclosed as an ancillary file in the arXiv version of this paper.

Remark 7.6. Notice that in the Python script, in order to obtain the result, it was easier to work with the projection (as in Remark 7.3) of $\mathcal{U}_{\omega_1+\omega_3}(-9)$ to the left orthogonal of $\langle \mathcal{B}, \mathcal{B}(1), \dots, \mathcal{B}(17) \rangle$. Then Q is easily obtained as the projection of $F(9)$ to the left of $\langle \mathcal{B}(1), \dots, \mathcal{B}(8) \rangle$. Similarly, if F' is the projection of $\mathcal{U}_{\omega_1+\omega_3}(-8)$ to the left of $\langle \mathcal{B}, \mathcal{B}(1), \dots, \mathcal{B}(17) \rangle$, then $L_{\mathcal{B}(1)}Q(1)$ is obtained as the projection of $F'(9)$ to the left of $\langle \mathcal{B}(1), \dots, \mathcal{B}(8) \rangle$. Since these operations preserve (numerical) orthogonality, we prove Proposition 7.5 using F and F' .

Data availability

No data was used for the research described in the article.

References

- [1] Alexander A. Beilinson, Coherent sheaves on \mathbf{P}^n and problems in linear algebra, *Funkc. Anal. Prilozh.* 12 (3) (1978) 68–69.
- [2] Pieter Belmans, Grassmannian.info — a periodic table of (generalised) grassmannians, <https://grassmannian.info>, 2025.
- [3] Pieter Belmans, Alexander Kuznetsov, Maxim Smirnov, Derived categories of the Cayley plane and the coadjoint Grassmannian of type F, *Transform. Groups* 28 (1) (2023) 9–34.
- [4] Anders Skovsted Buch, Andrew Kresch, Harry Tamvakis, Quantum Pieri rules for isotropic Grassmannians, *Invent. Math.* 178 (2) (2009) 345–405, MR 2545685.
- [5] Christian Böhning, Derived categories of coherent sheaves on rational homogeneous manifolds, *Doc. Math.* 11 (2006) 261–331.
- [6] Pierre-Emmanuel Chaptut, Laurent Manivel, Nicola Perrin, Quantum cohomology of minuscule homogeneous spaces, *Transform. Groups* 13 (1) (2008) 47–89.
- [7] Pierre-Emmanuel Chaptut, Laurent Manivel, Nicolas Perrin, Quantum cohomology of minuscule homogeneous spaces III. Semi-simplicity and consequences, *Can. J. Math.* 62 (6) (2010) 1246–1263.
- [8] Dragos Deliu, Homological projective duality for $Gr(3, 6)$, Ph.D. thesis, University of Pennsylvania, 2011.
- [9] Ivan Dimitrov, Mike Roth, Cup products of line bundles on homogeneous varieties and generalized PRV components of multiplicity one, *Algebra Number Theory* 11 (4) (2017) 767–815.
- [10] Daniele Faenzi, Laurent Manivel, On the derived category of the Cayley plane II, *Proc. Am. Math. Soc.* 143 (3) (2015) 1057–1074.
- [11] Anton V. Fonarev, Minimal Lefschetz decompositions of the derived categories for Grassmannians, *Izv. Ross. Akad. Nauk, Ser. Mat.* 77 (5) (2013) 203–224.
- [12] Anton V. Fonarev, Full exceptional collections on Lagrangian Grassmannians, *Int. Math. Res. Not.* (2) (2022) 1081–1122.
- [13] Lyalya Guseva, On the derived category of $IGr(3; 8)$, *Mat. Sb.* 211 (7) (2020) 24–59.
- [14] Mikhail M. Kapranov, On the derived categories of coherent sheaves on some homogeneous spaces, *Invent. Math.* 92 (3) (1988) 479–508.
- [15] Alexander Kuznetsov, Alexander Polishchuk, Exceptional collections on isotropic Grassmannians, *J. Eur. Math. Soc.* 18 (3) (2016) 507–574.
- [16] Alexander Kuznetsov, Maxim Smirnov, On residual categories for Grassmannians, *Proc. Lond. Math. Soc.* (3) 120 (5) (2020) 617–641.
- [17] Alexander Kuznetsov, Maxim Smirnov, Residual categories for (co)adjoint Grassmannians in classical types, *Compos. Math.* 157 (6) (2021) 1172–1206.
- [18] Alexander Kuznetsov, Hyperplane sections and derived categories, *Izv. Ross. Akad. Nauk, Ser. Mat.* 70 (3) (2006) 23–128.
- [19] Alexander G. Kuznetsov, Hyperplane sections and derived categories, *Izv. Ross. Akad. Nauk, Ser. Mat.* 70 (3) (2006) 23–128. MR 2238172.
- [20] Alexander Kuznetsov, Homological projective duality, *Publ. Math. Inst. Hautes Études Sci.* (105) (2007) 157–220. MR 2354207.
- [21] Alexander Kuznetsov, Exceptional collections for Grassmannians of isotropic lines, *Proc. Lond. Math. Soc.* (3) 97 (1) (2008) 155–182.
- [22] Joseph M. Landsberg, Laurent Manivel, The projective geometry of Freudenthal’s magic square, *J. Algebra* 239 (2) (2001) 477–512.
- [23] Laurent Manivel, On the derived category of the Cayley plane, *J. Algebra* 330 (2011) 177–187.
- [24] Eckhard Meinrenken, Clifford Algebras and Lie Theory, first ed., *Modern Surveys in Mathematics*, Springer, Berlin, Heidelberg, 2013.
- [25] Alexander Polishchuk, Alexander Samokhin, Full exceptional collections on the Lagrangian Grassmannians $LG(4, 8)$ and $LG(5, 10)$, *J. Geom. Phys.* 61 (10) (2011) 1996–2014.
- [26] Roger Richardson, Gerhard Röhrle, Robert Steinberg, Parabolic subgroups with abelian unipotent radical, *Invent. Math.* 110 (3) (1992) 649–671. MR 1189494.

- [27] Alexander Samokhin, Some remarks on the derived categories of coherent sheaves on homogeneous spaces, *J. Lond. Math. Soc.* (2) 76 (1) (2007) 122–134.
- [28] Maxim Smirnov, Residual categories of Grassmannians, arXiv:2212.01580 [math.AG], 2022.
- [29] Maxim Smirnov, On the derived category of the adjoint Grassmannian of type F, arXiv:2107.07814 [math.AG], 2023.
- [30] Alexander Samokhin, Wilberd van der Kallen, Highest weight category structures on $rep(b)$ and full exceptional collections on generalized flag varieties over \mathbb{z} , arXiv:2407.13653 [math.AG], 2024.
- [31] Marc A.A. van Leeuwen, Arej M. Cohen, Bert Lisser, LiE, a package for lie group computations, *Computer Algebra*, Nederland, Amsterdam, 1992.