

## The effect of traditional cupping on pain and mechanical thresholds in patients with chronic nonspecific neck pain: a randomised controlled pilot study

Romy Lauche, Holger Cramer, Claudia Hohmann, Kyung-Eun Choi, Thomas Rampp, Felix Joyonto Saha, Frauke Musial, Jost Langhorst, Gustav Dobos

### Angaben zur Veröffentlichung / Publication details:

Lauche, Romy, Holger Cramer, Claudia Hohmann, Kyung-Eun Choi, Thomas Rampp, Felix Joyonto Saha, Frauke Musial, Jost Langhorst, and Gustav Dobos. 2012. "The effect of traditional cupping on pain and mechanical thresholds in patients with chronic nonspecific neck pain: a randomised controlled pilot study." *Evidence-Based Complementary and Alternative Medicine* 2012: 429718. <https://doi.org/10.1155/2012/429718>.

## Research Article

# The Effect of Traditional Cupping on Pain and Mechanical Thresholds in Patients with Chronic Nonspecific Neck Pain: A Randomised Controlled Pilot Study

Romy Lauche,<sup>1</sup> Holger Cramer,<sup>1</sup> Claudia Hohmann,<sup>1</sup> Kyung-Eun Choi,<sup>1</sup> Thomas Ramm,<sup>1</sup> Felix Joyonto Saha,<sup>1</sup> Frauke Musial,<sup>2</sup> Jost Langhorst,<sup>1</sup> and Gustav Dobos<sup>1</sup>

<sup>1</sup> Chair of Complementary and Integrative Medicine, University of Duisburg-Essen, Knappschafts-Krankenhaus, Am Deimelsberg 34a, 45276 Essen, Germany

<sup>2</sup> Department of Community Medicine, The National Research Centre in Complementary and Alternative Medicine (NAFKAM), Faculty of Health Science, University of Tromsø, 9037 Tromsø, Norway

Correspondence should be addressed to Romy Lauche, r.lauche@kliniken-essen-mitte.de

Received 18 August 2011; Revised 7 September 2011; Accepted 13 September 2011

Academic Editor: Gerhard Litscher

Copyright © 2012 Romy Lauche et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Introduction.** Cupping has been used since antiquity in the treatment of pain conditions. In this pilot study, we investigated the effect of traditional cupping therapy on chronic nonspecific neck pain (CNP) and mechanical sensory thresholds. **Methods.** Fifty CNP patients were randomly assigned to treatment (TG,  $n = 25$ ) or waiting list control group (WL,  $n = 25$ ). TG received a single cupping treatment. Pain at rest (PR), pain related to movement (PM), quality of life (SF-36), Neck Disability Index (NDI), mechanical detection (MDT), vibration detection (MDT), and pressure pain thresholds (PPT) were measured before and three days after a single cupping treatment. Patients also kept a pain and medication diary (PaDi, MeDi) during the study. **Results.** Baseline characteristics were similar in the two groups. After cupping TG reported significantly less pain (PR:  $-17.9$  mm VAS, 95%CI  $-29.2$  to  $-6.6$ ; PM:  $-19.7$ , 95%CI  $-32.2$  to  $-7.2$ ; PaDi:  $-1.5$  points on NRS, 95%CI  $-2.5$  to  $-0.4$ ; all  $P < 0.05$ ) and higher quality of life than WL (SF-36, Physical Functioning: 7.5, 95%CI 1.4 to 13.5; Bodily Pain: 14.9, 95%CI 4.4 to 25.4; Physical Component Score: 5.0, 95%CI 1.4 to 8.5; all  $P < 0.05$ ). No significant effect was found for NDI, MDT, or VDT, but TG showed significantly higher PPT at pain-areas than WL (in lg(kPa); pain-maximum: 0.088, 95%CI 0.029 to 0.148, pain-adjacent: 0.118, 95%CI 0.038 to 0.199; both  $P < 0.01$ ). **Conclusion.** A single application of traditional cupping might be an effective treatment for improving pain, quality of life, and hyperalgesia in CNP.

## 1. Introduction

Neck pain, that is, pain between the occipital bone, the thoracic vertebra, and the extensions to the shoulder joint [1], is a major health-related socioeconomic problem and the lifetime prevalence is approximately 48.5% [2]. Neck pain can be caused by trauma, inflammatory diseases, or degeneration of the spine; however, most patients suffer from simple or non-specific neck pain, which is mainly caused by mechanical factors such as sprain and strains [3]. The aetiology of non-specific neck pain is not yet understood in detail, but different factors have been shown to contribute to the development and persistence of neck pain. They do not only include poor posture [3] and high physical load

[4], but also poor psychological health [1, 5], stress [6], low socioeconomic status [7], and smoking [8, 9]. Usually non-specific neck pain resolves within three to six months; but 14% of the patients will suffer from recurrent or persistent pain [10]. If neck pain persists for more than 3 months, it is considered chronic neck pain [11].

Besides the pain and the related impairment in daily activities, chronic neck pain is also associated with functional changes. For example dysfunctional microcirculation of the trapezius muscle [12, 13] has been reported as well as motor control disturbances of the neck musculature [14]. Mechanical hyperalgesia, that is, increased response to painful mechanical stimulation has also been shown in chronic non-specific neck pain [15–17]; an effect which might be related

to active trigger points, present in chronic non-specific neck pain patients but not in healthy controls [18]. However, it is still unknown if this process is restricted to the cervical area [16] or widespread [19]. Hyperalgesia in chronic non-specific neck pain also shows different patterns and seems to rely on different mechanisms than hyperalgesia in acute [19] and traumatic neck pain [15], respectively.

Although there is only limited evidence for these treatments, conventional treatment options include the prescription of nonsteroidal anti-inflammatory drugs [20], physical therapy [21, 22] or exercise [23, 24]. According to the literature [20, 25] and treatment guidelines [25] pharmacological therapy cannot be recommended, the same is true for manual therapy [26], or massages [21]. Dynamic and isometric exercises as part of physical therapy have also been proven to be only moderately effective in the long term [25]. Due to the limited treatment patients seek alternative treatment options, especially those patients with more intense pain [27, 28] and those who have not experienced improvements under conventional treatment [29].

Traditional cupping or wet cupping has been used in the treatment of pain and many other complaints for millennia [30]. A glass cup is utilized to create suction over a painful area after incisions are made to the skin. By doing so, it is hypothesised that “congested” blood is sucked out of the skin thereby increasing blood and lymphatic circulation and relieving painful muscle tension [30, 31].

Within the last years the interest in traditional cupping has emerged and there is growing evidence that cupping might be effective in various pain conditions [32–37]. Michalsen et al. [37] for example, found that a single traditional cupping treatment at the trapezius muscle was effective in relieving the symptoms of the carpal tunnel syndrome as well as associated neck pain. Lüdtkke et al. [36] investigated the effect of traditional cupping in Brachialgia parasthetica nocturna, that is, numb, tingling, and painful sensations in fingers or hands during the night. A single treatment significantly reduced symptoms and the associated neck pain and no adverse events were observed. Farhadi et al. [34] found significantly reduced pain, functional disability, and pain medication in patients with low-back pain three months after traditional cupping compared to standard care. Cupping might further be effective in migraine and tension-type head-ache [32] and postherpetic pain [38]. However, despite growing evidence there is yet no RCT to investigate the effectiveness of traditional cupping in the treatment of chronic non-specific neck pain.

The aim of this pilot study was to test the efficacy of a single traditional cupping treatment in patients with chronic non-specific neck pain. Besides pain ratings we determined mechanical thresholds at pain-related and control areas to serve as more objective pain markers. We hypothesised that patients in the treatment group would report less pain at T2 compared to the waiting list control group.

## 2. Methods

**2.1. Patients.** The study protocol was approved by the institutional review board of the University Duisburg-Essen

Medical Institutions (no. 09–3985). Between July 2009 and July 2010 50 patients aged 18 to 75 who suffered from neck pain for at least three months in a row with a minimum of 40 mm intensity on a 100 mm visual analogue scale (VAS) were included in the study. A specific inclusion criterion was based on the recommendations for traditional cupping [30, 39, 40]. Accordingly, patients show so-called plethora or overabundance. These terms refer to different signs and symptoms such as voluminous gelosis of the subskin, which indicates local blood congestion, swelling, and adhesions of the connective tissue in the neck region. A strong constitution, for example, high level of vitality, and high blood pressure, were further indicators for traditional cupping. Patients with blank myogelosis, that is, hyperirritable areas of skeletal muscle associated with small palpable nodules in taut bands of muscle fibres together with lowered microcirculation, were referred for dry cupping.

Patients were included only if neck pain was clearly identified to be of mechanical origin and specific causes for their neck pain had been excluded in the medical history either by an orthopaedist or a neurologist. Specific causes included traumatic neck pain (e.g., WAD), inflammatory or malignant disease, congenital malformation of the spine, radicular symptoms such as radiating pain, paresis, prickling, or tingling, invasive treatments within the last 4 weeks, surgery to the spine within the last year, and corticosteroid or opioid treatment. Further exclusion criteria were pregnancy, serious acute or chronic organic diseases such as diabetes or cancer, mental disorders, and haemorrhagic tendency or anticoagulation treatment. Nonsteroidal pain medication and physiotherapy were allowed if the treatment regimen was not altered for four weeks before and continued during the study. This ensured that statistical evaluation of the effects of cupping treatment was not influenced by alterations in medications or physiotherapy during the study phase.

All patients were recruited through advertisements in local newspapers and screened two times. First inclusion and exclusion criteria were checked in a standardised telephone interview, then the patients underwent a physical and neurological examination by the study physician at their first appointment.

### 2.2. Outcome Measures

**2.2.1. Pain.** Pain at rest (PR) and maximal pain related to movement (PM, provoked pain by neck flexion, neck extension, lateral neck flexion, and neck rotation in either direction) [41] were recorded on a VAS graded from 0 (no pain at all) to 100 mm (worst pain imaginable). The minimal clinical important difference (MCID) for the VAS, a highly reliable instrument to measure pain intensity [42], is a reduction of 30%, which is the equivalent of a moderate pain reduction [43]. For PM the direction that elicited highest pain report was chosen for analysis. Baseline and postintervention pain scores were recorded at T1 and T2. Additionally patients kept a pain (PaDi) and medication diary (MeDi) from day 0 (7 days prior to T1) until T2, where they rated their pain three times daily on a 11-point numeric rating scale (NRS ranging from 0 = no pain to 10 = worst

pain imaginable) and made notes of concurrent medication and treatments.

**2.2.2. Questionnaires.** Self-rated disability due to neck pain was assessed with the Neck Disability Index (NDI) [44], a 10-item questionnaire representing everyday activities. The MCID for the NDI is 10% improvement for uncomplicated neck pain [45]. The health-related quality of life was quantified by the German version of the Medical Outcomes Study Short Form-36 (SF-36) [46, 47]. The SF-36 provides a detailed health profile on the basis of eight health dimensions as well as sum scores for physical and mental health. Two versions were used in the study, the standard version (4-week time frame) for baseline assessment at T1 and the acute recall version (1-week time frame) at T2. The latter version was used because it was considered more sensitive to recent changes in health status [48].

**2.2.3. General Health Outcome.** Within the SF-36 the General Health outcome was recorded on a 5-point Likert scale that ranged from “My health is much better than before treatment” to “My health is much worse than before treatment.”

**2.2.4. Mechanical Sensory and Pain Thresholds.** Sensory testing included determination of mechanical detection threshold (MDT), vibration detection threshold (VDT), and pressure pain threshold (PPT) and was conducted in the following areas: the site of maximal pain (pain-maximum), adjacent to the pain maximum (pain-adjacent), hand and foot on the right side. Pain-maximum and pain-adjacent were determined for each patient individually. First, the patient was given a diagram of the body on which to mark the most painful spot in the neck region. This spot, defined as pain-maximum, was verified by physical examination. The second spot, defined as adjacent to the painful area (pain-adjacent) was defined outside the painful area, that is, patients did not report pain in that area. Again physical examination was used to confirm the location. Both spots were marked in the pain diagram for precise replication of the measurements at T2. Thresholds were also determined at control areas, that is, right hand and right foot, in order to estimate reliability of measurements. All sensory measures were determined and calculated according to the standardised protocol for the quantitative sensory testing (QST) by Rolke et al. [49, 50], and MDT and PPT were logarithmised to reach normal distribution [49].

Mechanical detection threshold was measured with a set of von Frey filaments (Aesthesiometer, SOMEDIC, Sweden) that exert forces between 0.26 and 1080 mN. The threshold was determined by the method of limits, whereby the stimulus intensity is decreased until the patient can no longer perceive the touch and is then increased until the patient first perceives the touch again. Five series of descending and ascending stimulus intensities were made at pain-maximum, pain-adjacent, on the dorsa of the right hand and the right foot. The final threshold was the log-transformed geometric mean of these five series [49].

VDT was quantified by a Rydel Seiffer tuning fork (64 Hz, 8/8 scale). It was placed over a bony prominence, for example, on the spinal process, the styloid process of ulna and the lateral malleolus and left there until the subject could not feel the vibration anymore. The arithmetic mean of three series was taken the individual vibration detection threshold [49].

PPT was measured by a pressure algometer (Algometer, SOMEDIC, Sweden) at pain-maximum, pain-adjacent, the thenar eminence, and the instep. It exerts forces up to 2000 kPa when used with a probe area of 1 cm<sup>2</sup>. The pressure pain threshold was measured in three ramps of increasing pressure intensities of ca. 50 kPa/s until the subject signalled the first sensation of pain in addition to the pressure sensation. The log-transformed arithmetic mean of these three series was taken the individual pressure pain threshold [49].

To evaluate the reliability of the sensory threshold measurements, the retest reliabilities were determined at the control areas in the control group participants (WL,  $N = 23$ ). Pearson's correlation coefficients were  $r = 0.35$  for MDT hand ( $P = 0.09$ ),  $r = 0.66$  for MDT foot ( $P = 0.001$ ),  $r = 0.79$  for PPT hand ( $P = 0.00001$ ),  $r = 0.71$  for PPT foot ( $P = 0.0001$ ),  $r = 0.56$  for VDT hand ( $P = 0.005$ ), and  $r = 0.73$  for VDT foot ( $P = 0.0001$ ). The average correlation coefficients was  $r = 0.63$  which indicates sufficient reliability.

**2.2.5. Safety.** All participants were asked to report any adverse events during the study period. The questionnaires relating to T2 also included an open question about relevant experiences and adverse events.

**2.2.6. Expectation.** After randomisation all patients had to self-rate their expectations towards cupping therapy on a VAS ranging from 0 = “not effective at all” to 100 mm = “most effective.”

**2.3. Intervention: Traditional Cupping Technique.** Based on data from previous studies on traditional cupping [36, 37] and clinical experience, a single cupping treatment was considered sufficient. Cupping was performed by the study physician, who was trained in cupping and regularly performed cupping in a clinical setting. Patients were asked to lay topless on the massage couch. The study physician used the patients pain diagram (see Section 2.2.4) and physical examination to identify the areas of pain and the voluminous geloses of the subskin, which most commonly were found at the descending parts of the trapezius muscle.

The cupping procedure involved the following steps: the skin was disinfected; superficial incisions were made with a disposable microlancet at the areas of pain and voluminous geloses; double-walled glass cups (2–6 glasses with diameters from 25 to 50 mm) were held inverted over an open flame to heat the air inside; the glass cup was placed on the incision. The air inside the cup cooled down and created a vacuum which sucked blood out through the incisions. The glasses were removed after 10 to 15 minutes, and the skin was




FIGURE 1: Study design.

disinfected and a plaster was applied. However, since bleeding generally stopped during treatment, this was only a precaution. Patients were asked not to take a bath or go swimming within the next 48 hours to prevent delays in wound healing. After some minutes of rest patients were free to leave.

**2.4. Study Design.** After the telephone interview potential participants were invited to be assessed on whether they were eligible for the study. The study physician also informed them about the study details. Written informed consent was obtained and patients were then randomly assigned to either a treatment group or a waiting list control group by means of sequentially numbered, sealed opaque envelopes, prepared by the study coordinator, who was neither involved in treatment nor in measurement. Patients were handed out the pain and medication diary (PaDi, MeDi) and measurement and treatment appointments were scheduled. Figure 1 illustrates the study design.

At baseline assessment (T1) study participants filled out the following questionnaires: pain at rest (PR), pain related to movement (PM), Neck Disability Index (NDI), and quality of life (SF-36). At last mechanical thresholds, that is, mechanical detection threshold (MDT), vibration detection threshold (VDT), and pressure pain threshold (PPT) were determined. At the end of T1 the treatment group received a single traditional cupping treatment whereas the waiting list control group received no treatment. Three days later participants returned for postintervention assessment (T2). They again filled out the questionnaires and underwent sensory testing. After they had completed the postintervention assessment, the wait-list control group was offered the cupping treatment.

**2.5. Statistical Analyses.** Treatment and waiting list control group were compared using chi-square analysis for discrete data and independent *t*-tests for continuous data on demographics, pain history, and baseline variables. For each outcome measure except the pain diary the results of the intervention were compared by analyses of covariance (ANCOVA) taking the post treatment measurement (T2) as a dependent and group as a between-subject factor. Respective baseline value of the outcome (T1) served as a covariate. This approach was chosen according to Vickers and Altman [51]. The intention-to-treat principle was used in this study,

however, since all drop outs were lost before T1 missing data could not be replaced by taking the last observation forward.

The pain diary (PaDi) was analysed by means of a repeated measurement ANCOVA. Within the statistical model the group variable served as between-subject factor, the post intervention measures as dependent factors, and the average pain in the week before T1 as covariate. Medication and concurrent treatments (MeDi) were continuously recorded in the diary and converted into relative amount of days under medication or treatment.

The General Health outcome was analysed by means of the Mann-Whitney *U* test.

Because of the pilot character of the study the level of statistical significance was not adjusted. An alpha of 0.05 was chosen for all analyses.

### 3. Results

**3.1. CONSORT Flowchart.** After the first telephone screening, 122 patients were invited for further evaluation. 50 of them fulfilled the study criteria and agreed to participate in the study.

Three patients in the treatment group and two in the waiting list control group resigned from participation before T1, no data could be collected from these patients. Final statistical analyses were conducted on 22 patients in the treatment group and on 23 patients in the waiting list control group. Figure 2 shows a flowchart of patient recruitment.

**3.2. Sample Characteristics.** All baseline values were comparable between the two groups, see Table 1. Two-thirds of participants in the study were female, the average age was 54.8 (TG) and 57.2 (WL). Study patients suffered for a very long time from neck pain; on average they reported 12.0 (TG) and 10.4 (WL) years of pain. The average pain intensity was rated 44.9 (TG) and 42.6 (WL). Expectation was comparable between the groups; therefore it was not included in further analysis.

Pre- and postintervention scores and estimated differences are presented in Table 2 and described in detail below.

**3.3. Pain.** Analysis of pain at rest (PR) shows a significant group difference at T2. TG reported 17.9 mm less pain on the

TABLE 1: Baseline demographic and clinical characteristics of trial groups.

Sociodemographic and clinical characteristics	TG (N = 22) mean ± SD	WL (N = 23) mean ± SD	P
Age (years)	54.8 ± 9.6	57.2 ± 9.4	0.393
Sex (F/M)	18/7	16/9	0.544
BMI (kg/m <sup>2</sup> )	28.9 ± 5.6	27.1 ± 4.3	0.203
Pain at rest (PR)	44.9 ± 18.2	42.6 ± 17.8	0.810
Average neck pain at baseline (PaDi)	4.8 ± 1.1	4.6 ± 1.4	0.552
History of neck pain (years)	12.0 ± 10.3	10.4 ± 11.5	0.618
Expected effectiveness of cupping therapy (VAS from 0 = not effective at all to 100 = highly effective)	72.8 ± 18.9	68.3 ± 20.5	0.448

TABLE 2: Outcomes of subjective measures at T1 and T2.

	T1		T2		Estimated difference at T2 diff TG versus WL* (95% CI)	ANCOVA		
	TG (n = 22) (mean ± SD)	WL (n = 23) (mean ± SD)	TG (n = 22) (mean ± SD)	WL (n = 23) (mean ± SD)		Df	F	P
Pain at rest (PR)	44.9±18.2	42.6±17.8	28.5 ± 23.9	45.7 ± 16.4	-17.9 (-29.2 to -6.6)	44	10.2	<b>0.003</b>
Maximal pain related to movement (PM)	53.9±25.7	65.6±22.1	29.1 ± 20.9	53.8 ± 26.1	-19.7 (-32.2 to -7.2)	44	10.1	<b>0.003</b>
Neck Disability Index (NDI)	29.9 ± 11.8	31.1 ± 9.1	24.5 ± 13.5	29.0 ± 9.3	-3.6 (-8.7 to 1.6)	44	2.0	0.168
SF-36 Physical Functioning	74.5 ± 19.1	71.3 ± 20.7	80.0 ± 15.3	70.2 ± 19.2	7.5 (1.4 to 13.5)	44	6.2	<b>0.017</b>
SF-36 Role Physical	39.8 ± 37.5	39.1 ± 41.9	58.0 ± 41.8	51.1 ± 38.8	6.4 (-12.0 to 24.8)	44	0.5	0.483
SF-36 Bodily Pain	37.8 ± 9.3	39.7 ± 9.1	53.1 ± 22.9	39.3 ± 11.4	14.9 (4.4 to 25.4)	44	8.2	<b>0.007</b>
SF-36 General Health Perception	62.2 ± 14.2	64.0 ± 19.3	64.0 ± 14.8	61.3 ± 20.7	4.1 (-3.3 to 11.5)	44	1.3	0.268
SF-36 Vitality	59.5 ± 21.0	53.5 ± 19.6	61.4 ± 21.4	53.5 ± 23.8	2.1 (-5.1 to 9.2)	44	0.3	0.561
SF-36 Social Function	70.5 ± 25.7	69.6 ± 24.7	79.0 ± 26.6	73.9 ± 26.9	4.4 (-6.8 to 15.6)	44	0.6	0.434
SF-36 Role Emotional	81.8 ± 36.7	71.0 ± 39.3	81.8 ± 33.7	76.8 ± 39.5	-0.1 (-19.8 to 19.6)	44	0.0	0.991
SF-36 Mental Health	72.4 ± 15.9	68.2 ± 18.3	69.6 ± 21.4	68.5 ± 22.4	-3.4 (-10.7 to 4.0)	44	0.9	0.358
SF-36 Physical Component Score	37.8 ± 7.8	38.7 ± 8.6	43.3 ± 8.5	39.0 ± 7.4	5.0 (1.4 to 8.5)	44	7.8	<b>0.008</b>
SF-36 Mental Component Score	51.8 ± 10.8	48.7 ± 11.3	50.4 ± 11.7	49.8 ± 13.6	-2.1 (-7.1 to 3.0)	44	0.7	0.415

\* Group differences and P values from an ANCOVA model with 2 groups, baseline values as covariate.




FIGURE 2: CONSORT flowchart of recruitment and study conditions.

VAS than WL. There was also a significant group difference on maximal pain related to movement (PM). While both groups were comparable in their pain ratings at T1, TG reported 19.7 mm less movement-related pain than the WL at T2.

Pain diary (PaDi) shows a sudden decline in pain ratings in TG at day 2, that is, the day after cupping therapy whereas it remained relatively stable in WL (Figure 3). A repeated measures ANCOVA revealed a significant interaction time × group ( $F = 5.22$ ,  $Df = 3/98$ ,  $\epsilon = 0.002$ ,  $P = 0.005$ ). Post hoc analyses confirmed a significant group difference at day 2 ( $\Delta -1.5$ , 95%CI  $-2.5$  to  $-0.4$ ,  $P = 0.008$ ) and single comparisons within TG also showed significant difference between baseline and day 2 ( $\Delta -0.9$ , 95% CI  $-1.7$  to  $-0.2$ ,  $P = 0.014$ ).

The majority of the patients went without any concurrent treatment in the week before T1 (medication: 60.0%, physiotherapeutic treatment: 91.1%). Those who did, used medication in  $27.8 \pm 22.2$  and physiotherapy in  $39.3 \pm 33.8$  of the days. The use of medication and concurrent treatments during the study was not further analysed.




FIGURE 3: Pain ratings (pain diary, NRS, mean  $\pm$  SD) decreased in TG at the day after cupping. \* $P < 0.05$ .

**3.4. Questionnaires.** No significant differences at T2 were found for the Neck Disability Index (NDI). The same was true for the Mental Component Score (SF-36) and the following subscales of the SF-36: Role Physical, Vitality, Social Function, Role Emotional, and General Health perceptions. On the other hand, significant differences occurred in the subscales Physical Functioning, Bodily Pain, and the Physical Component Score. At T2 TG reported significantly higher values on these scales indicating higher quality of life. Analysis of the General Health Outcome (SF-36) revealed a significant group difference with a significant higher rank for the TG (Mann Whitney  $U$  test; mean rank TG: 18.8; WL: 27.0;  $U = 160.5$ ;  $P = 0.019$ ) indicating more positive ratings than WL. In detail 11 patients of 22 in TG rated their health at least somewhat better than before, only two did so in WL. The majority in WL rated their health about the same as before (18 of 23). Three patients in each group even reported worse health (see also “safety issues”).

**3.5. Mechanical Sensory and Pain Thresholds.** MDT, VDT, and PPT for each group are listed in Table 3. Statistical analyses revealed no group differences for MDT or VDT, but for PPT. Significant differences in pressure pain thresholds were found at the pain-maximum and the pain-adjacent, but not at the control areas. Figure 4 displays the course of PPT at the pain-related areas.

**3.6. Safety.** Although most patients tolerated the treatment very well, adverse events were observed in some of the patients. One patient reported that the procedure itself was painful, other adverse events including slight reactions such as circulatory instability in the first minutes after treatment, tension headaches, a migraine attack, a reappearing tinnitus or wound healing itches. All of these adverse events were minor and transient.

However, two patients experienced more serious adverse events. As a result an ad hoc safety board was constituted to evaluate these adverse events and decide on further actions. The safety board was initiated by the principal investigator and consisted of the study physicians, the senior physicians of the Clinic for Complementary and Integrative Medicine, the head of the research group, an external statistician, and an external scientist, whose area of expertise is in safety issues and medical ethics. Two cases were presented and evaluated: one patient returned four days after treatment and reported worsened neck pain, a strong headache, and constant ear noises. The study physician examined the patient and diagnosed a cervical spine blockage. She was referred to an orthopedic for further diagnosis treatment. Later inquiries revealed that the symptoms had lasted for 2 to 3 weeks and improved subsequently. The neck pain, however, was neither better nor worse than before she participated in the study. Another patient complained of dizziness, nausea, and body misperception directly after treatment, so she had to lie down directly after treatment. Blood pressure and pulse measurement revealed normal circulatory function. The study physician diagnosed a transitory vagal reaction caused by the treatment and recommended her to rest until symptoms were resolved. After three hours lying and another hour sitting and walking the patient had mostly recovered. After examination the patient was sent home and a new appointment some days later was made. The patient then reported that the dizziness and nausea were fully resolved, but that the neck pain had worsened. The study physician offered her another treatment against the neck pain, which she refused. Later inquiries revealed that the pain had decreased within two weeks. The safety commission evaluated both incidents as adverse events, but not of a serious kind. Further actions as a consequence of occurrence of the adverse events involved obligatory follow-up check of patients in WL within two days after treatment. No adverse events were reported for WL after treatment.

## 4. Discussion

**4.1. Principal Findings.** To our knowledge, this is the first RCT where the effect of a single application of traditional cupping on chronic non-specific neck pain is investigated. Patients treated with cupping therapy showed significant improvements in their symptoms. Pain at rest (PR), maximal pain related to movement (PM), and bodily pain (SF-36) were reduced after a single cupping treatment. Pain diary (PaDi) showed a significant decline in pain ratings already on the day after cupping. According to the quality of life questionnaires (SF-36), the cupping treatment also significantly decreased Bodily Pain and improved Physical Functioning as well as the Physical Component Score.

Cupping also showed an effect on one of the nonsubjective parameters, the pressure-pain threshold (PPT), which is thought to reflect the functional status of (altered) pain perception. Pressure pain thresholds at pain-related areas increased or remained stable over time in the TG whereas patients of the WL control group became sensitised at those areas.

TABLE 3: Mechanical detection and pain thresholds at T1 and T2 (mean  $\pm$  SD).

		T1		T2		Estimated difference at T2 diff TG versus WL* (95% CI)	ANCOVA		
		TG ( <i>n</i> = 22) (mean $\pm$ SD)	WL ( <i>n</i> = 23) (mean $\pm$ SD)	TG ( <i>n</i> = 22) (mean $\pm$ SD)	WL ( <i>n</i> = 23) (mean $\pm$ SD)		<i>Df</i>	<i>F</i>	<i>P</i>
MDT in log(mN)	Pain- maximum	0.425 $\pm$ 0.427	0.443 $\pm$ 0.418	0.446 $\pm$ 0.508	0.411 $\pm$ 0.433	0.047 (−0.185 to 0.278)	44	0.686	
	Pain-sdjaent	0.290 $\pm$ 0.360	0.223 $\pm$ 0.374	0.382 $\pm$ 0.390	0.219 $\pm$ 0.477	0.124 (−0.094 to 0.341)	44	0.257	
VDT in X/8	Pain-maximum	6.061 $\pm$ 1.542	5.986 $\pm$ 1.135	6.061 $\pm$ 1.398	5.956 $\pm$ 1.075	0.024 (−0.482 to 0.529)	44	0.447	
	Pain-adjacent	5.288 $\pm$ 1.527	5.601 $\pm$ 1.162	5.515 $\pm$ 1.186	5.580 $\pm$ 1.401	0.199 (−0.366 to 0.764)	44	0.217	
PPT in log(kPa)	Pain-maximum	2.349 $\pm$ 0.169	2.357 $\pm$ 0.192	2.381 $\pm$ 0.149	2.299 $\pm$ 0.192	0.088, (0.029 to 0.148)	44	<b>0.005</b>	
	Pain-adjacent	2.396 $\pm$ 0.203	2.418 $\pm$ 0.200	2.423 $\pm$ 0.195	2.321 $\pm$ 0.204	0.118 (0.038 to 0.199)	44	<b>0.005</b>	

Baseline values were comparable between the groups. \*Differences were estimated by an ANCOVA model with 2 groups and the respective baseline values as covariate.




FIGURE 4: Course of pressure pain thresholds at pain-maximum and pain-adjacent (mean  $\pm$  SD) \**P* < 0.05.

**4.2. Interpretation.** In this study various pain measures such as pain at rest (PR), pain related to movement (PM), and pain diary (PaDi) data differed significantly between the TG and the WL after cupping. Thus, a single traditional cupping treatment appears to be effective in treating chronic non-specific neck pain. Since changes in the VAS and the NDI were also strongly correlated ( $r = 0.49$ ,  $P = 0.001$ ,  $N = 45$ ), pain relief appears to be associated with reduced impairment. However, there were no significant differences in NDI at T2 and the estimated difference was fewer than 10 points of improvement, which is the minimum clinical important change (MCIC) for the NDI [45]. This might have been due to the already low NDI scores at the beginning or due to the short followup. Interestingly, the pain diary ratings indicate that cupping has immediate effects. That is, the effects of traditional cupping are present already on the day after cupping treatment. This conforms to clinical observations, in which traditional cupping often shows dramatic and immediate effects on pain and other complaints.

Furthermore, Physical Functioning and the Physical Component Score (SF-36) changed significantly. These

changes are impressive since the post intervention measurement was only four days after the treatment. However, the immediate pain relief and the changes on physical scales of the SF-36 suggest that traditional cupping might work on a very somatic level. Effects of cupping have also been found on pressure pain thresholds at pain-related areas. Although the differences in PPT were relatively small, they were found consistently, suggesting that cupping might exert its effects locally, probably at receptor level.

Different modes of actions might explain the effect of traditional cupping on chronic neck pain. They involve neural, haematological, immune, and psychological effects [34]. Stimulation of the skin causes several autonomous, hormonal, and immune reactions [52]; this also applies for injuries due to the incisions [53]. Blood vessels in the treated areas are dilated by release of vasodilators such as adenosine, noradrenaline, and histamine, which lead to increased blood circulation [54].

In the course of cupping treatment, blood and other interstitial fluids are drawn out from the skin by the vacuum. Traditional cupping is mainly used in patients

with local blood congestion, swelling, and adhesions of the connective tissue in the neck region. It has been assumed that these congestions contain inflammatory extravasations [30, 35] and toxins. Cupping might therefore take the pressure off the tissue and relieve the neck area from these toxic congestions, which also increases circulation and lymphatic flow. Since circulation has been shown to be dysfunctional in chronic neck pain patients [12], cupping might restore normal circulation. Increased circulation in turn improves oxygen supply and cell metabolism [30] reducing the amount of inflammatory or toxic substances. This might also explain the significant effects of cupping on pressure pain thresholds at pain-related areas. Muscle spasm, congestion, and restricted blood flow can cause ischemic pain [55]. Accumulated inflammatory substances in skin and tissue might further induce hypersensitivity to noxious stimuli [16, 56], which is reflected by lowered pressure pain thresholds [57]. Since traditional cupping is supposed to evacuate toxins and inflammatory agents from the affected area and to restore normal circulation, this might explain the local effects on pressure pain thresholds.

The blood volume loss together with the local vasodilation might further increase parasympathetic activity by somatosympathetic reflexes, which corresponds well with the observed self-reported relaxation. Despite the invasiveness of traditional cupping the treatment group felt very relaxed after cupping treatment, on average they rated relaxation at  $62.2 \pm 20.1$  mm on a 100 mm VAS from 0 = “not relaxed at all” to 100 mm = “very relaxed.” In the worst case the reflex might cause a vasovagal syncope, as observed in one patient.

**4.3. Limitations of the Study.** The interpretation of the results might be limited due to choice of the passive control group. We are aware of the fact that unspecific effects such as expectation, conditioning, or environmental effects may have contributed to the observed overall effect size [35]. However, to date there are no suitable sham devices [58], even though there is an urgent need for a suitable sham procedure. Sham cupping by means of adhesives to keep the glass in place have been tried, but in our experience even cupping naïve subjects are likely to discover the sham intervention, even more so in traditional cupping than in dry cupping. Another serious problem is the impracticability of experimental blinding the assessor due to superficial wounds and visible cupping marks. On the other hand, traditional cupping is applied commonly in clinical CAM settings and has been proven to be helpful in alleviating several pain conditions [34, 36, 37], and patients request this treatment. Therefore there is a need for clinical trials on the topic, evaluating the efficacy and safety of these procedures. In conclusion, instead of ignoring the fact that there is a patient request for this reasonably invasive procedure and that there is limited data on efficacy and safety available, we decided to run an RCT with the best possible methodological approach, even though we are aware of its limitations.

Expectation was high in the patients participating in this study, a fact which might reflect a selection bias. It is likely that only patients with high expectations took part in this

study. However, baseline values were comparable between the groups.

A further limitation is the rather mild baseline pain intensity. Pain intensities reported by the patients in this study were at the lower end of the inclusion criteria scale. Some patients even fell below the required pain intensity of 40 on the VAS. This can be regarded as a possible source of bias, since patients probably exaggerated their complaints during screening to ensure inclusion into the study. The ceiling effect due to the low baseline pain intensity, likely limited the possible absolute reduction in pain intensity. Furthermore, due to the same problem, the likelihood of aggravation was also high due to the natural course of disease.

**4.4. Strengths of the Study.** Despite the limitations we could observe a strong pain reduction (VAS) of approximately  $32.8 \pm 51.1\%$  in the treatment group, compared to  $24.6 \pm 88.7\%$  in the waiting list control group. This pain reduction is comparable to studies on dry cupping [59] or massage [60] but in contrast to these methods the effect in traditional cupping occurs almost immediately after treatment. The overall pain reduction is within the range of clinical significance [43]. Moreover, treatment effects were also observed on pressure pain threshold, the concept of which is less transparent for the participant and therefore less open to presumption and hypothesising, which may make the results less biased than simple pain ratings.

## 5. Conclusion

A single application of traditional cupping might be effective in the treatment of chronic non-specific neck pain. Further studies are necessary to confirm these results and to evaluate the effectiveness of cupping compared to standard treatments. Studies investigating the effects of repeated traditional cupping interventions in different intervals and long-term observations are needed as well. Although measurements of sensory thresholds give possible hints on the physiology of pain processing, further investigations aiming at the mechanisms of action are necessary, too. However, the results of this first study and the patients' experiences with cupping therapy support the assumption that cupping might be a safe and effective treatment for chronic non-specific neck pain.

## Conflict of Interests

All authors disclose any commercial association that might create a conflict of interest in connection with the submitted paper. There is especially no competing financial interest for any of the authors.

## Acknowledgments

This paper was supported by a grant from the Karl and Veronica Carstens Foundation. The authors especially thank Henning Albrecht, Ph.D., and Rainer Lüdtke, M.S.c., for participating in the safety board.

## References

- [1] H. Merskey and N. Bogduk, *Classification of Chronic Pain: Description Of Chronic Pain Syndromes and Definitions of Pain Terms*, IASP Press, Seattle, Wash, USA, 1994.
- [2] R. Fejer, K. O. Kyvik, and J. Hartvigsen, "The prevalence of neck pain in the world population: a systematic critical review of the literature," *European Spine Journal*, vol. 15, no. 6, pp. 834–848, 2006.
- [3] A. Binder, "The diagnosis and treatment of nonspecific neck pain and whiplash," *Europa Medicophysica*, vol. 43, no. 1, pp. 79–89, 2007.
- [4] T. Skov, V. Borg, and E. Ørhede, "Psychosocial and physical risk factors for musculoskeletal disorders of the neck, shoulders, and lower back in salespeople," *Occupational and Environmental Medicine*, vol. 53, no. 5, pp. 351–356, 1996.
- [5] S. J. Linton, "A review of psychological risk factors in back and neck pain," *Spine*, vol. 25, no. 9, pp. 1148–1156, 2000.
- [6] K. Karjalainen, A. Malmivaara, M. van Tulder et al., "Multidisciplinary biopsychosocial rehabilitation for neck and shoulder pain among working age adults," *Cochrane Database of Systematic Reviews*, no. 2, Article ID CD002194, 2003.
- [7] S. M. Siivola, S. Levoska, K. Latvala, E. Hoskio, H. Vanharanta, and S. Keinänen-Kiukaanniemi, "Predictive factors for neck and shoulder pain: a longitudinal study in young adults," *Spine*, vol. 29, no. 15, pp. 1662–1669, 2004.
- [8] S. Hogg-Johnson, G. van der Velde, L. J. Carroll et al., "The burden and determinants of neck pain in the general population. Results of the bone and joint decade 2000–2010 task force on neck pain and its associated disorders," *Journal of Manipulative and Physiological Therapeutics*, vol. 32, no. 2, pp. S46–S60, 2009.
- [9] S. M. McLean, S. May, J. Klaber-Moffett, D. M. Sharp, and E. Gardiner, "Risk factors for the onset of non-specific neck pain: a systematic review," *Journal of Epidemiology and Community Health*, vol. 64, no. 7, pp. 565–572, 2010.
- [10] J. A. Schoffermann and M. E. Koestler, "Whiplash and neck pain-related disability," in *Handbook of Complex Occupational Disability Claims: Early Risk Identification, Intervention and Prevention*, I. Z. Schultz and R. J. Gatchel, Eds., pp. 203–216, Springer, New York, NY, USA, 2005.
- [11] I. Jensen and K. Harms-Ringdahl, "Strategies for prevention and management of musculoskeletal conditions. Neck pain," *Best Practice and Research: Clinical Rheumatology*, vol. 21, pp. 93–108, 2007.
- [12] R. Larsson, P. A. Öberg, and S. E. Larsson, "Changes of trapezius muscle blood flow and electromyography in chronic neck pain due to trapezius myalgia," *Pain*, vol. 79, no. 1, pp. 45–50, 1999.
- [13] V. Strøm, C. Røe, and S. Knardahl, "Work-induced pain, trapezius blood flux, and muscle activity in workers with chronic shoulder and neck pain," *Pain*, vol. 144, no. 1–2, pp. 147–155, 2009.
- [14] D. Falla, "Unravelling the complexity of muscle impairment in chronic neck pain," *Manual Therapy*, vol. 9, no. 3, pp. 125–133, 2004.
- [15] D. Scott, G. Jull, and M. Sterling, "Widespread sensory hypersensitivity is a feature of chronic whiplash-associated disorder but not chronic idiopathic neck pain," *Clinical Journal of Pain*, vol. 21, no. 2, pp. 175–181, 2005.
- [16] R. La Touche, C. Fernández-de-las-Peñas, J. Fernández-Carnero, S. Díaz-Parreño, A. Paris-Aleman, and L. Arendt-Nielsen, "Bilateral mechanical-pain sensitivity over the trigeminal region in patients with chronic mechanical neck pain," *Journal of Pain*, vol. 11, no. 3, pp. 256–263, 2010.
- [17] W. Jänig, "Grundlagen von reflextherapien," in *Naturheilverfahren und Unkonventionelle Medizinische Richtungen*, M. Bühring and F. H. Kremer, Eds., pp. 1–104, Springer, Berlin, Germany, 2005.
- [18] C. Fernández-de-las-Peñas, C. Alonso-Blanco, and J. C. Miangolarra, "Myofascial trigger points in subjects presenting with mechanical neck pain: a blinded, controlled study," *Manual Therapy*, vol. 12, no. 1, pp. 29–33, 2007.
- [19] K. Javanshir, R. Ortega-Santiago, M. A. Mohseni-Bandpei, J. C. Miangolarra-Page, and C. Fernández-De-Las-Peñas, "Exploration of somatosensory impairments in subjects with mechanical idiopathic neck pain: a preliminary study," *Journal of Manipulative and Physiological Therapeutics*, vol. 33, no. 7, pp. 493–499, 2010.
- [20] P. M. Peloso, A. Gross, T. Haines, K. Trinh, C. H. Goldsmith, and S. Burnie, "Medicinal and injection therapies for mechanical neck disorders," *Cochrane Database of Systematic Reviews*, no. 3, Article ID CD000319, 2007.
- [21] J. Ezzo, B. G. Haraldsson, A. R. Gross et al., "Massage for mechanical neck disorders: a systematic review," *Spine*, vol. 32, no. 3, pp. 353–362, 2007.
- [22] A. R. Gross, T. Kay, M. Hondras et al., "Manual therapy for mechanical neck disorders: a systematic review," *Manual Therapy*, vol. 7, no. 3, pp. 131–149, 2002.
- [23] T. M. Kay, A. Gross, C. Goldsmith, P. L. Santaguida, J. Hoving, and G. Bronfort, "Exercises for mechanical neck disorders," *Cochrane Database of Systematic Reviews*, no. 3, Article ID CD004250, 2005.
- [24] J. Ylinen, "Physical exercises and functional rehabilitation for the management of chronic neck pain," *Europa Medicophysica*, vol. 43, no. 1, pp. 119–132, 2007.
- [25] P. Tugwell, "Philadelphia panel evidence-based clinical practice guidelines on selected rehabilitation interventions for neck pain," *Physical Therapy*, vol. 81, no. 10, pp. 1701–1717, 2001.
- [26] A. Gross, J. Miller, J. D'Sylva et al., "Manipulation or mobilisation for neck pain: a Cochrane Review," *Manual Therapy*, vol. 15, no. 4, pp. 315–333, 2010.
- [27] S. K. Ndao-Brumblay and C. R. Green, "Predictors of complementary and alternative medicine use in chronic pain patients," *Pain Medicine*, vol. 11, no. 1, pp. 16–24, 2010.
- [28] K. J. Sherman, D. C. Cherkin, and C. J. Hogeboom, "The diagnosis and treatment of patients with chronic low-back pain by Traditional Chinese Medical acupuncturists," *Journal of Alternative and Complementary Medicine*, vol. 7, no. 6, pp. 641–650, 2001.
- [29] A. K. Kanodia, A. T. R. Legedza, R. B. Davis, D. M. Eisenberg, and R. S. Phillips, "Perceived benefit of Complementary and Alternative Medicine (CAM) for back pain: a national survey," *Journal of the American Board of Family Medicine*, vol. 23, no. 3, pp. 354–362, 2010.
- [30] I. Chirali, *Traditional Chinese Medicine Cupping Therapy*, Elsevier Churchill Livingstone, Philadelphia, Pa, USA, 2007.
- [31] A. Manz, *The Art of Cupping*, Thieme, Stuttgart, Germany, 2009.
- [32] A. Ahmadi, D. C. Schwebel, and M. Rezaei, "The efficacy of wet-cupping in the treatment of tension and migraine headache," *American Journal of Chinese Medicine*, vol. 36, no. 1, pp. 37–44, 2008.
- [33] H. Cao, J. Liu, and G. T. Lewith, "Traditional Chinese medicine for treatment of fibromyalgia: a systematic review of randomized controlled trials," *Journal of Alternative and Complementary Medicine*, vol. 16, no. 4, pp. 397–409, 2010.

- [34] K. Farhadi, D. C. Schwebel, M. Saeb, M. Choubsaz, R. Mohammadi, and A. Ahmadi, "The effectiveness of wet-cupping for nonspecific low back pain in Iran: a randomized controlled trial," *Complementary Therapies in Medicine*, vol. 17, no. 1, pp. 9–15, 2009.
- [35] J. I. Kim, M. S. Lee, D. H. Lee, K. Boddy, and E. Ernst, "Cupping for treating pain: a systematic review," *Evidence-Based Complementary and Alternative Medicine*, vol. 2011, Article ID 467014, 7 pages, 2011.
- [36] R. Lütke, U. Albrecht, R. Stange, and B. Uehleke, "Brachialgia paraesthetica nocturna can be relieved by "wet cupping"—results of a randomised pilot study," *Complementary Therapies in Medicine*, vol. 14, no. 4, pp. 247–253, 2006.
- [37] A. Michalsen, S. Bock, R. Lütke et al., "Effects of traditional cupping therapy in patients with carpal tunnel syndrome: a randomized controlled trial," *Journal of Pain*, vol. 10, no. 6, pp. 601–608, 2009.
- [38] H. Cao, C. Zhu, and J. Liu, "Wet cupping therapy for treatment of herpes zoster: a systematic review of randomized controlled trials," *Alternative therapies in health and medicine*, vol. 16, no. 6, pp. 48–54, 2010.
- [39] J. Abele, *Das Schröpfen: Eine Bewährte Alternative Heilmethode*, Urban & Fischer, München, Germany, 5th edition, 2003.
- [40] G. Bachmann and F. Pecker, *Die Schröpfkopfbehandlung*, Haug, Heidelberg, Germany, 3rd edition, 1978.
- [41] D. Irnich, N. Behrens, H. Molzen et al., "Randomised trial of acupuncture compared with conventional massage and "sham" laser acupuncture for treatment of chronic neck pain," *British Medical Journal*, vol. 322, no. 7302, pp. 1574–1577, 2001.
- [42] P. E. Bijur, W. Silver, and E. J. Gallagher, "Reliability of the visual analog scale for measurement of acute pain," *Academic Emergency Medicine*, vol. 8, no. 12, pp. 1153–1157, 2001.
- [43] R. H. Dworkin, D. C. Turk, K. W. Wyrwich et al., "Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations," *Journal of Pain*, vol. 9, no. 2, pp. 105–121, 2008.
- [44] H. Vernon and S. Mior, "The neck disability index: a study of reliability and validity," *Journal of Manipulative and Physiological Therapeutics*, vol. 14, pp. 409–415, 1991.
- [45] J. C. MacDermid, D. M. Walton, S. Avery et al., "Measurement properties of the neck disability index: a systematic review," *Journal of Orthopaedic and Sports Physical Therapy*, vol. 39, no. 5, pp. 400–416, 2009.
- [46] M. Bullinger, I. Kirchberger, and J. Ware, "The German SF-36 health survey translation and psychometric testing of a generic instrument for the assessment of health-related quality of life," *Journal of Public Health*, vol. 3, no. 1, pp. 21–36, 1995.
- [47] M. Bullinger and I. Kirchberger, *SF-36. Fragebogen zum Gesundheitszustand. Handanweisung*, Hogrefe, Göttingen, Germany, 1998.
- [48] S. D. Keller, M. S. Bayliss, J. E. Ware Jr., M. A. Hsu, A. M. Damiano, and T. F. Goss, "Comparison of responses to SF-36 health survey questions with one-week and four-week recall periods," *Health Services Research*, vol. 32, no. 3, pp. 367–384, 1997.
- [49] R. Rolke, W. Magerl, K. A. Campbell et al., "Quantitative sensory testing: a comprehensive protocol for clinical trials," *European Journal of Pain*, vol. 10, no. 1, pp. 77–88, 2006.
- [50] R. Rolke, R. Baron, C. Maier et al., "Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values," *Pain*, vol. 123, no. 3, pp. 231–243, 2006.
- [51] A. J. Vickers and D. G. Altman, "Analysing controlled trials with baseline and follow up measurements," *British Medical Journal*, vol. 323, no. 7321, pp. 1123–1124, 2001.
- [52] I. Lund and T. Lundberg, "Are minimal, superficial or sham acupuncture procedures acceptable as inert placebo controls?" *Acupuncture in Medicine*, vol. 24, no. 1, pp. 13–15, 2006.
- [53] A. Sato, "Neural mechanisms of autonomic responses elicited by somatic sensory stimulation," *Neuroscience and Behavioral Physiology*, vol. 27, no. 5, pp. 610–621, 1997.
- [54] M. Ernst and M. H. M. Lee, "Sympathetic effects of manual and electrical acupuncture of the Tsusanli knee point: comparison with the Hoku hand point sympathetic effects," *Experimental Neurology*, vol. 94, no. 1, pp. 1–10, 1986.
- [55] P. Weerapong, P. A. Hume, and G. S. Kolt, "The mechanisms of massage and effects on performance, muscle recovery and injury prevention," *Sports Medicine*, vol. 35, no. 3, pp. 235–256, 2005.
- [56] V. Johnston, N. L. Jimmieson, G. Jull, and T. Souvlis, "Quantitative sensory measures distinguish office workers with varying levels of neck pain and disability," *Pain*, vol. 137, no. 2, pp. 257–265, 2008.
- [57] R. D. Treede, R. Rolke, K. Andrews, and W. Magerl, "Pain elicited by blunt pressure: neurobiological basis and clinical relevance," *Pain*, vol. 98, no. 3, pp. 235–240, 2002.
- [58] E. Ernst, "Testing traditional cupping therapy," *Journal of Pain*, vol. 10, no. 6, p. 555, 2009.
- [59] R. Lauche, H. Cramer, K. -E. Choi et al., "The influence of a series of five dry cupping treatments on pain and mechanical thresholds in patients with chronic non-specific neck pain—a randomised controlled pilot study," *BMC Complementary and Alternative Medicine*, vol. 11, article 63, 2011.
- [60] K. J. Sherman, D. C. Cherkin, R. J. Hawkes, D. L. Miglioretti, and R. A. Deyo, "Randomized trial of therapeutic massage for chronic neck pain," *Clinical Journal of Pain*, vol. 25, no. 3, pp. 233–238, 2009.