

Short-term exposure to fine particulate matter and nitrogen dioxide and mortality in 4 countries

Yiqun Ma, Federica Nobile, Anne Marb, Robert Dubrow, Massimo Stafoggia, Susanne Breitner, Patrick L. Kinney, Kai Chen

Angaben zur Veröffentlichung / Publication details:

Ma, Yiqun, Federica Nobile, Anne Marb, Robert Dubrow, Massimo Stafoggia, Susanne Breitner, Patrick L. Kinney, and Kai Chen. 2024. "Short-term exposure to fine particulate matter and nitrogen dioxide and mortality in 4 countries." *JAMA Network Open* 7(3): e2354607.
<https://doi.org/10.1001/jamanetworkopen.2023.54607>.

Nutzungsbedingungen / Terms of use:

CC BY 4.0



Short-Term Exposure to Fine Particulate Matter and Nitrogen Dioxide and Mortality in 4 Countries

Yiqun Ma, MPhil; Federica Nobile, MSc; Anne Marb, MSc; Robert Dubrow, MD, PhD; Massimo Stafoggia, PhD; Susanne Breitner, ScD; Patrick L. Kinney, ScD; Kai Chen, PhD

Abstract

IMPORTANCE The association between short-term exposure to air pollution and mortality has been widely documented worldwide; however, few studies have applied causal modeling approaches to account for unmeasured confounders that vary across time and space.

OBJECTIVE To estimate the association between short-term changes in fine particulate matter (PM_{2.5}) and nitrogen dioxide (NO₂) concentrations and changes in daily all-cause mortality rates using a causal modeling approach.

DESIGN, SETTING, AND PARTICIPANTS This cross-sectional study used air pollution and mortality data from Jiangsu, China; California; central-southern Italy; and Germany with interactive fixed-effects models to control for both measured and unmeasured spatiotemporal confounders. A total of 8 963 352 deaths in these 4 regions from January 1, 2015, to December 31, 2019, were included in the study. Data were analyzed from June 1, 2021, to October 30, 2023.

EXPOSURE Day-to-day changes in county- or municipality-level mean PM_{2.5} and NO₂ concentrations.

MAIN OUTCOMES AND MEASURES Day-to-day changes in county- or municipality-level all-cause mortality rates.

RESULTS Among the 8 963 352 deaths in the 4 study regions, a 10-μg/m³ increase in daily PM_{2.5} concentration was associated with an increase in daily all-cause deaths per 100 000 people of 0.01 (95% CI, 0.001-0.01) in Jiangsu, 0.03 (95% CI, 0.004-0.05) in California, 0.10 (95% CI, 0.07-0.14) in central-southern Italy, and 0.04 (95% CI, 0.02-0.05) in Germany. The corresponding increases in mortality rates for a 10-μg/m³ increase in NO₂ concentration were 0.04 (95% CI, 0.03-0.05) in Jiangsu, 0.03 (95% CI, 0.01-0.04) in California, 0.10 (95% CI, 0.05-0.15) in central-southern Italy, and 0.05 (95% CI, 0.04-0.06) in Germany. Significant effect modifications by age were observed in all regions, by sex in Germany (eg, 0.05 [95% CI, 0.03-0.06] for females in the single-pollutant model of PM_{2.5}), and by urbanicity in Jiangsu (0.07 [95% CI, 0.04-0.10] for rural counties in the 2-pollutant model of NO₂).

CONCLUSIONS AND RELEVANCE The findings of this cross-sectional study contribute to the growing body of evidence that increases in short-term exposures to PM_{2.5} and NO₂ may be associated with increases in all-cause mortality rates. The interactive fixed-effects model, which controls for unmeasured spatial and temporal confounders, including unmeasured time-varying confounders in different spatial units, can be used to estimate associations between changes in short-term exposure to air pollution and changes in health outcomes.

JAMA Network Open. 2024;7(3):e2354607. doi:10.1001/jamanetworkopen.2023.54607

Key Points

Question What are the associations between short-term changes in fine particulate matter (PM_{2.5}) and nitrogen dioxide (NO₂) concentrations and changes in daily all-cause mortality rates?

Findings This cross-sectional study of more than 8.9 million deaths found that a 10-μg/m³ increase in daily PM_{2.5} concentrations was associated with increases in daily all-cause deaths per 100 000 people of 0.01 in Jiangsu, China, 0.03 in California, 0.10 in central-southern Italy, and 0.04 in Germany; corresponding increases in mortality rates for the same increase in NO₂ concentrations were 0.04, 0.03, 0.10, and 0.05, respectively.

Meaning The findings of this study add to the growing evidence that increases in short-term exposures to PM_{2.5} and NO₂ may be associated with increases in mortality rates.

+ [Invited Commentary](#)

+ [Supplemental content](#)

Author affiliations and article information are listed at the end of this article.

Open Access. This is an open access article distributed under the terms of the CC-BY License.

JAMA Network Open. 2024;7(3):e2354607. doi:10.1001/jamanetworkopen.2023.54607

March 1, 2024 1/13

Introduction

The association between short-term exposure to air pollution and mortality has been widely documented worldwide.¹⁻⁴ Most previous epidemiological studies have investigated the effects of short-term exposure to air pollution on mortality using time-series analyses.¹⁻³ By contrasting each spatial unit to itself over time, time-series studies avoid the confounding effects of spatial and time-invariant variables by design; however, results from time-series analyses may be biased due to unmeasured temporal confounding.⁵⁻⁷ Alternative causal modeling approaches that account for unmeasured time-varying confounders in different spatial units offer an attractive alternative for studying the association between short-term air pollution exposure and mortality.

Fixed-effects models, particularly 2-way fixed-effects (TWFE) models, have been widely applied for causal inference in econometrics and other social sciences.⁸ In the field of air pollution epidemiology, TWFE models have been recently applied, first in studies of long-term air pollution exposure⁹⁻¹¹ and most recently to estimate the effects of short-term fine particulate matter (PM_{2.5}) exposure on hospitalizations.⁷ By introducing indicators for each spatial unit and each time unit, TWFE models can control for all spatial confounders that vary across space but not across the study timescales (eg, geographical and socioeconomic factors) and all temporal confounders that vary by time but not space (eg, day of week and seasonality).^{9,12} However, this approach assumes no unmeasured confounders that display different temporal variations across spatial units (time-varying spatial unit effects), an assumption that is often violated.¹³ To relax this assumption of the TWFE model, the more flexible interactive fixed-effects (IFE) model was developed. By decomposing the unmeasured time-varying spatial unit-specific confounders into heterogeneous effects of common trends, the IFE model can potentially control for such confounders in the regression.¹³⁻¹⁵ To the best of our knowledge, this study is the first application of the IFE approach to model the causal effects of short-term exposure to air pollution on mortality.

Substantial heterogeneity in the association between air pollution concentrations and daily mortality exists across countries and regions,¹⁻³ especially between developed and developing countries and between regions with low and high levels of pollution.^{1,3} A multicountry approach has the potential to comprehensively describe the effects of air pollution in regions with different air pollution levels and socioeconomic statuses. This study focused on 4 study regions: Jiangsu Province, China (hereinafter Jiangsu); California; central-southern Italy; and Germany. We specifically examined PM_{2.5} and nitrogen dioxide (NO₂) as 2 major traffic- and/or health-related air pollutants. Using data from these 4 regions from January 1, 2015, to December 31, 2019, we estimated the association between changes in daily PM_{2.5} and NO₂ concentrations and changes in daily mortality rates using an IFE approach. Potential effect modifications by sex, age, and urbanicity were further examined.

Methods

This cross-sectional study used anonymized daily county- or municipality-level mortality records and was approved by the Yale Institutional Review Board. The need for informed consent was waived owing to the use of publicly available deidentified data. This study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline.

Mortality Rate

We collected daily all-cause mortality data from 2015 to 2019 by sex and age in each spatial unit (county in Jiangsu, California, and Germany and municipality in central-southern Italy) of each region (eAppendix 1 in [Supplement 1](#)). Based on the population size of each spatial unit (eAppendix 1 in [Supplement 1](#)), we calculated daily spatial unit-level all-cause mortality rates for the whole population and population subgroups by sex (male and female) and age (0-74 and ≥75 years). In central-southern Italy, to have a sufficient number of deaths in each included municipality, we

included in the analysis only the 669 municipalities with a population of 10 000 or greater of a total of 3250 municipalities. This population cutoff was only applied to central-southern Italy because all counties in the other 3 regions satisfied this population criterion.

Air Pollution and Meteorological Variables

Daily concentrations of PM_{2.5} and NO₂ at 1 × 1-km² resolution from 2015 to 2019 were estimated by spatiotemporal models for Jiangsu,^{16,17} central-southern Italy,^{18,19} and Germany^{20,21} (eAppendix 2 in Supplement 1). In brief, the models incorporated variables from satellite observations, chemical transport model simulations, land use and/or land cover characteristics, and meteorological data as predictive factors and applied advanced machine learning algorithms to model daily air pollution concentrations. The modeled air pollution data were aggregated to daily county- or municipality-level concentrations using the geographic mean and matched to the mortality data.

To the best of our knowledge, there was no available high-resolution daily spatiotemporal model for PM_{2.5} and NO₂ covering 2015 to 2019 for California at the time of our study, so we relied on the records of air quality monitoring sites managed by the US Environmental Protection Agency Air Quality System.²² The 32 California counties with monitoring sites for both PM_{2.5} and NO₂ were included in the analysis (of 58 counties), including data from 108 sites for PM_{2.5} and 115 sites for NO₂. For counties in California with multiple monitoring sites, we calculated county-level mean air pollution concentrations and used these in our analyses.

Hourly air temperature data at 0.1° × 0.1° resolution were extracted from the ERA5-Land reanalysis dataset²³ for all 4 regions. Like the air pollution data, we calculated the mean air temperature data for each day in each spatial unit. We also calculated the daily dew point temperature and relative humidity in each region (eAppendix 3 in Supplement 1).

Statistical Analysis

Data were analyzed from June 1, 2021, to October 30, 2023. The IFE model assumes that the unmeasured spatial unit effects that change over time have a factor structure and that all regressors and factors are stationary. Additionally, the IFE model allows a weak serial and cross-sectional correlation.¹⁴ In our study, the IFE model was expressed as $\Delta \text{mortality rate}_{i,t} = \mu + \alpha_i + \beta \Delta \text{air pollution}_{i,t} + ns(\Delta \text{temperature}_{i,t}, df = 5) + v_{i,t} + \epsilon_{i,t}$, where $v_{i,t} = \sum_{l=1}^d \lambda_{i,l} f_{l,t}$, the outcome variable $\Delta \text{mortality rate}_{i,t}$ is the change in all-cause mortality rate from day $t - 1$ to day t in spatial unit i , and $\Delta \text{air pollution}_{i,t}$ is the change in mean concentration of PM_{2.5} or NO₂ from day $t - 1$ to day t in spatial unit i . The day-to-day change in air temperature on the same lag day as the air pollution variable ($\Delta \text{temperature}_{i,t}$) was included as a flexible natural cubic spline with 5 df . First-order differences of the observed time series were taken to remove the long-term and seasonal trend of mortality rate, air pollution, and meteorological factors and to meet the stationarity and normality assumptions of the model. The α_i refers to time-invariant spatial unit effects, which can help control for the effects of spatial confounders that do not vary over time. The μ is the intercept; $\epsilon_{i,t}$ is the error term; and $v_{i,t}$ is the unmeasured time-varying spatial unit effect, which is decomposed into d common time-varying factors $f_{l,t}$, with corresponding unobserved spatial unit-level loading parameters $\lambda_{i,l}$.^{24,25} The number of factors (d) was selected following the criteria proposed by Bai and Ng²⁴ (eAppendix 4 in Supplement 1). The coefficient of changes in air pollution (β) represents the expected change in the daily mortality rate for each unit change in the daily air pollution level. Heteroskedasticity and autocorrelation-consistent standard errors (SEs) were calculated to account for potential heteroskedasticity and autocorrelation in the error terms. The sample R code for our main IFE model can be found in eAppendix 5 in Supplement 1.

To account for the potential confounding effects from the other air pollutant, we also used 2-pollutant models in which both PM_{2.5} and NO₂ were included. We explored the lag pattern in the associations of PM_{2.5} and NO₂ changes with changes in daily mortality rate on the current day and the previous 2 days using both single lag days (lag0 to lag2) and cumulative lag days (lag01 to lag02),

which was commonly used in previous studies.¹⁻³ The IFE analyses were performed with R software, version 4.1.3 (R Project for Statistical Computing) using the package phtt.²⁵

We conducted subgroup analyses to estimate potential differences in the effects of air pollution according to sex (male and female) and age (0-74 years and ≥ 75 years). In addition, we performed stratified analyses to examine the potential effect modification by urbanicity (urban and rural). The classification criteria of urban and rural areas in each study region are described in eAppendix 6 in Supplement 1. We tested the statistical difference in effect estimates between groups by calculating the z score as $(\hat{Q}_1 - \hat{Q}_2) / \sqrt{(SE_1)^2 + (SE_2)^2}$, where $\hat{Q}_1$ and $\hat{Q}_2$ are the estimates and SE_1 and SE_2 are their respective standard errors.²⁶ Statistical significance was evaluated at $P < .05$ (2-sided test).

To evaluate the model specification, we performed a randomization test in which we randomized the PM_{2.5} or NO₂ concentration 2000 times across all 1825 days in each spatial unit. Such a randomization test is widely applied to detect temporal dependence in panel models due to model misspecification.²⁷ In addition, we performed several sensitivity analyses to test the robustness of our results: (1) we additionally adjusted for relative humidity or dew point temperature in the model; (2) we applied a traditional fixed-effects model, which does not consider unmeasured time-varying spatial unit effects; and (3) we used 4 *df* or 6 *df* instead of 5 *df* in the natural cubic spline of air temperature. Furthermore, we compared the results from the IFE model with those from a traditional 2-stage time-series model (eAppendix 7 in Supplement 1).^{2,3}

Results

Description of Mortality Rate and Air Pollution Exposure

This study covered a total of 8 963 352 deaths, including 2 633 920 in Jiangsu (45.0% female and 55.0% male; 60.0% aged ≥ 75 years), 1 237 862 in California (48.3% female and 51.7% male; 58.7% aged ≥ 75 years), 1 227 482 in central-southern Italy (51.8% female and 48.2% male; 73.5% aged ≥ 75 years), and 3 864 088 in Germany (52.1% female and 47.9% male; 72.5% aged ≥ 75 years) from 2015 to 2019. The descriptive statistics of daily mortality rate, PM_{2.5} and NO₂ concentrations, and air temperature in each region are presented in Table 1. No data were missing in this study. The mean (SD) spatial unit-level PM_{2.5} and NO₂ concentrations were the highest in Jiangsu (PM_{2.5}, 50.8 [27.8] $\mu\text{g}/\text{m}^3$; NO₂, 32.2 [1.28] $\mu\text{g}/\text{m}^3$). The mean (SD) PM_{2.5} concentration was the lowest in California (9.9 [9.2] $\mu\text{g}/\text{m}^3$), and the mean (SD) NO₂ concentration was the lowest in Germany (12.2 [7.8] $\mu\text{g}/\text{m}^3$) (Table 1). The spatial distribution of PM_{2.5} and NO₂ in the 4 study regions is displayed in Figure 1.

Association Between Short-Term Air Pollution Exposure and Mortality Rate

Figure 2 shows the estimated associations between changes in daily PM_{2.5} and NO₂ concentrations and changes in daily mortality rate (per 100 000 people) from both single- and 2-pollutant models in each region. In single-pollutant models, increases in PM_{2.5} and NO₂ concentrations were associated with increases in mortality rates in all 4 study regions. We define the main lag for each pollutant in each region as the lag with the greatest effect size from the single-pollutant model; the estimated coefficients for all lags are presented in eTable 1 in Supplement 1. In the single-pollutant models for PM_{2.5} in Jiangsu and Germany, the greatest effect estimates were observed for lag01; a 10- $\mu\text{g}/\text{m}^3$ increase in 2-day mean concentration of PM_{2.5} was associated with increases in daily all-cause deaths per 100 000 people of 0.01 (95% CI, 0.001-0.01) in Jiangsu and 0.04 (95% CI, 0.02-0.05) in Germany. In California and central-southern Italy, the highest estimate for PM_{2.5} in the single-pollutant model was found for lag02; a 10- $\mu\text{g}/\text{m}^3$ increase in 3-day mean PM_{2.5} concentrations was associated with increases in daily all-cause deaths per 100 000 people of 0.03 (95% CI, 0.004-0.05) in California and 0.10 (95% CI, 0.07-0.14) in central-southern Italy. In the single-pollutant models, the largest effect estimates for a 10- $\mu\text{g}/\text{m}^3$ increase in NO₂ concentration were 0.04 (95% CI, 0.03-0.05) deaths per 100 000 people in Jiangsu (lag02), 0.03 (95% CI, 0.01-0.04) deaths per 100 000 people in California (lag02), 0.10 (95% CI, 0.05-0.15) deaths per 100 000 people in central-southern Italy (lag02), and 0.05 (95% CI, 0.04-0.06) deaths per 100 000 people in Germany (lag01). The results

from 2-pollutant models were generally consistent with those from single-pollutant models, though the estimated coefficients were relatively smaller (Figure 2 and eTable 1 in Supplement 1).

Effect Modification by Sex, Age, and Urbanicity

We used the lags with the largest effect sizes in each region to examine potential effect modification. In subgroup analyses by sex, we found a significantly larger effect size in the association between

Table 1. Descriptive Statistics of Daily County- or Municipality-Level Mortality Rate and Environmental Data, 2015 to 2019

Study region	Mean (SD)	Median (IQR) [range]
Jiangsu, China ^a		
All-cause mortality rate ^b		
All-group	1.9 (0.7)	1.8 (0.9) [0.0 to 11.2]
Male	2.0 (1.0)	1.9 (1.2) [0.0 to 11.4]
Female	1.7 (0.9)	1.6 (1.1) [0.0 to 13.5]
Aged 0-74 y	0.8 (0.4)	0.7 (0.5) [0.0 to 5.9]
Aged ≥75	26.7 (13.5)	24.6 (16.1) [0.0 to 210.7]
Environmental factors		
PM _{2.5} concentration, µg/m ³	50.8 (27.8)	44.0 (31.8) [6.0 to 254.4]
NO ₂ concentration, µg/m ³	32.2 (12.8)	29.8 (15.1) [6.6 to 122.5]
Air temperature, °C	16.1 (9.1)	17.1 (15.7) [-10.0 to 34.8]
California ^c		
All-cause mortality rate ^b		
All-group	2.0 (0.8)	1.9 (0.8) [0.0 to 9.5]
Male	2.0 (1.1)	1.9 (1.1) [0.0 to 14.7]
Female	1.9 (1.1)	1.8 (1.0) [0.0 to 10.4]
Aged 0-74 y	0.9 (0.6)	0.8 (0.6) [0.0 to 6.7]
Aged ≥75 y	19.8 (10.0)	18.7 (10.4) [0.0 to 108.5]
Environmental factors		
PM _{2.5} concentration, µg/m ³	9.9 (9.2)	8.0 (6.4) [0.0 to 411.7]
NO ₂ concentration, µg/m ³	15.7 (11.4)	12.8 (13.6) [0.0 to 108.2]
Air temperature, °C	16.3 (7.0)	15.7 (10.1) [-5.0 to 40.0]
Central-southern Italy ^d		
All-cause mortality rate ^b		
All-group	2.7 (3.9)	0.0 (4.7) [0.0 to 69.9]
Male	2.7 (5.5)	0.0 (3.4) [0.0 to 106.0]
Female	2.7 (5.3)	0.0 (3.5) [0.0 to 79.0]
Aged 0-74 y	0.8 (2.2)	0.0 (0.0) [0.0 to 43.6]
Aged ≥75 y	19.2 (32.7)	0.0 (31.6) [0.0 to 645.7]
Environmental factors		
PM _{2.5} concentration, µg/m ³	12.6 (6.0)	11.3 (5.7) [1.3 to 99.3]
NO ₂ concentration, µg/m ³	12.4 (7.3)	10.1 (7.6) [1.8 to 81.2]
Air temperature, °C	15.9 (6.8)	15.5 (10.6) [-12.7 to 35.3]
Germany ^e		
All-cause mortality rate ^b		
All-group	2.6 (1.6)	2.4 (1.9) [NA]
Male	2.5 (2.1)	2.2 (2.5) [NA]
Female	2.7 (2.2)	2.4 (2.5) [NA]
Aged 0-74 y	0.8 (0.9)	0.7 (1.2) [NA]
Aged ≥75 y	16.3 (10.9)	15.3 (13.3) [NA]
Environmental factors		
PM _{2.5} concentration, µg/m ³	10.0 (7.1)	8.2 (6.8) [0.5 to 141.7]
NO ₂ concentration, µg/m ³	12.2 (7.8)	10.1 (8.8) [0.6 to 83.9]
Air temperature, °C	10.2 (7.4)	10.0 (11.9) [-17.9 to 31.8]

Abbreviations: NA, not available; NO₂, nitrogen dioxide; PM_{2.5}, fine particulate matter.

^a Includes 82 counties with a total population of 77 955 026 persons.

^b Reported per 100 000 population.

^c Includes the 32 out of 58 counties with Environmental Protection Agency air quality monitoring stations, with a total population of 37 290 255.


^d Includes 669 of 3250 municipalities with a population of 10 000 or greater, with a total population of 24 545 330.

^e Includes 401 counties with a total population of 82 735 005. Minimum and maximum values of mortality rates in Germany were not allowed to be released according to the information protection policies of the German Statistical Offices.

changes in daily PM_{2.5} concentration and changes in daily mortality rate among females compared with males in Germany in both single-pollutant (0.05 [95% CI, 0.03-0.06] deaths per 100 000 people; *P* = .03) and 2-pollutant (0.04 [95% CI, 0.02-0.05] deaths per 100 000 people; *P* = .01) models (**Table 2**). No statistically significant differences between sexes were observed for PM_{2.5} in other regions or for NO₂.

In subgroup analyses by age, changes in daily PM_{2.5} concentration and changes in daily mortality rate were found to be significantly greater among people 75 years and older than among those aged 0 to 74 years in central-southern Italy for the single-pollutant model (1.00 [95% CI, 0.65-1.35] deaths per 100 000 people; *P* < .001) and the 2-pollutant model (0.90 [95% CI, 0.46-1.33] deaths per 100 000 people; *P* < .001) and in Germany for the single-pollutant model (0.21 [95% CI, 0.13-0.29] deaths per 100 000 people; *P* < .001) and the 2-pollutant model (0.11 [95% CI, 0.02-0.20] deaths per 100 000 people; *P* = .03). Consistently, for NO₂, the association was greater among people 75 years and older compared with those aged 0 to 74 years in all 4 regions in single-pollutant models (0.46 [95% CI, 0.23-0.69] deaths per 100 000 people in Jiangsu; 0.40 [95% CI, 0.19-0.61] deaths per 100 000 people in California; 0.98 [95% CI, 0.45-1.51] deaths per 100 000 people in central-southern Italy; and 0.31 [95% CI, 0.22-0.40] deaths per 100 000 people in Germany) (*P* < .001 for all) and in Jiangsu (0.61 [95% CI, 0.30-0.91] deaths per 100 000 people), California (0.39 [95% CI, 0.17-0.60] deaths per 100 000 people), and Germany (0.26 [95% CI, 0.17-0.36] deaths per 100 000 people) in 2-pollutant models (*P* < .001 for all) (Table 2).

Figure 1. Daily Mean Fine Particulate Matter (PM_{2.5}) and Nitrogen Dioxide (NO₂) Concentrations in Each Spatial Unit in All Study Regions



The blank areas in California represent counties that lack air quality monitoring stations for both PM_{2.5} and NO₂; the blank areas in central-southern Italy represent municipalities with a population of less than 10 000 individuals.


In stratified analyses by urbanicity, we found no statistically significant differences in the association between changes in daily PM_{2.5} or NO₂ concentration and changes in daily mortality rate between urban and rural areas except for NO₂ in Jiangsu, where the difference was significant in rural compared with urban areas in the 2-pollutant model (0.07 [95% CI, 0.04-0.10] deaths per 100 000 people; $P = .02$) (Figure 3 and eTable 2 in Supplement 1).

Randomization Test, Sensitivity Analysis, and Comparative Analysis

As shown in eFigure 1 in Supplement 1, the distributions of the model coefficients when the changes in PM_{2.5} or NO₂ concentrations were randomly assigned 2000 times across days were centered at approximately 0 in all 4 regions, and the coefficient estimates from our main model fell substantially outside these distributions in all study regions. This indicates that the estimated associations between air pollution and mortality in our study were unlikely driven by temporal dependence due to a misspecification of the model.

Sensitivity analyses showed that our results remained robust when additionally adjusting for dew point temperature and relative humidity using traditional fixed-effects models and alternative degrees of freedom for air temperature (eTable 3 in Supplement 1). The estimated unmeasured time-varying spatial unit effects in the main IFE model for the main lag in Jiangsu and California are visualized in eFigure 2 in Supplement 1. We observed 1 unmeasured factor that displayed different temporal variations across spatial units in Jiangsu and 5 in California. However, these factors may not

Figure 2. Estimated Change in Daily Mortality Rate Associated With a 10- $\mu\text{g}/\text{m}^3$ Increase in Fine Particulate Matter (PM_{2.5}) or Nitrogen Dioxide (NO₂) Concentration



Single- and 2-pollutant models on different lag days in each study region are shown. The error bars represent the 95% CIs. Lag0 to lag2 represent single lag days; lag01 to lag02, cumulative lag days.

be directly interpretable because they are, at best, linear transformations of the true factors. No such factors were detected by the IFE model in central-southern Italy and Germany.

The results from the traditional 2-stage time-series analysis are shown in eFigure 3 in Supplement 1. While the estimates from the IFE model and those from the time-series analysis should not be directly compared because of their different interpretations, the positive estimates from each analysis reinforce the findings from the IFE model. Both suggested that short-term exposure to PM_{2.5} or NO₂ was associated with an increase in all-cause mortality.

Discussion

Using the IFE model, a causal modeling approach, we found that short-term increases in PM_{2.5} and NO₂ concentrations were associated with increases in daily mortality rate in Jiangsu, California, central-southern Italy, and Germany. The IFE model has the advantage of controlling for unobserved confounding factors that vary over time within each spatial unit. This study showcases the

Table 2. Estimated Changes in Daily Mortality Rate Associated With a 10-μg/m³ Increase in PM_{2.5} and NO₂ Concentrations by Sex and Age^a

Subgroup	Single-pollutant model				Two-pollutant model			
	PM _{2.5}		NO ₂		PM _{2.5}		NO ₂	
	Estimated changes in daily mortality rate (95% CI) ^b	P value ^c	Estimated changes in daily mortality rate (95% CI) ^b	P value ^c	Estimated changes in daily mortality rate (95% CI) ^b	P value ^c	Estimated changes in daily mortality rate (95% CI) ^b	P value ^c
Jiangsu, China								
Sex								
Male	0.01 (0.003 to 0.01)		0.04 (0.03 to 0.05)		−0.0006 (−0.01 to 0.004)		0.05 (0.03 to 0.06)	
Female	0.01 (0.001 to 0.01)	.51	0.04 (0.03 to 0.05)	.83	−0.004 (−0.01 to 0.0004)	.31	0.05 (0.04 to 0.07)	.39
Age, y								
0–74	0.008 (−0.0009 to 0.002)		0.01 (0.01 to 0.02)		−0.002 (−0.004 to 0.00006)		0.02 (0.01 to 0.02)	
≥75	0.06 (−0.01 to 0.13)	.10	0.46 (0.23 to 0.69)	<.001	−0.01 (−0.10 to 0.07)	.78	0.61 (0.30 to 0.91)	<.001
California								
Sex								
Male	0.03 (−0.003 to 0.06)		0.02 (−0.01 to 0.04)		0.02 (−0.01 to 0.05)		0.01 (−0.01 to 0.04)	
Female	0.03 (0.003 to 0.06)	.79	0.04 (0.01 to 0.06)	.25	0.02 (−0.01 to 0.05)	.94	0.03 (0.01 to 0.05)	.27
Age, y								
0–74	0.02 (0.01 to 0.03)		0.004 (−0.004 to 0.01)		0.02 (0.01 to 0.03)		−0.0009 (−0.01 to 0.01)	
≥75	0.20 (−0.08 to 0.47)	.20	0.40 (0.19 to 0.61)	<.001	0.06 (−0.22 to 0.34)	.78	0.39 (0.17 to 0.60)	<.001
Central-southern Italy								
Sex								
Male	0.09 (0.03 to 0.14)		0.08 (0.01 to 0.16)		0.08 (0.01 to 0.14)		0.02 (−0.07 to 0.11)	
Female	0.12 (0.07 to 0.17)	.37	0.12 (0.05 to 0.19)	.50	0.11 (0.04 to 0.17)	.55	0.04 (−0.06 to 0.13)	.83
Age, y								
0–74	0.01 (−0.01 to 0.04)		0.03 (−0.0006 to 0.06)		0.002 (−0.02 to 0.03)		0.03 (−0.01 to 0.07)	
≥75	1.00 (0.65 to 1.35)	<.001	0.98 (0.45 to 1.51)	<.001	0.90 (0.46 to 1.33)	<.001	0.28 (−0.37 to 0.93)	.45
Germany								
Sex								
Male	0.02 (0.01 to 0.04)		0.05 (0.04 to 0.07)		0.004 (−0.01 to 0.02)		0.05 (0.03 to 0.07)	
Female	0.05 (0.03 to 0.06)	.03	0.05 (0.03 to 0.07)	.79	0.04 (0.02 to 0.05)	.01	0.03 (0.01 to 0.05)	.19
Age, y								
0–74	0.01 (0.01 to 0.02)		0.02 (0.01 to 0.02)		0.01 (0.001 to 0.01)		0.01 (0.01 to 0.02)	
≥75	0.21 (0.13 to 0.29)	<.001	0.31 (0.22 to 0.40)	<.001	0.11 (0.02 to 0.20)	.03	0.26 (0.17 to 0.36)	<.001

Abbreviations: NO₂, nitrogen dioxide; PM_{2.5}, fine particulate matter.

^a In subgroup analyses, we used the main lag day (lag01 to lag02 represent cumulative lag days) for each pollutant in each region (Jiangsu: PM_{2.5} lag01, NO₂ lag02; California: PM_{2.5} lag02, NO₂ lag02; central-southern Italy: PM_{2.5} lag02, NO₂ lag02; and Germany: PM_{2.5} lag01, NO₂ lag01).


^b Calculated per 100 000 population.

^c We tested the statistical differences in effect estimates between males and females and between those aged 0 to 74 years and those 75 years or older based on the z score calculated using the coefficients and standard errors for different groups.

applicability of the IFE approach across diverse regions, including China, the US, and Europe, encompassing varying levels of air pollution concentrations and socioeconomic statuses.

Our estimated associations between short-term changes in air pollution and changes in daily mortality rates in 4 countries are generally consistent with findings from previous studies using time-series analysis.¹⁻³ For example, in the multicountry and multicity studies,^{2,3} a 10- $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} concentration on lag01 or in NO₂ concentration on lag1 was found to be associated with a 0.68% or 0.46% increase in daily all-cause mortality, respectively. In a study based on 272 cities in China, a 10- $\mu\text{g}/\text{m}^3$ increase in the 2-day mean NO₂ concentration was associated with an increase of 0.9% in daily total nonaccidental mortality.¹ However, the estimates of our study cannot be directly compared with those from time-series analyses due to different interpretations of the model coefficients. The coefficient in our model estimates the change in the daily mortality rate for each 10- $\mu\text{g}/\text{m}^3$ change in the daily air pollution level; the coefficient does not directly provide information about the association between the absolute level of air pollution and the absolute level of mortality. Nevertheless, the consistency between the results from traditional time-series analyses and the novel IFE approach provides robust support for an association between short-term exposure to PM_{2.5} and NO₂ and mortality.

Figure 3. Estimated Change in Daily Mortality Rate Associated With a 10- $\mu\text{g}/\text{m}^3$ Increase in Fine Particulate Matter (PM_{2.5}) or Nitrogen Dioxide (NO₂) Concentration in Urban and Rural Areas



Single- and 2-pollutant models in urban and rural areas in each study region are shown. The error bars represent the 95% CIs. The lag with the strongest pollutant-mortality association for each pollutant in each region was used in the stratified analysis (Jiangsu: PM_{2.5} lag01, NO₂ lag02; California: PM_{2.5} lag02, NO₂ lag02; central-southern Italy: PM_{2.5} lag02, NO₂ lag02; Germany: PM_{2.5} lag01, NO₂ lag01). We tested the statistical differences in effect estimates between urban and rural areas by calculating *P* values based on the *z* score derived from the coefficients and SEs for different subgroups. *P* values were noted only for statistically significant between-group differences.

In addition, the results from our main IFE models were similar to those from traditional fixed-effects models, which do not take into account unmeasured time-varying effects at the spatial unit level (eTable 3 in [Supplement 1](#)). This suggests that even though we observed such unmeasured spatial unit-specific temporal confounding (as seen in the model for the main lags in Jiangsu and California) (eFigure 2 in [Supplement 1](#)), its role in the estimated air pollution-mortality association was minimal. This observation further bolsters our confidence in traditional time-series analysis. Despite its inherent limitations in adjusting for unmeasured temporal confounding, the results it produces remain valid.

Although short-term exposure to NO₂ has been linked to mortality and morbidity outcomes by numerous epidemiological studies,^{28,29} questions remain about the causal nature of this association.³⁰⁻³² As stated in the Integrated Science Assessment by the US Environmental Protection Agency, current evidence for short-term NO₂ exposure is "suggestive of, but not sufficient to infer, a causal relationship."³⁰ Because NO₂ and other combustion-derived air pollutants such as PM_{2.5} can be coemitted from traffic and other sources,^{33,34} it has been speculated that NO₂ only serves as a surrogate for other traffic-related air pollutants.³⁴⁻³⁶ This uncertainty not only calls into question the weight of epidemiological evidence on the health effects of NO₂ but also complicates policy making for air quality regulations.³⁴ In our study, the NO₂ association remained after adjustment for PM_{2.5} in the 2-pollutant model in all 4 study regions, suggesting that NO₂ may have its own independent effect on mortality that is not explained by PM_{2.5} levels.

We found substantial effect modification by age in all study regions. The effect modification of associations between short-term changes in air pollution and changes in daily mortality rate was significantly larger among people 75 years and older than among those aged 0 to 74 years. This association among older people reflects both the high mortality rate (Table 1) and the high susceptibility in the older population as found in previous studies.³⁷

This study encompasses regions from 4 distinct countries spanning different air pollution levels and diverse socioeconomic backgrounds. Substantial heterogeneity in the associations between short-term changes in air pollution and changes in daily mortality rate was observed across study regions. These heterogeneous results could be explained by different chemical compositions of particulate matter, population characteristics (such as age structure) affecting susceptibility, regional climate, socioeconomic factors, and other factors.³⁸ In addition, in central-southern Italy, the spatial unit was the municipality, which was much smaller than the counties used in other study regions. This resulted in fewer daily mortality counts in each spatial unit and, consequently, greater uncertainties in central-southern Italy. The difference in spatial scale also introduced different degrees of exposure measurement error; using larger spatial units such as counties might dilute the actual effects.

Limitations

This study has several limitations. First, using county- or municipality-level air pollution and mortality data, we were unable to capture variations within individual spatial units, which may be particularly pronounced in larger counties or municipalities. Second, the air pollution exposures in Jiangsu, central-southern Italy, and Germany were modeled and therefore subject to uncertainty. In California, our reliance on records from air quality monitoring stations did not yield full spatial coverage. Third, although the IFE model relaxed the assumptions of the traditional TWFE model, it has its own assumptions, and violation of these assumptions might bias the results. In addition, we assumed a linear association between air pollution and mortality, but this assumption may oversimplify the actual relationship. Studies with finer spatial resolution, wider spatial coverage, and more accurate exposure assessment are needed to further investigate the potentially nonlinear causal relationship between short-term air pollution exposure and health outcomes.

Conclusions

In this cross-sectional study, after controlling for both measured and unmeasured spatiotemporal confounders, we found that increases in short-term PM_{2.5} and NO₂ concentrations were associated with

increases in all-cause mortality rates. This study contributed to the growing body of evidence on the potentially detrimental health effects of short-term PM_{2.5} and NO₂ exposure. The IFE model can be used to estimate associations between changes in short-term exposure to air pollution and changes in daily health outcomes.

ARTICLE INFORMATION

Accepted for Publication: November 22, 2023.

Published: March 1, 2024. doi:10.1001/jamanetworkopen.2023.54607

Open Access: This is an open access article distributed under the terms of the [CC-BY License](#). © 2024 Ma Y et al. *JAMA Network Open*.

Corresponding Author: Kai Chen, PhD, Department of Environmental Health Sciences, Yale School of Public Health, 60 College St, New Haven, CT 06520 (kai.chen@yale.edu).

Author Affiliations: Department of Environmental Health Sciences, Yale School of Public Health, New Haven, Connecticut (Ma, Dubrow, Chen); Yale Center on Climate Change and Health, Yale School of Public Health, New Haven, Connecticut (Ma, Dubrow); Department of Epidemiology, Lazio Region Health Service ASL Roma 1, Rome, Italy (Nobile, Stafoggia); Chair of Epidemiology, Institute for Medical Information Processing, Biometry, and Epidemiology, Faculty of Medicine, Ludwig-Maximilians-Universität München, Munich, Germany (Marb, Breitner); Institute of Epidemiology, Helmholtz Zentrum München-German Research Center for Environmental Health, Neuherberg, Germany (Marb, Breitner); Department of Environmental Health, Boston University School of Public Health, Boston, Massachusetts (Kinney).

Author Contributions: Mss Ma and Marb had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Marb, Dubrow, Stafoggia, Breitner, Kinney, Chen.

Acquisition, analysis, or interpretation of data: Ma, Nobile, Marb, Dubrow, Breitner, Chen.

Drafting of the manuscript: Ma, Nobile, Marb.

Critical review of the manuscript for important intellectual content: Marb, Dubrow, Stafoggia, Breitner, Kinney, Chen.

Statistical analysis: Ma, Nobile, Marb, Stafoggia, Breitner, Chen.

Obtained funding: Marb, Breitner, Chen.

Administrative, technical, or material support: Nobile.

Supervision: Kinney, Chen.

Conflict of Interest Disclosures: Ms Ma reported receiving grant funding from the Health Effects Institute (HEI) during the conduct of the study. Ms Nobile reported receiving grant funding from the HEI during the conduct of the study. Ms Marb reported receiving grant funding from the HEI during the conduct of the study. Dr Dubrow reported receiving grant funding from the HEI during the conduct of the study. Dr Breitner reported receiving grant funding from the HEI during the conduct of the study. Dr Chen reported receiving grant funding from the HEI during the conduct of the study. No other disclosures were reported.

Funding/Support: This work was supported by grant 4984-RFA20-1B/21-7 from the HEI. The research described in this article was conducted under contract with the HEI, an organization jointly funded by award No. R-82811201 from the US Environmental Protection Agency (EPA) and certain motor vehicle and engine manufacturers.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Disclaimer: The contents of this article do not necessarily reflect the views of the HEI or its sponsors, nor do they necessarily reflect the views and policies of the EPA or motor vehicle and engine manufacturers.

Data Sharing Statement: See [Supplement 2](#).

REFERENCES

1. Chen R, Yin P, Meng X, et al. Associations between ambient nitrogen dioxide and daily cause-specific mortality: evidence from 272 Chinese cities. *Epidemiology*. 2018;29(4):482-489. doi:10.1097/EDE.0000000000000829
2. Meng X, Liu C, Chen R, et al. Short term associations of ambient nitrogen dioxide with daily total, cardiovascular, and respiratory mortality: multilocation analysis in 398 cities. *BMJ*. 2021;372(534):n534. doi:10.1136/bmj.n534

3. Liu C, Chen R, Sera F, et al. Ambient particulate air pollution and daily mortality in 652 cities. *N Engl J Med*. 2019;381(8):705-715. doi:10.1056/NEJMoa1817364
4. Lu F, Xu D, Cheng Y, et al. Systematic review and meta-analysis of the adverse health effects of ambient PM_{2.5} and PM₁₀ pollution in the Chinese population. *Environ Res*. 2015;136:196-204. doi:10.1016/j.envres.2014.06.029
5. Schwartz J, Bind MA, Koutrakis P. Estimating causal effects of local air pollution on daily deaths: effect of low levels. *Environ Health Perspect*. 2017;125(1):23-29. doi:10.1289/EHP232
6. Wei Y, Wang Y, Wu X, et al. Causal effects of air pollution on mortality rate in Massachusetts. *Am J Epidemiol*. 2020;189(11):1316-1323. doi:10.1093/aje/kwaa098
7. Guo Y, Wu Y, Ye T, Zhang L, Johnson A, Li S. A causal modelling framework for short-term effects of PM_{2.5} on hospitalisations: a nationwide time series study in Brazil. *Environ Int*. 2023;171:107688. doi:10.1016/j.envint.2022.107688
8. Cunningham S. *Causal Inference: The Mixtape*. Yale University Press; 2021.
9. Wang Y, Kloog I, Coull BA, Kosheleva A, Zanobetti A, Schwartz JD. Estimating causal effects of long-term PM_{2.5} exposure on mortality in New Jersey. *Environ Health Perspect*. 2016;124(8):1182-1188. doi:10.1289/ehp.1409671
10. Renzi M, Forastiere F, Schwartz J, Davoli M, Michelozzi P, Stafoggia M. Long-term PM₁₀ exposure and cause-specific mortality in the Latium Region (Italy): a difference-in-differences approach. *Environ Health Perspect*. 2019;127(6):67004. doi:10.1289/EHP3759
11. Schwartz J, Wei Y, Yitshak-Sade M, Di Q, Dominici F, Zanobetti A. A national difference in differences analysis of the effect of PM_{2.5} on annual death rates. *Environ Res*. 2021;194:110649. doi:10.1016/j.envres.2020.110649
12. Goodman-Bacon A. Difference-in-differences with variation in treatment timing. *J Econometrics*. 2021;225(2):254-277. doi:10.1016/j.jeconom.2021.03.014
13. Liu L, Wang Y, Xu Y. A practical guide to counterfactual estimators for causal inference with time-series cross-sectional data. *Am J Pol Sci*. Published online August 2, 2022. doi:10.1111/ajps.12723
14. Bai J. Panel data models with interactive fixed effects. *Econometrica*. 2009;77(4):1229-1279. doi:10.3982/ECTA6135
15. Xu Y. Generalized synthetic control method: causal inference with interactive fixed effects models. *Polit Anal*. 2017;25(1):57-76. doi:10.1017/pan.2016.2
16. Huang C, Hu J, Xue T, Xu H, Wang M. High-resolution spatiotemporal modeling for ambient PM_{2.5} exposure assessment in China from 2013 to 2019. *Environ Sci Technol*. 2021;55(3):2152-2162. doi:10.1021/acs.est.0c05815
17. Huang C, Sun K, Hu J, Xue T, Xu H, Wang M. Estimating 2013-2019 NO₂ exposure with high spatiotemporal resolution in China using an ensemble model. *Environ Pollut*. 2022;292(pt A):118285. doi:10.1016/j.envpol.2021.118285
18. Stafoggia M, Cattani G, Ancona C, Ranzi A. Exposure assessment of air pollution in Italy 2016-2019 for future studies on air pollution and COVID-19. Article in Italian. *Epidemiol Prev*. 2020;44(5-6)(suppl 2):161-168. doi:10.19191/EP20.5-6.S2.115
19. Stafoggia M, Bellander T, Bucci S, et al. Estimation of daily PM₁₀ and PM_{2.5} concentrations in Italy, 2013-2015, using a spatiotemporal land-use random-forest model. *Environ Int*. 2019;124:170-179. doi:10.1016/j.envint.2019.01.016
20. Felming J, Stern R. *Data Assimilation on the Basis of Optimal Interpolation for the Mapping of Immission Loads: Description of the Methodology and Practical Application for 2002*. Book in German. Institut für Meteorologie, Troposphärische Umweltforschung, Freie Universität Berlin; 2004.
21. German Environment Agency. Cartographic representation of the areal immission load in Germany by combination of measurements and calculations: 1999-2003. January 2006. Accessed November 1, 2021. <https://www.umweltbundesamt.de/publikationen/kartografische-darstellung-flaechenhaften>
22. United States Environmental Protection Agency. Outdoor air quality data. Updated July 5, 2023. Accessed June 1, 2021. <https://www.epa.gov/outdoor-air-quality-data/download-daily-data>
23. Muñoz Sabater J. ERA5-Land hourly data from 1950 to present. Climate Data Store. 2019. Accessed November 1, 2021. <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=overview>
24. Bai J, Ng S. Determining the number of factors in approximate factor models. *Econometrica*. 2002;70(1):191-221. doi:10.1111/1468-0262.00273
25. Bada O, Liebl D. phtt: Panel data analysis with heterogeneous time trends in R. *J Stat Softw*. 2014;59(6):1-33. doi:10.18637/jss.v059.i06
26. Chen K, Dubrow R, Breitner S, et al. Triggering of myocardial infarction by heat exposure is modified by medication intake. *Nat Cardiovasc Res*. 2022;1(8):727-731. doi:10.1038/s44161-022-00102-z

27. Benschaul-Tolonen A. Local industrial shocks and infant mortality. *Econ J (Lond)*. 2019;129(620):1561-1592.
28. Mills IC, Atkinson RW, Kang S, Walton H, Anderson HR. Quantitative systematic review of the associations between short-term exposure to nitrogen dioxide and mortality and hospital admissions. *BMJ Open*. 2015;5(5):e006946. doi:10.1136/bmjopen-2014-006946
29. Wang M, Li H, Huang S, et al. Short-term exposure to nitrogen dioxide and mortality: a systematic review and meta-analysis. *Environ Res*. 2021;202:111766. doi:10.1016/j.envres.2021.111766
30. US Environmental Protection Agency. Integrated science assessment (ISA) for oxides of nitrogen, oxides of sulfur and particulate matter ecological criteria (final report, 2020). EPA/600/R-20/278. Updated December 6, 2022. Accessed June 1, 2023. <https://assessments.epa.gov/risk/document/&deid=349473>
31. World Health Organization. *Review of Evidence on Health Aspects of Air Pollution: REVIHAAP Project: Technical Report*. WHO Regional Office for Europe; 2021.
32. Seaton A, Dennekamp M. Hypothesis: ill health associated with low concentrations of nitrogen dioxide—an effect of ultrafine particles? *Thorax*. 2003;58(12):1012-1015. doi:10.1136/thorax.58.12.1012
33. Brunekreef B, Strak M, Chen J, et al. Mortality and morbidity effects of long-term exposure to low-level PM_{2.5}, BC, NO₂, and O₃: an analysis of European cohorts in the ELAPSE Project. *Res Rep Health Eff Inst*. 2021;2021(208):1-127.
34. Forastiere F, Peters A. Invited perspective: the NO₂ and mortality dilemma solved? almost there! *Environ Health Perspect*. 2021;129(12):121304. doi:10.1289/EHP10286
35. Alotaibi R, Bechle M, Marshall JD, et al. Traffic related air pollution and the burden of childhood asthma in the contiguous United States in 2000 and 2010. *Environ Int*. 2019;127:858-867. doi:10.1016/j.envint.2019.03.041
36. Mills IC, Atkinson RW, Anderson HR, Maynard RL, Strachan DP. Distinguishing the associations between daily mortality and hospital admissions and nitrogen dioxide from those of particulate matter: a systematic review and meta-analysis. *BMJ Open*. 2016;6(7):e010751. doi:10.1136/bmjopen-2015-010751
37. Shumake KL, Sacks JD, Lee JS, Johns DO. Susceptibility of older adults to health effects induced by ambient air pollutants regulated by the European Union and the United States. *Aging Clin Exp Res*. 2013;25(1):3-8. doi:10.1007/s40520-013-0001-5
38. Samet JM. Air pollution risk estimates: determinants of heterogeneity. *J Toxicol Environ Health A*. 2008;71(9-10):578-582. doi:10.1080/15287390801997666

SUPPLEMENT 1.

eAppendix 1. Sources of Mortality and Population Data in Each Region

eAppendix 2. Air Pollution Spatiotemporal Models for Jiangsu, China, Central-Southern Italy, and Germany

eAppendix 3. Relative Humidity Calculation in Each Region

eAppendix 4. Selection of the Number of Common Time-Varying Factors in the Unmeasured Time-Varying Spatial Unit Effect in the Interactive Fixed-Effects Model

eAppendix 5. Sample R Code for the Main Interactive Fixed-Effects Model

eAppendix 6. Classification of Urban and Rural Areas in Each Study Region

eAppendix 7. Traditional 2-Stage Time-Series Analysis

eFigure 1. Randomization Test for the Main Lag in Each Study Region

eFigure 2. Plot of Estimated Common Time-Varying Factors and Unmeasured Time-Varying Spatial Unit Effects in the Main Single-Pollutant Models for PM_{2.5} or NO₂ for Jiangsu, China, and California

eFigure 3. Association Between Air Pollution and Mortality Estimated From Interactive Fixed-Effects Model and Traditional 2-Stage Time-Series Analysis

eTable 1. Estimated Change in Daily Mortality Rate (per 100 000 People) Associated With a 10-μg/m³ Increase in PM_{2.5} or NO₂ Concentration (95% CI) by Lag

eTable 2. Estimated Change in Daily Mortality Rate (per 100 000 People) Associated With a 10-μg/m³ Increase in PM_{2.5} or NO₂ Concentration by Urbanicity (95% CI)

eTable 3. Estimated Changes in Daily Mortality Rate (per 100 000 People) Associated With a 10-μg/m³ Increase in PM_{2.5} and NO₂ Concentrations in Sensitivity Analyses

eReferences.

SUPPLEMENT 2.

Data Sharing Statement