

## Exact many-electron ground states on diamond and triangle Hubbard chains

Zsolt Gulácsi, Arno P. Kampf, Dieter Vollhardt

### Angaben zur Veröffentlichung / Publication details:

Gulácsi, Zsolt, Arno P. Kampf, and Dieter Vollhardt. 2008. "Exact many-electron ground states on diamond and triangle Hubbard chains." *Progress of Theoretical Physics Supplement* 176: 1–21. <https://doi.org/10.1143/ptps.176.1>.

### Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

**Deutsches Urheberrecht**

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



## Exact Many-Electron Ground States on Diamond and Triangle Hubbard Chains

Zsolt GULÁCSI,<sup>1,2</sup> Arno KAMPF<sup>1</sup> and Dieter VOLLHARDT<sup>1</sup>

<sup>1</sup>*Theoretical Physics III, Center for Electronic Correlations and Magnetism,  
Institute of Physics, University of Augsburg, D-86135 Augsburg, Germany*

<sup>2</sup>*Department of Theoretical Physics, Faculty of Natural Sciences,  
University of Debrecen, H-4010 Debrecen, Hungary*

We construct exact ground states of interacting electrons on triangle and diamond Hubbard chains. The construction requires (i) a rewriting of the Hamiltonian into positive semidefinite form, (ii) the construction of a many-electron ground state of this Hamiltonian, and (iii) the proof of the uniqueness of the ground state. This approach works in any dimension, requires no integrability of the model, and only demands sufficiently many microscopic parameters in the Hamiltonian which have to fulfill certain relations. The scheme is first employed to construct exact ground state for the diamond Hubbard chain in a magnetic field. These ground states are found to exhibit a wide range of properties such as flat-band ferromagnetism and correlation induced metallic, half-metallic or insulating behavior, which can be tuned by changing the magnetic flux, local potentials, or electron density. Detailed proofs of the uniqueness of the ground states are presented. By the same technique exact ground states are constructed for triangle Hubbard chains and a one-dimensional periodic Anderson model with nearest-neighbor hybridization. They permit direct comparison with results obtained by variational techniques for  $f$ -electron ferromagnetism due to a flat band in  $\text{CeRh}_3\text{B}_2$ .

### §1. Introduction

Electronic correlations in solids can result in a high sensitivity of these systems to small changes in external parameters such as temperature, pressure, magnetic field, or doping.<sup>1)–3)</sup> This may then lead to strongly non-linear responses such as large resistivity and volume changes, a strong thermoelectric response, colossal magnetoresistance, and high- $T_c$  superconductivity. The complexity displayed by electronically correlated materials is not only the topic of fascinating fundamental research, but is also significant for technological applications, e.g., in sensors, switches, and in spintronics.

Even highly simplified models of interacting electrons are known to display a dazzling diversity of phases, e.g., metallic, insulating, and many types of long-range ordered states.<sup>4),5)</sup> Theoretical investigations of correlation effects almost always make some kind of approximation which cannot, in general, explain the full complexity of a model in detail. In this situation exact solutions of correlation models are particularly important and useful.

In this paper we discuss a general strategy for the construction of exact many-electron ground states of lattice models such as the Hubbard model. The approach is based on three distinct steps: (i) the rewriting of the Hamiltonian in positive semidefinite form, (ii) the construction of exact ground states for this Hamiltonian, and (iii) the proof of the uniqueness of the ground states constructed thereby.

This strategy was recently employed to construct a class of exact ground states of a three-dimensional periodic Anderson model (PAM), including the conventional PAM on regular Bravais lattices at and above  $3/4$  filling,<sup>6),7)</sup> and to the diamond Hubbard chain in a magnetic field.<sup>8)</sup> In both cases a wide range of features such as metallic, non-Fermi liquid, and insulating phases in the case of the PAM,<sup>6),7)</sup> and flat-band ferromagnetism, correlation-induced metallic, half-metallic, or insulating behavior in the case of the diamond Hubbard chain<sup>8)</sup> was found. These exact results prove that the phase diagram of even seemingly simple quantum mechanical models of many-particle systems can indeed be surprisingly complex.

The construction of exact ground states by casting the Hamiltonian into positive semidefinite form was originally applied by Brand and Gieseke to the Hubbard model and the PAM on special perovskite-type decorated lattices in dimensions  $d \geq 2$  at  $U = \infty$ .<sup>9)</sup> The procedure was generalized by Strack<sup>10)</sup> to the PAM and the extended Emery model on regular lattices in low dimensions,<sup>11)</sup> and to the extended Hubbard model. The method was subsequently used to derive rigorous criteria for the stability of saturated ferromagnetism in general Hubbard-type models.<sup>12)</sup> The possibility of superconductivity in the extended Hubbard model and its strong coupling limit was also investigated.<sup>13)–15)</sup> Furthermore, next-nearest neighbor terms in the Hamiltonian were shown to cause charge density waves, degenerate non-magnetic localized phases, and phase separation.<sup>16)</sup> Introducing operators to rewrite the Hamiltonian in positive semidefinite form which are defined on the entire unit cell of a  $d$ -dimensional lattice (e.g., on plaquettes in  $d = 2$  or cubes in  $d = 3$ ) rather than only on bonds, Orlik and Gulácsi<sup>17)</sup> investigated the PAM on regular two- and three-dimensional lattices at  $U = \infty$ , finding insulating non-magnetic phases. This method was used by Sarasua and Continentino<sup>18)</sup> to analyze the one-dimensional PAM in more detail. Gurin and Gulácsi<sup>19)</sup> and Gulácsi<sup>20)</sup> demonstrated that it is also possible to investigate the PAM in  $d = 2, 3$  at  $U < \infty$ . The exact ground states constructed by Gulácsi and Vollhardt for the PAM at  $3/4$ <sup>6)</sup> and  $1/4$ <sup>7)</sup> gave the first mathematical proof for the presence of ferromagnetism in this model in dimension  $d = 3$ . These authors also detected a strong anomaly in the compressibility of the system at the boundary of the parameter region where the non-magnetic localized phase is stable — a feature which is typical for  $f$  electron systems such as Cerium. In this connection the uniqueness of the constructed states was proved explicitly for the first time.

Most recently, the present authors constructed exact ground states of interacting electrons on the diamond Hubbard chain in a magnetic field.<sup>8)</sup> This model was shown to exhibit a wide range of properties such as flat-band ferromagnetism and correlation induced metallic, half-metallic or insulating behavior which can be tuned by changing the magnetic flux, local potentials, or the electron density.

In this paper we present details of the construction of exact many-electron ground states on the diamond Hubbard chain and of their physical properties.<sup>8)</sup> In particular, in Appendices A and B we provide details of the mathematical proof of the uniqueness of the ground-state solutions for this model. We also construct exact ground states on triangle Hubbard chains and show that within a particular subspace of parameters the triangle Hubbard chain may be mapped to a one-dimensional PAM.

The exact solutions obtained for this model can be applied to investigate the correlated electron material  $\text{CeRh}_3\text{B}_2$ , for which Kono and Kuramoto<sup>21)</sup> had constructed an approximate solution.

The paper is structured as follows: In §2 we discuss the general strategy for the construction of exact many-electron ground states. Exact ground states are constructed for the diamond Hubbard chain in §3. Details of the proof of the uniqueness of these ground states are presented in Appendices A and B. The technique is employed in §4 to obtain exact ground states also for the triangle Hubbard chain, and the results are applied to explain properties of  $\text{CeRh}_3\text{B}_2$ .

## §2. Construction of exact many-electron ground states: General strategy

Our construction of exact many-electron ground states is based on the following three steps.

### 2.1. Step 1: Rewriting the Hamiltonian in positive semidefinite form

In the first step the many-electron Hamiltonian is cast into positive semidefinite form. This means that one rewrites the Hamiltonian in terms of a number of positive semidefinite operators  $\hat{P}_n$  as

$$\hat{H} = \hat{H}_0 + \hat{H}_U = \sum_{n=1}^L \hat{P}_n + E_g \equiv \hat{H}' + E_g, \quad (2.1)$$

where  $E_g$  is the ground-state energy. Here (2.1) is supposed to be an operator identity, involving no approximations. The positive semidefinite nature of  $\hat{P}_n$  is expressed by the relation  $\langle \Psi | \hat{P}_n | \Psi \rangle \geq 0$ , which must be satisfied for all  $|\Psi\rangle$ . Due to this inequality, the minimal eigenvalue of  $\hat{P}_n$  is zero. This property, i.e., the existence of a well-defined lower bound, namely zero, in the spectrum of  $\hat{H}' = \hat{H} - E_g$  in (2.1) is the corner stone of the procedure. Simple examples for a positive semidefinite operator are  $\hat{P}_n = \hat{\Omega}^\dagger \hat{\Omega}$ , or  $\hat{P}_n = \hat{\Omega} \hat{\Omega}^\dagger$ , or  $\hat{P} = \sum_i \hat{P}_i$ , where the positive semidefinite operator  $\hat{P}_i = \hat{n}_{i,\uparrow} \hat{n}_{i,\downarrow} - (\hat{n}_{i,\uparrow} + \hat{n}_{i,\downarrow}) + 1$  assumes its minimum eigenvalue if at least one electron is present on the site  $i$ .

The scalar  $E_g$  in (2.1), i.e., the ground state energy, is usually a complicated function of the parameters entering in  $\hat{H}$ . It should be noted that the decomposition of  $\hat{H}$  in positive semidefinite form is, in general, not unique and can be obtained in different ways. Each transformation is valid only in certain parts  $\mathcal{R}_p$  of the parameter space.

### 2.2. Step 2: Construction of the exact many-electron ground states

After the transformation of a given Hamiltonian  $\hat{H}$  into a particular positive semidefinite form valid in a parameter space region  $\mathcal{R}_p$ , the exact ground state of  $\hat{H}$  needs to be constructed. Since  $\hat{H}' = \sum_{n=1}^L \hat{P}_n$ , and the minimal eigenvalue of each  $\hat{P}_n$  is zero, this requires the construction of the most general state  $|\Psi_g\rangle$  which

satisfies

$$\hat{P}_n |\Psi_g\rangle = 0, \quad (2.2)$$

for all  $n$ . Consequently,

$$\hat{H} |\Psi_g\rangle = E_g |\Psi_g\rangle \quad (2.3)$$

holds, where  $|\Psi_g\rangle$  is the ground state, with  $E_g$  as the ground state energy in  $\mathcal{R}_p$ . The actual construction of  $|\Psi_g\rangle$  depends on the explicit expression of the operators  $\hat{P}_n$  in (2.1). Therefore each case must be individually analyzed. Nevertheless the following guideline may be applied (here we restrict ourselves to fermionic particles):

i) For operators  $\hat{P}_n = \hat{\Omega}^\dagger \hat{\Omega}$  the ground state has, in general, the form  $|\phi_g\rangle = \hat{\Omega}^\dagger |0\rangle$ , where  $|0\rangle$  is the vacuum, and the anti-commutation relation  $\{\hat{\Omega}, \hat{\Omega}^\dagger\} = 0$  holds. Indeed, in this case one has  $\hat{P}_n |\phi_g\rangle = 0$ . To construct the global ground state one then tries to extend this property to all operators  $\hat{P}_n$ .

ii) Likewise, for operators  $\hat{P}_n = \hat{\Omega} \hat{\Omega}^\dagger$  the ground state construction makes use of  $(\hat{\Omega}^\dagger)^2 = 0$  and starts from states of the form  $|\psi_g\rangle = \hat{\Omega}^\dagger |0\rangle$ , since in this case  $\hat{P}_n |\psi_g\rangle = 0$ . To obtain  $|\Psi_g\rangle$  from  $|\psi_g\rangle$  the property  $\hat{P}_n |\psi_g\rangle = 0$ , valid for a particular  $n$ , must be extended to all operators  $\hat{P}_n$ .

### 2.3. Step 3: Proof of the uniqueness of the ground states

Finally the constructed ground state has to be shown to be unique, i.e., one must prove that there exist no other linearly independent eigenstates of  $\hat{H}$  in  $\mathcal{R}_p$  with the same energy  $E_g$ . This step employs the property of the *kernel* of  $\hat{H}'$  which is defined as the Hilbert subspace of states  $|\Phi\rangle$  with  $\hat{H}' |\Phi\rangle = 0$ , e.g.

$$\ker(\hat{H}') := \{ |\Phi\rangle \mid \hat{H}' |\Phi\rangle = 0 \}. \quad (2.4)$$

Since  $\hat{H}' = \sum_{n=1}^L \hat{P}_n$ , one has  $\ker(\hat{H}') = \bigcap_{n=1}^L \ker(\hat{P}_n)$ . To prove the uniqueness one must show that  $|\Psi_g\rangle$  spans  $\ker(\hat{H}')$ . For this it is necessary and sufficient to prove that

- (i)  $|\Psi_g\rangle$  is an element of the intersection of the kernels of all positive semidefinite operators  $\hat{P}_n$ , i.e.,  $|\Psi_g\rangle \in \ker(\hat{H}') = \bigcap_{n=1}^L \ker(\hat{P}_n)$ , and that
- (ii) all states  $|\Psi\rangle \in \bigcap_{n=1}^L \ker(\hat{P}_n)$  can be written in terms of the constructed ground state  $|\Psi_g\rangle$ .

We note that if condition (ii) is not satisfied the constructed ground state is not unique, implying that other, linearly independent states with the same lowest energy  $E_g$  exist. The uniqueness of the ground states constructed in this paper is proved in Appendices A and B.

### 2.4. Discussion

The procedure described above works, in principle, in any dimension and for any model, independent of the concept of integrability. On the other hand, a decomposition of  $\hat{H}$  into positive semidefinite form and a subsequent construction of the ground state may not always be possible — especially for Hamiltonians with a very

simple structure. Indeed, the procedure works the better the more microscopic parameters (e.g., hopping amplitudes, bands, etc.) enter in  $\hat{H}$ . This is due to the fact that in a high-dimensional parameter space it is easier to find a subspace  $\mathcal{R}_p$  where a relation between microscopic parameters is found, for which the transformation of the Hamiltonian into positive semidefinite form holds.

### §3. Exact many-electron ground states on diamond Hubbard chains

The first exact many-electron ground states for the diamond Hubbard chain in an external magnetic field have been deduced by Gulácsi, Kampf, and Vollhardt.<sup>8)</sup> Here we provide further details of the construction process and the physical properties of this system.

The diamond Hubbard chain is shown in Fig. 1. This chain has 3 sites per unit cell, hence 3 sublattices, denoted by the sublattice index  $s = 1, 2, 3$ , and therefore 3 bands. One denotes by  $N, N_c$ , and  $n = N/(3N_c)$  the number of electrons, cells, and electron density, respectively. The Hamiltonian of the chain  $\hat{H} = \hat{H}_0 + \hat{H}_U$  is given by

$$\hat{H}_0 = \sum_{\sigma} \sum_{i=1}^{N_c} \left\{ [t e^{i\frac{\delta}{2}} (\hat{c}_{i+r_2, \sigma}^{\dagger} \hat{c}_{i, \sigma} + \hat{c}_{i+a, \sigma}^{\dagger} \hat{c}_{i+r_2, \sigma} + \hat{c}_{i+r_1, \sigma}^{\dagger} \hat{c}_{i+a, \sigma} + \hat{c}_{i, \sigma}^{\dagger} \hat{c}_{i+r_1, \sigma}) + t_{\perp} \hat{c}_{i+r_2, \sigma}^{\dagger} \hat{c}_{i+r_1, \sigma} + t_{\parallel} \hat{c}_{i+a, \sigma}^{\dagger} \hat{c}_{i, \sigma} + \text{H.c.}] + \varepsilon \sum_{s=1,2} \hat{n}_{i+r_s, \sigma} \right\}, \quad (3.1)$$

$$\hat{H}_U = U \sum_{i=1}^{N_c} \sum_{s=1}^3 \hat{n}_{i+r_s, \uparrow} \hat{n}_{i+r_s, \downarrow}. \quad (3.2)$$

In the external magnetic field the hopping matrix elements acquire the Peierls phase factor with  $\delta = 2\pi\Phi/\Phi_0$  and  $\Phi_0 = hc/e$  is the flux quantum. Here we have chosen the vector potential  $\mathbf{A} \parallel \mathbf{a}$ , and the field dependent hopping amplitudes follow from  $t_{j,j'}(\mathbf{B}) = t_{j,j'}(0) \exp[(2i\pi/\Phi_0) \int_j^{j'} \mathbf{A} \cdot d\mathbf{l}]$ . For one flux quantum per unit cell (triangle)  $\delta = \pi$ . The Zeeman term is not explicitly included in  $\hat{H}$ , but all deduced ground states have also been analyzed




Fig. 1. Diamond Hubbard chain; the hopping matrix elements on different bonds are indicated, as well as the sites on the sublattices  $s = 1$  and  $2$  with potential  $\varepsilon$ .




Fig. 2. Schematic view of three particular bare band structures for (a)  $t_{\perp} = t_{\parallel} = 0$ ,  $\delta = \pi/2$ , (b)  $t_{\parallel} > 0$ ,  $\varepsilon = -t_{\perp} + t_{\perp}^{-1}/2$ ,  $\delta = \pi$ , and (c)  $t_{\parallel} = 0$ ,  $\varepsilon t_{\perp} \cos \delta = t_{\perp}^2 - \cos^2 \delta$ .

in the presence of the Zeeman term.

Fourier transforming  $\hat{H}_0$  one finds  $\hat{H}_0 = \sum_{\mathbf{k}, \sigma} \sum_{s,s'=1}^3 M_{s,s'}(\mathbf{k}) \hat{c}_{s,\mathbf{k},\sigma}^\dagger \hat{c}_{s',\mathbf{k},\sigma}$ , where the matrix elements  $M_{s,s'}$ , which determine the free-electron dispersion, are given in Ref. 8). In the following all energies will be given in units of  $2t$ . Examples for bare band structures are schematically shown in Fig. 2.

### 3.1. Flat-band ferromagnetism




Fig. 3. Diamond Hubbard chain at  $t_\perp = t_\parallel = 0$ . The thin line depicts the localized eigenstate corresponding to  $\hat{C}_{3,i,\sigma}^\dagger$ .

We first consider the parameters  $t_\perp = t_\parallel = 0, \delta = \frac{\pi}{2}$ , in which case the single-electron band structure consists of three flat bands (see Fig. 2(a)). This corresponds to the “Aharonov-Bohm cage” limit analyzed by Vidal et al.<sup>22)</sup> for the two electron problem with  $\varepsilon = 0$  (see also Fig. 3). Based on their finding that excited singlet eigenstates are localized for  $U = 0$  but delocalized for  $U > 0$ , Vidal et al. conjectured that such delocalized states should emerge

also at finite electron densities. Below we deduce indeed exact ground states for the diamond Hubbard chain at finite densities.

Following the strategy outlined in §2 one first transforms the kinetic part of the Hamiltonian into positive semidefinite form by introducing new canonical fermionic operators  $\hat{C}_{\nu,i,\sigma}$  in real space (for details see Ref. 8)); these operators represent localized Wannier eigenstates (see Fig. 3). With the energies  $E_2 = \varepsilon$ ,  $E_{2\pm 1} = (\varepsilon \mp \sqrt{\varepsilon^2 + 4})/2$  one obtains

$$\hat{H}_0 = \sum_{\sigma, i=1}^{N_c} \sum_{\nu=1}^3 E_\nu \hat{C}_{\nu,i,\sigma}^\dagger \hat{C}_{\nu,i,\sigma}. \quad (3.3)$$

Since  $\hat{C}_{\nu,i,\sigma}^\dagger \hat{C}_{\nu,i,\sigma}$  and  $\hat{H}_U$  are positive semidefinite operators the ground state of  $\hat{H}$  for  $N_c \geq N$ ,  $U > 0$  becomes

$$|\Psi_g^I(N)\rangle = \prod_{i=1}^N \hat{C}_{3,i,\sigma_i}^\dagger |0\rangle, \quad E_g^I = E_3 N. \quad (3.4)$$

For  $n = 1/3$  one obtains a fully saturated ferromagnetic ground state in the lowest band (see Fig. 4) which represents an explicit realization of Mielke-Tasaki’s flat band ferromagnetism.<sup>23)</sup> The strategy for the proof of the uniqueness of this ground state is discussed in Appendix A. For  $n < 1/3$  only the Wannier states with a spatial overlap have the same spin, the (highly degenerate) ground state hence consists of ferromagnetic clusters of arbitrary spin orientation. Since  $\hat{H}_U |\Psi_g^I(N)\rangle = 0$  and the kinetic part of  $\hat{H}$  is diagonal in real space, the ground state (3.4) is localized. The explicit proof of the uniqueness of (3.4) for  $n < 1/3$  is presented in Appendix B.

The flat-band ferromagnetism found here realizes early ideas of Gutzwiller<sup>24)</sup> and Kanamori<sup>25)</sup> from 1963 about the origin of ferromagnetism. They argued that the ferromagnetic orientation of the electronic spins is due to the fact that the local interaction is thereby completely suppressed thus lowering the energy greatly. However, they were not able to compute the associated loss in kinetic energy of this highly correlated state. In the present case of flat-band ferromagnetism the latter effect poses no problem since the kinetic energy is zero anyway.

We note that at  $U > 0$  the lowest band is flat only for  $\delta = \pi/2$ , but is dispersive for  $\delta = \pi$ , or  $\delta = 0$ . In the latter situation the system is most probably conducting for incommensurate fillings. Hence one encounters here a magnetic field induced metal-insulator transition.

### 3.2. Correlated half-metal

The previous results suggest that itinerant states are easier to realize at  $\delta \neq \pi/2$ . To analyze this problem we investigate below the special case  $t_{\parallel} > 0$ ,  $\epsilon = -t_{\perp} + t_{\perp}^{-1}/2$ , and  $\delta = \pi$ ; the single-electron band structure has a dispersive lowest band and two upper flat bands (see Fig. 2(b)).

#### 3.2.1. Transformation of $\hat{H}$ into positive semidefinite form

For the rewriting of  $\hat{H}_0$  one defines first the non-canonical fermionic operators

$$\hat{A}_{i,\sigma} = a_1 \hat{c}_{i,\sigma} + a_2 \hat{c}_{i+r_2,\sigma} + a_3 \hat{c}_{i+a,\sigma} + a_4 \hat{c}_{i+r_1,\sigma}, \quad (3.5)$$

which fulfill  $(\hat{A}_{i,\sigma})^2 = 0$  and  $\{\hat{A}_{i,\sigma}, \hat{A}_{j,\sigma}^{\dagger}\} \neq \delta_{i,j}$ . Based on (3.5) one obtains

$$\begin{aligned} \hat{A}_{i,\sigma}^{\dagger} \hat{A}_{i,\sigma} &= (a_2^* a_1 \hat{c}_{i+r_2,\sigma}^{\dagger} \hat{c}_{i,\sigma} + a_3^* a_2 \hat{c}_{i+a,\sigma}^{\dagger} \hat{c}_{i+r_2,\sigma} + a_4^* a_3 \hat{c}_{i+r_1,\sigma}^{\dagger} \hat{c}_{i+a,\sigma} + a_1^* a_4 \hat{c}_{i,\sigma}^{\dagger} \hat{c}_{i+r_1,\sigma} \\ &\quad + a_2^* a_4 \hat{c}_{i+r_2,\sigma}^{\dagger} \hat{c}_{i+r_1,\sigma} + a_3^* a_1 \hat{c}_{i+a,\sigma}^{\dagger} \hat{c}_{i,\sigma} + \text{H.c.}) \\ &\quad + |a_1|^2 \hat{n}_{i,\sigma} + |a_2|^2 \hat{n}_{i+r_2,\sigma} + |a_3|^2 \hat{n}_{i+a,\sigma} + |a_4|^2 \hat{n}_{i+r_1,\sigma}. \end{aligned} \quad (3.6)$$

With the requirement  $-\sum_{i,\sigma} \hat{A}_{i,\sigma}^{\dagger} \hat{A}_{i,\sigma} = \hat{H}_0$  one arrives at the matching conditions

$$\begin{aligned} a_2^* a_1 &= a_3^* a_2 = a_4^* a_3 = a_1^* a_4 = -t e^{i\frac{\delta}{2}}, & a_2^* a_4 &= -t_{\perp}, & a_3^* a_1 &= -t_{\parallel}, \\ |a_1|^2 + |a_3|^2 &= \epsilon + |a_2|^2 = \epsilon + |a_4|^2, \end{aligned} \quad (3.7)$$

from which

$$\hat{A}_{i,\sigma} = \sqrt{t_{\parallel}} [\hat{c}_{i,\sigma} - \hat{c}_{i+a,\sigma} - 2t_{\perp} e^{i\frac{\delta}{2}} (\hat{c}_{i+r_1,\sigma} - \hat{c}_{i+r_2,\sigma})] \quad (3.8)$$




Fig. 4. The ferromagnetic flat-band solution obtained at  $t_{\perp} = t_{\parallel} = 0$ ; the ground state result does not provide information about higher lying states as indicated by the question mark.



follows. Based on (3.6)–(3.8),  $\hat{H}_0$  is rewritten as

$$\hat{H}_0 = - \sum_{\mathbf{i}, \sigma} \hat{A}_{\mathbf{i}, \sigma}^\dagger \hat{A}_{\mathbf{i}, \sigma} = + \sum_{\mathbf{i}, \sigma} \hat{A}_{\mathbf{i}, \sigma} \hat{A}_{\mathbf{i}, \sigma}^\dagger - 2N_c \sum_{m=1}^4 |a_m|^2. \quad (3.9)$$

For the interaction part  $\hat{H}_U$ , one uses

$$\hat{H}_U = U \sum_{\mathbf{i}}^{3N_c} \hat{n}_{\mathbf{i}, \uparrow} \hat{n}_{\mathbf{i}, \downarrow} = U \hat{P} + U \hat{N} - U N_c, \quad \hat{P} = \sum_{\mathbf{i}} \hat{P}_{\mathbf{i}}. \quad (3.10)$$

$\hat{N}$  is the particle number operator, and  $\hat{P}_{\mathbf{i}} = (\hat{n}_{\mathbf{i}, \uparrow} - 1)(\hat{n}_{\mathbf{i}, \downarrow} - 1)$  is a positive semidefinite operator, which gives zero for at least one electron on site  $\mathbf{i}$  and 1 for an unoccupied site  $\mathbf{i}$ . Consequently,  $\hat{H}$  has been transformed into the positive semidefinite form

$$\hat{H} = \sum_{\mathbf{i}, \sigma} \hat{A}_{\mathbf{i}, \sigma} \hat{A}_{\mathbf{i}, \sigma}^\dagger + U \hat{P} + E_g^{II}, \quad (3.11)$$

where  $E_g^{II} = (\epsilon + U + t_\perp)N - N_c(3U + 4t_\perp + 1/t_\perp)$ .

### 3.2.2. Construction of the ground state

As discussed in §2.2, the ground state  $|\Psi_g\rangle$  for  $U > 0$  is deduced such to satisfy  $\hat{A}_{\mathbf{i}, \sigma}^\dagger |\Psi_g\rangle = 0$  and  $\hat{P} |\Psi_g\rangle = 0$ , from where indeed

$$\hat{H} |\Psi_g\rangle = E_g |\Psi_g\rangle \quad (3.12)$$

follows. Since  $(\hat{A}_{\mathbf{i}, \sigma}^\dagger)^2 = 0$  one finds

$$|\Psi_g^{II}(4N_c)\rangle \propto \prod_{\mathbf{i}} \hat{A}_{\mathbf{i}, -\sigma}^\dagger \hat{A}_{\mathbf{i}, \sigma}^\dagger |0\rangle. \quad (3.13)$$

In order to satisfy  $\hat{P} |\Psi_g\rangle = 0$  as well, which requires one electron on each site, besides the contribution from (3.13) one has to introduce in the ground state wavefunction also the operator

$$\hat{F}_\sigma^\dagger = \prod_{\mathbf{i}} [\hat{c}_{\mathbf{i}+\mathbf{r}_{s_{i,1}}, \sigma}^\dagger \hat{c}_{\mathbf{i}+\mathbf{r}_{s_{i,2}}, \sigma}^\dagger], \quad (3.14)$$

which creates two electrons with fixed spin  $\sigma$  on arbitrary sites of each unit cell. Hence, the unnormalized ground state becomes

$$|\Psi_g^{II}(4N_c)\rangle = c \left[ \prod_{\mathbf{i}} \hat{A}_{\mathbf{i}, -\sigma}^\dagger \hat{A}_{\mathbf{i}, \sigma}^\dagger \right] \hat{F}_\sigma^\dagger |0\rangle, \quad (3.15)$$

where  $c$  is a normalization constant. The contribution in (3.13) creates one  $\sigma$  electron, and an electron with spin  $-\sigma$  in each unit cell, while  $\hat{F}_\sigma^\dagger$  creates two electrons with spin  $\sigma$  on arbitrary sites of each unit cell. Consequently the ground state in

(3.15) corresponds to  $N = 4N_c$ ,  $n = 4/3$ ,  $n_\sigma = N_\sigma/3N_c = 1$ ,  $n_{-\sigma} = 1/3$ . The proof of the uniqueness of (3.15) is presented in detail in Appendix A. In conclusion, (3.15) is the unique ground state.

Concerning the physical properties of the ground state one observes that in (3.15) each lattice site is occupied by one  $\sigma$ -electron; hence, the  $\sigma$ -electrons are localized. The  $-\sigma$  electrons are spatially extended, but localized for  $N_c \rightarrow \infty$ . This is inferred from the ground-state expectation value of the hopping term  $\Gamma_{\mathbf{r},-\sigma} = \langle \hat{c}_{\mathbf{j},-\sigma}^\dagger \hat{c}_{\mathbf{j}+\mathbf{r},-\sigma} + \text{H.c.} \rangle$ , which in the limit  $N_c \rightarrow \infty$ , for  $r/a = m$ , is calculated as

$$\Gamma_{m,-\sigma} = \frac{(-1)^m}{\sqrt{1+1/t_\perp}} e^{-\frac{m}{\xi_{-\sigma}}}. \quad (3.16)$$

The one-particle localization length  $\xi_{-\sigma}$  in (3.16) increases almost linearly with  $1/t_\perp$ .<sup>8)</sup>

### 3.2.3. Solution for $n > 4/3$

For electron densities  $n > 4/3$  there are  $N > 4N_c$  electrons. No  $\sigma$ -electrons but  $\Delta N$   $-\sigma$ -electrons can be added to the system such that  $n_\sigma = 1$ ,  $n_{-\sigma} = 1/3 + \Delta N/N_c$ . The ground state then becomes

$$|\Psi_g^{II}(4N_c + \Delta N)\rangle = \prod_{\alpha=1}^{\Delta N} \hat{c}_{n_\alpha, \mathbf{k}_\alpha, -\sigma}^\dagger |\Psi_g^{II}(4N_c)\rangle, \quad (3.17)$$

where  $n_\alpha$  can be arbitrarily chosen from  $s = 1, 2, 3$ . Equation (3.17) contains plane wave-type contributions from the  $-\sigma$ -electrons. Consequently, the additional  $\Delta N$   $-\sigma$  electrons are itinerant. Indeed, this follows explicit from the expectation value of the hopping term  $\Gamma_{\mathbf{r},-\sigma}$ . For  $|\mathbf{r}/a| \gg 1$ ,  $\Delta N = 1$ ,  $s = 1$ , one finds  $\Gamma_{\mathbf{r},-\sigma}/\Gamma_{\mathbf{r},-\sigma}^{(1)} = 1 - [2 + (1 - \cos ak)/t_\perp]^{-1}$ . Here  $\Gamma_{\mathbf{r},-\sigma}^{(1)}$  is the plane wave result and  $k$  is the norm of the momentum of the electron added above  $n = 4/3$ . We note that the momenta  $\mathbf{k}_\alpha$  in (3.17) are arbitrary. Hence it is not possible to define a Fermi surface, and the ground state therefore corresponds to a non-Fermi liquid.

In the density range  $4/3 < n < 5/3$  the ground state has  $3N_c$  immobile  $\sigma$  electrons and  $N_c$  electrons with spin  $-\sigma$  confined to their localized Wannier functions. For this reason the ground state has a high geometric degeneracy; its uniqueness is discussed in Appendix B.

Only the  $\Delta N$  conducting  $-\sigma$  electrons are itinerant, leading to metallic behavior with a low carrier density and a low spin polarization. Since the conduction through this correlated half-metal involves only electrons of one spin species, such a system may serve as a spin-valve device. Since  $U > 0$  is required for this property one encounters here a correlation induced localization-delocalization transition to a half-metal. The magnetization behaves as  $M \propto (1 - \Delta N/N_c) \rightarrow 0$  for  $\Delta N \rightarrow N_c$  in a finite magnetic field. The transition can also be induced by tuning the local potential  $\varepsilon$ . At  $n = 5/3$  the ground state becomes nonmagnetic and localized in the thermodynamic limit, while at  $n > 5/3$  one finds a nonmagnetic but conducting ground state.

### 3.3. Exact ground states for general magnetic flux

It is also possible to construct exact ground states for general magnetic flux values including also the  $\mathbf{B} = \mathbf{0}$  case, for example at  $\delta \in (-\pi/2, \pi/2)$  with  $t_{\parallel} = 0$ ,  $t_{\perp} < 0$ ,  $b \equiv -\cos \delta/t_{\perp}$ ,  $\varepsilon = b - b^{-1}$ , and  $n \geq 5/3$ ; <sup>8)</sup> the single-electron band structure has two dispersive bands and an upper flat band (see Fig. 2(c)). For these hopping and potential parameters one obtains a localized nonmagnetic ground state for  $\delta = 0$  over an extended range of densities  $n \geq 5/3$ . By contrast, at finite magnetic fields the ground state is a nonsaturated ferromagnet. This localized state at  $n = 5/3$  is insulating, but gapless and itinerant for  $n > 5/3$ . For the latter densities the majority spin  $\sigma$ -electrons are immobile and only the  $-\sigma$  electrons are itinerant. Therefore, by varying the external magnetic field and the sublattice potential one can tune from a localized, nonmagnetic ground state to a nonsaturated ferromagnet in the density range  $n \geq 5/3$ ; the ferromagnetic state is insulating at  $n = 5/3$ , but conducting for  $n > 5/3$ .

We finally note the properties derived above do not depend on the Zeeman coupling. Only the solution in §3.1 for  $n < 1/3$  is altered by the Zeeman term, whose presence leads to a fully spin aligned ferromagnetic solution.

## §4. Exact many-electron ground states on triangle Hubbard chains

The procedure for the construction of exact ground states on the diamond Hubbard chain is now applied to a Hubbard chain composed of corner-sharing triangles (see Fig. 5). Triangles are well-known to frustrate non-ferromagnetic magnetic order. Hence interacting electrons on lattices with triangles as structural sub-units are expected to favor ferromagnetic ground states.




Fig. 5. Triangle Hubbard chain.

The triangle Hubbard chain in Fig. 5 has two sites per cell which leads to two energy bands. The number of electrons, cells, and electron density are here denoted by  $N$ ,  $N_c$  and  $n = N/(2N_c)$ , respectively. This model was previously investigated in detail by Penc et al.,<sup>26)</sup> Fazekas et al.,<sup>27)</sup> Müller-Hartmann<sup>28)</sup> and Derzhko et al.<sup>29)</sup> with

a variety of analytical and numerical techniques.

The Hamiltonian of the triangle chain  $\hat{H}^{tri} = \hat{H}_0^{tri} + \hat{H}_U^{tri}$  has the form

$$\begin{aligned} \hat{H}_0^{tri} = \sum_{\sigma} \sum_{i=1}^{N_c} [ & (t \hat{c}_{i,\sigma}^{\dagger} \hat{c}_{i+a,\sigma} + t' (\hat{c}_{i,\sigma}^{\dagger} \hat{c}_{i+r,\sigma} + \hat{c}_{i+a,\sigma}^{\dagger} \hat{c}_{i+r,\sigma}) + \text{H.c.}) \\ & + \varepsilon_0 \hat{c}_{i,\sigma}^{\dagger} \hat{c}_{i,\sigma} + \varepsilon_1 \hat{c}_{i+r,\sigma}^{\dagger} \hat{c}_{i+r,\sigma} ], \\ \hat{H}_U^{tri} = \sum_i [ & U_0 \hat{n}_{i,\uparrow} \hat{n}_{i,\downarrow} + U_1 \hat{n}_{i+r,\uparrow} \hat{n}_{i+r,\downarrow} ]. \end{aligned} \quad (4.1)$$

The model parameters are explained in Fig. 5. In the following we will construct

exact ground states in the subspace of parameters defined by the relation  $(t')^2/t = \varepsilon_1 - \varepsilon_0 + 2t$ ,  $\varepsilon_1 - \varepsilon_0 > -2t$ ,  $t > 0$ , for which the lowest bare band is flat.

#### 4.1. Ground states for the triangle Hubbard chain

##### 4.1.1. Ground state for $U_0, U_1 > 0$

Employing the operators

$$\hat{A}_{\mathbf{i},\sigma} = \sqrt{t}[\hat{c}_{\mathbf{i},\sigma} + \hat{c}_{\mathbf{i}+\mathbf{a},\sigma} + (t'/t)\hat{c}_{\mathbf{i}+\mathbf{r},\sigma}], \quad (4.2)$$

the Hamiltonian (4.1) can be transformed into positive semidefinite form:

$$\hat{H}^{tri} = \sum_{\mathbf{i},\sigma} \hat{A}_{\mathbf{i},\sigma}^\dagger \hat{A}_{\mathbf{i},\sigma} + K\hat{N} + H_U^{tri}. \quad (4.3)$$

Here  $\hat{N}$  is the particle number operator and  $K = \varepsilon_0 - 2t$ . To construct the ground state one has to find operators  $\hat{B}_{\mathbf{i},\sigma}^\dagger$  which satisfy  $\{\hat{A}_{\mathbf{i},\sigma}, \hat{B}_{\mathbf{i}',\sigma'}^\dagger\} = 0$  for all values of the indices. The ground state then has the form

$$|\Psi^{tri}(N)\rangle = \prod_{\mathbf{i}}^N \hat{B}_{\mathbf{i},\sigma_i}^\dagger |0\rangle, \quad (4.4)$$

with  $\hat{B}_{\mathbf{i},\sigma}^\dagger = [\hat{c}_{\mathbf{i}-\mathbf{a}+\mathbf{r},\sigma}^\dagger + \hat{c}_{\mathbf{i}+\mathbf{r},\sigma}^\dagger - (t'/t)\hat{c}_{\mathbf{i},\sigma}^\dagger]$ ,  $N \leq N_c$ . For  $n < 1/2$  there exist ferromagnetic clusters, and only operators  $\hat{B}_{\mathbf{i},\sigma_i}^\dagger$  which touch have the same spin index. For  $n = 1/2$  all clusters touch such that a fully saturated ferromagnetic state is obtained as shown in Fig. 6(a) for the  $U_1 > 0$  case. The ground state corresponds to a particular realization of flat-band ferromagnetism,<sup>23)</sup> for which Derzhko et al.<sup>29)</sup> calculated the thermodynamic properties.




Fig. 6. Ground-state configuration of the electrons for (a)  $U_0, U_1 > 0$ , and (b)  $U_0 > 0, U_1 = 0$ . The ground-state results do not provide information about higher lying states as indicated by the question marks.

#### 4.1.2. Ground state for $U_0 > 0, U_1 = 0$

The ground state has the same form as (4.4), but now it is *non*-magnetic even for  $n = 1/2$  (see Fig. 6(b)). This is due to the fact that, although the connectivity condition is fulfilled (i.e.,  $\hat{B}_{\mathbf{i},\sigma}^\dagger$  and  $\hat{B}_{\mathbf{i}+\mathbf{a},\sigma}^\dagger$  overlap at site  $\mathbf{i}+\mathbf{r}$ ) there is no interaction at the connectivity point ( $U_1 = 0$ ). The fulfillment of the connectivity condition for the Wannier functions is therefore not sufficient for the ground state to be ferromagnetic. In addition a finite Hubbard repulsion is needed *at* the connectivity points to guarantee a correlation between the electrons.

#### 4.1.3. Ground state for $U_0 = 0, U_1 > 0$

The ground state has again the form (4.4), but now it is a fully saturated ferromagnetic state at  $n = 1/2$  even for  $U_0 = 0$ , since the Hubbard interaction is non-zero at the connectivity points ( $U_1 > 0$ ). We thus see that for flat-band ferromagnetism to occur the Hubbard interaction is needed *only* at the connectivity points. The ferromagnetic states discussed in §§4.1.1 and 4.1.3. are localized in the thermodynamic limit.

### 4.2. Mapping to the one-dimensional periodic Anderson model (PAM)

By the following change of notation in (4.1):

$$\hat{c}_{\mathbf{i}\sigma} \rightarrow \hat{d}_{\mathbf{i},\sigma}, \quad \hat{c}_{\mathbf{i}+\mathbf{r},\sigma} \rightarrow \hat{f}_{\mathbf{i},\sigma}, \quad t' \rightarrow V, \quad \varepsilon_1 \rightarrow E_f, \quad U_1 \rightarrow U, \quad (4.5)$$

and setting  $\varepsilon_0 = U_0 = 0$  the triangle Hubbard chain transforms into a one-dimensional PAM with zero on-site hybridization and nearest-neighbor hybridization  $V$ :

$$\begin{aligned} \hat{H}^{PAM} = & \sum_{\sigma} \sum_{\mathbf{i}=1}^{N_c} (t \hat{d}_{\mathbf{i},\sigma}^\dagger \hat{d}_{\mathbf{i}+\mathbf{a},\sigma} + \text{H.c.}) + E_f \hat{f}_{\mathbf{i}+\mathbf{r},\sigma}^\dagger \hat{f}_{\mathbf{i}+\mathbf{r},\sigma} \\ & + V (\hat{d}_{\mathbf{i},\sigma}^\dagger \hat{f}_{\mathbf{i}+\mathbf{r},\sigma} + \hat{d}_{\mathbf{i}+\mathbf{a},\sigma}^\dagger \hat{f}_{\mathbf{i}+\mathbf{r},\sigma} + \text{H.c.}) + U \sum_{\mathbf{i}} \hat{n}_{\mathbf{i}+\mathbf{r},\uparrow}^f \hat{n}_{\mathbf{i}+\mathbf{r},\downarrow}^f. \end{aligned} \quad (4.6)$$

A lowest bare flat band is obtained for  $V^2/t = E_f + 2t$ ,  $E_f > -2t$ ,  $t > 0$ .

#### 4.2.1. Exact ground state for the one-dimensional PAM

In the parameter regime  $V^2/t = E_f + 2t$ ,  $E_f > -2t$ ,  $t > 0$  the exact ground state for the one-dimensional PAM is obtained from (4.4) as

$$|\Psi_g^{PAM}(N = N_c)\rangle = \prod_{\mathbf{i}=1}^{N_c} \left[ \hat{f}_{\mathbf{i}-\mathbf{a}+\mathbf{r},\sigma}^\dagger + \hat{f}_{\mathbf{i}+\mathbf{r},\sigma}^\dagger - \frac{V}{t} \hat{d}_{\mathbf{i},\sigma}^\dagger \right] |0\rangle. \quad (4.7)$$

#### 4.2.2. Itinerant ferromagnetism of finite chains

In analogy with (3.16) the localization length  $\xi$  of the  $d$ -electrons can be calculated from the expression for the expectation value of the hopping term  $\Gamma_{\mathbf{r}}$ . Its dependence on the nearest-neighbor hybridization strength  $V$  is given by

$$\xi = \frac{a}{\ln[1 + \frac{V^2}{2t^2}(1 - \sqrt{1 + \frac{4t^2}{V^2}})]}, \quad (4.8)$$

as shown in Fig. 7. Here  $a$  is the distance between sites at the base of the triangle chain (see Fig. 5). Although flat-band ferromagnetism in the thermodynamic limit usually corresponds to localized electrons, the localization length can be so large that it exceeds the length of a finite sample. In realistic situations the exact ground state of the finite chain therefore shows *itinerant ferromagnetism*.

#### 4.3. Application to $f$ -electron ferromagnetism in $\text{CeRh}_3\text{B}_2$

As pointed out by Kono and Kuramoto<sup>21)</sup> the one-dimensional PAM, (4-6), may be employed to model the  $4f$ -electron compound  $\text{CeRh}_3\text{B}_2$ . This material is ferromagnetic below  $T_c = 120$  K. Its properties are interesting for several reasons: (i) the ferromagnetic state cannot be explained by a RKKY interaction, (ii) the  $4f$  moment has the remarkably small value of  $0.45 \mu_B$  compared to the free  $\text{Ce}^{3+}$  moment of  $2.14 \mu_B$ , and (iii) the Curie temperature  $T_c$  has the highest value among all known Ce compounds with non-magnetic elements.

The lattice structure of the class of rare-earth (RE) compounds  $\text{RERh}_3\text{B}_2$  was determined by Yamada et al.<sup>30)</sup> A cut through the  $(a, c)$ -plane of the Cerium system ( $\text{RE} = \text{Ce}$ ) is shown in Fig. 8. The remarkable difference between the distances of the Ce atoms along the  $a$ - and the  $c$ -direction, respectively, makes this material a quasi one-dimensional two-band system.<sup>31)</sup> For this reason Kono and Kuramoto<sup>21)</sup> modeled  $\text{CeRh}_3\text{B}_2$  by the triangle PAM chain shown in Fig. 9. The parameters entering the model are  $t = 0.34$  eV,  $V = 0.24$  eV,  $E_f = -0.714$  eV,  $U = 7$  eV,  $n = 0.55$ , as obtained from band-structure calculations and X-ray absorption spectroscopy (we note that the density  $n = 0.55$  corresponds to twice the value, i.e.,  $n = 1.1$ , in the notation of Ref. 21)).

##### 4.3.1. Variational results for the one-dimensional PAM

Kono and Kuramoto<sup>21)</sup> computed the band structure of  $\text{CeRh}_3\text{B}_2$  by constructing a Gutzwiller variational wave function and using variational Monte Carlo to determine the ground state. They found the lowest band of  $\text{CeRh}_3\text{B}_2$  to be almost perfectly flat and fully polarized, and the second band dispersive and almost unpolarized. It should be stressed that the shape of the lowest quasiparticle band of interacting electrons depends on  $U$ .

##### 4.3.2. Exact ground state of the one-dimensional PAM

We can use the exact ground state of the one-dimensional PAM, (4-7), to make direct contact with the  $f$ -electron ferromagnetism found in  $\text{CeRh}_3\text{B}_2$ . Namely, this ground state represents a saturated flat-band ferromagnetic state of the  $f$ -electrons for electron density  $n = 1/2$  and arbitrary  $U > 0$ , provided the condition  $V^2/t =$




Fig. 7. Localization length  $\xi$  of the  $d$ -electrons of the one-dimensional PAM, (4-6), as a function of the nearest-neighbor hybridization strength  $V$ .




Fig. 8. Cut through the  $(a, c)$ -plane of the lattice structure of  $\text{CeRh}_3\text{B}_2$  based on the parameters determined in Ref. 30).




Fig. 9. One-dimensional PAM used to model  $\text{CeRh}_3\text{B}_2$ .<sup>21)</sup>

$E_f + 2t$ ,  $E_f > -2t$ ,  $t > 0$  holds. The condition is indeed fulfilled for the parameter values  $t = 0.34$  eV,  $V = 0.23$  eV,  $E_f = -0.52$  eV, which are very close to the experimentally determined values. In this case the lowest band is perfectly flat for all  $U > 0$ . Apparently the ferromagnetism of the  $f$ -electrons in  $\text{CeRh}_3\text{B}_2$  is due to the presence of an (almost) flat lowest band.

#### 4.3.3. Magnetic moments

The experimentally observed magnetic moment of the  $f$ -electrons is  $m_f = 0.45 \mu_B$ .<sup>32)</sup> The variational result<sup>21)</sup> yields  $m_f = 0.94 \mu_B$ , while the exact ground state discussed above leads to a value of  $m_f = 0.68 \mu_B$  which is even closer to the experimental value. Both theoretical values are somewhat higher than the experimental  $f$  moment. This may be due to the fact that the theoretical approaches do not include the spin-orbit interaction which causes an antiferromagnetic coupling between the  $d$ - and the  $f$ -electrons.

## §5. Conclusions

Despite its simplicity, the diamond Hubbard chain displays remarkably complex physical properties with insulating, conducting, and fully or partially spin polarized states. The constructed exact many-electron ground states revealed flat-band ferromagnetism, correlated half-metallic behavior, and metal-insulator transitions in the diamond Hubbard chain. The virtue of tuning fundamentally different ground states through external magnetic fields, site selective potentials, or the electron density, points to new prospects for designing electronic devices, which allow to switch between different ground states, including states that operate as a spin valve. We applied the general strategy for the construction of exact ground states also to triangle Hubbard chains. The mapping of the triangle Hubbard chain to the one-dimensional periodic Anderson model within a restricted parameter set provided a theory for

$f$ -electron ferromagnetism in  $\text{CeRh}_3\text{B}_2$ , which is based on the physical properties of an exact ground state.

Further applications of the construction of exact ground states by casting the Hamiltonian into positive semidefinite form are possible for model systems with selected geometries in artificially designed structures. In particular, we constructed exact ground states for pentagon Hubbard chains. These results will be presented elsewhere.

### Acknowledgements

We thank F. Mila, Y. Kuramoto, and F. C. Zhang for discussions regarding the uniqueness of the solutions. Support by the Hungarian Research Fund through Contracts No. OTKA-T48782, 76821, the Alexander von Humboldt Foundation, and the Deutsche Forschungsgemeinschaft through SFB 484 is gratefully acknowledged.

### Appendix A

#### — Uniqueness of Ground States with $N = pN_c$ Where $p$ Is an Integer —

Here we prove the uniqueness of those exact ground states constructed in this paper for which the electron number  $N$  is an integer multiple of the number of unit cells  $N_c$ . The proof proceeds as discussed in §2.3 and is exemplified by proving the uniqueness of the ground state  $|\Psi_g^{II}(4N_c)\rangle$ , (3·15), of the diamond Hubbard chain. In this case  $N = 4N_c$ , i.e.,  $p = 4$ , and the electron density has the value  $n = 4/3$ . The Hamiltonian, the unnormalized ground state, and the  $\hat{A}_{i,\sigma}$  operators entering therein are given by (3·11), (3·15), and (3·8), respectively. To prove the uniqueness one first has to determine the kernel of  $\hat{H}' = \hat{H} - E_g^{II} = \sum_{\sigma} \sum_{i=1}^{N_c} \hat{A}_{i,\sigma} \hat{A}_{i,\sigma}^{\dagger} + U\hat{P}$ . The form of the Hamiltonian  $\hat{H}'$  implies that its kernel is simply given by the intersection of the kernels of the individual terms, i.e.,  $\ker(\hat{H}') = \ker(\sum_{\sigma} \sum_{i=1}^{N_c} \hat{A}_{i,\sigma} \hat{A}_{i,\sigma}^{\dagger}) \cap \ker(\hat{P})$ , where  $\hat{P}$  is an operator, which assumes its minimum eigenvalue 0, if there is at least one electron at each site.

To prove the uniqueness of the ground state  $|\Psi_g^{II}(4N_c)\rangle$ , (3·15), it is now necessary to demonstrate that it spans  $\ker(\hat{H}')$ . According to the discussion in §2.3 the proof consists of two parts. First one has to establish that

(i)  $|\Psi_g^{II}(4N_c)\rangle$  is an element of the intersection of the kernels of the positive semidefinite operators entering in  $\hat{H}'$ .

Then one must show that

(ii) all states from the kernel of  $\hat{H}'$  can be written in the form of the ground state  $|\Psi_g^{II}(4N_c)\rangle$ .

The first part of the proof, (i), is covered by the following Theorems A-1 to A-3. The second part of the proof, (ii), then follows immediately (see below).

To prove (i) we note that the first term in  $\hat{H}'$  is a sum of positive semidefinite operators  $\hat{A}_{i,\sigma} \hat{A}_{i,\sigma}^{\dagger}$ . The ground state of  $\hat{H}'$  must therefore be an element of each kernel of all the operator  $\hat{A}_{i,\sigma} \hat{A}_{i,\sigma}^{\dagger}$ , for all  $i, \sigma$ . Thus we first determine the kernel of



$\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger$  (Appendix A.1), then the kernel of the sum  $\sum_\sigma \sum_{i=1}^{N_c} \hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger$  (Appendix A.2), and finally the kernel of the full Hamiltonian  $\hat{H}'$  (Appendix A.3).

#### A.1. The kernel of $\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger$

**Theorem A-1:** The kernel of  $\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger$  is spanned by states of the form

$$|\Psi\rangle = \hat{A}_{i,\sigma}^\dagger \hat{W}^\dagger |0\rangle, \quad (\text{A.1})$$

where  $\hat{W}^\dagger$  is an arbitrary operator.

*Proof of Theorem A-1*

a) From  $(\hat{A}_{i,\sigma}^\dagger)^2 = 0$  it follows that  $\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger|\Psi\rangle = 0$ . This implies that  $|\Psi\rangle \in \ker(\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger)$ .

b) In order to show that all elements of  $\ker(\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger)$  can be written in the form  $|\Psi\rangle = \hat{A}_{i,\sigma}^\dagger \hat{W}^\dagger |0\rangle$  we consider a state  $|\Phi\rangle = \hat{Y}^\dagger |0\rangle \in \ker(\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger)$  and show that  $|\Phi\rangle$  has the form  $|\Psi\rangle = \hat{A}_{i,\sigma}^\dagger \hat{W}^\dagger |0\rangle$ .

Making use of  $\{\hat{A}_{i,\sigma}, \hat{A}_{i,\sigma}^\dagger\} = 8t_\perp^2 t_\parallel + 2t_\parallel =: a_{i,i}^\sigma > 0$  and  $\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger|\Phi\rangle = 0$  we can write

$$|\Phi\rangle = \hat{Y}^\dagger |0\rangle = \frac{1}{a_{i,i}^\sigma} (\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger + \hat{A}_{i,\sigma}^\dagger\hat{A}_{i,\sigma})\hat{Y}^\dagger |0\rangle = \hat{A}_{i,\sigma}^\dagger \left( \frac{1}{a_{i,i}^\sigma} \hat{A}_{i,\sigma}\hat{Y}^\dagger \right) |0\rangle. \quad (\text{A.2})$$

With the identification  $\hat{W}^\dagger := \frac{1}{a_{i,i}^\sigma} \hat{A}_{i,\sigma}\hat{Y}^\dagger$  we have therefore verified that indeed

$$|\Phi\rangle = \hat{A}_{i,\sigma}^\dagger \hat{W}^\dagger |0\rangle = |\Psi\rangle. \quad (\text{A.3})$$

This completes the proof of Theorem A-1.

#### A.2. The kernel of $\sum_\sigma \sum_{i=1}^{N_c} \hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger$

The kernel of  $\sum_\sigma \sum_{i=1}^{N_c} \hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger$  has the form

$$\ker \left( \sum_\sigma \sum_{i=1}^{N_c} \hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger \right) = \bigcap_{\sigma=\uparrow,\downarrow} \bigcap_{i=1}^{N_c} \ker(\hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger). \quad (\text{A.4})$$

**Theorem A-2:**  $\ker(\sum_\sigma \sum_{i=1}^{N_c} \hat{A}_{i,\sigma}\hat{A}_{i,\sigma}^\dagger)$  is spanned by vectors of the form

$$|\Psi\rangle = \left[ \prod_\sigma \prod_{i=1}^{N_c} \hat{A}_{i,\sigma}^\dagger \right] \hat{W}^\dagger |0\rangle, \quad (\text{A.5})$$

where the operator  $\hat{W}^\dagger$  is arbitrary.

Theorem A-2 follows directly from the linear independence of the operators  $\hat{A}_{i,\sigma}^\dagger$  (that is,  $\hat{A}_{i,\sigma}^\dagger \neq \sum_{j \neq i} \alpha_{j,\sigma} \hat{A}_{j,\sigma}^\dagger$  holds for arbitrary coefficients  $\alpha_{j,\sigma}$  due to the fact that the operators  $\hat{c}_{i+r_1,\sigma}^\dagger, \hat{c}_{i+r_2,\sigma}^\dagger$  enter only in  $\hat{A}_{i,\sigma}^\dagger$  and not in  $\hat{A}_{j,\sigma}^\dagger$ , for  $j \neq i$ ).

### A.3. The kernel of $\hat{H}'$

The kernel of  $\hat{H}'$  is given by  $\ker(\hat{H}') = \ker(\sum_{\sigma} \sum_{i=1}^{N_c} \hat{A}_{i,\sigma} \hat{A}_{i,\sigma}^{\dagger}) \cap \ker(\hat{P})$ , where  $\hat{P}$  is an operator, which assumes its minimum eigenvalue 0, if there is at least one electron at each site. Hence,  $\ker(\hat{H}')$  is spanned by vectors of the form in Eq. (A-1), where now the operator  $\hat{W}^{\dagger}$  is no longer arbitrary, but must have a *specific* form to satisfy the above-mentioned requirement imposed by  $\ker(\hat{P})$ .

**Theorem A-3:** For  $N = 4N_c$  the operator  $\hat{W}^{\dagger}$  is given by  $\hat{W}^{\dagger} = \hat{F}_{\sigma}^{\dagger}$ , where  $\hat{F}_{\sigma}^{\dagger}$  creates two electrons with fixed spin  $\sigma$  in each unit cell.

*Proof of Theorem A-3.*

The proof proceeds in two main steps: we show that  $\hat{W}^{\dagger} = \hat{F}_{\sigma}^{\dagger}$  is 1) a possible choice, and 2) the unique choice.

1) Let us denote by  $\mathbf{T}_i$  the unit cell (= triangle), which contains the sites  $i, i + \mathbf{r}_1, i + \mathbf{r}_2$ . The state  $\hat{F}_{\sigma}^{\dagger}|0\rangle$  has one empty site in each unit cell  $\mathbf{T}_i$ . We now analyze how these empty sites are filled by the product  $\prod_{i=1}^{N_c} \hat{A}_{i,\sigma}^{\dagger}$ . We first observe that  $\hat{A}_{i,\sigma}^{\dagger}$  acts on all sites in  $\mathbf{T}_i$ , which implies that the empty site in  $\mathbf{T}_i$  can indeed be filled by applying  $\hat{A}_{i,\sigma}^{\dagger}$ . There are  $N_c$  empty sites and  $N_c$  operators  $\hat{A}_{i,\sigma}^{\dagger}$ ; thus, the product  $\prod_{i=1}^{N_c} \hat{A}_{i,\sigma}^{\dagger}$  inserts  $N_c$  electrons with spin  $\sigma$  into the system. Since  $(\hat{c}_{j,\sigma}^{\dagger})^2 = 0$ , the state  $\prod_{i=1}^{N_c} \hat{A}_{i,\sigma}^{\dagger} \hat{F}_{\sigma}^{\dagger}|0\rangle$  has *one*  $\sigma$ -electron on all  $3N_c$  sites of the lattice. Hence,  $\hat{F}_{\sigma}^{\dagger}$  is a possible choice for  $\hat{W}^{\dagger}$  in constructing the states contained in  $\ker(\hat{H}')$ .

2) We now flip the spin  $\sigma$  of one electron in  $\hat{F}_{\sigma}^{\dagger}$  into  $-\sigma$ , and denote the new operator  $\hat{W}^{\dagger}$  by  $\hat{F}_{\sigma}^{\dagger}$ . In this case, a single unit cell, say  $\mathbf{T}_{i_o}$ , will have one empty site, one electron with spin  $\sigma$  on another site, and one electron with spin  $-\sigma$  on the third site. Then the operator  $\hat{A}_{i_o,\sigma}^{\dagger}$  can create one electron with spin  $\sigma$  on the site, which already contains the electron with spin  $-\sigma$ , thereby creating a double occupancy. One empty site will remain in  $\mathbf{T}_{i_o}$ . For  $i \neq i_o$ , the operators  $\hat{A}_{i,\sigma}^{\dagger}$  can introduce their electron into the empty sites of  $\mathbf{T}_i$ . Hence one empty site can remain in the system. The state  $|\chi\rangle = \prod_{i=1}^{N_c} \hat{A}_{i,\sigma}^{\dagger} \hat{F}_{\sigma}^{\dagger}|0\rangle$  therefore has at least one term with an empty site and contains one  $-\sigma$  electron. Acting now with  $\prod_{i=1}^{N_c} \hat{A}_{i,-\sigma}^{\dagger}$  on  $|\chi\rangle$  adds  $N_c$  electrons with spin  $-\sigma$  on  $(3N_c - 1)$  sites. The new state  $|\Phi\rangle = \prod_{i=1}^{N_c} \hat{A}_{i,-\sigma}^{\dagger} |\chi\rangle$  will therefore have at least one term with an empty site, and is thus not an element of  $\ker(\hat{P})$  and  $\ker(\hat{H})$ . This shows that  $\hat{W}^{\dagger} = \hat{F}_{\sigma}^{\dagger}$  is the *unique* choice. This completes the proof of Theorem A-3.

Theorems A-1, A-2 and A-3 prove that  $|\Psi_g^{II}(4N_c)\rangle$ , (3-15), is indeed an element of the intersection of the kernels of the positive semidefinite operators entering in  $\hat{H}'$ . This completes part (i) of the proof of the uniqueness discussed in §2.3.

Since all vectors from  $\ker(\hat{H}')$  can be written in the form  $|\Psi_g^{II}(4N_c)\rangle$ , (3-15), and since this state represents the constructed ground state  $|\Psi_g\rangle$ , it follows immediately that *all* vectors in  $\ker(\hat{H}')$  can be written in the form (3-15). Therefore the ground state (3-15) is the *unique* ground state of the Hamiltonian  $\hat{H}$ , (3-11).

The proofs of the uniqueness of the ground state (3-4) at density  $n = 1/3$ , corresponding to  $p = 1$ , and of solution III in Ref. 8) at density  $n = 5/3$  ( $p = 5$ )

proceed similarly.

## Appendix B

### —— Uniqueness of Ground-State Solutions with $N = pN_c$ for Non-Integer $p$ ——

#### B.1. Geometric degeneracies

For non-integer  $p$  as in the case of (3·4) for  $n < 1/3$ , and (3·17) for  $n > 4/3$  (as well as in the case of solution III for  $n > 5/3$  of Ref. 8)), the ground states have a high geometric degeneracy. In these cases there exist linearly independent eigenstates  $|\Psi_g^\gamma\rangle$  with the same ground-state energy. They may be denoted by a degeneracy index  $\gamma = 1, 2, \dots, M_d$ , where  $M_d$  is the degree of degeneracy. The ground state then becomes  $|\Psi_g\rangle = \sum_{\gamma=1}^{M_d} \alpha_\gamma |\Psi_g^\gamma\rangle$ , where apart from the normalization condition, the coefficients  $\alpha_\gamma$  are arbitrary. For these ground states the proof of uniqueness proceeds analogously to that described in Appendix A. Namely, one has to prove that  $|\Psi_g\rangle$  spans  $\ker(\hat{H}')$ , but now, given the specific structure of  $|\Psi_g\rangle$ , it has to be verified that

- 1) all linearly independent states  $|\Psi_g^\gamma\rangle$  are elements of the intersection of the kernels of the positive semidefinite operators  $\hat{P}_n$ , i.e.,  $|\Psi_g^\gamma\rangle \in \ker(\hat{H}') = \bigcap_{n=1}^L \ker(\hat{P}_n)$ , and
- 2) all states  $|\Psi_g\rangle \in \bigcap_{n=1}^L \ker(\hat{P}_n)$  can be written in the form of the constructed ground state, e.g. as a linear combination of the vectors  $|\Psi_g^\gamma\rangle$ .

As an example we now prove the uniqueness of the ground state  $|\Psi_g^I(N)\rangle$ , (3·4), for  $n < 1/3$ . The proof is presented for fixed  $N$ , namely  $N = N_c - 1$ . For other  $N$  values the proof proceeds similarly. The first part of the proof, (i), (see §2.3) is covered by the following Theorems B-1 and B-2. The second part of the proof, (ii), then follows immediately (see below).

#### B.2. Uniqueness of the ground state (3·4) for $N = N_c - 1$ particles

The transformed Hamiltonian  $\hat{H} = \hat{H}_0 + \hat{H}_U$  is determined by  $\hat{H}_0$  and  $\hat{H}_U$  as given by (3·3) and (3·2), respectively. Furthermore, one has  $\hat{N} = \sum_{i,\sigma} \sum_{\nu=1}^3 \hat{C}_{\nu,i,\sigma}^\dagger \hat{C}_{\nu,i,\sigma}$ , where  $\hat{C}_{\nu,i,\sigma}$  are genuine canonical Fermi operators, and  $E_3 < E_1, E_2$ . Since  $\hat{N}$  is a constant of motion, the Hamiltonian can be written as  $\hat{H} = \sum_{i,\sigma} [(E_1 - E_3) \hat{C}_{1,i,\sigma}^\dagger \hat{C}_{1,i,\sigma} + (E_2 - E_3) \hat{C}_{2,i,\sigma}^\dagger \hat{C}_{2,i,\sigma}] + \hat{H}_U + E_g^I$ , where  $E_g^I = E_3 N$  and  $(E_1 - E_3), (E_2 - E_3) > 0$ . Consequently,  $\hat{H}' := \hat{H} - E_g^I$  is a positive semidefinite operator:  $\hat{H}' = \hat{P}_1 + \hat{P}_2$ , where  $\hat{P}_1 = \sum_{i,\sigma} [(E_1 - E_3) \hat{C}_{1,i,\sigma}^\dagger \hat{C}_{1,i,\sigma} + (E_2 - E_3) \hat{C}_{2,i,\sigma}^\dagger \hat{C}_{2,i,\sigma}]$ , and  $\hat{P}_2 = \hat{H}_U$ . The ground state is given by (3·4).

##### B.2.1. The kernel of $\hat{P}_1$

**Theorem B-1:** The kernel of  $\hat{P}_1$  is spanned by states of the form

$$|\Psi(\{\mathbf{i}\}, \{\sigma_i\})\rangle = \prod_{i=1}^{N_c-1} \hat{C}_{3,i,\sigma_i}^\dagger |0\rangle, \quad (\text{B} \cdot 1)$$

where the product is taken over arbitrary  $N_c - 1$  sites, and  $\sigma_i$  is arbitrary. Note that  $|\Psi(\{\mathbf{i}\}, \{\sigma_i\})\rangle$  depends on the set of sites  $\{\mathbf{i}\}$ , and the set of spin orientations  $\{\sigma_i\}$ .  
*Proof of Theorem B-1.*

The proof proceeds in two steps: a) one first shows that Eq. (B.1) is a possible choice, and b) then verifies that it is the unique choice.

a) The initial  $6N_c$  canonical Fermi operators  $\hat{c}_{\mathbf{i}+\mathbf{r}_s, \sigma}$ , where  $s = 1, 2, 3$  is the sublattice index, have been transformed by a linear transformation into the  $6N_c$  operators  $\hat{C}_{\nu, \mathbf{i}, \sigma_i}$ , which satisfy canonical fermionic anti-commutation rules. Since the set of the new operators  $\{\hat{C}_{\nu, \mathbf{i}, \sigma_i}\}$  substitutes the set  $\{\hat{c}_{\mathbf{i}+\mathbf{r}_s, \sigma}\}$ , all operators in the system acting on states defined by arbitrary  $(\mathbf{i}, \sigma_i)$  can be expressed in terms of the operators  $\hat{C}_{\nu, \mathbf{i}, \sigma_i}$ .

Given these conditions, an operator  $\hat{X}^\dagger$  which satisfies

$$\{\hat{C}_{1, \mathbf{i}, \sigma_i}, \hat{X}^\dagger\} = \{\hat{C}_{2, \mathbf{i}, \sigma_i}, \hat{X}^\dagger\} = 0 \quad (\text{B.2})$$

for all indices  $\mathbf{i}, \sigma_i$ , must have the form  $\hat{X}^\dagger = \prod_j \prod_{\sigma_j} \hat{C}_{3, \mathbf{j}, \sigma_j}^\dagger$ . Indeed, the anti-commutation relations (B.2) hold only for this type of  $\hat{X}^\dagger$  operator since an arbitrary  $\hat{X}^\dagger$  operator can be expressed in terms of  $\hat{C}_{\nu, \mathbf{j}, \sigma_j}$  operators, and only  $\hat{C}_{3, \mathbf{j}, \sigma_j}^\dagger$  anticommutes with all  $\hat{C}_{1, \mathbf{i}, \sigma_i}$  and  $\hat{C}_{2, \mathbf{i}, \sigma_i}$ .

b) If  $\hat{X}^\dagger$  contains at least one  $\hat{C}_{\nu, \mathbf{j}, \sigma_j}^\dagger$  operator with  $\nu = 1, 2$ , Eq. (B.2) no longer holds, hence Eq. (B.1) is the unique choice. This proves Theorem B.1.

We furthermore observe that due to the fermionic operator nature of  $\hat{C}_{3, \mathbf{i}, \sigma_i}$ , the deduced elements of  $\ker(\hat{P}_1)$  from Eq. (B.1) are not only linearly independent but also orthogonal:

$$\langle \Psi(\{\mathbf{i}\}, \{\sigma_i\}) | \Psi(\{\mathbf{i}'\}, \{\sigma'_i\}) \rangle = \delta_{\{\mathbf{i}\}, \{\mathbf{i}'\}} \delta_{\{\sigma_i\}, \{\sigma'_i\}}. \quad (\text{B.3})$$

### B.2.2. The kernel of $\hat{H}'$

Since  $\ker(\hat{H}') = \ker(\hat{P}_1) \cap \ker(\hat{P}_2)$ , the states in  $\ker(\hat{H}')$  have the form (B.1), but the specific set of indices  $(\{\mathbf{i}\}, \{\sigma_i\})$  has to be chosen such that these states are also contained in  $\ker(\hat{P}_2)$ . This condition requires the proof of the following theorem.

**Theorem B-2:** The kernel of  $\hat{H}'$  is spanned by states of the form

$$|\Psi(\mathbf{j})\rangle = |\Psi(\{\mathbf{i}\})\rangle = \prod_{\mathbf{i}=1}^{N_c-1} \hat{C}_{3, \mathbf{i}, \sigma}^\dagger |0\rangle, \quad (\text{B.4})$$

where  $\sigma$  is fixed, and the product extends over  $N_c - 1$  arbitrary sites.

Since the product in Eq. (B.4) is taken over  $N_c - 1$  sites of the sublattice  $s = 3$ , the vector  $|\Psi(\{\mathbf{i}\})\rangle$  can be denoted by the specific site  $\mathbf{j}$  excluded from the product, i.e.  $|\Psi(\{\mathbf{i}\})\rangle \equiv |\Psi(\mathbf{j})\rangle$ . This site  $\mathbf{j}$  plays the role of the degeneracy index  $\gamma$  in Appendix B.1.

*Proof of Theorem B-2.*

The theorem is proved in three steps: First one has to show that all  $\sigma_i$  spin indices in (B.4) must have the same value, secondly it has to be verified that the deduced

states in  $\ker(\hat{H}')$  are linearly independent, and thirdly one needs to prove that the set of the deduced states is complete.

a) All states in  $\ker(\hat{P}_2)$  require the absence of any double occupancy. Therefore, all indices  $\sigma_i$  must be fixed to the same value  $\sigma$ . The double occupancy cannot be avoided otherwise, since in  $N_c$  unit cells  $N_c - 1$  operators  $\hat{C}_{3,i,\sigma}^\dagger$  share a common site.

b) Due to the relation (B.3) the vectors  $|\Psi(\mathbf{j})\rangle$  are linearly independent, since they are orthogonal  $\langle\Psi(\mathbf{j})|\Psi(\mathbf{j}')\rangle = \delta_{\mathbf{j},\mathbf{j}'}$ .

c) Obviously, the degree of degeneracy of a macroscopic many-body state is given by the number of equivalent configurations. Due to part a), apart from the trivial degeneracy with respect to the total spin direction, one finds  $M_d = N_c$ , because there are  $N_c$  possibilities for placing  $N_c - 1$  identical objects on  $N_c$  sites. Consequently,  $\ker(\hat{H}')$  is an  $M_d = N_c$  dimensional Hilbert subspace. The index  $\mathbf{j}$  in the states of the form (B.4) refers to  $N_c$  possible different unit cells. Furthermore, the vectors  $|\Psi(\mathbf{j})\rangle$  are linearly independent, hence the set  $\{|\Psi(\mathbf{j})\rangle\}$  forms a (orthonormalized) basis of  $\ker(\hat{H}')$ . Therefore,  $\{|\Psi(\mathbf{j})\rangle\}$  spans  $\ker(\hat{H}')$ , and the ground state has the general form  $|\Psi_g\rangle = \sum_{\mathbf{j}=1}^{N_c} \alpha_{\mathbf{j}} |\Psi(\mathbf{j})\rangle$ , with arbitrary coefficients  $\alpha_{\mathbf{j}}$ . This proves Theorem B-2.

Since  $\{|\Psi(\mathbf{j})\rangle\}$  is a basis of  $\ker(\hat{H}')$ , all states  $|\phi\rangle \in \ker(\hat{H}')$  can be written in the form  $|\Psi_g\rangle$  of part c). Therefore (3.4) is the unique ground state of the Hamiltonian  $\hat{H}$  consisting of the two terms (3.3) and (3.2).

For  $N = N_c$  the set  $\{\mathbf{j}\}$  is an empty set, and the upper limit set to  $N_c$  in Eq. (B.4) provides the unique ground state at density  $n = 1/3$  (see (3.4) for  $N = N_c$ ). A similar strategy for  $N < N_c - 1$  allows one to prove the uniqueness of the ground-state solutions for all electron densities  $n < 1/3$  in (3.4). Similar proofs are expected to hold in the case of the ground state  $|\Psi_g^{II}(4N_c + \Delta N)\rangle$ , (3.17), for  $n > 4/3$  ( $p > 4$ ) and solution III for  $n > 5/3$  ( $p > 5$ ) constructed in Ref. 8).

## References

- 1) M. Imada, A. Fujimori and Y. Tokura, Rev. Mod. Phys. **70** (1998), 1039.
- 2) Y. Tokura and N. Nagaosa, Science **288** (2000), 462.
- 3) E. Dagotto, Science **309** (2005), 257.
- 4) *The Hubbard Model*, ed. A. Montorsi (World Scientific, Singapore, 1992).
- 5) H. Yao, W.-F. Tsai and S. A. Kivelson, Phys. Rev. B **76** (2007), 161104(R).
- 6) Z. Gulácsi and D. Vollhardt, Phys. Rev. Lett. **91** (2003), 186401.
- 7) Z. Gulácsi and D. Vollhardt, Phys. Rev. B **72** (2005), 075130.
- 8) Z. Gulácsi, A. Kampf and D. Vollhardt, Phys. Rev. Lett. **99** (2007), 026404.
- 9) U. Brandt and A. Giesekeus, Phys. Rev. Lett. **68** (1992), 2648.
- 10) R. Strack and D. Vollhardt, Phys. Rev. Lett. **70** (1993), 2637; Physica B **199-200** (1994), 273.
- 11) R. Strack, Phys. Rev. Lett. **70** (1993), 833.
- 12) R. Strack and D. Vollhardt, Phys. Rev. Lett. **72** (1994), 3425; J. Low Temp. Phys. **99** (1995), 385.
- 13) A. Montorsi and D. K. Campbell, Phys. Rev. B **53** (1996), 5153.
- 14) L. G. Sarason, Phys. Rev. B **75** (2007), 054504.
- 15) R. B. Laughlin, Philos. Mag. B **86** (2006), 1165.
- 16) P. Gurin and Z. Gulácsi, Philos. Mag. B **76** (1997), 827.
- 17) I. Orlik and Z. Gulácsi, Philos. Mag. Lett. **78** (1998), 177.
- 18) L. G. Sarason and M. A. Continentino, Phys. Rev. B **65** (2002), 233107.
- 19) P. Gurin and Z. Gulácsi, Phys. Rev. B **64** (2001), 045118 [Errata; **65** (2002), 129901].
- 20) Z. Gulácsi, Phys. Rev. B **66** (2002), 165109; Eur. Phys. J. B **30** (2002), 295.

- 21) H. K. Kono and Y. Kuramoto, J. Phys. Soc. Jpn. **75** (2006), 084706.
- 22) J. Vidal, B. Doucot, R. Mosseri and P. Butaud, Phys. Rev. Lett. **85** (2000), 3906.
- 23) A. Mielke and H. Tasaki, Commun. Math. Phys. **158** (1993), 341.
- 24) M. C. Gutzwiller, Phys. Rev. Lett. **10** (1963), 159.
- 25) J. Kanamori, Prog. Theor. Phys. **30** (1963), 275.
- 26) K. Penc, H. Shiba, F. Mila and T. Tsukagoshi, Phys. Rev. B **54** (1996), 4056.
- 27) P. Fazekas, Philos. Mag. B **76** (1997), 797.
- 28) E. Müller-Hartmann, J. Low Temp. Phys. **99** (1995), 349.
- 29) O. Derzhko, A. Honecker and J. Richter, Phys. Rev. B **76** (2007), 220402(R).
- 30) M. Yamada, Y. Obiraki, T. Okubo, T. Shiromoto, Y. Kida, M. Shiimoto, H. Kohara, T. Yamamoto, D. Honda, A. Galatanu, Y. Haga, T. Takeuchi, K. Sugiyama, R. Settai, K. Kindo, S. K. Dhar, H. Harima and Y. Onuki, J. Phys. Soc. Jpn. **73** (2004), 2266.
- 31) T. Okubo, M. Yamada, A. Thamizhavel, S. K. Y. Inada, R. Settai, H. Harima, K. Takegahara, A. Galatanu, E. Yamamoto and Y. Onuki, J. of Phys.: Cond. Mat. **15** (2003), L721.
- 32) A. Galatanu, E. Yamamoto, T. Okubo, M. Yamada, A. Thamizhavel, T. Takeuchi, K. Sugiyama, Y. Inada and Y. Onuki, J. of Phys.: Cond. Mat. **15** (2003), S2187.