

**A COMPARISON OF A POSTERIORI ERROR ESTIMATORS
FOR MIXED FINITE ELEMENT DISCRETIZATIONS
BY RAVIART-THOMAS ELEMENTS**

BARBARA I. WOHLMUTH AND RONALD H.W. HOPPE

ABSTRACT. We consider mixed finite element discretizations of linear second order elliptic boundary value problems with respect to an adaptively generated hierarchy of possibly highly nonuniform simplicial triangulations. In particular, we present and analyze four different kinds of error estimators: a residual based estimator, a hierarchical one, error estimators relying on the solution of local subproblems and on a superconvergence result, respectively. Finally, we examine the relationship between the presented error estimators and compare their local components.

1. INTRODUCTION

We consider the following boundary value problem for a linear second order elliptic differential operator

$$(1.1) \quad \begin{aligned} u &:= -\operatorname{div}(a\nabla u) + bu = f && \text{in } \Omega, \\ u &= 0 && \text{on } \Gamma := \partial\Omega, \end{aligned}$$

where Ω stands for a bounded, polygonal domain in the Euclidean space \mathbb{R}^2 and $f \in L^2(\Omega)$. Furthermore, we assume $a = (a_{ij})_{i,j=1}^2$ to be a symmetric, matrix-valued function with $a_{ij} \in L^\infty(\Omega)$, $1 \leq i, j \leq 2$, and $b \in L^\infty(\Omega)$ satisfying

$$(1.2) \quad \begin{aligned} \alpha_0 |\xi|^2 &\leq \sum_{i,j=1}^2 a_{ij}(x) \xi_i \xi_j \leq \alpha_1 |\xi|^2, \\ \xi &\in \mathbb{R}^2, \quad 0 < \alpha_0 \leq \alpha_1, \quad 0 \leq b_0 \leq b(x) \leq b_1, \end{aligned}$$

for almost all $x \in \Omega$. The local bounds on a subset $D \subset \Omega$ are denoted by α_i^D , β_i^D , $0 \leq i \leq 1$. For simplicity, we have chosen homogeneous Dirichlet boundary data, but all subsequent results can be easily applied to more general boundary conditions.

In many applications, the flux $\mathbf{j} := -a\nabla u$ is more important than the primal variable u . Therefore, the original problem (1.1) is transformed into a first order system by introducing the auxiliary variable \mathbf{j} . The natural ansatz space for the flux is

$$H(\operatorname{div}; \Omega) := \{ \mathbf{q} \in (L^2(\Omega))^2 \mid \operatorname{div} \mathbf{q} \in L^2(\Omega) \},$$

Received by the editor September 6, 1996 and, in revised form, August 11, 1997 and January 6, 1998.

1991 *Mathematics Subject Classification*. Primary 65F10, 65N30, 65N50, 65N55.

Key words and phrases. Mixed finite elements, a posteriori error estimation, adaptive grid refinement.

which is a Hilbert space with respect to the inner product

$$(\mathbf{p}, \mathbf{q})_{\text{div}} := (\mathbf{p}, \mathbf{q})_0 + (\text{div } \mathbf{p}, \text{div } \mathbf{q})_0$$

and the associated norm $\|\cdot\|_{\text{div}} := (\cdot, \cdot)_{\text{div}}^{1/2}$. As usual we denote by $(\cdot, \cdot)_k$, $k \geq 0$, the standard inner product on $H^k(\Omega)$ and $(H^k(\Omega))^2$, while $|\cdot|_k$, $\|\cdot\|_k$ stand for the associated seminorms and norms, respectively.

Then, the weak formulation associated with (1.1) gives rise to the following saddle point problem:

Find $(\mathbf{j}, u) \in H(\text{div}; \Omega) \times L^2(\Omega)$ such that

$$(1.3) \quad \begin{aligned} a(\mathbf{j}, \mathbf{q}) + b(\mathbf{q}, u) &= 0, & \mathbf{q} \in H(\text{div}; \Omega), \\ b(\mathbf{j}, v) - c(u, v) &= -l(v), & v \in L^2(\Omega), \end{aligned}$$

where the bilinear forms $a(\cdot, \cdot)$, $b(\cdot, \cdot)$, $c(\cdot, \cdot)$ and the functional $l(\cdot)$ are given by

$$\begin{aligned} a(\mathbf{p}, \mathbf{q}) &:= \int_{\Omega} a^{-1} \mathbf{p} \cdot \mathbf{q} \, dx, & \mathbf{p}, \mathbf{q} \in H(\text{div}; \Omega), \\ b(\mathbf{q}, v) &:= - \int_{\Omega} \text{div } \mathbf{q} v \, dx, & \mathbf{q} \in H(\text{div}; \Omega), v \in L^2(\Omega), \\ c(u, v) &:= \int_{\Omega} b u v \, dx, & u, v \in L^2(\Omega), \\ l(v) &:= \int_{\Omega} f v \, dx, & v \in L^2(\Omega). \end{aligned}$$

The existence and uniqueness of the solution of system (1.3) are well known (cf. e.g. [12, §II, Thm. 1.2]). Associated with the bilinear form $a(\cdot, \cdot)$ is the norm $\|\cdot\|_{\text{div}}$, where $\|\mathbf{q}\|_{\text{div}}^2 := \int_{\Omega} a^{-1} \mathbf{q} \mathbf{q} \, dx + \int_{\Omega} \text{div } \mathbf{q} \text{div } \mathbf{q} \, dx$. Compared with the norm $\|\cdot\|_{\text{div}}$, the norm $\|\cdot\|_{\text{div}}$ is weighted by a^{-1} and plays the same role as the energy norm in the primal formulation. Both norms are equivalent due to the positive definiteness of a .

The mixed finite element approach is based on (1.3). Here, we use Raviart-Thomas finite elements with respect to a simplicial triangulation \mathcal{T}_h of Ω . The sets of vertices and edges are denoted by $\mathcal{P}_h := \mathcal{P}_h^0 \cup \mathcal{P}_h^\Gamma$, $\mathcal{E}_h := \mathcal{E}_h^0 \cup \mathcal{E}_h^\Gamma$ where \mathcal{P}_h^0 , \mathcal{E}_h^0 refer to the interior vertices and edges and \mathcal{P}_h^Γ , \mathcal{E}_h^Γ to those located on $\Gamma = \partial\Omega$. Furthermore, $P_k(D)$, $D \subseteq \Omega$, $k \geq 0$, stands for the set of polynomials of degree $\leq k$ on D .

For the discretization of the flux $\mathbf{j} \in H(\text{div}; \Omega)$ we choose the Raviart-Thomas ansatz space

$$RT_k(\Omega; \mathcal{T}_h) := \{\mathbf{q}_h \in H(\text{div}; \Omega) \mid \mathbf{q}_h|_T \in RT_k(T), T \in \mathcal{T}_h\},$$

where $RT_k(T)$, $T \in \mathcal{T}_h$, stands for the Raviart-Thomas element

$$RT_k(T) := P_k(T)^2 + P_k(T)\mathbf{x}, \quad \mathbf{x} = (x_1, x_2)^T.$$

The degrees of freedom of $RT_k(T)$ are given by the following moments:

$$\int_{e_i} \mathbf{n} \cdot \mathbf{q}_h p \, d\sigma, \quad p \in P_k(e_i), \quad \int_T \mathbf{p} \cdot \mathbf{q}_h \, dx, \quad \mathbf{p} \in P_{k-1}(T),$$

where \mathbf{n} is the outer normal on ∂T and e_i , $1 \leq i \leq 3$, are the edges of T . The discrete ansatz space for the primal variable $u \in L^2(\Omega)$ associated with $RT_k(\Omega; \mathcal{T}_h)$ is given by piecewise polynomial functions:

$$W_k(\Omega; \mathcal{T}_h) := \{v_h \in L^2(\Omega) \mid v_h|_T \in P_k(T), T \in \mathcal{T}_h\}.$$

Thus, the natural requirement $\operatorname{div} RT_k(\Omega; \mathcal{T}_h) = W_k(\Omega; \mathcal{T}_h)$ is guaranteed. Now, the lowest order mixed discretization of system (1.3) can be written as follows:

Find $(\mathbf{j}_h, u_h) \in RT_0(\Omega; \mathcal{T}_h) \times W_0(\Omega; \mathcal{T}_h)$ such that the following discrete saddle point problem is satisfied

$$(1.4) \quad \begin{aligned} a(\mathbf{j}_h, \mathbf{q}_h) + b(\mathbf{q}_h, u_h) &= 0, & \mathbf{q}_h &\in RT_0(\Omega; \mathcal{T}_h), \\ b(\mathbf{j}_h, v_h) - c(u_h, v_h) &= -l(v_h), & v_h &\in W_0(\Omega; \mathcal{T}_h). \end{aligned}$$

Note that the Babuška-Brezzi condition is fulfilled and that system (1.4) admits a unique solution (cf., e.g., [12, §II, Prop. 2.11]).

Throughout the following we refer to $(\mathbf{j}, u) \in H(\operatorname{div}; \Omega) \times L^2(\Omega)$ as the unique solution of the mixed variational problem (1.3) and to $(\mathbf{j}_h, u_h) \in RT_0(\Omega; \mathcal{T}_h) \times W_0(\Omega; \mathcal{T}_h)$ as the lowest order Raviart-Thomas approximation satisfying (1.4). Further, we denote by $(\tilde{\mathbf{j}}_h, \tilde{u}_h) \in RT_0(\Omega; \mathcal{T}_h) \times W_0(\Omega; \mathcal{T}_h)$ an available computed approximation obtained by means of an appropriate iterative solution process.

In particular, we advocate multilevel iterative solvers that work on a hierarchy $(\mathcal{T}_k)_{k=0}^j$ of simplicial triangulations of Ω generated by the well known refinement process due to Bank et al. [5]. The refinement strategy is such that a triangle $T \in \mathcal{T}_k$, $k \geq 0$, either remains unrefined, or is subdivided into four congruent subtriangles, or is bisected into two subtriangles. Following the refinement rules in [5], each triangle $T \in \mathcal{T}_k$, $k \geq 0$, is geometrically similar either to an element of \mathcal{T}_0 or to a bisected triangle of \mathcal{T}_0 . The diameter of T , $T \in \mathcal{T}_k$, is denoted by h_T , and h_e stands for the length of the edge $e \in \mathcal{E}_k$. Then, the regularity of the sequence $(\mathcal{T}_k)_{k=0}^j$ guarantees the existence of constants $0 < \kappa_0 \leq \kappa_1$ such that

$$(1.5) \quad \kappa_0 h_{e_i}^2 \leq |T| \leq \kappa_1 h_{e_i}^2, \quad 1 \leq i \leq 3, \quad T \in \mathcal{T}_k,$$

where $|T|$ is the area of T . Moreover, due to the local quasiuniformity of $(\mathcal{T}_k)_{k=0}^j$ there exist constants $\kappa_D, \kappa_K > 0$ such that for $T \in \mathcal{T}_k$

$$(1.6) \quad \operatorname{card}\{e \in \mathcal{E}_k, e \cap \partial T \neq \emptyset\} \leq \kappa_K, \quad \operatorname{card}\{T' \in \mathcal{T}_k, \partial T' \cap \partial T \neq \emptyset\} \leq \kappa_D.$$

We assume that the iterative approximation $(\tilde{\mathbf{j}}_h, \tilde{u}_h) \in RT_0(\Omega; \mathcal{T}_k) \times W_0(\Omega; \mathcal{T}_k)$ satisfies the second equation of the discrete saddle point problem exactly. This can be achieved, for instance, by using the algorithm proposed by Ewing and Wang [17] for a vanishing Helmholtz term $b \equiv 0$, which was later generalized in [19] and [31] to the case of nonvanishing b .

Reliable and efficient a posteriori error estimators are an indispensable tool for efficient adaptive algorithms. We refer to the pioneering work done by Babuška and Rheinboldt [3, 4], Eriksson, Johnson and Hansbo [16, 22, 23], and to the recent survey articles by Bornemann et al. [8] and Verfürth [27, 30] (cf., e.g., [22, 26, 33]). Following the classification of Verfürth [27], we generalize the standard concepts for a posteriori error estimators and present four different types. In the mixed setting, it takes an extra effort to develop an adequate error estimator compared with the standard primal formulation.

We shall derive a posteriori error estimators for the total error $\mathbf{e}_j := \mathbf{j} - \tilde{\mathbf{j}}_h$ in the flux measured in a weighted norm of the flux space $H(\operatorname{div}; \Omega)$, the total error $e_u := u - \tilde{u}_h$ in the primal variable measured in the L^2 -norm, and the total error in both the flux and the primal variable. Denoting the total error to be estimated by ϵ , an estimator η is said to be *efficient* if there exists a constant $\gamma > 0$, independent of the refinement level such that $\gamma\eta \leq \epsilon$, whereas η is called *reliable* if there exists another constant $\Gamma \geq \gamma$, independent of the refinement level, such that $\epsilon \leq \Gamma\eta$. In

this paper, we shall consider a posteriori error estimators that are both efficient and reliable, i.e., estimators satisfying

$$\gamma\eta \leq \epsilon \leq \Gamma\eta.$$

The paper is organized as follows:

In Section 2, the different types of a posteriori error estimators are introduced and the main results of this paper are summarized. For details we refer to Sections 3-6, where the error estimators are discussed thoroughly and upper and lower bounds for the total error are established.

In Section 3, we investigate a residual based error estimator. This kind of error estimator is based on the dual norm of the residual (cf., e.g., [3, 4, 7, 28, 27, 29]). Due to a Helmholtz decomposition of the ansatz space for the flux into subspaces of solenoidal and weakly irrotational vector fields, the corresponding continuous defect problem can be split into two independent subproblems. The first subproblem is associated with the divergence free part of the flux space and can be treated as in the conforming primal formulation, whereas the second subproblem gives rise to an indefinite saddle point problem.

In Sections 4 and 5, we present two types of hierarchical error estimators that are strongly related and require an adequate saturation assumption (cf., e.g., [6, 14, 15, 21, 25, 27]). In particular, for the derivation of the first hierarchical error estimator, which is dealt with in Section 4, we start from an approximation of the defect problem in a higher order ansatz space followed by a localization in terms of an appropriate hierarchical two-level splitting. For the construction of the other hierarchical error estimator, in Section 5 we proceed the other way around and begin with a suitable localization of the defect problem involving local subproblems that are solved by a hierarchical splitting of an elementwise higher order ansatz. In each case, we propose an estimator for the $H(\operatorname{div}; \Omega)$ -norm of the error in the flux variable, as well as an estimator for the combined error in the flux and in the primal variable.

In Section 6, we consider an error estimator for the primal variable in the L^2 -norm. This error estimator is motivated by a superconvergence result for the finite element approximation of u . It is obtained by a comparison of the piecewise constant approximation of the primal variable with a higher order finite element solution arising from a modified nonconforming approach. Finally, we show that the difference between the piecewise constant and the nonconforming approximation is equivalent to a formulation that can be obtained by using some local averaging techniques (cf., e.g., [11, 24, 32, 34]).

In Section 7, we discuss the relationship between these error estimators and prove their equivalence up to higher order terms. We note that the error estimators under consideration are constructed by means of their elementwise contributions according to

$$(1.7) \quad \eta = \sum_{T \in \mathcal{T}_k} \eta_T.$$

Two estimators $\eta^{(1)}$ and $\eta^{(2)}$ are said to be equivalent (locally equivalent), if there exist constants $0 < \delta \leq \Delta$ ($0 < \delta_T \leq \Delta_T$, $T \in \mathcal{T}_k$), independent of the refinement level, such that

$$\delta\eta^{(2)} \leq \eta^{(1)} \leq \Delta\eta^{(2)} \quad (\delta_T\eta_T^{(2)} \leq \eta_T^{(1)} \leq \Delta_T\eta_T^{(2)}, \quad T \in \mathcal{T}_k).$$

In view of (1.7), local equivalence implies equivalence but the converse does not necessarily hold true. Neglecting higher order terms, we obtain local equivalence of the residual based error estimator and the hierarchical one. Using this result, we investigate the estimator based on the solution of local subproblems in more detail, and establish local equivalence with the hierarchical estimator in the case of an appropriate modification of the discrete Dirichlet boundary data. As far as the estimator relying on superconvergence results is concerned, we cannot expect equivalence with the other ones. However, adding two additional terms allows us to prove equivalence, whereas no local equivalence can be established.

2. DEFINITION OF THE ERROR ESTIMATORS AND MAIN RESULTS

In the context of standard primal variational problems, a posteriori error estimators are well established. The recent survey articles of Verfürth [27] and Bornemann et al. [8] give an excellent comparison of different kinds of error estimators in the conforming setting. These concepts have been generalized to nonconforming finite element discretizations by Crouzeix-Raviart elements of lowest order in [21, 31]. For mixed finite element methods, there only exists some work of Braess et al. [9], Braess and Verfürth [10] and Verfürth [28] concerning residual based error estimators and indicators. Here, we use the same techniques as in the conforming case, but the investigation of the estimator for the mixed setting is much more complicated and requires some additional tools. In case of the residual based error estimator, we assume that the coefficient matrix a is a piecewise constant diagonal matrix and b is piecewise constant.

It can be easily seen that the total error $(\mathbf{j}_e, u_e) := (\mathbf{j} - \tilde{\mathbf{j}}_h, u - \tilde{u}_h)$ satisfies the continuous variational problem

$$(2.1) \quad \begin{aligned} a(\mathbf{j} - \tilde{\mathbf{j}}_h, \mathbf{q}) + b(\mathbf{q}, u - \tilde{u}_h) &= r(\mathbf{q}), & \mathbf{q} \in H(\operatorname{div}; \Omega), \\ b(\mathbf{j} - \tilde{\mathbf{j}}_h, v) - c(u - \tilde{u}_h, v) &= -(f - \Pi_0 f, v)_0, & v \in L^2(\Omega), \end{aligned}$$

where the residual r is given by $r(\mathbf{q}) := -a(\tilde{\mathbf{j}}_h, \mathbf{q}) - b(\mathbf{q}, \tilde{u}_h)$, $\mathbf{q} \in H(\operatorname{div}; \Omega)$, and $\Pi_0 f$ denotes the L^2 -projection of f onto $W_0(\Omega; \mathcal{T}_k)$.

The basic idea behind the construction of a residual based error estimator for the total error $\|u - \tilde{u}_h\|_0 + \|\mathbf{j} - \tilde{\mathbf{j}}_h\|_{\operatorname{div}}$ is to use a Helmholtz decomposition of $H(\operatorname{div}; \Omega)$. We obtain an error estimator which can be easily calculated by means of the available finite element approximation $(\tilde{\mathbf{j}}_h, \tilde{u}_h)$. In contrast to the hierarchical error estimator, no additional subproblem has to be solved. This is the main advantage of the residual based error estimator $\hat{\eta}_R$ which is determined by its local contributions:

$$(2.2) \quad \begin{aligned} \hat{\eta}_R^2 &:= \sum_{T \in \mathcal{T}_k} \hat{\eta}_{R;T}^2, \\ \hat{\eta}_{R;T}^2 &:= \|f - \Pi_0 f\|_{0;T}^2 + h_T^2 \|a^{-1} \tilde{\mathbf{j}}_h\|_{0;T}^2 + \sum_{i=1}^3 w_i \alpha_{e_i} h_{e_i} \|[a^{-1} \tilde{\mathbf{j}}_h \cdot \mathbf{t}_{e_i}]_J\|_{0;e_i}^2. \end{aligned}$$

Here, the weighting factors α_{e_i} and w_i are defined by $\alpha_{e_i} := \frac{1}{4}(\alpha_0^{T_1} + \alpha_0^{T_2} + \alpha_1^{T_1} + \alpha_1^{T_2})$ and $w_i := 1/2$, if $e_i = \partial T_1 \cap \partial T_2$ is an interior edge, and by $\alpha_{e_i} := \frac{1}{2}(\alpha_0^{T_e} + \alpha_1^{T_e})$ and $w_i := 1$, if $e_i \subset \partial T_e \cap \partial \Omega$. The jump $[\cdot]_J$ on e has to be defined as

$$[a^{-1} \tilde{\mathbf{j}}_h \cdot \mathbf{t}_e]_J := (a^{-1} \tilde{\mathbf{j}}_h \cdot \mathbf{t}_e|_{T_i} - a^{-1} \tilde{\mathbf{j}}_h \cdot \mathbf{t}_e|_{T_a}), \quad \partial T_i \cap \partial T_a = e,$$




FIGURE 2.1. Orientation of \mathbf{n}_e and \mathbf{t}_e

if e is an interior edge, and

$$[a^{-1}\tilde{\mathbf{j}}_h \cdot \mathbf{t}_e]_J := (a^{-1}\tilde{\mathbf{j}}_h \cdot \mathbf{t}_e|_{T_i}), \quad \partial T_i \cap \partial\Omega \supset e,$$

if $e \in \partial\Omega$. For the orientation of \mathbf{t}_e , we refer to Figure 2.1.

This error estimator provides an upper and a lower bound for the total error $\|u - \tilde{u}_h\|_0 + \|\mathbf{j} - \tilde{\mathbf{j}}_h\|_{\text{div}}$, if the gridsize and the iteration error are small enough.

Theorem 2.1. *There exist constants $h_{\max} > 0$ and $c_R, C_R, C_{it} > 0$, independent of the refinement level, such that for all $h \leq h_{\max}$*

$$c_R \hat{\eta}_R^2 \leq \|u - \tilde{u}_h\|_0^2 + \|\mathbf{j} - \tilde{\mathbf{j}}_h\|_{\text{div}}^2 \leq C_R \hat{\eta}_R^2 + C_{it} (\|u_h - \tilde{u}_h\|_0^2 + \|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\text{div}}^2).$$

In the case of standard conforming finite element discretizations, the hierarchical basis error estimator has been investigated by Deuffhard, Leinen, Yserentant [14]. An excellent overview is given by Bornemann et al. [8]. Recently, this concept has been generalized by Achchab et al. [1] for the mixed setting, but no easily accessible local error estimator is proposed. Here, the introduction of the error estimator is based on the principle of defect correction in higher order ansatz spaces. By means of appropriate localization and decoupling techniques of the flux ansatz space, we obtain an easily computable, efficient and reliable a posteriori error estimator for the flux error and the total error. The hierarchical error estimator presented in this section has been investigated in [19, 20, 31]. We summarize the main ideas and the basic results; for details we refer to the papers cited above.

The hierarchical basis error estimator is based on a discrete defect problem considered on appropriately chosen higher dimensional ansatz spaces. There are two different approaches. Either we consider the same mixed ansatz spaces associated with a finer triangulation, e.g., obtained by uniform refinement from the actual one, or we use higher order mixed ansatz spaces providing improved a priori estimates. Here, we restrict ourselves to the second approach, and consider the Raviart-Thomas ansatz space $RT_1(\Omega; \mathcal{T}_k)$. In contrast to the residual based error estimator, the continuous defect problem (2.1) will not be considered. Instead, to obtain an appropriate approximation of (2.1) we use the higher dimensional Raviart-Thomas ansatz space $RT_1(\Omega; \mathcal{T}_k)$ for the flux and the ansatz space $W_1(\Omega; \mathcal{T}_k)$ of piecewise linear functions for the primal variable. We restrict ourselves to the discrete saddle point problem which requires the computation of a pair $(e_j, e_u) \in RT_1(\Omega; \mathcal{T}_k) \times W_1(\Omega; \mathcal{T}_k)$ such that

$$(2.3) \quad \begin{aligned} a(e_j, \mathbf{q}) + b(\mathbf{q}, e_u) &= r(\mathbf{q}), & \mathbf{q} \in RT_1(\Omega; \mathcal{T}_k), \\ b(e_j, v) - c(e_u, v) &= -(f - \Pi_0 f, v)_0, & v \in W_1(\Omega; \mathcal{T}_k). \end{aligned}$$

Denoting by $(\mathbf{j}_{RT_1}, u_{RT_1})$ the solution of the discrete variational problem (1.4) on $RT_1(\Omega; \mathcal{T}_k) \times W_1(\Omega; \mathcal{T}_k)$, the introduction and the analysis of the error estimator

are based on the following saturation assumptions:

$$(S1 \text{ a}) \quad \|\mathbf{j} - \mathbf{j}_{RT_1}\|_{\text{div}} \leq \beta_k \|\mathbf{j} - \mathbf{j}_h\|_{\text{div}}, \quad \beta_k > 0,$$

$$(S1 \text{ b}) \quad \|\mathbf{j} - \mathbf{j}_{RT_1}\|_{\text{div}}^2 + \|u - u_{RT_1}\|_0^2 \leq \hat{\beta}_k (\|\mathbf{j} - \mathbf{j}_h\|_{\text{div}}^2 + \|u - u_h\|_0^2), \quad \hat{\beta}_k > 0,$$

with $\beta_k \leq \beta_\infty < 1$ and $\hat{\beta}_k \leq \hat{\beta}_\infty < 1$. These saturation assumptions are motivated by the well known a priori error estimates for $\mathbf{j} - \mathbf{j}_h$, $\mathbf{j} - \mathbf{j}_{RT_1}$ and $u - u_h$, $u - u_{RT_1}$ (see, e.g., [12]). In particular, the saturation assumption (S1 a) implies both an upper and a lower bound for the total error in terms of the $\|\cdot\|_{\text{div}}$ -norms of e_j and the iteration error $\mathbf{j}_h - \tilde{\mathbf{j}}_h$:

$$(2.4) \quad \begin{aligned} (1 + \beta_\infty)^{-1} (\|e_j\|_{\text{div}} - \beta_\infty \|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\text{div}}) &\leq \|\mathbf{j} - \tilde{\mathbf{j}}_h\|_{\text{div}}, \\ (1 - \beta_\infty)^{-1} (\|e_j\|_{\text{div}} + \beta_\infty \|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\text{div}}) &\geq \|\mathbf{j} - \tilde{\mathbf{j}}_h\|_{\text{div}}. \end{aligned}$$

Therefore, only the solution of (2.3) has to be examined. The approximation of (e_j, e_u) is based on the hierarchical two-level splitting of the mixed ansatz spaces $RT_1(\Omega; \mathcal{T}_k)$ and $W_1(\Omega; \mathcal{T}_k)$. By means of the L^2 -projection Π of $W_1(\Omega; \mathcal{T}_k)$ onto $W_0(\Omega; \mathcal{T}_k)$ and the interpolation operator $\rho : RT_1(\Omega; \mathcal{T}_k) \rightarrow RT_0(\Omega; \mathcal{T}_k)$ given locally by

$$\int_e \mathbf{n}_e \cdot (\rho \mathbf{q}) \, d\sigma = \int_e \mathbf{n}_e \cdot \mathbf{q} \, d\sigma, \quad e \in \mathcal{E}_k,$$

we obtain a hierarchical splitting of $W_1(\Omega; \mathcal{T}_k)$ and $RT_1(\Omega; \mathcal{T}_k)$ according to

$$(2.5) \quad W_1(\Omega; \mathcal{T}_k) = W_0(\Omega; \mathcal{T}_k) \oplus \widehat{W}_1(\Omega; \mathcal{T}_k),$$

$$(2.6) \quad RT_1(\Omega; \mathcal{T}_k) = RT_0(\Omega; \mathcal{T}_k) \oplus \widehat{RT}_1(\Omega; \mathcal{T}_k),$$

where $\widehat{W}_1(\Omega; \mathcal{T}_k) := (\text{Id} - \Pi) W_1(\Omega; \mathcal{T}_k)$ and $\widehat{RT}_1(\Omega; \mathcal{T}_k) := (\text{Id} - \rho) RT_1(\Omega; \mathcal{T}_k)$. The hierarchical surplus in the flux, $\widehat{RT}_1(\Omega; \mathcal{T}_k)$, can be further decomposed into a divergence free part $\widehat{RT}_1^0(\Omega; \mathcal{T}_k)$ and its complement $\widehat{RT}_1^1(\Omega; \mathcal{T}_k)$:

$$(2.7) \quad \widehat{RT}_1(\Omega; \mathcal{T}_k) = \widehat{RT}_1^0(\Omega; \mathcal{T}_k) \oplus \widehat{RT}_1^1(\Omega; \mathcal{T}_k),$$

where

$$\begin{aligned} \widehat{RT}_1^0(\Omega; \mathcal{T}_k) &:= \mathbf{curl} \hat{S}_2(\Omega; \mathcal{T}_k), \\ \widehat{RT}_1^1(\Omega; \mathcal{T}_k) &:= \{\mathbf{q} \in \widehat{RT}_1(\Omega; \mathcal{T}_k) \mid \mathbf{n}_e \cdot \mathbf{q}|_e = 0, \, e \in \mathcal{E}_k\}. \end{aligned}$$

Here, $\hat{S}_2(\Omega; \mathcal{T}_k)$ stands for the space of quadratic bubble functions associated with the midpoints of the edges. This space is given in terms of the hierarchical two-level splitting $S_2(\Omega; \mathcal{T}_k) = S_1(\Omega; \mathcal{T}_k) \oplus \hat{S}_2(\Omega; \mathcal{T}_k)$, where $S_1(\Omega; \mathcal{T}_k)$ and $S_2(\Omega; \mathcal{T}_k)$ refer to the conforming ansatz spaces associated with the standard P1 and P2 approximations. The structure of the decomposition (2.6) and (2.7) is symbolized in Figure 2.2. The splitting (2.7) is somewhat similar to the Helmholtz decomposition of the $H(\text{div}; \Omega)$ ansatz space that will be used in case of the residual based error estimator (see Section 3 below). But in contrast to this Helmholtz decomposition, the ansatz spaces $\widehat{RT}_1^0(\Omega; \mathcal{T}_k)$ and $\widehat{RT}_1^1(\Omega; \mathcal{T}_k)$ are not orthogonal with respect to the bilinear form $a(\cdot, \cdot)$. Furthermore, it is easy to see that $\widehat{RT}_1^1(\Omega; \mathcal{T}_k)$ can be written




FIGURE 2.2. Degrees of freedom of the subspaces of the higher order Raviart-Thomas space

as the direct sum of local two-dimensional subspaces $\widehat{RT}_1^1(T)$ which correspond to the “interior” degrees of freedom of $RT_1(T)$:

$$\widehat{RT}_1^1(\Omega; \mathcal{T}_k) = \bigoplus_{T \in \mathcal{T}_k} \widehat{RT}_1^1(T), \quad \widehat{RT}_1^1(T) := \left\{ \mathbf{q} \in \widehat{RT}_1^1(\Omega; \mathcal{T}_k) \mid \mathbf{q}|_{\Omega \setminus T} = 0 \right\}.$$

Due to the special structure of $W_1(\Omega; \mathcal{T}_k)$, the subspace $\widehat{W}_1(\Omega; \mathcal{T}_k)$ can be decomposed into the direct sum of local two-dimensional subspaces:

$$\widehat{W}_1(\Omega; \mathcal{T}_k) = \bigoplus_{T \in \mathcal{T}_k} \widehat{W}_1(T), \quad \widehat{W}_1(T) := \left\{ v \in \widehat{W}_1(\Omega; \mathcal{T}_k) \mid v|_{\Omega \setminus T} = 0 \right\}.$$

For the definition of the error estimator we have to consider two different types of local low dimensional variational problems associated with the two parts $\widehat{RT}_1^0(\Omega; \mathcal{T}_k)$ and $\widehat{RT}_1^1(\Omega; \mathcal{T}_k)$ of the hierarchical surplus. For each edge of the triangulation we have to solve a single equation

$$(2.8) \quad a(\mathbf{curl} \varphi_e, \mathbf{curl} \Phi_e) = r(\mathbf{curl} \Phi_e),$$

where Φ_e is the quadratic bubble function associated with the edge e and $\varphi_e \in \text{span}\{\Phi_e\}$. The second variational problem is associated with the elements T . For each element we have to consider a 4×4 saddle point problem: Find $(\tilde{e}_{\mathbf{j}_1}^1, \tilde{e}_{u_1}) \in \widehat{RT}_1^1(T) \times \widehat{W}_1(T)$ such that

$$(2.9) \quad \begin{aligned} a|_T(\tilde{e}_{\mathbf{j}_1}^1, \mathbf{q}) + b|_T(\mathbf{q}, \tilde{e}_{u_1}) &= r|_T(\mathbf{q}), & \mathbf{q} \in \widehat{RT}_1^1(T), \\ b|_T(\tilde{e}_{\mathbf{j}_1}^1, v) - c|_T(\tilde{e}_{u_1}, v) &= (f, v)_{0;T}, & v \in \widehat{W}_1(T). \end{aligned}$$

The solutions of (2.8) and (2.9) lead to the local definition of an a posteriori flux-oriented error estimator η_H :

$$(2.10) \quad \begin{aligned} \eta_H^2 &:= \sum_{T \in \mathcal{T}_k} \eta_{H;T}^2, \\ \eta_{H;T}^2 &:= \|\tilde{e}_{\mathbf{j}_1}^1\|_{\text{div};T}^2 + \sum_{i=1}^3 w_i \|\mathbf{curl} \varphi_{e_i}\|_{\text{div}}^2, \quad T \in \mathcal{T}_k, \end{aligned}$$

where the weighting factor w_i is defined as in the case of the residual based error estimator.

Theorem 2.2. *Under the saturation assumption (S1 a) with $0 < \beta_k < \beta_\infty < 1$ there exist constants $c_{\text{hier}}, C_{\text{hier}} > 0$ and $\gamma_{\text{hier}}, \Gamma_{\text{hier}} > 0$, independent of the*

refinement level, such that

$$(2.11) \quad \begin{aligned} (1 + \beta_\infty)^{-1} (c_{hier}\eta_H - \gamma_{hier}\|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\text{div}}) &\leq \|\mathbf{j} - \tilde{\mathbf{j}}_h\|_{\text{div}}, \\ (1 - \beta_\infty)^{-1} (C_{hier}\eta_H + \Gamma_{hier}\|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\text{div}}) &\geq \|\mathbf{j} - \tilde{\mathbf{j}}_h\|_{\text{div}}. \end{aligned}$$

Next, we consider an error estimator η_L based on the solution of local subproblems that is strongly related to the hierarchical error estimator. In the standard conforming setting, this kind of error estimator is due to Bank and Weiser [6]. For nonconforming techniques we refer to [21]. It relies on a defect correction with respect to a higher order ansatz space and an appropriate localization based on a hierarchical two-level splitting. It turns out that the estimator can be computed elementwise by the solution of local Dirichlet problems similar to the original global one (cf. [21]). In contrast to the standard conforming and nonconforming setting, the boundary data cannot be obtained by a simple averaging of the computed approximation.

For simplicity, we only consider the discretization error $(\mathbf{j} - \mathbf{j}_h, u - u_h)$ and not the total error $(\mathbf{j} - \tilde{\mathbf{j}}_h, u - \tilde{u}_h)$. Note that the results can be generalized to the total error. This time, we state the defect problem as a local Dirichlet problem for each element $T \in \mathcal{T}_k$:

$$(2.12) \quad \begin{aligned} a|_T(\mathbf{j} - \mathbf{j}_h, \mathbf{q}) + b|_T(\mathbf{q}, u - u_h) &= - \int_{\partial T} u \mathbf{n} \cdot \mathbf{q} \, d\sigma + r|_T(\mathbf{q}), & \mathbf{q} \in H(\text{div}; T), \\ b|_T(\mathbf{j} - \mathbf{j}_h, v) - c|_T(u - u_h, v) &= -(f - \Pi_0 f, v)_{0;T}, & v \in L^2(T). \end{aligned}$$

As in the case of the hierarchical error estimator, we only look for an adequate approximation of the solution of (2.12). The original ansatz space $H(\text{div}; T) \times L^2(T)$ will be replaced by $RT_1(T) \times P_1(T)$, and the Dirichlet data u by some appropriate approximation u_D which can be easily calculated from (\mathbf{j}_h, u_h) . A possible choice of u_D will be discussed in Section 5. In contrast to the hierarchical error estimator, we need an additional saturation assumption concerning the approximation of the Dirichlet data. We refer to $RT_1^{-1}(\Omega; \mathcal{T}_k)$ as the nonconforming ansatz space, where

$$RT_l^{-1}(\Omega; \mathcal{T}_k) := \{\mathbf{q} \in (L^2(\Omega))^2 \mid \mathbf{q}|_T \in RT_l(T), \quad T \in \mathcal{T}_k\}, \quad 0 \leq l \leq 1,$$

and to λ_{RT_1} as the Lagrange multiplier which is a piecewise linear function on the edges uniquely determined by means of $\mathbf{q} \in RT_1^{-1}(\Omega; \mathcal{T}_k)$,

$$\sum_{T \in \mathcal{T}_k} \int_{\partial T} \lambda_{RT_1} \mathbf{n} \cdot \mathbf{q} \, d\sigma = - \sum_{T \in \mathcal{T}_k} \int_T a^{-1} \mathbf{j}_{RT_1} \cdot \mathbf{q} \, dx + \sum_{T \in \mathcal{T}_k} \int_T \text{div} \, \mathbf{q} u_{RT_1} \, dx.$$

Then, introducing the weighted norm $\|\cdot\|_{0; \hat{\mathcal{E}}_k^{-1}}$ according to

$$\|v\|_{0; \hat{\mathcal{E}}_k^{-1}} := \left(\sum_{e \in \hat{\mathcal{E}}_k} h_e^{-1} \alpha_e \|v\|_{0; e}^2 \right)^{1/2}, \quad v \in L^2(\mathcal{E}_k),$$

we assume

$$(S2) \quad \|\lambda_{RT_1} - [u_D]_A\|_{0; \hat{\mathcal{E}}_k^{-1}} \leq \beta_{2; k} \|\mathbf{j} - \mathbf{j}_h\|_{\text{div}}, \quad \beta_{2; k} > 0,$$

to hold with $\beta_{2; k} \leq \beta_{2; \infty} < \infty$, where the average $[v]_A$ on e , $v \in \prod_{T \in \mathcal{T}_k} H^1(T)$, is given by $[v]_A := \frac{1}{2}(v|_{T_i} + v|_{T_a})$, $\partial T_i \cap \partial T_a = e$, if e is an interior edge, and by $[v]_A := \frac{1}{2}(v|_{T_i})$, $\partial T_i \cap \partial \Omega \supset e$, if $e \subset \partial \Omega$. In general, assumption (S2) is

motivated by an adequate a priori error estimate or by some equivalence results (see subsection 7.2). Because of the saturation assumptions (S1 a) and (S2), the previous simplifications are justified. We consider the discretized defect problem

$$(2.13) \quad \begin{aligned} a|_T(\hat{e}_j|_T, \mathbf{q}) + b|_T(\mathbf{q}, \hat{e}_u|_T) &= - \int_{\partial T} [u_D]_A \mathbf{n} \cdot \mathbf{q} \, d\sigma + r|_T(\mathbf{q}), & \mathbf{q} \in RT_1(T), \\ b|_T(\hat{e}_j|_T, v) - c|_T(\hat{e}_u|_T, v) &= -(f - \Pi_0, v)_{0;T}, & v \in P_1(T). \end{aligned}$$

Then $\hat{e}_j := (\hat{e}_j|_T)_{T \in \mathcal{T}_k}$ does guarantee lower and upper bounds for the error in the flux.

Theorem 2.3. *Under the saturation assumptions (S1 a) with $0 < \beta_k < \beta_\infty < 1$ and (S2) with $0 < \beta_{2;k} < \beta_{2;\infty} < \infty$ there exist constants $c_{loc}, C_{loc} > 0$, independent of the refinement level, such that*

$$(2.14) \quad c_{loc} \|\hat{e}_j\|_{\text{div}} \leq \|\mathbf{j} - \mathbf{j}_h\|_{\text{div}} \leq (1 - \beta_\infty)^{-1} C_{loc} \|\hat{e}_j\|_{\text{div}}.$$

Finally, we use the same hierarchical splittings and decoupling techniques as before. For each element we thus obtain three scalar equations and one saddle point problem that have to be solved. The error estimator η_L is defined by

$$(2.15) \quad \begin{aligned} \eta_L^2 &:= \sum_{T \in \mathcal{T}_k} \eta_{L;T}^2, \\ \eta_{L;T}^2 &:= \|\hat{e}_{\mathbf{j}_1}^1\|_{\text{div};T}^2 + \sum_{i=1}^3 a|_T(\mathbf{curl} \varphi_{e_i;T}, \mathbf{curl} \varphi_{e_i;T}), \quad T \in \mathcal{T}_k, \end{aligned}$$

where $\hat{e}_{\mathbf{j}_1}^1$ and $\mathbf{curl} \varphi_{e_i;T}$ are the solutions of the local problems on the element T .

Theorem 2.4. *Let the saturation assumptions (S1 a) with $\beta_\infty < 1$ and (S2) with $\beta_{2;\infty} < \infty$ be satisfied. Then, there exist constants $0 < c_{loc} \leq C_{loc}$, independent of the refinement level, such that*

$$(2.16) \quad c_{loc} \eta_L \leq \|\mathbf{j} - \mathbf{j}_h\|_{\text{div}} \leq C_{loc} \eta_L.$$

In Section 5, we will show that $\hat{e}_{\mathbf{j}_1}^1 = \tilde{e}_{\mathbf{j}_1}^1$, and thus the first part of the error estimator is exactly the same as in the case of the hierarchical basis error estimator.

Finally, in Section 6 we propose an estimator η_S for the error in the primal variable that can be motivated by a superconvergence result for the technique of mixed hybridization. In contrast to the hierarchical basis error estimator η_H and the error estimator η_L based on the solution of local subproblems, we do not have to solve additional defect problems. In the standard conforming case, error estimators obtained by some postprocessing of the approximation have been introduced by Zienkiewicz and Zhu in [32, 34] and have been further analyzed by Rodriguez [24]. In the mixed setting there is some work of Brandts [11]. In contrast to the conforming situation, we will be able to prove the equivalence between η_S and the L^2 -norm of the difference of \tilde{u}_h and some higher order finite element approximation which turns out to be the solution of a modified nonconforming variational approach. Therefore, we have a strong relationship between the error estimator and a discrete defect problem which is solved in a higher order ansatz space. Details will be given

in Section 6. The error estimator η_S is given as follows:

$$(2.17) \quad \begin{aligned} \eta_S^2 &:= \sum_{T \in \mathcal{T}_k} \eta_{S;T}^2, \\ \eta_{S;T}^2 &:= \sum_{i=1}^3 w_i h_{e_i}^2 ([\tilde{u}_h]_J|_{e_i})^2, \end{aligned}$$

where the weighting factor w_i is defined as before.

Theorem 2.5. *Let $\tilde{u}_h \in W_0(\Omega; \mathcal{T}_k)$ be an approximation of the primal variable u obtained by an iterative solution process for system (1.4). Then, there exist constants $0 < \sigma_0 \leq \sigma_1$ and $0 < C_0 < C_1$, depending only on the shape regularity of \mathcal{T}_k and the ellipticity constants in (1.2), such that*

$$(2.18) \quad \sigma_0 \eta_S - C_0 \|u_h - \tilde{u}_h\|_0 \leq \|u - \tilde{u}_h\|_0 \leq \sigma_1 \eta_S + C_1 \|u_h - \tilde{u}_h\|_0.$$

We emphasize that the computation of the estimator only involves the jumps of the piecewise constant approximation for the primal variable across the inner element boundaries.

In [11], Brandts introduced an a posteriori error estimator for the flux in the L^2 -norm based on a higher order recovering of the flux. If we add this estimator and $\|f - \Pi_0 f\|_0$ to η_S , it turns out that we again obtain an error estimator $\hat{\eta}_S$ for the total error.

The error estimator in [11] is defined as

$$\|\mathbf{j}_h - K\mathbf{j}_h\|_0,$$

where the linear operator $K : RT_0(\Omega; \mathcal{T}_k) \rightarrow (CR(\Omega; \mathcal{T}_k))^2$ is locally given by its value at the midpoint m_e of the edges of the triangulation

$$(2.19a) \quad K\mathbf{p}(m_e) := \frac{1}{2} (\mathbf{p}|_{T_1} + \mathbf{p}|_{T_2})(m_e), \quad T_1 \cap T_2 = e \in \mathcal{E}_k,$$

if e is an interior edge, and by

$$(2.19b) \quad \mathbf{t} \cdot (K\mathbf{p})(m_e) := 0, \quad \mathbf{n} \cdot (K\mathbf{p})(m_e) := \mathbf{n} \cdot \mathbf{p}|_{T_e}, \quad e \subset \partial T_e,$$

if e is an edge on the boundary $\partial\Omega$.

The error estimator $\hat{\eta}_S$ is then given by

$$(2.20) \quad \begin{aligned} \hat{\eta}_S^2 &:= \sum_{T \in \mathcal{T}_k} \hat{\eta}_{S;T}^2, \\ \hat{\eta}_{S;T}^2 &:= \eta_{S;T}^2 + \|\mathbf{j}_h - K\mathbf{j}_h\|_{0;T}^2 + \|f - \Pi_0 f\|_{0;T}^2. \end{aligned}$$

In subsection 7.3, it will be shown that $\hat{\eta}_S$ and the residual based error estimator $\hat{\eta}_R$ are globally equivalent.

3. THE RESIDUAL BASED ERROR ESTIMATOR

First, we summarize some technical results which are an indispensable tool for the investigation of the residual based error estimator. We consider the projection operator $P_C : L^2(\Omega) \rightarrow S_1(\Omega; \mathcal{T}_k)$ with respect to a discrete L^2 -norm due to Clément [13]. In contrast to the Lagrangian interpolation operator, this operator can be applied to discontinuous functions. Denoting by $\tilde{D}_p := \bigcup\{T \in \mathcal{T}_k \mid p \in \partial T\}$ the union of all triangles containing the vertex p and by λ_p the linear conforming nodal basis function, i.e., $\lambda_p(p') = \delta_{p,p'}$, it is defined by

$$P_C v := \sum_{p \in \mathcal{P}_k} (q_p(v))(p) \lambda_p, \quad v \in L^2(\Omega),$$

where $q_p(v) \in P_1(\tilde{D}_p)$, $p \in \mathcal{P}_k$, is the projection of v onto the space $P_1(\tilde{D}_p)$,

$$(q_p, q)_{0, \tilde{D}_p} = (v, q)_{0, \tilde{D}_p}, \quad q \in P_1(\tilde{D}_p).$$

P_C has the following properties:

Lemma 3.1. *Let $v \in H^1(\Omega)$, $T \in \mathcal{T}_k$ and $e \in \mathcal{E}_k$, $k \geq 0$. Then, there exist constants $C_{0;C}$, $C_{1;C}$, $\hat{C}_{0;C} > 0$, independent of the refinement level, such that*

$$(3.1a) \quad \|v - P_C v\|_{0;T} \leq h_T C_{0;C} |v|_{1; \tilde{D}_T},$$

$$(3.1b) \quad \|\hat{a}^{1/2} \nabla P_C v\|_{0;T} \leq C_{1;C} \|\hat{a}^{1/2} \nabla v\|_{0; \tilde{D}_T},$$

$$\hat{a} = (\hat{a}_{ij})_{i,j=1}^2, \quad \hat{a}_{11} := (a^{-1})_{22}, \quad \hat{a}_{22} := (a^{-1})_{11}, \quad \hat{a}_{12} = \hat{a}_{21} := 0,$$

$$(3.1c) \quad \|v - P_C v\|_{0;e} \leq \sqrt{\alpha_e h_e} \hat{C}_{0;C} \|\hat{a}^{-1/2} \nabla v\|_{0; \tilde{D}_e}$$

where $\tilde{D}_e := \bigcup \{T \in \mathcal{T}_k \mid e \cap \partial T \neq \emptyset\}$, $\tilde{D}_T := \bigcup \{T' \in \mathcal{T}_k \mid \partial T' \cap \partial T \neq \emptyset\}$, $\alpha_e := \frac{1}{4} (\alpha_0^{T_1} + \alpha_0^{T_2} + \alpha_1^{T_1} + \alpha_1^{T_2})$, $\partial T_1 \cap \partial T_2 = e \subset \Omega$ and $\alpha_e := \frac{1}{2} (\alpha_0^{T_e} + \alpha_1^{T_e})$, $e \subset \partial T_e \cap \partial \Omega$.

For the proof we refer to [13] and remark that the matrix \hat{a} is strictly positive definite.

We shall also take advantage of the following approximation property of the L^2 -projection Π_0 onto the space of piecewise constant functions:

$$(3.2) \quad \|v - \Pi_0 v\|_{0;T} \leq C_{proj} h_T |v|_{1;T}, \quad v \in H^1(T), \quad T \in \mathcal{T}_k,$$

where C_{proj} stands for a positive constant independent of the refinement level.

We further need two different types of bubble functions Φ_T , $T \in \mathcal{T}_k$, and Φ_e , $e \in \mathcal{E}_k$, associated with the element T and the edge e , respectively. Denoting by $\lambda_{p_i}^T$, $p_i \in \mathcal{P}_k \cap \partial T$, $1 \leq i \leq 3$, the barycentric coordinates of $T \in \mathcal{T}_k$ and by $p_i^e \in \mathcal{P}_k \cap e$, $1 \leq i \leq 2$, the vertices of the edge $e \subset \partial T$, they are defined as follows:

$$(3.3) \quad \Phi_T := 27 \lambda_{p_1}^T \lambda_{p_2}^T \lambda_{p_3}^T, \quad \Phi_e := 4 \lambda_{p_1^e}^T \lambda_{p_2^e}^T.$$

Lemma 3.2. (i) *Let $\mathbf{p} \in (P_1(T))^2$, $T \in \mathcal{T}_k$, $k \geq 0$. Then, there exist constants $C_{0;D}$, $C_{1;D} > 0$ independent of $T \in \mathcal{T}_k$ such that*

$$(3.4a) \quad \|\mathbf{p}\|_{0;T}^2 \leq C_{0;D}^2 \int_T \mathbf{p} \cdot \Phi_T \mathbf{p} \, dx,$$

$$(3.4b) \quad \|\operatorname{div}(\Phi_T \mathbf{p})\|_{0;T} \leq h_T^{-1} C_{1;D} \|\mathbf{p}\|_{0;T}.$$

(ii) *Let $p \in P_1(e)$, $e \in \mathcal{E}_k$, $k \geq 0$. Then, there exist constants $C_{0;K}$, $C_{1;K} > 0$, independent of $e \in \mathcal{E}_k$, such that*

$$(3.5a) \quad \|p\|_{0;e}^2 \leq C_{0;K}^2 \int_e p^2 \Phi_e \, d\sigma,$$

$$(3.5b) \quad \|\hat{a}^{-1/2} \nabla (p^{D_e} \Phi_e)\|_{0;D_e} \leq (\alpha_e h_e)^{-1/2} C_{1;K} \|p\|_{0;e},$$

$$D_e := \bigcup \{T \in \mathcal{T}_k \mid e \subset \partial T\},$$

where p^{D_e} defines a prolongation of p on Ω , $e_T \subset \partial T$ fixed:

$$(3.6) \quad p^{D_e}(x) := \begin{cases} 0, & x \notin D_e, \\ p(x_e), & x \in T \subset D_e \text{ and } x_e \in e, (x - x_e) \parallel e_T \neq e. \end{cases}$$

The proof is an easy consequence of the affine equivalence of the elements and the inverse inequality for polynomials. Note that the constants depend only on the initial triangulation and on the ratio of the local upper and lower bounds of a . The same type of result is used in [27].

The investigation of the a posteriori error estimator will be provided in several steps. In a first step, we decompose the flux ansatz space $H(\operatorname{div}; \Omega)$ into a solenoidal and a weakly irrotational part:

$$(3.7) \quad H(\operatorname{div}; \Omega) = H^0(\operatorname{div}; \Omega) \oplus H^1(\operatorname{div}; \Omega),$$

where $H^0(\operatorname{div}; \Omega) := \{\mathbf{q} \in H(\operatorname{div}; \Omega) \mid \operatorname{div} \mathbf{q} = 0\}$ and the orthogonal complement is defined by $H^1(\operatorname{div}; \Omega) := \{\mathbf{q} \in H(\operatorname{div}; \Omega) \mid a(\mathbf{q}, \mathbf{q}^0) = 0, \mathbf{q}^0 \in H^0(\operatorname{div}; \Omega)\}$. According to this splitting, the flux error can be written as $\mathbf{j}_e = \mathbf{j}_e^0 + \mathbf{j}_e^1$, where $\mathbf{j}_e^0 \in H^0(\operatorname{div}; \Omega)$ and $\mathbf{j}_e^1 \in H^1(\operatorname{div}; \Omega)$. Note that this representation is unique. Now, we study the subspaces $H^0(\operatorname{div}; \Omega)$ and $H^1(\operatorname{div}; \Omega)$ in more detail.

Lemma 3.3. *The following properties of $H^0(\operatorname{div}; \Omega)$ and $H^1(\operatorname{div}; \Omega)$ hold true:*

- (i) *For each element $\mathbf{q} \in H^0(\operatorname{div}; \Omega)$ there exists a scalar function $\phi \in H^1(\Omega)$ such that*

$$(3.8) \quad \mathbf{q} = \operatorname{curl} \phi := \begin{pmatrix} \phi_y \\ -\phi_x \end{pmatrix}.$$

- (ii) *On $H^1(\operatorname{div}; \Omega)$ the following norm equivalence holds true:*

$$(3.9) \quad c_{\operatorname{div}} \|\operatorname{div} \mathbf{q}\|_0 \leq \|\mathbf{q}\|_{\operatorname{div}} \leq C_{\operatorname{div}} \|\operatorname{div} \mathbf{q}\|_0$$

with constants $0 < c_{\operatorname{div}} < C_{\operatorname{div}}$ independent of $\mathbf{q} \in H^1(\operatorname{div}; \Omega)$.

Proof. The proof of the first assertion can be found in [18, Thm. 3.4], whereas the second assertion is an easy consequence of [12, Prop. 1.2]. \square

The construction of the a posteriori error estimator is mainly based on the preceding splitting of the flux ansatz space. The variational problem (2.1) consists of two independent subproblems. The first subproblem is associated with the solenoidal subspace and gives rise to a positive definite problem:

$$a(\mathbf{j}_e^0, \mathbf{q}) = r(\mathbf{q}), \quad \mathbf{q} \in H^0(\operatorname{div}; \Omega).$$

Introducing the weighted norm $\|\cdot\|_{0, \mathcal{E}_k}$ according to

$$\|v\|_{0, \mathcal{E}_k}^2 := \sum_{e \in \mathcal{E}_k} \alpha_e h_e \|v\|_{0, e}^2, \quad v \in L^2(\mathcal{E}_k),$$

we will show that $\|[a^{-1} \tilde{\mathbf{j}}_h \cdot \mathbf{t}_e]_J\|_{0, \mathcal{E}_k}$ yields sharp upper and lower bounds for the solenoidal part of the flux error, provided the iteration error is small enough. The existence of a nonvanishing Helmholtz term and the indefiniteness of the saddle point problem (2.1) do not influence the construction of the bounds.

Lemma 3.4. *There exist constants $c_{\mathbf{j}_e^0}, C_{\mathbf{j}_e^0}, C_{it}^0 > 0$, independent of the refinement level, such that*

$$c_{\mathbf{j}_e^0} \|[a^{-1} \tilde{\mathbf{j}}_h \cdot \mathbf{t}_e]_J\|_{0, \mathcal{E}_k} \leq \|\mathbf{j}_e^0\|_{\operatorname{div}} \leq C_{\mathbf{j}_e^0} \|[a^{-1} \tilde{\mathbf{j}}_h \cdot \mathbf{t}_e]_J\|_{0, \mathcal{E}_k} + C_{it}^0 \|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\operatorname{div}},$$

where $c_{\mathbf{j}_e^0} := (\sqrt{3}C_{0;K}^2 C_{1;K})^{-1}$, $C_{\mathbf{j}_e^0} := \sqrt{\kappa_K} \hat{C}_{0;C}$ and $C_{it}^0 := \sqrt{\kappa_D} C_{1;C}$.

Proof. Following the same lines as in the conforming setting [27], we evaluate the residual as a continuous linear functional restricted to $H^0(\operatorname{div}; \Omega)$. Let $\mathbf{q} = \mathbf{curl} \phi \in H^0(\operatorname{div}; \Omega)$; then

$$\begin{aligned}
r(\mathbf{q}) &= -a(\tilde{\mathbf{J}}_h, \mathbf{q}) = -\int_{\Omega} a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{q} \, dx = -\int_{\Omega} a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{curl} \phi \, dx \\
&= \sum_{T \in \mathcal{T}_k} \int_T \left(a_{1,1}^{-1} (\tilde{\mathbf{J}}_h)_{1,y} - a_{2,2}^{-1} (\tilde{\mathbf{J}}_h)_{2,x} \right) \phi \, dx \\
&\quad - \sum_{T \in \mathcal{T}_k} \int_{\partial T} \left(a_{1,1}^{-1} (\tilde{\mathbf{J}}_h)_1 \mathbf{n}_y - a_{2,2}^{-1} (\tilde{\mathbf{J}}_h)_2 \mathbf{n}_x \right) \phi \, d\sigma \\
&= \sum_{T \in \mathcal{T}_k} \int_{\partial T} (a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}) \phi \, d\sigma = \sum_{e \in \mathcal{E}_k} \int_e ([a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J) \phi \, d\sigma.
\end{aligned}$$

By means of Clément's projection P_C and the fact that $a(\mathbf{j}_h, \mathbf{q}) = 0$ for $\mathbf{q} \in RT_0(\Omega; \mathcal{T}_k) \cap H^0(\operatorname{div}; \Omega)$, we obtain an upper bound for the solenoidal part of the flux error. Let $\mathbf{j}_e^0 = \mathbf{curl} \psi$. Then, observing (1.6), (3.1b) and (3.1c), an upper bound for $\|\mathbf{j}_e^0\|_{\operatorname{div}}$ results from

(3.10)

$$\begin{aligned}
\|\mathbf{j}_e^0\|_{\operatorname{div}}^2 &= r(\mathbf{j}_e^0) = \sum_{e \in \mathcal{E}_k} \int_e ([a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J) \psi \, d\sigma \\
&= \sum_{e \in \mathcal{E}_k} \int_e ([a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J) (\psi - P_C \psi) \, d\sigma + \sum_{e \in \mathcal{E}_k} \int_e [a^{-1} (\tilde{\mathbf{J}}_h - \mathbf{j}_h) \cdot \mathbf{t}_e]_J P_C \psi \, d\sigma \\
&\leq \sum_{e \in \mathcal{E}_k} \left(\alpha_e h_e \int_e [a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J^2 \, d\sigma \right)^{1/2} \left(\frac{1}{\alpha_e h_e} \int_e (\psi - P_C \psi)^2 \, d\sigma \right)^{1/2} \\
&\quad + a(\mathbf{j}_h - \tilde{\mathbf{J}}_h, \mathbf{curl} P_C \psi) \\
&\leq \|\mathbf{j}_e^0\|_{\operatorname{div}} \left(\sqrt{\kappa_K} \hat{C}_{0,C} \| [a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J \|_{0;\mathcal{E}_k} + C_{1,C} \sqrt{\kappa_D} \|\tilde{\mathbf{J}}_h - \mathbf{j}_h\|_{\operatorname{div}} \right).
\end{aligned}$$

On the other hand, taking into account (3.5a) and (3.5b), a lower bound for $\|\mathbf{j}_e^0\|_{\operatorname{div}}$ can be established by means of the quadratic bubble functions Φ_e associated with the midpoints of the edges (cf. (3.3)):

$$\begin{aligned}
C_{0,K}^{-2} \| [a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J \|_{0;e}^2 &\leq \int_e [a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J^2 \Phi_e \, d\sigma \\
&= \int_e [a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J \cdot ([a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J \Phi_e) \, d\sigma \\
&= - \int_{D_e} \mathbf{curl} \cdot \left([a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J^{D_e} \Phi_e \right) \cdot (a^{-1} (\tilde{\mathbf{J}}_h - \mathbf{j})) \, dx \\
&\leq \| a^{-1/2} \mathbf{j}_e^0 \|_{0;D_e} \cdot \| \hat{a}^{-1/2} \nabla \left([a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J^{D_e} \Phi_e \right) \|_{0;D_e} \\
&\leq (\alpha_e h_e)^{-1/2} C_{1,K} \| a^{-1/2} \mathbf{j}_e^0 \|_{0;D_e} \cdot \| [a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J \|_{0;e},
\end{aligned}$$

whence

$$(3.11) \quad \| [a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J \|_{0;\mathcal{E}_k} \leq \sqrt{3} C_{0,K}^2 C_{1,K} \|\mathbf{j}_e^0\|_{\operatorname{div}}.$$

We recall that the extension $[a^{-1} \tilde{\mathbf{J}}_h \cdot \mathbf{t}_e]_J^{D_e}$ is defined according to (3.6). The assertion is an immediate consequence of (3.10) and (3.11). \square

In order to obtain a sharp estimate for the total flux error, in a second step we have to consider \mathbf{j}_e^1 . It is easy to see that in the special case of a vanishing Helmholtz term

$$\int_{\Omega} \operatorname{div}(\mathbf{j} - \tilde{\mathbf{j}}_h)v \, dx = \int_{\Omega} (f - \Pi_0 f)v \, dx, \quad v \in L^2(\Omega)$$

whence $\|\operatorname{div}(\mathbf{j} - \tilde{\mathbf{j}}_h)\|_0 = \|f - \Pi_0 f\|_0$. Because $\operatorname{div} \mathbf{j}_e = \operatorname{div} \mathbf{j}_e^1$, (3.9) readily provides a lower and an upper bound for $\|\mathbf{j}_e^1\|_{\operatorname{div}}$.

However, in the general case $b \neq 0$ the situation is more complicated, and the errors $\|u_e\|_0$ in the primal variable u and in the divergence part of the flux error $\|\mathbf{j}_e^1\|_{\operatorname{div}}$ are coupled.

Lemma 3.5. *The following inequalities hold true:*

$$\begin{aligned} \|\operatorname{div} \mathbf{j}_e^1\|_{0;T} &\leq \|f - \Pi_0 f\|_{0;T} + b_1^T \|u_e\|_{0;T}, \\ \|f - \Pi_0 f\|_{0;T} &\leq b_1^T \|u_e\|_{0;T} + \|\operatorname{div} \mathbf{j}_e^1\|_{0;T}. \end{aligned}$$

Proof. The second equation of the variational problem (2.1) states that

$$\int_{\Omega} \operatorname{div} \mathbf{j}_e v \, dx + \int_{\Omega} b u_e v \, dx = \int_{\Omega} (f - \Pi_0 f)v \, dx, \quad v \in L^2(\Omega),$$

and we conclude by a straightforward application of the triangle inequality. \square

Finally, we will focus our attention on the error in the primal variable with respect to the L^2 -norm $\|u - \tilde{u}_h\|_0$. For that purpose, we consider the L^2 -projection of u onto the space of piecewise constant functions and use the following result:

$$(3.12) \quad \|\Pi_0 u - u_h\|_0^2 \leq C_{sup}^2 \sum_{T \in \mathcal{T}_k} h_T^2 \|a^{-1}(\mathbf{j} - \mathbf{j}_h)\|_{\operatorname{div};T}^2,$$

where the constant C_{sup} only depends on the geometry of the initial triangulation and on the ratio of the local bounds of the coefficients in (1.1). In case of the Poisson equation, this result is well established (cf., e.g., [2, 12]). In the general case, it can be proved assuming a discrete H^2 -regularity and using some duality techniques (cf., e.g., [12, Remark 2.16]).

Lemma 3.6. *There exist constants c_{u_e} , $c_{\mathbf{j}_e}$, C_{u_e} , $C_{\mathbf{j}_e}$, $C_{it}^1 > 0$, independent of the refinement level, such that*

$$\begin{aligned} c_{u_e} \|u_e\|_0 &\leq \|a^{-1} \tilde{\mathbf{j}}_h\|_{0;\mathcal{T}_k} + c_{\mathbf{j}_e} \|a^{-1} \mathbf{j}_e\|_{\operatorname{div};\mathcal{T}_k} \\ &\quad + C_{it}^1 (\|u_h - \tilde{u}_h\|_0 + \|a^{-1}(\mathbf{j}_h - \tilde{\mathbf{j}}_h)\|_{\operatorname{div};\mathcal{T}_k}), \\ \|a^{-1} \tilde{\mathbf{j}}_h\|_{0;\mathcal{T}_k} &\leq (C_{u_e} \|u_e\|_0 + C_{\mathbf{j}_e} \|a^{-1} \mathbf{j}_e\|_{0;\mathcal{T}_k}), \end{aligned}$$

where the weighted norms are given by $\|\cdot\|_{0;\mathcal{T}_k}^2 := \sum_{T \in \mathcal{T}_k} h_T^2 \|\cdot\|_{0;T}^2$ and $\|\cdot\|_{\operatorname{div};\mathcal{T}_k}^2 := \sum_{T \in \mathcal{T}_k} h_T^2 \|\cdot\|_{\operatorname{div};T}^2$.

Proof. Since $\mathbf{j} = -a\nabla u$, in view of (3.3) and (3.12) we obtain

$$\begin{aligned} \|u - \tilde{u}_h\|_0 &\leq \|u - \Pi_0 u\|_0 + \|\Pi_0 u - \tilde{u}_h\|_0 \\ &\leq C_{proj} \left(\sum_{T \in \mathcal{T}_k} h_T^2 \|\nabla u\|_{0;T}^2 \right)^{1/2} + C_{sup} \|a^{-1}(\mathbf{j} - \mathbf{j}_h)\|_{\text{div};\mathcal{T}_k} + \|u_h - \tilde{u}_h\|_0 \\ &\leq C_{proj} (\|a^{-1}\tilde{\mathbf{j}}_h\|_{0;\mathcal{T}_k} + \|a^{-1}(\mathbf{j} - \tilde{\mathbf{j}}_h)\|_{0;\mathcal{T}_k}) \\ &\quad + C_{sup} (\|a^{-1}(\mathbf{j} - \tilde{\mathbf{j}}_h)\|_{\text{div};\mathcal{T}_k} + \|a^{-1}(\mathbf{j}_h - \tilde{\mathbf{j}}_h)\|_{\text{div};\mathcal{T}_k}) + \|u_h - \tilde{u}_h\|_0. \end{aligned}$$

The constants $c_{\mathbf{j}_e}$, c_{u_e} and $C_{it}^1 > 0$ are defined as $c_{\mathbf{j}_e} := 1 + C_{sup} C_{proj}^{-1}$, $c_{u_e} := C_{proj}^{-1}$ and $C_{it}^1 := C_{proj}^{-1} \max(1, C_{sup})$.

It remains to establish an upper bound for $\|a^{-1}\tilde{\mathbf{j}}_h\|_{0;\mathcal{T}_k}$. This can be achieved using the cubic bubble function ϕ_T (cf. (3.3)) and observing (3.4a), (3.4b):

$$\begin{aligned} C_{0;D}^{-2} \int_T |a^{-1}\tilde{\mathbf{j}}_h|^2 dx &\leq \int_T (a^{-1}\tilde{\mathbf{j}}_h + \nabla u) \cdot (\phi_T a^{-1}\tilde{\mathbf{j}}_h) dx - \int_T \nabla u \cdot (\phi_T a^{-1}\tilde{\mathbf{j}}_h) dx \\ &= \int_T \text{div}(\phi_T a^{-1}\tilde{\mathbf{j}}_h)(u - \tilde{u}_h) dx + \int_T a^{-1}(\tilde{\mathbf{j}}_h - \mathbf{j}) \cdot (\phi_T a^{-1}\tilde{\mathbf{j}}_h) dx \\ &\leq \|\tilde{u}_h - u\|_{0;T} \cdot \|\text{div}(\phi_T a^{-1}\tilde{\mathbf{j}}_h)\|_{0;T} + \|a^{-1}(\tilde{\mathbf{j}}_h - \mathbf{j})\|_{0;T} \cdot \|\phi_T a^{-1}\tilde{\mathbf{j}}_h\|_{0;T} \\ &\leq (h_T^{-1} C_{1;D}) \|\tilde{u}_h - u\|_{0;T} + \|a^{-1}(\tilde{\mathbf{j}}_h - \mathbf{j})\|_{0;T} \|a^{-1}\tilde{\mathbf{j}}_h\|_{0;T}. \end{aligned}$$

This local inequality holds true for all elements $T \in \mathcal{T}_k$. Therefore, we obtain the global estimate

$$\|a^{-1}\tilde{\mathbf{j}}_h\|_{0;\mathcal{T}_k} \leq C_{u_e} \|\tilde{u}_h - u\|_0 + C_{\mathbf{j}_e} \|a^{-1}(\mathbf{j} - \tilde{\mathbf{j}}_h)\|_{0;\mathcal{T}_k},$$

where $C_{u_e} := \sqrt{2} C_{0;D}^2 C_{1;D}$ and $C_{\mathbf{j}_e} := \sqrt{2} C_{0;D}^2$. \square

If we take into account the definition of the residual based error estimator given by (2.2) and the results of Lemmas 3.4, 3.5, and 3.6, the proof of Theorem 2.1 can be readily given:

Proof of Theorem 2.1. By means of Lemmas 3.4, 3.5, and 3.6, we get

$$\begin{aligned} \|f - \Pi_0 f\|_0^2 &+ \|[a^{-1}\tilde{\mathbf{j}}_h \mathbf{t}_e]_J\|_{0;\mathcal{E}_k} + \|a^{-1}\tilde{\mathbf{j}}_h\|_{0;\mathcal{T}_k} \\ &\leq 2(b_1^2 + C_{u_e}^2) \|u_e\|_0^2 + \left(c_{\mathbf{j}_e}^{-2} + 2h_0^2 \alpha_0^{-1} C_{\mathbf{j}_e}^2 \right) \|\mathbf{j}_e^0\|_{\text{div}}^2 \\ &\quad + 2 \max\left(1, h_0^2 \alpha_0^{-1} C_{\mathbf{j}_e}^2\right) \|\mathbf{j}_e^1\|_{\text{div}}^2. \end{aligned}$$

Hence, the first inequality of Theorem 2.1 holds true with

$$c_R^{-1} := \max\left(2, \left(c_{\mathbf{j}_e}^{-2} + 2h_0^2 \alpha_0^{-1} C_{\mathbf{j}_e}^2\right), 2(b_1^2 + C_{u_e}^2)\right).$$

To establish the upper bound for the error, we need an additional assumption on h . Assuming

$$h_0 \leq h_{\max} := \left(\frac{3}{\alpha_0} \left(\frac{c_{\mathbf{j}_e}}{c_{u_e}} \right)^2 \max(1, \alpha_0^{-1}) (1 + 4C_{div}^2 b_1^2 \max(1, \alpha_0^{-1})) \right)^{-1/2},$$

by means of Lemmas 3.4, 3.5, and 3.6 we obtain

$$\begin{aligned}
\|u_e\|_0^2 + \|\mathbf{j}_e\|_{\text{div}}^2 &= \|u_e\|_0^2 + \|\mathbf{j}_e^0\|_{\text{div}}^2 + \|\mathbf{j}_e^1\|_{\text{div}}^2 \\
&\leq 4C_{\mathbf{j}_e^0}^2 \|[a^{-1}\tilde{\mathbf{j}}_h \cdot \mathbf{t}_e]_J\|_{0;\mathcal{E}_k}^2 + 4C_{\text{div}}^2 \max(1, \alpha_0^{-1}) \|f - \Pi_0 f\|_0^2 \\
&\quad + 3c_{u_e}^{-2} (1 + 4b_1^2 C_{\text{div}}^2 \max(1, \alpha_0^{-1})) \|a^{-1}\tilde{\mathbf{j}}_h\|_{0;\mathcal{T}_k}^2 + 4(C_{it}^0)^2 \|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\text{div}}^2 \\
&\quad + 3(c_{u_e}^{-1} C_{it}^1)^2 (1 + 4b_1^2 C_{\text{div}}^2 \max(1, \alpha_0^{-1})) (\|u_h - \tilde{u}_h\|_0 + \|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\text{div};\mathcal{T}_k})^2.
\end{aligned}$$

Then, the upper bound holds true with

$$C_R := \max\left(3c_{u_e}^{-2} (1 + 4b_1^2 C_{\text{div}}^2 \max(1, \alpha_0^{-1})), 4C_{\mathbf{j}_e^0}^2, 4C_{\text{div}}^2 \max(1, \alpha_0^{-1})\right)$$

and

$$\begin{aligned}
C_{it} := 2 \max &\left(3 \left(\frac{C_{it}^1}{c_{u_e}}\right)^2 (1 + 4b_1^2 C_{\text{div}}^2 \max(1, \alpha_0^{-1})), \right. \\
&\left. 2(C_{it}^0)^2 + \left(\frac{C_{it}^1}{c_{\mathbf{j}_e}} \frac{\max(1, \alpha_1)}{\max(1, \alpha_0^{-1})}\right)^2\right).
\end{aligned}$$

□

Remark. Note that we even get an upper bound if

$$(3.13) \quad h_0 < C_\infty := \left(\sqrt{\frac{6}{\alpha_0}} b_1 C_{\text{div}} \frac{c_{\mathbf{j}_e}}{c_{u_e}} \max(1, \alpha_0^{-1})\right)^{-1}.$$

In case $b \equiv 0$, we define $C_\infty := \infty$, and thus we have no limitation on h_0 . If h_0 tends to C_∞ , the upper bounds in the theorem tend to ∞ as well.

4. THE HIERARCHICAL BASIS ERROR ESTIMATOR

As indicated in Section 2, the idea behind the hierarchical type a posteriori error estimator consists in an approximation of the error equation with respect to the higher order mixed ansatz spaces $RT_1(\Omega; \mathcal{T}_k)$ and $W_1(\Omega; \mathcal{T}_k)$, followed by an appropriate localization in terms of a hierarchical two-level splitting. In particular, introducing the local spaces

$$\widehat{RT}_1^0(T) := \left\{ \mathbf{p} \in RT_1(T) \mid \mathbf{p} = \mathbf{q}|_T, \mathbf{q} \in \widehat{RT}_1^0(\Omega; \mathcal{T}_k) \right\}, \quad T \in \mathcal{T}_k, k \geq 0,$$

and $\widehat{RT}_1(T) := \widehat{RT}_1^0(T) \oplus \widehat{RT}_1^1(T)$, where $\widehat{RT}_1^1(T)$ is given as in Section 2, the following strengthened Cauchy-Schwarz inequalities (4.1) will play an important role for the derivation of the error estimator:

(4.1)

$$\begin{aligned}
a|_T(\mathbf{q}^0, \mathbf{q}^1)^2 &\leq \eta_0^2 a|_T(\mathbf{q}^0, \mathbf{q}^0) a|_T(\mathbf{q}^1, \mathbf{q}^1), \quad \mathbf{q}^0 \in RT_0(T), \mathbf{q}^1 \in \widehat{RT}_1(T), \\
a|_T(\hat{\mathbf{q}}^0, \hat{\mathbf{q}}^1)^2 &\leq \eta_1^2 a|_T(\hat{\mathbf{q}}^0, \hat{\mathbf{q}}^0) a|_T(\hat{\mathbf{q}}^1, \hat{\mathbf{q}}^1), \quad \hat{\mathbf{q}}^\nu \in \widehat{RT}_1^\nu(T), 0 \leq \nu \leq 1,
\end{aligned}$$

with $\eta_\nu^2 < 1$, $0 \leq \nu \leq 1$, being independent of the refinement level. The inequalities (4.1) are an easy consequence of the affine equivalence of the Raviart-Thomas elements and the shape regularity of the triangulations.

The system (2.3) cannot be solved locally, and therefore the approximation (e_j, e_u) is not suited for an easily computable error estimator. Consequently, the main idea of our proposed error estimator consists in the replacement of the original

bilinear form $a(\cdot, \cdot)$ by a modified bilinear form $\tilde{a}(\cdot, \cdot)$. According to the splitting of $RT_1(\Omega; \mathcal{T}_k)$ the vector fields $\mathbf{q}, \mathbf{p} \in RT_1(\Omega; \mathcal{T}_k)$ are decomposed as follows:

$$\mathbf{q} = \mathbf{q}^0 + \hat{\mathbf{q}}^0 + \hat{\mathbf{q}}^1, \quad \mathbf{p} = \mathbf{p}^0 + \hat{\mathbf{p}}^0 + \hat{\mathbf{p}}^1,$$

where $\mathbf{q}^0, \mathbf{p}^0 \in RT_0(\Omega; \mathcal{T}_k)$, $\hat{\mathbf{q}}^0, \hat{\mathbf{p}}^0 \in \widehat{RT}_1^0(\Omega; \mathcal{T}_k)$ and $\hat{\mathbf{q}}^1, \hat{\mathbf{p}}^1 \in \widehat{RT}_1^1(\Omega; \mathcal{T}_k)$. Then, the modified bilinear form $\tilde{a}(\mathbf{q}, \mathbf{p})$ is defined as follows:

$$\tilde{a}(\mathbf{q}, \mathbf{p}) := a(\mathbf{q}^0, \mathbf{p}^0) + a(\hat{\mathbf{q}}^0, \hat{\mathbf{p}}^0) + a(\hat{\mathbf{q}}^1, \hat{\mathbf{p}}^1).$$

It is easy to see that $\tilde{a}(\cdot, \cdot)$ is orthogonal with respect to the decomposition of the ansatz space of the flux. A simple consequence of the strengthened Cauchy-Schwarz inequalities (4.1) and Young's inequality is the equivalence of the bilinear forms $a(\cdot, \cdot)$ and $\tilde{a}(\cdot, \cdot)$:

$$(4.2) \quad c_{RT}^2 \tilde{a}|_T(\mathbf{q}, \mathbf{q}) \leq a|_T(\mathbf{q}, \mathbf{q}) \leq C_{RT}^2 \tilde{a}|_T(\mathbf{q}, \mathbf{q}), \quad \mathbf{q} \in RT_1(T),$$

with constants $0 < c_{RT} \leq C_{RT}$ independent of $T \in \mathcal{T}_k$, $k \geq 1$. A detailed proof of (4.2) can be found in [32, Lemma 4.6]. Now, we consider a modified discrete variational problem which is obtained from (2.3) if we replace the bilinear form $a(\cdot, \cdot)$ by $\tilde{a}(\cdot, \cdot)$: Find $(\tilde{e}_j, \tilde{e}_u) \in RT_1(\Omega; \mathcal{T}_k) \times W_1(\Omega; \mathcal{T}_k)$ such that

$$(4.3) \quad \begin{aligned} \tilde{a}(\tilde{e}_j, \mathbf{q}) + b(\mathbf{q}, \tilde{e}_u) &= r(\mathbf{q}), & \mathbf{q} \in RT_1(\Omega; \mathcal{T}_k), \\ b(\tilde{e}_j, v) - c(\tilde{e}_u, v) &= -(f - \Pi_0 f, v)_0, & v \in W_1(\Omega; \mathcal{T}_k). \end{aligned}$$

The following theorem states the equivalence of the solutions of the saddle point problems (2.3) and (4.3). By means of (2.4) and Theorem 2.2, $\|\tilde{e}_j\|_{\text{div}}$ provides sharp upper and lower bounds for the error in the flux.

Theorem 4.1. *The solutions (e_j, e_u) and $(\tilde{e}_j, \tilde{e}_u) \in RT_1(\Omega; \mathcal{T}_k) \times W_1(\Omega; \mathcal{T}_k)$ of the discrete variational problems (2.3) and (4.3) are equivalent in the sense that there exist constants $0 < c_{j,RT} \leq C_{j,RT}$ and $c_{d,RT}, c_{u,RT}, C_{d,RT}, C_{u,RT} > 0$, independent of \mathcal{T}_k , $k \geq 1$, such that*

$$(4.4a) \quad c_{j,RT}^2 \tilde{a}(\tilde{e}_j, \tilde{e}_j) \leq a(e_j, e_j) \leq C_{j,RT}^2 \tilde{a}(\tilde{e}_j, \tilde{e}_j),$$

$$(4.4b) \quad \|\text{div } \tilde{e}_j\|_0 \leq \|\text{div } e_j\|_0 + c_{d,RT} a(e_j, e_j)^{1/2},$$

$$\|\text{div } e_j\|_0 \leq \|\text{div } \tilde{e}_j\|_0 + C_{d,RT} \tilde{a}(\tilde{e}_j, \tilde{e}_j)^{1/2},$$

$$(4.4c) \quad \|\tilde{e}_u\|_0 \leq \|e_u\|_0 + c_{u,RT} a(e_j, e_j)^{1/2},$$

$$\|e_u\|_0 \leq \|\tilde{e}_u\|_0 + C_{u,RT} \tilde{a}(\tilde{e}_j, \tilde{e}_j)^{1/2}.$$

Proof. In view of (2.3) and (4.3) we compare the solutions e_j and \tilde{e}_j and obtain

$$a(e_j - \tilde{e}_j, e_j - \tilde{e}_j) + c(e_u - \tilde{e}_u, e_u - \tilde{e}_u) = \tilde{a}(\tilde{e}_j, e_j - \tilde{e}_j) - a(\tilde{e}_j, e_j - \tilde{e}_j)$$

as well as

$$\tilde{a}(e_j - \tilde{e}_j, e_j - \tilde{e}_j) + c(e_u - \tilde{e}_u, e_u - \tilde{e}_u) = \tilde{a}(e_j, e_j - \tilde{e}_j) - a(e_j, e_j - \tilde{e}_j).$$

Observing (4.2) and the fact that $c(\cdot, \cdot)$ is positive semidefinite, by straightforward calculations we get

$$\begin{aligned} a(e_j - \tilde{e}_j, e_j - \tilde{e}_j)^{1/2} &\leq \left(C_{RT} + \frac{1}{c_{RT}} \right) \tilde{a}(\tilde{e}_j, \tilde{e}_j)^{1/2}, \\ \tilde{a}(e_j - \tilde{e}_j, e_j - \tilde{e}_j)^{1/2} &\leq \left(C_{RT} + \frac{1}{c_{RT}} \right) a(e_j, e_j)^{1/2}. \end{aligned}$$

The triangle inequality applied to $a(e_j, e_j)^{1/2}$ and $\tilde{a}(\tilde{e}_j, \tilde{e}_j)^{1/2}$ proves (4.4a) with $C_{j;RT} := 2C_{RT} + c_{RT}^{-1}$ and $c_{j;RT} := (C_{RT} + 2c_{RT}^{-1})^{-1}$.

Recalling that $\operatorname{div} RT_1(\Omega; \mathcal{T}_k) = W_1(\Omega; \mathcal{T}_k)$ and

$$b(\tilde{e}_j - e_j, v) = c(\tilde{e}_u - e_u, v), \quad v \in W_1(\Omega; \mathcal{T}_k),$$

we see that (4.4b) follows from

$$\begin{aligned} \|\operatorname{div}(\tilde{e}_j - e_j)\|_0 &= \sup_{\substack{v \in W_1(\Omega; \mathcal{T}_k) \\ v \neq 0}} \frac{b(\tilde{e}_j - e_j, v)}{\|v\|_0; \Omega} \leq \sqrt{\beta_1} c(e_u - \tilde{e}_u, e_u - \tilde{e}_u)^{1/2} \\ &= \sqrt{\beta_1} (\tilde{a}(e_j - \tilde{e}_j, \tilde{e}_j) - a(e_j - \tilde{e}_j, e_j))^{1/2} \\ &\leq \sqrt{\beta_1} \sqrt{2(C_{RT} + c_{RT}^{-1})} \tilde{a}(\tilde{e}_j, \tilde{e}_j)^{1/4} a(e_j, e_j)^{1/4}. \end{aligned}$$

By means of the triangle inequality and (4.4a) we conclude with that $C_{d;RT}^2 := 2\beta_1 C_{j;RT} (C_{RT} + c_{RT}^{-1})$ and $c_{d;RT}^2 := 2c_{j;RT}^{-1} \beta_1 (C_{RT} + c_{RT}^{-1})$.

For the proof of (4.4c) we note that for all $\mathbf{q} \in RT_1(\Omega; \mathcal{T}_k)$ we have

$$(4.5) \quad b(\mathbf{q}, e_u - \tilde{e}_u) = \tilde{a}(\tilde{e}_j, \mathbf{q}) - a(e_j, \mathbf{q}).$$

Taking into account the inequality

$$\|v\|_0 \leq \beta_{\operatorname{div}} \sup_{\substack{\mathbf{q} \in RT_1(\Omega; \mathcal{T}_k) \\ \mathbf{q} \neq 0}} \frac{b(\mathbf{q}, v)}{\|\mathbf{q}\|_{\operatorname{div}}}, \quad v \in W_1(\Omega; \mathcal{T}_k),$$

and (4.2), (4.4a) and (4.5), we obtain

$$\begin{aligned} \|e_u - \tilde{e}_u\|_0 &\leq \alpha_0^{-1/2} \beta_{\operatorname{div}} (1 + c_{RT}^{-1} c_{j;RT}^{-1}) a(e_j, e_j)^{1/2}, \\ \|e_u - \tilde{e}_u\|_0 &\leq \alpha_0^{-1/2} \beta_{\operatorname{div}} (C_{j;RT} + c_{RT}^{-1}) \tilde{a}(\tilde{e}_j, \tilde{e}_j)^{1/2}. \end{aligned}$$

□

A simple consequence of Theorem 4.1 is the existence of constants $0 < c_{a-1} \leq C_{a-1}$ such that

$$(4.6) \quad c_{a-1}^2 (\tilde{a}(\tilde{e}_j, \tilde{e}_j) + \|\operatorname{div} \tilde{e}_j\|_{0; \Omega}^2) \leq \|e_j\|_{\operatorname{div}}^2 \leq C_{a-1}^2 (\tilde{a}(\tilde{e}_j, \tilde{e}_j) + \|\operatorname{div} \tilde{e}_j\|_{0; \Omega}^2),$$

where $c_{a-1}^{-2} := \max(2, 2c_{d;RT}^2 + c_{j;RT}^{-2})$ and $C_{a-1}^2 := \max(2, 2C_{d;RT}^2 + C_{j;RT}^2)$.

To prove the upper and lower bounds for the error estimator defined by (2.10), we have to show the equivalence between $\tilde{a}(\tilde{e}_j, \tilde{e}_j) + \|\operatorname{div} \tilde{e}_j\|_{0; \Omega}^2$ and η_H^2 . A first step in this direction is to consider the saddle point problem (4.3) in more detail. According to the hierarchical splitting of the spaces $RT_1(\Omega; \mathcal{T}_k)$ and $W_1(\Omega; \mathcal{T}_k)$, (4.3) can be rewritten in terms of three independent subproblems. We decompose \tilde{e}_j and \tilde{e}_u as follows:

$$\begin{aligned} \tilde{e}_j &= \tilde{e}_{j_0} + \tilde{e}_{j_1}^0 + \tilde{e}_{j_1}^1, \quad \tilde{e}_{j_0} \in RT_0(\Omega; \mathcal{T}_k), \quad \tilde{e}_{j_1}^0 \in \widehat{RT}_1^0(\Omega; \mathcal{T}_k), \quad \tilde{e}_{j_1}^1 \in \widehat{RT}_1^1(\Omega; \mathcal{T}_k), \\ \tilde{e}_u &= \tilde{e}_{u_0} + \tilde{e}_{u_1}, \quad \tilde{e}_{u_0} \in W_0(\Omega; \mathcal{T}_k), \quad \tilde{e}_{u_1} \in \widehat{W}_1(\Omega; \mathcal{T}_k), \end{aligned}$$

and obtain three variational problems:

$$(4.7) \quad \begin{aligned} a(\tilde{e}_{j_0}, \mathbf{q}) + b(\mathbf{q}, \tilde{e}_{u_0}) &= r(\mathbf{q}), \quad \mathbf{q} \in RT_0(\Omega; \mathcal{T}_k), \\ b(\tilde{e}_{j_0}, v) - c(\tilde{e}_{u_0}, v) &= 0, \quad v \in W_0(\Omega; \mathcal{T}_k), \end{aligned}$$

$$(4.8) \quad a(\tilde{e}_{j_1}^0, \mathbf{q}) = r(\mathbf{q}), \quad \mathbf{q} \in \widehat{RT}_1^0(\Omega; \mathcal{T}_k),$$

$$(4.9) \quad \begin{aligned} a\left(\tilde{\mathbf{e}}_{\mathbf{j}_1}^1, \mathbf{q}\right) + b(\mathbf{q}, \tilde{e}_{u_1}) &= r(\mathbf{q}), & \mathbf{q} \in \widehat{RT}_1^1(\Omega; \mathcal{T}_k), \\ b\left(\tilde{\mathbf{e}}_{\mathbf{j}_1}^1, v\right) - c(\tilde{e}_{u_1}, v) &= -(f, v)_0, & v \in \widehat{W}_1(\Omega; \mathcal{T}_k). \end{aligned}$$

Again $r(\mathbf{q})$ denotes the residual. The solution $(\tilde{e}_{\mathbf{j}_0}, \tilde{e}_{u_0})$ of (4.7) is zero only in case of a vanishing iteration error $(\tilde{\mathbf{j}}_h - \mathbf{j}_h, \tilde{u}_h - u_h)$. An upper bound for $a(\tilde{e}_{\mathbf{j}_0}, \tilde{e}_{\mathbf{j}_0})$ and $\|\operatorname{div} \tilde{e}_{\mathbf{j}_0}\|_{0;\Omega}^2$ can be easily established by means of the iteration error.

The variational problems (4.7) and (4.9) are indefinite saddle point problems. On the other hand, (4.8) represents a symmetric and positive definite system which can be decoupled by well known standard techniques (cf. [14]). The bilinear form $a(\cdot, \cdot)$ applied to the discrete space $\widehat{RT}_1^0(\Omega; \mathcal{T}_k) \times \widehat{RT}_1^0(\Omega; \mathcal{T}_k)$ can be seen as a bilinear form $\hat{a}(\cdot, \cdot)$ on $\hat{S}_2(\Omega; \mathcal{T}_k) \times \hat{S}_2(\Omega; \mathcal{T}_k)$. Let $\mathbf{q}, \mathbf{p} \in \widehat{RT}_1^0(\Omega; \mathcal{T}_k)$ and $\phi, \psi \in \hat{S}_2(\Omega; \mathcal{T}_k)$ with $\mathbf{q} = \operatorname{curl} \phi$ and $\mathbf{p} = \operatorname{curl} \psi$; then

$$(4.10) \quad a(\mathbf{q}, \mathbf{p}) = \sum_{T \in \mathcal{T}_k} \int_T a^{-1} \operatorname{curl} \phi \operatorname{curl} \psi \, dx = \sum_{T \in \mathcal{T}_k} \int_T \hat{a} \nabla \phi \nabla \psi \, dx =: \hat{a}(\phi, \psi).$$

The matrix \hat{a} is defined by $\hat{a}_{11} := (a^{-1})_{22}$, $\hat{a}_{22} := (a^{-1})_{11}$ and $\hat{a}_{12} = \hat{a}_{21} := -(a^{-1})_{12}$, and has the same eigenvalues as a^{-1} . Each element $\phi \in \hat{S}_2(\Omega; \mathcal{T}_k)$ can be written as the direct sum of quadratic bubble functions associated with the edges of the triangulations according to $\phi = \sum_{e \in \mathcal{E}_k} \phi_e$, where ϕ_e , $e \in \mathcal{E}_k$, is a multiple of the nodal basis function Φ_e . The inequality

$$(4.11) \quad c_{\operatorname{curl}} \sum_{i=1}^3 \hat{a}|_T(\phi_{e_i}, \phi_{e_i}) \leq \hat{a}|_T(\phi, \phi) \leq C_{\operatorname{curl}} \sum_{i=1}^3 \hat{a}|_T(\phi_{e_i}, \phi_{e_i}), \quad T \in \mathcal{T}_k,$$

with constants $0 < c_{\operatorname{curl}} \leq C_{\operatorname{curl}}$ states that the global problem (4.8) can be replaced by local ones. The coupling between different bubble functions is neglected and a single equation has to be solved for each edge (cf. (2.8) of Section 2):

$$\hat{a}(\varphi_e, \Phi_e) = r(\operatorname{curl} \Phi_e).$$

Then, equivalence of $\sum_{e \in \mathcal{E}_k} \hat{a}(\varphi_e, \varphi_e)$ and $a(\tilde{\mathbf{e}}_{\mathbf{j}_1}^0, \tilde{\mathbf{e}}_{\mathbf{j}_1}^0)$ is guaranteed by (4.11).

It remains to examine (4.9). Due to the special structure of $\widehat{RT}_1^1(\Omega; \mathcal{T}_k)$ and $\widehat{W}_1(\Omega; \mathcal{T}_k)$, the global problem consists of independent local subproblems associated with the elements of the triangulation. For each element we have to solve the 4×4 saddle point problem (2.9), which can be further reduced to two 2×2 problems using an L^2 -orthogonal basis of $\widehat{W}_1(T)$.

For the proof of Theorem 2.2 it remains to be shown that the flux of the solution of subproblem (4.7) is bounded by the iteration error independently of the refinement level. For this purpose we reconsider the residual:

$$\begin{aligned} r(\tilde{e}_{\mathbf{j}_0}) &= a(\tilde{e}_{\mathbf{j}_0}, \tilde{e}_{\mathbf{j}_0}) + c(\tilde{e}_{u_0}, \tilde{e}_{u_0}) = -a(\tilde{\mathbf{j}}_h, \tilde{e}_{\mathbf{j}_0}) - b(\tilde{e}_{\mathbf{j}_0}, \tilde{u}_h) \\ &= a(\mathbf{j}_h - \tilde{\mathbf{j}}_h, \tilde{e}_{\mathbf{j}_0}) + b(\tilde{e}_{\mathbf{j}_0}, u_h - \tilde{u}_h) = a(\mathbf{j}_h - \tilde{\mathbf{j}}_h, \tilde{e}_{\mathbf{j}_0}) + c(\tilde{e}_{u_0}, u_h - \tilde{u}_h). \end{aligned}$$

In view of the equalities

$$c(u_h - \tilde{u}_h, u_h - \tilde{u}_h) = b(\mathbf{j}_h - \tilde{\mathbf{j}}_h, u_h - \tilde{u}_h) \quad \text{and} \quad \|\operatorname{div}(\tilde{e}_{\mathbf{j}_0})\|_{0;\Omega}^2 = b(\tilde{e}_{\mathbf{j}_0}, \operatorname{div}(\tilde{e}_{\mathbf{j}_0})),$$

we obtain

$$\begin{aligned} a(\tilde{e}_{\mathbf{j}_0}, \tilde{e}_{\mathbf{j}_0}) + c(\tilde{e}_{u_0}, \tilde{e}_{u_0}) &\leq \|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\operatorname{div}}^2, \\ \|\operatorname{div}(\tilde{e}_{\mathbf{j}_0})\|_{0;\Omega}^2 &\leq \beta_1 \|\mathbf{j}_h - \tilde{\mathbf{j}}_h\|_{\operatorname{div}}^2. \end{aligned}$$

Proof of Theorem 2.2. Because of (2.4), (4.6) and (4.11), the assertion (2.11) is a direct consequence of the definition of the error estimator with constants $C_{hier} := C_{a-1} \max(1, c_{curl}^{-1/2})$ and $\Gamma_{hier} := \beta_\infty \max(1, \sqrt{\beta_1}) + C_{a-1} \sqrt{1 + \beta_1}$ for the upper bounds of the error. The constants for the lower bounds are given by $c_{hier} := c_{a-1} \min(1, C_{curl}^{-1/2})$ and $\gamma_{hier} := \beta_\infty \max(1, \sqrt{\beta_1})$. \square

Remark. If we are interested in an error estimator for the error in the flux and in the primal variable, we have to take into account an additional term in the definition of the error estimator $\hat{\eta}_H$:

$$(4.12) \quad \begin{aligned} \hat{\eta}_H^2 &:= \sum_{T \in \mathcal{T}_k} \hat{\eta}_{H;T}^2, \\ \hat{\eta}_{H;T}^2 &:= \eta_{H;T}^2 + \|\tilde{e}_{u_1}\|_{0;T}^2, \quad T \in \mathcal{T}_k. \end{aligned}$$

It is easy to see that the saturation assumption (S1 b) and (4.4c) as well as (2.11) guarantee that the error estimator $\hat{\eta}_H$ provides sharp lower and upper bounds for the total error $(\|u - \tilde{u}_h\|_0^2 + \|\mathbf{j} - \tilde{\mathbf{j}}_h\|_{\text{div}}^2)^{1/2}$, if the iteration error is small enough.

5. AN ERROR ESTIMATOR BASED ON LOCAL SUBPROBLEMS

This section is devoted to the proof of Theorems 2.3 and 2.4.

Proof of Theorem 2.3. As a first step we consider $\hat{e}_j - e_j$. We recall that (e_j, e_u) denotes the discrete solution of (2.3). In general, the solution $\hat{e}_j \in RT_1^{-1}(\Omega; \mathcal{T}_k)$ is not contained in $H(\text{div}; \Omega)$. In the following, the bilinear forms $a(\cdot, \cdot)$ and $b(\cdot, \cdot)$ are extended to the nonconforming ansatz spaces in a natural way, i.e., the integrals over Ω are replaced by the sum of the integrals over the elements T of the triangulation \mathcal{T}_k :

$$\begin{aligned} a(\hat{e}_j - e_j, \hat{e}_j) &= -b(\hat{e}_j, \hat{e}_u - e_u) - \sum_{e \in \mathcal{E}_k} \int_e ([u_D]_A - \lambda_{RT_1}) [\mathbf{n}_e \cdot \hat{e}_j]_J d\sigma \\ &= -c(\hat{e}_u, \hat{e}_u - e_u) + (f - \Pi_0 f, \hat{e}_u - e_u)_0 \\ &\quad - \sum_{e \in \mathcal{E}_k} \int_e ([u_D]_A - \lambda_{RT_1}) [\mathbf{n}_e \cdot (\hat{e}_j - e_j)]_J d\sigma \\ &= -c(\hat{e}_u - e_u, \hat{e}_u - e_u) - b(e_j, \hat{e}_u - e_u) \\ &\quad - \sum_{e \in \mathcal{E}_k} \int_e ([u_D]_A - \lambda_{RT_1}) [\mathbf{n}_e \cdot (\hat{e}_j - e_j)]_J d\sigma \\ &\leq \|\text{div } e_j\|_0 \|\hat{e}_u - e_u\|_0 + \|[u_D]_A - \lambda_{RT_1}\|_{0; \hat{\mathcal{E}}_k^{-1}} \|\mathbf{n}_e \cdot (\hat{e}_j - e_j)\|_{0; \hat{\mathcal{E}}_k}. \end{aligned}$$

Note that the weighted norm $\|\cdot\|_{0; \hat{\mathcal{E}}_k}$ has the inverse weighting factor compared to $\|\cdot\|_{0; \hat{\mathcal{E}}_k^{-1}}$:

$$\|v\|_{0; \hat{\mathcal{E}}_k} := \left(\sum_{e \in \mathcal{E}_k} \alpha_e^{-1} h_e \|v\|_{0;e}^2 \right)^{1/2}, \quad v \in L^2(\mathcal{E}_k).$$

For $\|\hat{e}_u - e_u\|_0$ we have to establish an adequate upper bound. We obtain

$$(5.1) \quad \begin{aligned} \|e_u - \hat{e}_u\|_0 &\leq \beta_{\text{div}} \sup_{\substack{\mathbf{q} \in RT_1(\Omega; \mathcal{T}_k) \\ \mathbf{q} \neq 0}} \frac{b(\mathbf{q}, e_u - \hat{e}_u)}{\|\mathbf{q}\|_{\text{div}}} = \beta_{\text{div}} \sup_{\substack{\mathbf{q} \in RT_1(\Omega; \mathcal{T}_k) \\ \mathbf{q} \neq 0}} \frac{a(\mathbf{q}, \hat{e}_j - e_j)}{\|\mathbf{q}\|_{\text{div}}} \\ &\leq \alpha_0^{-1/2} \beta_{\text{div}} a(e_j - \hat{e}_j, e_j - \hat{e}_j)^{1/2}. \end{aligned}$$

Due to an inverse estimate for the Raviart-Thomas elements [12, 31], an upper bound for the weighted jump $\|[\mathbf{n}_e \cdot (\hat{e}_j - e_j)]_J\|_{0; \hat{\mathcal{E}}_k}$ is given by

$$\|[\mathbf{n}_e \cdot (\hat{e}_j - e_j)]_J\|_{0; \hat{\mathcal{E}}_k} \leq C_{\text{nor}} a(\hat{e}_j - e_j, \hat{e}_j - e_j)^{1/2},$$

where the constant $0 < C_{\text{nor}}$ is independent of the refinement level. Altogether, we arrive at an upper bound for $a(\hat{e}_j, \hat{e}_j)^{1/2}$:

$$a(\hat{e}_j, \hat{e}_j)^{1/2} \leq 2a(e_j, e_j)^{1/2} + \alpha_0^{-1/2} \beta_{\text{div}} \|\text{div } e_j\|_0 + C_{\text{nor}} \beta_{2; \infty} \|\mathbf{j} - \mathbf{j}_h\|_{\text{div}}.$$

It remains to establish an upper bound for $\|\text{div } \hat{e}_j\|_0$. According to the equality $b(e_j - \hat{e}_j, v) = c(e_u - \hat{e}_u, v)$, $v \in W_1(\Omega; \mathcal{T}_k)$, we get

$$\begin{aligned} \|\text{div } \hat{e}_j\|_0 &\leq \|\text{div } e_j\|_0 + \beta_1 \|e_u - \hat{e}_u\|_0 \\ &\leq \|\text{div } e_j\|_0 + \alpha_0^{-1/2} \beta_1 \beta_{\text{div}} a(\hat{e}_j - e_j, \hat{e}_j - e_j)^{1/2} \\ &\leq \alpha_0^{-1/2} \beta_1 \beta_{\text{div}} a(e_j, e_j)^{1/2} + (1 + \alpha_0^{-1} \beta_1 \beta_{\text{div}}^2) \|\text{div } e_j\|_0 \\ &\quad + \alpha_0^{-1/2} \beta_1 \beta_{\text{div}} C_{\text{nor}} \beta_{2; \infty} \|\mathbf{j} - \mathbf{j}_h\|_{\text{div}}. \end{aligned}$$

To summarize the results, $\|\hat{e}_j\|_{\text{div}}$ yields a lower bound for $\|\mathbf{j} - \mathbf{j}_h\|_{\text{div}}$:

$$C_{a-1}^e \left(1 + \beta_\infty + \frac{C_{a-1}^0}{C_{a-1}^e} \right) \|\mathbf{j} - \mathbf{j}_h\|_{\text{div}} \geq \|\hat{e}_j\|_{\text{div}}$$

where

$$C_{a-1}^e := \sqrt{2} \left(1 + \max \left(1 + \frac{\beta_1 \beta_{\text{div}}}{\sqrt{\alpha_0}}, \frac{\beta_1 \beta_{\text{div}}^2}{\alpha_0} + \frac{\beta_{\text{div}}}{\sqrt{\alpha_0}} \right) \right)$$

and $C_{a-1}^0 := C_{\text{nor}} \beta_{2; \infty} (1 + \alpha_0^{-1/2} \beta_1 \beta_{\text{div}})$.

To prove an upper bound for $\|\mathbf{j} - \mathbf{j}_h\|_{\text{div}}$, we again examine $a(e_j - \hat{e}_j, e_j - \hat{e}_j)$:

$$\begin{aligned} a(e_j - \hat{e}_j, e_j) &= b(e_j, \hat{e}_u - e_u) = c(e_u, \hat{e}_u - e_u) - (f - \Pi_0 f, \hat{e}_u - e_u)_0 \\ &= -c(\hat{e}_u - e_u, \hat{e}_u - e_u) + b(\hat{e}_j, \hat{e}_u - e_u), \\ a(e_j - \hat{e}_j, e_j - \hat{e}_j) &\leq -a(e_j - \hat{e}_j, \hat{e}_j) + \|\text{div } \hat{e}_j\|_0 \|\hat{e}_u - e_u\|_0. \end{aligned}$$

Using the upper bound (5.1) for $\|e_u - \hat{e}_u\|_0$, we obtain

$$a(e_j - \hat{e}_j, e_j - \hat{e}_j)^{1/2} \leq a(\hat{e}_j, \hat{e}_j)^{1/2} + \alpha_0^{-1/2} \beta_{\text{div}} \|\text{div } \hat{e}_j\|_0.$$

By means of the triangle inequality we get the upper bound in (2.14) with the constant

$$C_{\text{loc}} := 1 + \sqrt{2} (1 + \alpha_0^{-1} \beta_1^2 \beta_{\text{div}}^2) \max(1, \alpha_0^{-1} \beta_{\text{div}}).$$

□

The computation of (\hat{e}_j, \hat{e}_u) in (2.13) requires the solution of an 11×11 saddle point problem for each element. Therefore, the determination of \hat{e}_j is too expensive for a cheap error estimator. The rest of this section is devoted to the reduction of the computational amount.

As in the previous section, we replace the original bilinear form $a|_T(\cdot, \cdot)$ by the modified form $\tilde{a}|_T(\cdot, \cdot)$ and consider three local subproblems. First, it is easy to see that the solution of the subproblem associated with $RT_0(\Omega; \mathcal{T}_k) \times W_0(\Omega; \mathcal{T}_k)$ is equal to zero. Second, let $\hat{e}_{\mathbf{j}_1}^0|_T \in \widehat{RT}_1^0(T)$ be the unique solution of the symmetric, positive definite 3×3 system

$$(5.2) \quad a|_T(\hat{e}_{\mathbf{j}_1}^0, \mathbf{q}) = - \int_{\partial T} [u_D]_A \mathbf{n} \cdot \mathbf{q} \, d\sigma + r|_T(\mathbf{q}), \quad \mathbf{q} \in \widehat{RT}_1^0(T).$$

We can further reduce the amount of computation, if we replace the stiffness matrix in (5.2) by its spectrally equivalent diagonal matrix. Then, we have to solve one scalar equation for each edge $e_i \in \partial T$, $1 \leq i \leq 3$:

$$(5.3) \quad \int_T \hat{a} \nabla \varphi_{e_i; T} \nabla \Phi_{e_i} \, dx = - \int_{\partial T} [u_D]_A \mathbf{n} \cdot (\mathbf{curl} \Phi_{e_i}) \, d\sigma + r|_T(\mathbf{curl} \Phi_{e_i}).$$

We observe that, due to (4.11),

$$c_{\mathbf{curl}} a|_T(\hat{e}_{\mathbf{j}_1}^0, \hat{e}_{\mathbf{j}_1}^0) \leq \sum_{i=1}^3 a|_T(\mathbf{curl} \varphi_{e_i}, \mathbf{curl} \varphi_{e_i}) \leq C_{\mathbf{curl}} a|_T(\hat{e}_{\mathbf{j}}^0, \hat{e}_{\mathbf{j}}^0).$$

Finally, we have to take into account the solution $(\hat{e}_{\mathbf{j}_1}^1|_T, \hat{e}_u^1|_T) \in \widehat{RT}_1^1(T) \times \widehat{W}_1(T)$, $T \in \mathcal{T}_k$, of the following saddle point problem:

$$(5.4) \quad \begin{aligned} a|_T(\hat{e}_{\mathbf{j}_1}^1, \mathbf{q}) + b|_T(\mathbf{q}, \hat{e}_{u_1}) &= r|_T(\mathbf{q}), & \mathbf{q} \in \widehat{RT}_1^1(T), \\ b|_T(\hat{e}_{\mathbf{j}_1}^1, v) - c|_T(\hat{e}_{u_1}, v) &= -(f, v)_{0; T}, & v \in \widehat{W}_1(T). \end{aligned}$$

Proof of Theorem 2.4. We only have to show the equivalence of the norms $\|\hat{e}_{\mathbf{j}}\|_{\text{div}}^2$ and $\|\hat{e}_{\mathbf{j}_1}^0\|_{\text{div}}^2 + \|\hat{e}_{\mathbf{j}_1}^1\|_{\text{div}}^2$. Again, we replace the bilinear form $a|_T(\cdot, \cdot)$ by $\tilde{a}|_T(\cdot, \cdot)$. Using the modified residual $\tilde{r}(\mathbf{q}) := r(\mathbf{q}) - \sum_{e \in \mathcal{E}_k} \int_e [u_D]_A [\mathbf{n}_e \cdot \mathbf{q}]_J \, d\sigma$, we are in the situation of Theorem 4.1 with \tilde{r} instead of r . Then, the same techniques give the assertion. \square

Remark. An error estimator for the error $\|u - u_h\|_0^2 + \|\mathbf{j} - \mathbf{j}_h\|_{\text{div}}^2$ is given by

$$(5.5) \quad \begin{aligned} \hat{\eta}_L^2 &:= \sum_{T \in \mathcal{T}_k} \hat{\eta}_{L; T}^2, \\ \hat{\eta}_{L; T}^2 &:= \eta_{L; T}^2 + \|\hat{e}_{u_1}\|_{0; T}^2, \quad T \in \mathcal{T}_k. \end{aligned}$$

Since $(\hat{e}_{\mathbf{j}}^1, \hat{e}_u^1) = (\tilde{e}_{\mathbf{j}_1}^1, \tilde{e}_{u_1})$, the assertion is evident.

For the evaluation of the error estimators η_L and $\hat{\eta}_L$ we have to specify the local Dirichlet data u_D . A possible choice is to take a piecewise quadratic function $v_{\mathbf{j}}$, $v_{\mathbf{j}}|_T \in P_2(T)$, $T \in \mathcal{T}_k$. Let $v_{\mathbf{j}}$ be locally defined as the unique quadratic function such that

$$\nabla v_{\mathbf{j}}|_T = -a^{-1} \tilde{\mathbf{j}}_h|_T, \quad \Pi_0 v_{\mathbf{j}}|_T = \tilde{u}_h|_T, \quad T \in \mathcal{T}_k,$$

and take $u_D = v_{\mathbf{j}}$. In subsection 7.2, we will see that this definition guarantees the local equivalence between the hierarchical error estimator and the estimator based on the solution on local subproblems.

6. ERROR ESTIMATOR BASED ON A SUPERCONVERGENCE RESULT

The starting point for the construction of the error estimator is the following discrete nonconforming variational problem:

$$(6.1) \quad a_N(\Psi_{NC}, \eta) = (\Pi_0 f, \eta)_0, \quad \eta \in N(\Omega; \mathcal{T}_k),$$

where the bilinear form $a_N : N(\Omega; \mathcal{T}_k) \times N(\Omega; \mathcal{T}_k) \rightarrow \mathbb{R}$ is given by

$$a_N(\psi, \eta) := \sum_{T \in \mathcal{T}_k} \int_T \left(\hat{P}_{a^{-1}}(a \nabla \psi) \cdot \nabla \eta + b \Pi_0 \psi \cdot \Pi_0 \eta \right) dx, \quad \psi, \eta \in N(\Omega; \mathcal{T}_k),$$

and the nonconforming ansatz space $N(\Omega; \mathcal{T}_k)$ is the lowest order Crouzeix-Raviart space augmented by cubic bubble functions:

$$N(\Omega; \mathcal{T}_k) := CR(\Omega; \mathcal{T}_k) \oplus B(\Omega; \mathcal{T}_k).$$

Here, $CR(\Omega; \mathcal{T}_k) := \{v \in L^2(\Omega) \mid v|_T \in P_1(T), T \in \mathcal{T}_k, v|_{T_1}(m_e) = v|_{T_2}(m_e), \partial T_1 \cap \partial T_2 = e \subset \Omega, v|_T(m_e) = 0, e \subset \partial T \cap \partial \Omega\}$ and $B(\Omega; \mathcal{T}_k) := \{v \in L^2(\Omega) \mid v|_T \in P_3(T), v|_{\partial T} = 0, T \in \mathcal{T}_k\}$. The local operator Π_0 is the L^2 -projection onto $W_0(\Omega; \mathcal{T}_k)$, whereas $\hat{P}_{a^{-1}}$ denotes the orthogonal projection onto $RT_0^{-1}(\Omega; \mathcal{T}_k)$ with respect to the bilinear form $a(\cdot, \cdot)$. Then, the following equalities are true [2, 12]:

$$(6.2) \quad u_h = \Pi_0 \Psi_{NC}, \quad \lambda_h = \Pi_e \Psi_{NC}, \quad \mathbf{j}_h = -\hat{P}_{a^{-1}}(a \nabla \Psi_{NC}),$$

where $(\mathbf{j}_h, u_h, \lambda_h) \in RT_0^{-1}(\Omega; \mathcal{T}_k) \times W_0(\Omega; \mathcal{T}_k) \times M_0(\Omega; \mathcal{E}_k)$ denotes the unique solution of the mixed hybrid formulation of the variational problem (1.4):

$$(6.3) \quad \begin{aligned} a(\mathbf{j}_h, \mathbf{q}) + b(\mathbf{q}, u_h) + d(\lambda_h, \mathbf{q}) &= 0, & \mathbf{q} \in RT_0^{-1}(\Omega; \mathcal{T}_k), \\ b(\mathbf{j}_h, v) - c(u_h, v) &= -(f, v)_0, & v \in W_0(\Omega; \mathcal{T}_k), \\ d(\mu, \mathbf{j}_h) &= 0, & \mu \in M_0(\Omega; \mathcal{E}_k). \end{aligned}$$

Here, $M_0(\Omega; \mathcal{E}_k)$ is the ansatz space for the Lagrange multipliers

$$M_0(\Omega; \mathcal{E}_k) := \{\mu \in L^2(\mathcal{E}_k) \mid \mu|_e \in P_0(e), e \in \mathcal{E}_k, \mu|_e = 0, e \subset \partial \Omega\}$$

and Π_e stands for the L^2 -projection onto $M_0(\Omega; \mathcal{E}_k)$. Finally, the bilinear form $d_{RT} : M_0(\Omega; \mathcal{E}_k) \times RT_0^{-1}(\Omega; \mathcal{T}_k) \rightarrow \mathbb{R}$ is given by

$$d(\mu, \mathbf{q}) := \sum_{T \in \mathcal{T}_k} \int_{\partial T} \mu \mathbf{n} \cdot \mathbf{q} d\sigma, \quad \mu \in M_0(\Omega; \mathcal{E}_k), \quad \mathbf{q} \in RT_0^{-1}(\Omega; \mathcal{T}_k).$$

Now, we assume the existence of a constant $0 \leq \beta < 1$ such that

$$(6.4) \quad \|u - \hat{u}_{RT_0}\|_0 \leq \beta \|u - u_h\|_0,$$

with $\hat{u}_{RT_0} \in CR(\Omega; \mathcal{T}_k)$ being the nonconforming extension of λ_h . This saturation assumption is motivated by a superconvergence result that holds true in the case of mixed hybridization, stating that the nonconforming extension \hat{u}_{RT_0} of the multiplier λ_h does provide a better approximation of the primal variable u than the piecewise constant approximation u_h (see [2, 12]). It is easy to see that (6.4) gives rise to an upper and a lower bound for the discretization error of the primal variable u in the L^2 -norm:

$$(1 + \beta)^{-1} \|\hat{u}_{RT_0} - u_h\|_0 \leq \|u - u_h\|_0 \leq (1 - \beta)^{-1} \|\hat{u}_{RT_0} - u_h\|_0.$$

Up to now, the error estimator depends on u_h and \hat{u}_{RT_0} . If the original system (1.4) is solved, the nonconforming approximation \hat{u}_{RT_0} is not available without additional computation. We have to solve additional local problems to obtain \hat{u}_{RT_0} .

In the rest of this section, the equivalence between $\|u_h - \hat{u}_{RT_0}\|_0^2$ and a weighted sum of the squared jumps of u_h across the edges $e \in \mathcal{E}_k$ will be established.

Theorem 6.1. *Let $(\mathbf{j}_h, u_h, \lambda_h) \in RT_0^{-1}(\Omega; \mathcal{T}_k) \times W_0(\Omega; \mathcal{T}_k) \times M_0(\Omega; \mathcal{E}_k)$ be the unique solution of (6.3), and assume that $\hat{u}_{RT_0} \in CR(\Omega; \mathcal{T}_k)$ is the nonconforming extension of λ_h . Then, there exist constants $0 < \sigma_0 \leq \sigma_1$, depending only on the shape regularity of \mathcal{T}_k and the ellipticity constants in (1.2), such that*

$$(6.5) \quad \sigma_0 \left(\sum_{e' \in \mathcal{E}_k} h_e^2 ([u_h]_{J|e})^2 \right)^{1/2} \leq \|u_h - \hat{u}_{RT_0}\|_0 \leq \sigma_1 \left(\sum_{e' \in \mathcal{E}_k} h_e^2 ([u_h]_{J|e})^2 \right)^{1/2}.$$

Proof. A detailed proof of the theorem can be found in [19, Thm. 4.1], [32, Thm. 5.8]. Here, we will only sketch the main ideas. By straightforward computation, we obtain

$$(6.6) \quad c \|u_h - \hat{u}_{RT_0}\|_0^2 \leq \sum_{e \in \mathcal{E}_k} h_e^2 \left(2 ([u_h]_{A|e} - \lambda_h|_e)^2 + \frac{1}{2} ([u_h]_{J|e})^2 \right) \leq C \|u_h - \hat{u}_{RT_0}\|_0^2,$$

where the constants $0 < c \leq C$ are independent of the refinement level. As a direct consequence of (6.6), we obtain the lower bound in (6.5).

However, the proof of the upper bound is more involved. It is sufficient to show that

$$(6.7) \quad \sum_{e \in \mathcal{E}_k} h_e^2 ([u_h]_{A|e} - \lambda_h|_e)^2 \leq c \sum_{e \in \mathcal{E}_k} h_e^2 ([u_h]_{J|e})^2$$

with an appropriate positive constant c . As a first step, one can establish the following relationship between λ_h and the averages and jumps of u_h :

$$(6.8) \quad \lambda_h|_e = [u_h]_{A|e} - \sum_{e' \in \mathcal{E}_k} \frac{h_{e'}}{h_e} [u_h]_{J|e'} [\mathbf{n}_{e'} \cdot P_{a^{-1}}(\boldsymbol{\tau}_e)]_{A|e'},$$

where $P_{a^{-1}}$ denotes the global orthogonal projection onto $RT_0(\Omega; \mathcal{T}_k)$ with respect to the weighted L^2 -inner product $a(\cdot, \cdot)$, and $\boldsymbol{\tau}_e$ stands for a local function of $RT_0^{-1}(\Omega; \mathcal{T}_k)$ associated with the edge e ,

$$(6.9) \quad \boldsymbol{\tau}_e = \frac{1}{2} (\boldsymbol{\tau}_e^{T_{in}} + \boldsymbol{\tau}_e^{T_{out}}).$$

Here, $\boldsymbol{\tau}_e^{T_{in}}$, $\boldsymbol{\tau}_e^{T_{out}}$ are the nodal basis vector fields in $RT_0^{-1}(\Omega; \mathcal{T}_k)$ with support in T_{in} and T_{out} , respectively, given by

$$\mathbf{n} \cdot \boldsymbol{\tau}_e^T|_{e'} = \delta_{e,e'}, \quad e' \subset \partial T, \quad T \in \{T_{in}, T_{out}\}.$$

Note that $\boldsymbol{\tau}_e$ is not contained in $H(\text{div}; \Omega)$ for an interior edge. For the next step, one has to consider the projection $P_{a^{-1}}$ in more detail and to prove that the spectral radius of $P_{a^{-1}} P_{a^{-1}}^T$ is bounded independently of the refinement level. This can be achieved by considering the local matrix representations of the positive definite operator A associated with the bilinear form $a(\cdot, \cdot)$ and the natural embedding of $RT_0(\Omega; \mathcal{T}_k)$ into $RT_0^{-1}(\Omega; \mathcal{T}_k)$. \square

Proof of Theorem 2.5. It is easy to see that

$$|h_e| \| [w]_{J|e} \|_0^2 \leq \frac{2}{\kappa_0} \|w\|_{0; T_1 \cup T_2}^2, \quad T_1 \cap T_2 = e, \quad w \in W_0(\Omega; \mathcal{T}_k).$$

Due to the triangle inequality and Theorem 2.5 we obtain

$$\begin{aligned} \|u - \tilde{u}_h\|_0 &\leq \frac{\sigma_1}{1 - \beta} \left(\sum_{e' \in \mathcal{E}_k} h_e \|\tilde{u}_h\|_{0,e} \right)^{1/2} + \left(\frac{\sigma_1}{1 - \beta} \sqrt{6\kappa_0^{-1}} + 1 \right) \|u_h - \tilde{u}_h\|_0, \\ \|u - \tilde{u}_h\|_0 &\geq \frac{\sigma_0}{1 + \beta} \left(\sum_{e' \in \mathcal{E}_k} h_e \|\tilde{u}_h\|_{0,e} \right)^{1/2} - \left(\frac{\sigma_0}{1 + \beta} \sqrt{6\kappa_0^{-1}} + 1 \right) \|u_h - \tilde{u}_h\|_0, \end{aligned}$$

and hence the assertion is proved. \square

7. COMPARISON OF THE DIFFERENT ERROR ESTIMATORS

The error estimators $\hat{\eta}_R$, η_H , $\hat{\eta}_H$, η_L , $\hat{\eta}_L$, η_S and $\hat{\eta}_S$ have been investigated independently in Sections 3–6, respectively. In this section, we examine the relationships between their local contributions. For simplicity we restrict ourselves to the Poisson equation with homogeneous Dirichlet boundary data

$$(7.1) \quad -\Delta u = f \quad \text{in } \Omega \quad \text{and} \quad u = 0 \quad \text{on } \partial\Omega,$$

and we assume that the exact discrete solution is available.

7.1. Equivalence between the residual based and the hierarchical error estimator.

Theorem 7.1. *Let $\hat{\eta}_{R;T}$ and $\hat{\eta}_{H;T}$, $T \in \mathcal{T}_k$, be given by (2.2) and (4.12), respectively. Then, there exist constants $c_{R;H}$, $C_{R;H}$ and $\Gamma_{R;H} > 0$, independent of the refinement level and $T \in \mathcal{T}_k$, such that*

$$\begin{aligned} c_{R;H} \hat{\eta}_{H;T}^2 &\leq \hat{\eta}_{R;T}^2 \leq C_{R;H} \hat{\eta}_{H;T}^2 + \|f - \Pi_1 f\|_{0;T}^2 \\ &\quad + \Gamma_{R;H} h_T^2 \left(h_T^2 \|\Pi_0 f\|_{0;T}^2 + \sum_{i=1}^3 h_{e_i} \|\Pi_0 f\|_{0;e_i}^2 \right). \end{aligned}$$

Proof. We recall that the local components of $\hat{\eta}_R$ and $\hat{\eta}_H$ are given by

$$\begin{aligned} \hat{\eta}_{R;T}^2 &:= \|f - \Pi_0 f\|_{0;T}^2 + h_T^2 \|a^{-1} \tilde{\mathbf{j}}_h\|_{0;T}^2 + \sum_{i=1}^3 w_i \alpha_{e_i} h_{e_i} \| [a^{-1} \tilde{\mathbf{j}}_h \cdot \mathbf{t}_{e_i}]_J \|_{0;e_i}^2, \\ \hat{\eta}_{H;T}^2 &:= \|\tilde{e}_{\mathbf{j}_1}^1\|_T^2 + \sum_{i=1}^3 w_i \|\mathbf{curl} \varphi_{e_i}\|_{\text{div}}^2 + \|\tilde{e}_{u_1}\|_{0;T}^2, \quad T \in \mathcal{T}_k. \end{aligned}$$

In order to establish their equivalence we proceed in several steps. First, we determine an upper bound for $\hat{\eta}_{H;T}$. To this end we consider the three parts of $\hat{\eta}_{H;T}$ separately. It is easy to see that there exists a constant $c_1 > 0$, independent of the refinement level, such that

$$(7.2) \quad \|\mathbf{q}\|_{0;T} \leq c_1 h_T \|\text{div } \mathbf{q}\|_{0;T}, \quad \mathbf{q} \in \widehat{RT}_1^1(T), \quad T \in \mathcal{T}_k.$$

Inequality (7.2) guarantees an upper bound for $\|\tilde{e}_{\mathbf{j}_1}^1\|_T^2$:

$$(7.3) \quad \begin{aligned} \|\tilde{e}_{\mathbf{j}_1}^1\|_T^2 &= \|\tilde{e}_{\mathbf{j}_1}^1\|_{0;T}^2 + \|\text{div } \tilde{e}_{\mathbf{j}_1}^1\|_{0;T}^2 \\ &\leq (1 + c_1^2 h_T^2) \|\text{div } \tilde{e}_{\mathbf{j}_1}^1\|_{0;T}^2 \leq (1 + c_1^2 h_T^2) \|f - \Pi_0 f\|_{0;T}^2. \end{aligned}$$

On the other hand, an upper bound for the second term $\|\mathbf{curl} \varphi_{e_i}\|_{\text{div}}$ is obtained by

$$\begin{aligned}
(7.4) \quad \|\mathbf{curl} \varphi_{e_i}\|_{\text{div}} &= \sup_{\substack{\mathbf{q} \in \widehat{RT}_1^0(\Omega; \mathcal{T}_k) \\ \mathbf{q} \cdot \mathbf{n}|_e = 0, e \neq e_i}} \frac{\int_{\Omega} \mathbf{curl} \varphi_{e_i} \cdot \mathbf{q} \, dx}{\|\mathbf{q}\|_0} = \sup_{\substack{\mathbf{q} \in \widehat{RT}_1^0(\Omega; \mathcal{T}_k) \\ \mathbf{q} \cdot \mathbf{n}|_e = 0, e \neq e_i}} \frac{r(\mathbf{q})}{\|\mathbf{q}\|_0} \\
&\leq \left(\int_{e_i} [\mathbf{j}_h \cdot \mathbf{t}_{e_i}]_J^2 \, d\sigma \right)^{1/2} \cdot \|\nabla \Phi_{e_i}\|_0^{-1} \left(\int_{e_i} \Phi_{e_i}^2 \, d\sigma \right)^{1/2} \\
&\leq c_2 h_{e_i}^{1/2} \|\mathbf{j}_h \cdot \mathbf{t}_{e_i}\|_J,
\end{aligned}$$

where the constant c_2 depends only on the geometry of \mathcal{T}_0 . For an estimation of the third term, we note that $\tilde{e}_{u_1}|_T$ is an element of $\widehat{W}_1(T)$. Hence

$$\begin{aligned}
(7.5) \quad \|\tilde{e}_{u_1}\|_{0;T} &= \sup_{\substack{\mathbf{q} \in \widehat{RT}_1^1(T) \\ \mathbf{q} \neq 0}} \frac{\int_T \text{div} \mathbf{q} \tilde{e}_{u_1} \, dx}{\|\text{div} \mathbf{q}\|_{0;T}} = \sup_{\substack{\mathbf{q} \in \widehat{RT}_1^1(T) \\ \mathbf{q} \neq 0}} \frac{\int_T (\tilde{e}_{j_1}^1 + \mathbf{j}_h) \cdot \mathbf{q} \, dx}{\|\text{div} \mathbf{q}\|_{0;T}} \\
&\leq c_1 h_T \left(\|\tilde{e}_{j_1}^1\|_{0;T} + \|\mathbf{j}_h\|_{0;T} \right).
\end{aligned}$$

The right-hand sides in (7.3), (7.4) and (7.5) give rise to $\hat{\eta}_{R;T}^2$. Thus, we have derived an upper bound for $\hat{\eta}_{H;T}$, which in turn implies a lower bound for $\hat{\eta}_{R;T}$.

It remains to establish the upper bound for $\hat{\eta}_{R;T}^2$. Again, we consider the different parts of $\hat{\eta}_{R;T}^2$ separately. As a first step, we examine $h_T^2 \|\mathbf{j}_h\|_{0;T}^2$. Let $\mathbf{j}_d \in RT_0(T)$ be given by

$$\mathbf{j}_d = \frac{\Pi_0 f}{2} (\mathbf{x} - \mathbf{x}_c),$$

where \mathbf{x}_c denotes the center of gravity of T . Then, it is easy to see that

$$\text{div}(\mathbf{j}_h|_T - \mathbf{j}_d) = 0.$$

Note that there exists a unique $v \in P_1(T)$ such that $\Pi_0 v = 0$ and $\nabla v = \mathbf{j}_h|_T - \mathbf{j}_d$.

Further, let $\mathbf{p} \in \widehat{RT}_1^1(T)$ be uniquely given by means of $-\text{div} \mathbf{p} = v$. There exists a constant c_3 , independent of the refinement level and $T \in \mathcal{T}_k$, such that $\|w\|_{0;T} \geq c_3 h_T \|w\|_{1;T}$ for all $w \in P_1(T)$. Then, we obtain

$$c_3 h_T \|\mathbf{j}_h|_T - \mathbf{j}_d\|_{0;T} \|\text{div} \mathbf{p}\|_{0;T} \leq \|v\|_{0;T}^2 = - \int_T \text{div} \mathbf{p} v \, dx = \int_T (\mathbf{j}_h|_T - \mathbf{j}_d) \cdot \mathbf{p} \, dx,$$

whence

$$\begin{aligned}
\|\mathbf{j}_h - \mathbf{j}_d\|_{0;T} &\leq (c_3 h_T)^{-1} \sup_{\mathbf{q} \in \widehat{RT}_1^1(T)} \int_T (\mathbf{j}_h - \mathbf{j}_d) \cdot \mathbf{q} \, dx \|\text{div} \mathbf{q}\|_{0;T}^{-1} \\
&\leq c_1 c_3^{-1} \left(\|\mathbf{j}_d\|_{0;T} + \|\tilde{e}_{j_1}^1\|_{0;T} \right) + (c_3 h_T)^{-1} \|\tilde{e}_{u_1}\|_{0;T}.
\end{aligned}$$

On the other hand, we get

$$\|\mathbf{j}_d\|_{0;T} \leq \frac{1}{2} h_T \|\Pi_0 f\|_{0;T}.$$

Altogether, this results in an upper bound for $h_T^2 \|\mathbf{j}_h\|_{0;T}^2$:

$$h_T^2 \|\mathbf{j}_h\|_{0;T}^2 \leq 3 \left(\frac{1}{c_3} \right)^2 \|\tilde{e}_{u_1}\|_{0;T}^2 + 3 \left(\frac{c_1}{c_3} \right)^2 h_T^2 \|\tilde{e}_{j_1}^1\|_{0;T}^2 + \frac{3}{4} \left(1 + \frac{c_1}{c_3} \right)^2 h_T^4 \|\Pi_0 f\|_{0;T}^2.$$

Finally, we consider $\|\mathbf{j}_h \cdot \mathbf{t}_e\|_{0;e}$ and remark that $\mathbf{j}_h \cdot \mathbf{t}_e \in P_1(e)$ can be written as

$$[\mathbf{j}_h \cdot \mathbf{t}_e]_J(\mathbf{x}_s) = a_e + b_e(s - h_e/2), \quad s \in [0; h_e], \quad \mathbf{x}_s := \mathbf{p}_e + s \cdot \mathbf{t}_e,$$

where $a_e := [\mathbf{j}_h \cdot \mathbf{t}_e]_J(m_e)$ and $b_e := (d[\mathbf{j}_h \cdot \mathbf{t}_e]_J/ds)(m_e)$. Here, m_e denotes the midpoint of the edge e and \mathbf{p}_e is that vertex of e such that the other vertex is given by $\mathbf{p}_e + h_e \mathbf{t}_e$. Since $b_e = \nabla[\mathbf{j}_h \cdot \mathbf{t}_e]_J \cdot \mathbf{t}_e$, we get

$$b_e = \frac{1}{2} [\Pi_0 f]_J,$$

whence

$$h_e \|\mathbf{j}_h \cdot \mathbf{t}_e\|_{0;e}^2 = a_e^2 h_e^2 + \frac{1}{12} h_e^3 \|\Pi_0 f\|_{0;e}^2.$$

On the other hand,

$$\begin{aligned} \pm a_e &= \pm 3(2h_e)^{-1} \int [\mathbf{j}_h \cdot \mathbf{t}_e]_J \Phi_e d\sigma = \pm 3(2h_e)^{-1} r(\mathbf{curl} \Phi_e) \\ &\leq 3(2h_e)^{-1} |\Phi_e|_1 \|\mathbf{curl} \varphi_e\|_{\text{div}} \leq 3(2h_e)^{-1} c_4 \|\mathbf{curl} \varphi_e\|_{\text{div}}. \end{aligned}$$

Note that in view of

$$(7.6) \quad 0 < \hat{c}_4 \leq |\Phi_e|_1 \leq c_4, \quad e \in \mathcal{E}_k, \quad k \geq 0,$$

the positive constant $c_4 > 0$ depends only on the geometry of \mathcal{T}_0 . We thus get

$$h_e \|\mathbf{j}_h \cdot \mathbf{t}_e\|_{0;e}^2 \leq \frac{9}{4} c_4^2 \|\mathbf{curl} \varphi_e\|_{\text{div}}^2 + \frac{1}{12} h_e^3 \|\Pi_0 f\|_{0;e}^2.$$

Summarizing the preceding results, we establish an upper bound for $\hat{\eta}_{R;T}^2$:

$$\begin{aligned} \hat{\eta}_{R;T}^2 &\leq \max(1, 3c_1^2 c_3^{-2} h_T^2) \|\tilde{e}_{j_1}^1\|_T^2 + \frac{9}{4} c_4^2 \sum_{i=1}^3 w_i \|\mathbf{curl} \varphi_{e_i}\|_{\text{div}}^2 + 3c_3^{-2} \|\tilde{e}_{u_1}\|_{0;T}^2 \\ &\quad + \frac{3}{4} \left(1 + c_1 c_3^{-1} \right)^2 h_T^4 \|\Pi_0 f\|_{0;T}^2 + \frac{1}{12} \sum_{i=1}^3 w_i h_{e_i}^3 \|\Pi_0 f\|_{0;e_i}^2 + \|f - \Pi_1 f\|_{0;T}^2. \end{aligned}$$

□

The fourth, fifth and sixth terms in the upper bound of $\hat{\eta}_{R;T}$ are, in general, higher order perturbations of $\hat{\eta}_{R;T}$. Provided the grid size is small enough, they can be neglected.

7.2. Remarks on the error estimator based on the solution of local sub-problems. We will show that the estimators η_L and $\hat{\eta}_L$ based on the solution of local subproblems and the hierarchical basis error estimators η_H and $\hat{\eta}_H$ are locally equivalent.

Theorem 7.2. *Let $\eta_{H;T}$, $\eta_{L;T}$ and $\hat{\eta}_{H;T}$, $\hat{\eta}_{L;T}$, $T \in \mathcal{T}_k$, be given by (2.10), (2.15), (4.12) and (5.5), respectively. Then*

$$(7.7) \quad \eta_{H;T}^2 \leq \eta_{L;T}^2 + \frac{1}{2} \sum_{e=\partial T \cap \partial T_e} \eta_{L;T_e}^2, \quad \hat{\eta}_{H;T}^2 \leq \hat{\eta}_{L;T}^2 + \frac{1}{2} \sum_{e=\partial T \cap \partial T_e} \hat{\eta}_{L;T_e}^2.$$

where $e = \partial T \cap \partial T_e$ denotes an interior edge and T_e stands for the adjacent element sharing the edge e with T .

Moreover, if we choose $u_D = v_j$ with $\nabla v_j = \mathbf{j}_h$ and $[v_j]_A = 0$, $[\mathbf{t}_e \cdot \mathbf{j}_h]_J = 0$ for $e \subset \partial\Omega$ in (5.3), then there exist constants $C_{L;H}$, $\hat{C}_{L;H} > 0$, independent of the refinement level and $T \in \mathcal{T}_k$, such that

$$(7.8) \quad \eta_{L;T}^2 \leq C_{L;H} \eta_{H;T}, \quad \hat{\eta}_{L;T}^2 \leq \hat{C}_{L;H} \hat{\eta}_{H;T}.$$

Proof. The local contributions of η_H , $\hat{\eta}_H$ and η_L , $\hat{\eta}_L$ are given by

$$\begin{aligned} \eta_{H;T}^2 &= \|\hat{e}_{\mathbf{j}_1}^1\|_{\text{div};T}^2 + \sum_{i=1}^3 w_i \|\mathbf{curl} \varphi_{e_i}\|_{\text{div}}^2, & \hat{\eta}_{H;T}^2 &= \eta_{H;T}^2 + \|\hat{e}_{u_1}\|_{0;T}, \\ \eta_{L;T}^2 &= \|\hat{e}_{\mathbf{j}_1}^1\|_{\text{div};T}^2 + \sum_{i=1}^3 a|_T(\mathbf{curl} \varphi_{e_i;T}, \mathbf{curl} \varphi_{e_i;T}), & \hat{\eta}_{L;T}^2 &= \eta_{L;T}^2 + \|\hat{e}_{u_1}\|_{0;T}, \end{aligned}$$

where $w_i = 1$ if $e_i \subset \partial\Omega$ and $w_i = \frac{1}{2}$ if e_i is an interior edge.

We note that $(\hat{e}_{\mathbf{j}_1}^1, \hat{e}_{u_1}) = (\tilde{e}_{\mathbf{j}_1}^1, \tilde{e}_{u_1})$ and $\varphi_{\hat{e}_{\mathbf{j}_1}^1}^e = \varphi_{\tilde{e}_{\mathbf{j}_1}^1}^e$ if $e \subset \partial\Omega$. On the other hand, for an interior edge $e = \partial T \cap \partial T_e$ we get

$$\begin{aligned} a(\mathbf{curl} \varphi_e, \mathbf{curl} \varphi_e) &= r(\mathbf{curl} \varphi_e) \\ &= a|_T(\mathbf{curl} \varphi_{e;T}, \mathbf{curl} \varphi_e) + a|_{T_e}(\mathbf{curl} \varphi_{e;T}, \mathbf{curl} \varphi_e). \end{aligned}$$

Thanks to Young's inequality, this proves (7.7).

For the proof of (7.8), we choose $u_D = v_j$ in (5.3) and obtain

$$\begin{aligned} \int_T \nabla \varphi_{e;T} \cdot \nabla \Phi_e \, dx &= \int_{\partial T} [v_j]_A \nabla \Phi_e \cdot \mathbf{t} \, d\sigma - a|_T(\mathbf{j}_h, \mathbf{curl} \Phi_e) \\ &= \int_{\partial T} \mathbf{j}_h \cdot \mathbf{t} \Phi_e \, d\sigma - \int_{\partial T} [\nabla v_j \cdot \mathbf{t}]_A \Phi_e \, d\sigma = \pm \frac{1}{2} \int_e [\mathbf{j}_h \cdot \mathbf{t}_e]_J \Phi_e \, d\sigma \end{aligned}$$

for an interior edge. Consequently

$$\begin{aligned} a|_T(\mathbf{curl} \Phi_e, \mathbf{curl} \Phi_e) a|_T(\mathbf{curl} \varphi_{e;T}, \mathbf{curl} \varphi_{e;T}) &= \frac{1}{4} \left(\int_e [\mathbf{t} \cdot \mathbf{j}_h]_J \Phi_e \, d\sigma \right)^2 \\ &= \frac{1}{4} a(\mathbf{curl} \varphi_e, \mathbf{curl} \Phi_e)^2 = \frac{1}{4} a(\mathbf{curl} \varphi_e, \mathbf{curl} \varphi_e) a(\mathbf{curl} \Phi_e, \mathbf{curl} \Phi_e) \\ &\leq \frac{1}{4} (1 + C) a(\mathbf{curl} \varphi_e, \mathbf{curl} \varphi_e) a|_T(\mathbf{curl} \Phi_e, \mathbf{curl} \Phi_e), \end{aligned}$$

where the constant $C > 0$ only depends on the local geometry of \mathcal{T}_0 .

The preceding inequality implies (7.8) with $C_{L;H} = \hat{C}_{L;H} := \max(1, \frac{1}{2}(1 + C))$. \square

7.3. Remarks on the error estimator based on superconvergence. The estimator $\hat{\eta}_R$ guarantees sharp upper and lower bounds for the combination of the flux error and the error in the primal variable. Since η_S is an error estimator designed only for the L^2 -error in the primal variable, we cannot expect η_S to be equivalent to $\hat{\eta}_R$.

Theorem 7.3. Let $\hat{\eta}_R$ and η_S be given by (2.2) and (2.17), respectively. Then, there exist constants $0 < c_{S;R} \leq C_{S;R}$, independent of the refinement level, such that

$$(7.9) \quad c_{S;R} \eta_S^2 \leq \sum_{T \in \mathcal{T}_k} h_T^2 \|\mathbf{j}_h\|_{0;T}^2 \leq C_{S;R} \eta_S^2.$$

Proof. We first establish an upper bound for $h_e \| [u_h]_J \|_{0;e}^2$ if $e = \partial T_1 \cap \partial T_2$ is an interior edge. For that purpose we denote by $\mathbf{q}_e \in RT_0(\Omega; \mathcal{T}_k)$ the nodal basis field associated with e , i.e., $\mathbf{n}_e \cdot \mathbf{q}_e|_{e'} = \delta_{e,e'}$. It follows that

$$\begin{aligned} h_e \| [u_h]_J \|_{0;e}^2 &= h_e \int_e [u_h]_J [u_h]_J \mathbf{q}_e \cdot \mathbf{n}_e \, d\sigma = h_e \int_{\Omega} [u_h]_J \mathbf{j}_h \cdot \mathbf{q}_e \, dx \\ &\leq C (h_{T_1}^2 \|\mathbf{j}_h\|_{0;T_1}^2 + h_{T_2}^2 \|\mathbf{j}_h\|_{0;T_2}^2), \end{aligned}$$

where $C > 0$ is independent of $e \in \mathcal{E}_k$. Consequently

$$\eta_{S;T}^2 \leq C \sum_{T' \in D_T} h_{T'}^2 \|\mathbf{j}_h\|_{0;T'}^2, \quad T \in \mathcal{T}_k,$$

where $D_T := \{T' \in \mathcal{T}_k \mid \partial T' \cap \partial T \in \mathcal{E}_k\}$ is the union of all triangles sharing an edge with T . Summing over all $T \in \mathcal{T}_k$ and observing (1.6) gives the first inequality in (7.9).

On the other hand, to prove the second part of (7.9) we assume $\mathbf{q} \in RT_0(T)$ with $\mathbf{q} = \mathbf{j}_h|_T$, $T \in \mathcal{T}_k$. Then, we get

$$\begin{aligned} \|\mathbf{j}_h\|_{0;T}^2 &= \int_T \operatorname{div} \mathbf{q} u_h \, dx - \int_{\partial T} \lambda_h \mathbf{q} \cdot \mathbf{n} \, d\sigma = \int_{\partial T} (u_h|_T - \hat{u}_h) \mathbf{q} \cdot \mathbf{n} \, d\sigma \\ &\leq C h_T^{-1} \|u_h - \hat{u}_h\|_{0;T} \|\mathbf{j}_h\|_{0;T}, \end{aligned}$$

where $C > 0$ is independent of $T \in \mathcal{T}_k$. Summing over $T \in \mathcal{T}_k$ and following the reasoning in the proof of Theorem 2.5 gives the assertion. \square

We will now show that $\|\mathbf{j}_h - K\mathbf{j}_h\|_{0;T}$, $T \in \mathcal{T}_k$, is equivalent to some other part of $\hat{\eta}_{R;T}$. This, combined with the previous result, gives global equivalence of the error estimators $\hat{\eta}_S$ and $\hat{\eta}_R$.

Theorem 7.4. Let the operator K be given by (2.19a) and (2.19b). Then, there exist constants $0 < c_{K;R} \leq C_{K;R}$, depending only on the local geometry of \mathcal{T}_0 , such that

$$(7.10) \quad c_{K;R} \|\mathbf{j}_h - K\mathbf{j}_h\|_{0;T}^2 \leq \sum_{i=1}^3 w_i \|\operatorname{curl} \varphi_{e_i}\|_{\operatorname{div}}^2 \leq C_{K;R} \|\mathbf{j}_h - K\mathbf{j}_h\|_{0;T}^2, \quad T \in \mathcal{T}_k.$$

Proof. We consider a triangle $T \in \mathcal{T}_k$ with interior edges. Then (1.5), (7.6), and straightforward computations yield

$$\begin{aligned} \|\mathbf{j}_h - K\mathbf{j}_h\|_{0;T}^2 &= \frac{1}{3} |T| \sum_{i=1}^3 \left((\mathbf{t} \cdot (\mathbf{j}_h - K\mathbf{j}_h)|_T(m_{e_i}))^2 + (\mathbf{n} \cdot (\mathbf{j}_h - K\mathbf{j}_h)|_T(m_{e_i}))^2 \right) \\ &= \frac{1}{12} |T| \sum_{i=1}^3 ((\mathbf{j}_h)_J(m_{e_i}) \cdot \mathbf{t})^2 \leq \frac{3}{8} \kappa_1 c_4^2 \sum_{i=1}^3 w_i \|\operatorname{curl} \varphi_{e_i}\|_{\operatorname{div}}^2, \\ \|\mathbf{j}_h - K\mathbf{j}_h\|_{0;T}^2 &\geq \frac{3}{8} \kappa_0 c_4^2 \sum_{i=1}^3 w_i \|\operatorname{curl} \varphi_{e_i}\|_{\operatorname{div}}^2. \end{aligned}$$

\square

Corollary 7.5. *Let $\hat{\eta}_R$ and $\hat{\eta}_S$ be given by (2.2) and (2.20), respectively. Then, there exist constants $0 < \hat{c}_{S,R} \leq \hat{C}_{S,R}$, independent of the refinement level, such that*

$$(7.11) \quad \hat{c}_{S,R} \hat{\eta}_S^2 \leq \hat{\eta}_R^2 \leq \hat{C}_{S,R} \hat{\eta}_S^2.$$

Proof. The proof is an immediate consequence of Theorem 7.3 and Theorem 7.4. \square

We note that (7.11) only provides global equivalence of $\hat{\eta}_S$ and $\hat{\eta}_R$. However, local equivalence can be obtained, if we use $\|u_h - \hat{u}_h\|_0^2$ instead of $\sum_{e \in \mathcal{E}_k} h_e \| [u_h]_J \|_{0;e}^2$.

REFERENCES

- [1] B. ACHCHAB, A. AGOUZAL, J. BARANGER, AND J. MAITRE, *Estimateur d'erreur a posteriori hiérarchique. Application aux éléments finis mixtes*, Numer. Math. 80 (1998), 159–179. CMP 49:01
- [2] D. ARNOLD AND F. BREZZI, *Mixed and nonconforming finite element methods: Implementation, post-processing and error estimates*, RAIRO Modél. Math. Anal. Numér. 19 (1985), pp. 7–35. MR 87g:65176
- [3] I. BABUŠKA AND W. RHEINBOLDT, *Error estimates for adaptive finite element computations*, SIAM J. Numer. Anal., 15 (1978), pp. 736–754. MR 58:3400
- [4] ———, *A posteriori error estimates for the finite element method*, Int. J. Numer. Methods Eng., 12 (1978), pp. 1597–1615.
- [5] R. BANK, A. SHERMAN, AND A. WEISER, *Refinement algorithm and data structures for regular local mesh refinement*, in Scientific Computing, R. Stepleman et al., ed., vol. 44, IMACS North-Holland, Amsterdam, 1983, pp. 3–17. MR 85i:00014
- [6] R. BANK AND A. WEISER, *Some a posteriori error estimators for elliptic partial differential equations*, Math. Comp., 44 (1985), pp. 283–301. MR 86g:65207
- [7] R. BECKER AND R. RANNACHER, *Weighted a posteriori error control in FE methods*, Tech. Report 96 - 1, SFB 359 - Universität Heidelberg, January 1996.
- [8] F. BORNEMANN, B. ERDMANN, AND R. KORNUBER, *A posteriori error estimates for elliptic problems in two and three spaces dimensions*, SIAM J. Numer. Anal., 33 (1996), pp. 1188–1204. MR 98a:65161
- [9] D. BRAESS, O. KLAAS, R. NIEKAMP, E. STEIN, AND F. WOBSCAL, *Error indicators for mixed finite elements in 2-dimensional linear elasticity*, Comput. Methods Appl. Mech. Eng., 127, (1995), pp. 345–356. MR 96h:73045
- [10] D. BRAESS AND R. VERFÜRTH, *A posteriori error estimators for the Raviart-Thomas element*, SIAM J. Numer. Anal., 33 (1996), pp. 2431–2444. MR 97m:65201
- [11] J. BRANDTS, *Superconvergence and a posteriori error estimation for triangular mixed finite elements*, Numer. Math., 68 (1994), pp. 311–324. MR 97a:65162
- [12] F. BREZZI AND M. FORTIN, *Mixed and hybrid finite element methods*, Springer-Verlag, New York, 1991. MR 92d:65187
- [13] P. CLÉMENT, *Approximation by finite element functions using local regularization*, RAIRO Anal. Numér., 2 (1975), pp. 77–84. MR 53:4569
- [14] P. DEUFLHARD, P. LEINEN, AND H. YSERENTANT, *Concepts of an adaptive hierarchical finite element code*, IMPACT Comput. Sci. Engrg., 1 (1989), pp. 3–35.
- [15] R. DURAN AND R. RODRIGUEZ, *On the asymptotic exactness of Bank-Weiser's estimator*, Numer. Math., 62 (1992), pp. 297–303. MR 93e:65135
- [16] K. ERIKSSON AND C. JOHNSON, *An adaptive finite element method for linear elliptic problems*, Math. Comput., 50 (1988), pp. 361–383.
- [17] R. EWING AND J. WANG, *Analysis of multilevel decomposition iterative methods for mixed finite element methods*, RAIRO Modél. Math. Anal. Numér. 28 (1994), pp. 377–398. MR 95e:65099

- [18] R. HOPPE AND B. WOHLMUTH, *Adaptive multilevel techniques for mixed finite element discretizations of elliptic boundary value problems*, SIAM J. Numer. Anal., 34 (1997), pp. 1658–1681. MR **98e**:65095
- [19] ———, *Efficient numerical solution of mixed finite element discretizations by adaptive multilevel methods*, Appl. Math., 40 (1995), pp. 227–248. MR **96b**:65111
- [20] ———, *Multilevel iterative solution and adaptive mesh refinement for mixed finite element discretizations*, Appl. Num. Math., 23 (1997), pp. 97–117 MR **98g**:65109
- [21] ———, *Element-oriented and edge-oriented local error estimators for nonconforming finite element methods*, RAIRO Modél. Math. Anal. Numér. 30 (1996), pp. 237–263. MR **97e**:65124
- [22] C. JOHNSON, *Numerical Solution of Partial Differential Equations by the Finite Element Method*, no. 101, Cambridge University Press, Cambridge, 1987. MR **89b**:65003a
- [23] C. JOHNSON AND P. HANSBO, *Adaptive finite element methods in computational mechanics*, Comp. Meth. Appl. Mech. Eng., 101 (1992), pp. 143–181. MR **93m**:65157
- [24] R. RODRIGUEZ, *Some remarks on the Zienkiewicz-Zhu estimator*, Numer. Meth. for PDE, 10 (1994), pp. 625–636. MR **95e**:65103
- [25] T. STROUBOULIS AND J. ODEN, *A posteriori error estimation of finite element approximations in fluid dynamics*, Comp. Meth. Appl. Mech. Engrg., 78 (1990), pp. 201–242. MR **91a**:76066
- [26] B. SZABÓ AND I. BABUŠKA, *Finite Element Analysis*, Wiley, New York, 1991. MR **93f**:73001
- [27] R. VERFÜRTH, *A Review of A Posteriori Error Estimation and Adaptive Mesh-Refinement Techniques*, Wiley-Teubner, Chichester, 1996.
- [28] ———, *A posteriori error estimators for the Stokes equations*, Numer. Math., 55 (1989), pp. 309–325. MR **90d**:65187
- [29] ———, *A posteriori error estimates for nonlinear problems. Finite element discretizations of elliptic equations*, Math. Comput., 62 (1994), pp. 445–475. MR **94j**:65136
- [30] ———, *A posteriori error estimation and adaptive mesh-refinement techniques*, J. Comp. Appl. Math., 50 (1994), pp. 67–83. MR **95c**:65171
- [31] B. WOHLMUTH, *Adaptive Multilevel-Finite-Elemente Methoden zur Lösung elliptischer Randwertprobleme*, PhD thesis, TU München, 1995.
- [32] J. ZHU AND O. ZIENKIEWICZ, *Adaptive techniques in the finite element method*, Commun. Appl. Numer. Methods, 4 (1988), pp. 197–204.
- [33] O. ZIENKIEWICZ AND R. TAYLOR, *The Finite Element Method*, vol. 1, Mc Graw-Hill, London, 1989.
- [34] O. ZIENKIEWICZ AND J. ZHU, *A simple error estimator and adaptive procedure for practical engineering analysis*, J. Numer. Meth. Eng., 28 (1987), pp. 28–39.

MATH. INSTITUTE, UNIVERSITY OF AUGSBURG, D-86135 AUGSBURG, GERMANY
E-mail address: wohlmuth@math.uni-augsburg.de

MATH. INSTITUTE, UNIVERSITY OF AUGSBURG, D-86135 AUGSBURG, GERMANY
E-mail address: hoppe@math.uni-augsburg.de