

Globally Lipschitz minimizers for variational problems with linear growth

Lisa Beck, Miroslav Bulíček, Erika Maringová

Angaben zur Veröffentlichung / Publication details:

Beck, Lisa, Miroslav Bulíček, and Erika Maringová. 2018. "Globally Lipschitz minimizers for variational problems with linear growth." *ESAIM: Control, Optimisation and Calculus of Variations* 24 (4): 1395–1413. <https://doi.org/10.1051/cocv/2017065>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



GLOBALLY LIPSCHITZ MINIMIZERS FOR VARIATIONAL PROBLEMS WITH LINEAR GROWTH*

LISA BECK^{1,*}, MIROSLAV BULÍČEK² AND ERIKA MARINGOVÁ²

Abstract. We study the minimization of convex, variational integrals of linear growth among all functions in the Sobolev space $W^{1,1}$ with prescribed boundary values (or its equivalent formulation as a boundary value problem for a degenerately elliptic Euler–Lagrange equation). Due to insufficient compactness properties of these Dirichlet classes, the existence of solutions does not follow in a standard way by the direct method in the calculus of variations and in fact might fail, as it is well-known already for the non-parametric minimal surface problem. Assuming radial structure, we establish a necessary and sufficient condition on the integrand such that the Dirichlet problem is in general solvable, in the sense that a Lipschitz solution exists for any regular domain and all prescribed regular boundary values, *via* the construction of appropriate barrier functions in the tradition of Serrin’s paper [J. Serrin, *Philos. Trans. R. Soc. Lond., Ser. A* **264** (1969) 413–496].

Mathematics Subject Classification. 35A01, 35B65, 35J70, 49N60.

Received October 3, 2016. Accepted October 5, 2017.

1. INTRODUCTION

In this paper we are concerned with the existence of (unique) scalar-valued Lipschitz solutions to the Dirichlet problem

$$\begin{aligned} -\operatorname{div}(a(|\nabla u|)\nabla u) &= 0 && \text{in } \Omega, \\ u &= u_0 && \text{on } \partial\Omega, \end{aligned} \tag{1.1}$$

where $\Omega \subset \mathbb{R}^d$ is a bounded, regular domain and with regular prescribed boundary values u_0 . The focus is on coefficient functions $a \in C^1(\mathbb{R}^+)$, which on the one hand represent a radial structure condition and which on the other belong to a convex linear growth problem which is naturally formulated in the Sobolev space $W^{1,1}(\Omega)$, meaning that we work under the permanent assumption that the function $s \mapsto a(s)s$ is increasing and remains

Keywords and phrases. Variational problems, linear growth, Lipschitz minimizers, non-convex domains.

* *L. Beck is grateful for the support by M.O.P.S. of the Mathematical Institute at the University of Augsburg. M. Bulíček and E. Maringová were supported by the ERC-CZ project LL1202 financed by the Ministry of Education, Youth and Sports, Czech Republic. The support of the grant SVV-2016-260335 is also acknowledged. M. Bulíček is a member of the Nečas Center for Mathematical Modeling.*

¹ Institut für Mathematik, Universität Augsburg, Universitätsstr. 14, 86159 Augsburg, Germany.

² Mathematical Institute, Faculty of Mathematics and Physics, Charles University, Sokolovská 83, 186 75, Prague, Czech Republic.

*Corresponding author: lisa.beck@math.uni-augsburg.de

bounded. In this setting, the existence of a weak solution to the Dirichlet problem (1.1) is equivalent to the existence of a minimizer of a related (convex) variational integral in the Dirichlet class $u_0 + W_0^{1,1}(\Omega)$, and we may equivalently look for a function $u \in u_0 + W_0^{1,1}(\Omega)$ such that for all smooth, compactly supported test functions $\varphi \in \mathcal{D}(\Omega)$ we have

$$\int_{\Omega} F(|\nabla u|) \, dx \leq \int_{\Omega} F(|\nabla u_0 + \nabla \varphi|) \, dx, \quad (1.2)$$

where F and a are linked *via* the identity

$$F'(s) = a(s)s \quad \text{for all } s \in \mathbb{R}^+. \quad (1.3)$$

The minimal surface equation is clearly the most prominent example for such a Dirichlet problem, and other prototypic examples are given *via* the coefficient functions

$$a_p(s) := (1 + s^p)^{-\frac{1}{p}} \quad (1.4)$$

for $s \in \mathbb{R}^+$ and $p > 0$ (which for the specific case $p = 2$ just gives the minimal surface equation).

One peculiarity of linear growth problems is that, even if the equation (1.1) is monotone and the integrand $z \mapsto F(|z|)$ of the variational functional is convex, standard monotonicity methods and the direct method of the calculus of variations fail in general, since the Sobolev space $W^{1,1}(\Omega)$ is non-reflexive and hence has insufficient compactness properties. For the study of such linear growth problem of variational type, one common strategy is to extend in a first step the functional by lower semicontinuity (in the sense of Lebesgue and Serrin [15, 21]) to the larger space $BV(\Omega)$ of functions of bounded variation, *i.e.*, to consider, for fixed boundary values u_0 , the functional

$$w \mapsto \inf \left\{ \liminf_{n \rightarrow \infty} \int_{\Omega} F(|\nabla w_n|) \, dx : (w_n)_{n \in \mathbb{N}} \text{ in } u_0 + W_0^{1,1}(\Omega) \text{ with } w_n \rightarrow w \text{ in } L^1(\Omega) \right\}$$

with $w \in BV(\Omega)$. This extension also allows for an integral representation, see *e.g.* [13], which consists in the original functional evaluated for the absolutely continuous part of the measure derivative and penalization terms for non-vanishing singular measure derivative or non-attainment of the prescribed boundary values (note that the trace operator is in general not continuous with respect to weak-* convergence in $BV(\Omega)$). In a second step, one can then study minimizers of the extended functional in $BV(\Omega)$ (which can be interpreted as generalized minimizers of the original functional), which exist as a consequence of the direct method, applied in the space $BV(\Omega)$ equipped with the weak-* topology (observe that the lower semicontinuity of the extended functional in this topology was established by Reshetnyak [20]).

Returning to the original question, one can then investigate whether or not these generalized minimizers belong to $u_0 + W_0^{1,1}(\Omega)$ which amounts to excluding the singular measure derivative and to show attainment of the boundary values. However, this is in general not the case, meaning that in fact no minimizer in the space $u_0 + W_0^{1,1}(\Omega)$ might exist. This situation has been studied in full detail for the minimal surface equation, *i.e.* the non-parametric minimal surface problem. A by now classical result by Miranda [19] states that for any locally pseudoconvex domain Ω and any continuous prescribed boundary values u_0 a unique minimizer $u \in \mathcal{C}(\overline{\Omega}) \cap \mathcal{C}^2(\Omega)$ exists. Moreover, this result is sharp, and neither the convexity assumption nor the regularity assumption on u_0 can be considerably weakened in order to guarantee the existence of minimizers in $u_0 + W_0^{1,1}(\Omega)$. For a broad overview on results concerning the existence, uniqueness and regularity of these non-parametric minimal surfaces, the connection to parametric minimal surfaces, and detailed proofs we refer to Giusti's monograph [14].

Concerning higher regularity of such generalized minimizers, let us mention briefly that the essential difficulty is the degeneration of the second order derivatives of the integrand $f(z) = F(|z|)$, namely that the ratio between its largest and smallest eigenvalue blows up in the limit $|z| \rightarrow \infty$. This phenomenon does not only happen for functionals exhibiting linear growth, but also nearly linear growth (with the model case $F(s) = s \log(1 + s)$, *cp.* [18]). In these cases the role of the so-called μ -ellipticity condition was investigated, which quantifies the

ellipticity degeneration (and which in our case would basically mean $F''(s) \geq cs^{-\mu}$ for $s \geq 1$). As the main outcome it was observed, first by Fuchs and Mingione [11] for nearly linear growth problem and by Bildhauer and Fuchs in a series of works [4–7] for linear growth problems, that $C_{\text{loc}}^1(\Omega)$ regularity of generalized minimizers can still be established, also for the vectorial case with radial structure, provided that μ is sufficiently small. Specifically in our situation, this applies for $\mu \in (1, 3)$ (see [4], Thm. 2.7, but also [17], Thm. B and [2], Thm. 1.3), while in the limit case $\mu = 3$ (as for the area functional) only $W^{1,1}(\Omega)$ regularity is known, with improved $L \log L$ integrability of the gradients (see [4], Thm. 2.5 and [1], Cor. 1.13). However, these paper focus primarily on regularity of generalized minimizers, and attainment of the prescribed boundary values u_0 is in fact not expected in this general setting, as highlighted above.

In the present paper, we proceed with an alternative (and also very classical) strategy, which directly addresses the minimization problem (or equivalently the Dirichlet problem), without passing through the relaxed formulation. Indeed, the goal is to characterize integrands F (or equivalently coefficient functions a), in terms of properties of F only, such that the minimization or Dirichlet problem admits a solution in $u_0 + W_0^{1,1}(\Omega)$, for any (regular) domain Ω and boundary values u_0 . In fact, necessary and sufficient conditions for the solvability of the Dirichlet problem in the planar case $d = 2$ for the second order elliptic equation

$$\mathcal{A}(Du) \cdot \nabla^2 u = 0 \quad (1.5)$$

(with the convention $\mathcal{A}(z) \cdot \tilde{z} := \sum_{i,j=1}^d \mathcal{A}_{ij}(z) \tilde{z}_{ij}$) were already investigated by Bernstein [3], in terms of the Bernstein genre g defined *via* the validity of

$$c|z|^{2-g} \leq \frac{\mathcal{A}(z) \cdot (z \otimes z)}{\text{tr } \mathcal{A}(z)} \leq C|z|^{2-g} \quad \text{for all } z \in \mathbb{R}^d \setminus B_1(\mathbf{0})$$

for some constants $c \leq C$ (if well-defined), and then generalized by Leray [16]. One can easily calculate that the Bernstein genre of the equation for the prototypic coefficients a_p from (1.4) is given by p (hence, the minimal surface equation is of Bernstein genre 2). Bernstein's discovery in [3] was that the question of the solvability of the Dirichlet problem splits into the two classes of genre $g \leq 1$ and $g > 1$. While for the first class the Dirichlet problem is in general solvable, one needs to impose curvature restriction on the second class (as the pseudoconvexity condition mentioned before in Miranda's result). An extension to the higher-dimensional case $d \geq 2$ and a systematic treatment of more general non-uniformly elliptic equations of the form (1.5) was given later by Serrin [22]. He defined the equation to be *regularly elliptic* if

$$\frac{\mathcal{A}(z) \cdot (z \otimes z)}{\text{tr } \mathcal{A}(z)} \geq \Phi(|z|) \quad \text{for all } z \in \mathbb{R}^d \setminus B_1(\mathbf{0}) \quad (1.6)$$

holds for some increasing function $\Phi \in \mathcal{C}(\mathbb{R}^+)$ satisfying

$$\int_1^\infty \Phi(t) t^{-2} dt = \infty. \quad (1.7)$$

Obviously, equations with a well-defined Bernstein genre g are regularly elliptic if and only if $g \leq 1$. Furthermore, our specific equation (1.1) is regularly elliptic in particular if the left-hand side in (1.6) is increasing and if we have in addition

$$\int_1^\infty \frac{a'(t)t + a(t)}{a'(t)t + da(t)} dt = \int_1^\infty \frac{F''(t)}{a'(t)t + da(t)} dt = \infty.$$

The relevance of the structure condition (1.7) consists in the fact that it implies *a priori* estimates for the gradient of solutions on the boundary, *via* the construction of so-called *global barriers functions*, and interior *a priori* bounds follow in turn. Since the existence of solutions to the Dirichlet problem can be reduced to the proof of *a priori* estimates, Serrin obtained as a consequence that regularly elliptic Dirichlet problems allow for a solution, for arbitrary (regular) domains and prescribed boundary values (while for non-regularly elliptic

equations it is in general again necessary to impose restrictions on the domain, *e.g.* on the curvature of the boundary $\partial\Omega$.

The main result of the present article concerns the solvability of the Dirichlet problem in $u_0 + W_0^{1,1}(\Omega)$ and higher regularity of the solution in the sense of Lipschitz continuity, which we prove simultaneously, following the strategy of Serrin's work [22]. We here work under a radial structure condition, which allows for an easier construction of barriers, and we work under a bounded oscillation assumption, which in some sense acts as a substitute for the monotonicity assumption of the function Φ introduced above. We then obtain a necessary and sufficient condition for the solvability of the Dirichlet problem, in terms of an integral condition as in (1.7), and the precise statement is the following:

Theorem 1.1. *Let $F \in \mathcal{C}^2(\mathbb{R}^+)$ be a strictly convex function with $\lim_{s \rightarrow 0} F'(s) = 0$ which satisfies, for some constants $C_1, C_2 > 0$,*

$$\begin{aligned} C_1 s - C_2 &\leq F(s) \leq C_2(1 + s) && \text{for all } s \in \mathbb{R}^+, \\ \frac{F''(s)}{F''(t)} &\leq C_2 && \text{for all } s \geq 1 \text{ and } t \in [s/2, 2s]. \end{aligned} \quad (1.8)$$

Then the following statements are equivalent:

- (i) *For arbitrary domains Ω of class \mathcal{C}^1 satisfying an exterior ball condition and arbitrary prescribed boundary values $u_0 \in \mathcal{C}^{1,1}(\bar{\Omega})$ there exists a unique function $u \in \mathcal{C}^{0,1}(\bar{\Omega})$ solving (1.1).*
- (ii) *The function F satisfies*

$$\int_1^\infty t F''(t) dt = \infty. \quad (1.9)$$

The proof of Theorem 1.1 will be divided into two parts. In Section 2 we first deal with the failure of the existence of Lipschitz solutions to some regular boundary value problem of the type (1.1) if the assumption (1.9) is not satisfied, by an adaptation of an example constructed by Finn [10] for the minimal surface equation. We here emphasize that this non-existence result concerns the *general* solvability of the Dirichlet problem, and in fact, some restricted solvability results, for specific (non-convex) domains and boundary values, might still be true even if (1.9) is violated, see for instance ([8], Thm. 2.1). The rest of the paper is then devoted to the proof of existence of Lipschitz solutions if assumption (1.9) holds. To this end, we provide in Section 3 some auxiliary lemmata, before proceeding in Section 4 to the main proof, which consists of a number of steps. First, since the existence of solutions cannot be addressed directly, we perform an approximation of the original Dirichlet problem by a family of Dirichlet problems exhibiting a quadratic growth condition (thus, admitting solutions in $u_0 + W_0^{1,2}(\Omega)$ by the direct method). Then, the passage to the limit yields a Lipschitz solution with boundary values u_0 (and not only a generalized solution) if we can show uniform $W^{1,\infty}(\Omega)$ estimates for the solutions of the approximate problems. This is achieved by the construction of appropriate barrier functions, in the tradition of Serrin's paper [22], and concludes the proof of Theorem 1.1. Finally, it is worth mentioning that the integral condition (1.9) is trivially satisfied if F satisfies the aforementioned μ -ellipticity condition studied by Bildhauer and Fuchs for some $\mu \in (1, 2]$. Hence, in the scalar setting (to which the barrier techniques are limited), every Dirichlet problem is solvable according to Theorem 1.1, which provides a connection to the first approach to the minimization problem *via* relaxation described at the beginning of the introduction. Moreover, we have recovered that the Dirichlet problem in the prototypic example with coefficients (1.4) is in general solvable for arbitrary regular domains and boundary values if and only if $p \leq 1$ holds.

Remark 1.2. Concerning the assumptions on the domain and the function F , let us note:

- (i) A domain Ω satisfies the exterior ball condition if there exists a number $r_0 > 0$ such that for every point $\mathbf{x}_0 \in \partial\Omega$ there is a ball $B_{r_0}(\hat{\mathbf{x}}_0)$ with $\overline{B_{r_0}(\hat{\mathbf{x}}_0)} \cap \bar{\Omega} = \{\mathbf{x}_0\}$. Convexity or $\mathcal{C}^{1,1}$ -regularity of the domain are sufficient for the exterior ball condition (see *e.g.* [9], Thm. 1.9), thus, Theorem 1.1 holds in particular for all convex domains of class \mathcal{C}^1 and for arbitrary domains of class $\mathcal{C}^{1,1}$.

- (ii) The assumption $\lim_{s \rightarrow 0} F'(s) = 0$ ensures the ellipticity of equation (1.1) and the strict convexity of the associated minimization problem. For convenience, we can further assume that $\lim_{s \rightarrow 0} F''(s) = 0$ holds, which implies that $s \mapsto F(|s|)$ is of class $\mathcal{C}^2(\mathbb{R})$. In fact, if this was not the case, we could regularize it by a standard ε -mollifying kernel, which, for small $\varepsilon > 0$, preserves the structure assumptions (1.8) and (1.9). The result of Theorem 1.1 is then recovered in the limit $\varepsilon \searrow 0$, since all constants appearing in the derivation of a Lipschitz bound on the solution do only depend on those, and most crucially not on F'' for small values.
- (iii) For simplicity, but without loss of generality, we can restrict ourselves in the whole paper to functions F which satisfy

$$F(0) = 0 \quad \text{and} \quad \lim_{s \rightarrow \infty} \frac{F(s)}{s} = \lim_{s \rightarrow \infty} F'(s) = 1, \quad (1.10)$$

since the Dirichlet problem is invariant under addition and multiplication by a constant to F . Notice for the second relation that the function $s \mapsto F'(s)$ is monotonically increasing by convexity of F and therefore has a limit as $s \rightarrow \infty$. Moreover, strict positivity and finiteness of this limit follow from (1.8) and L'Hôpital's rule, which also shows that the two limits in (1.10) coincide.

- (iv) With a change of variables (and the normalization from the previous remark), we have

$$\int_1^\infty t F''(t) dt = \int_{F'(1)}^1 (F')^{-1}(s) ds.$$

Since the latter integral can be rewritten *via* the conjugate function

$$F^*(s^*) := \sup_{s \in \mathbb{R}^+} \{ss^* - F(s)\} \quad \text{for } s^* \in \mathbb{R}^+$$

(which appears in the dual formulation of the Dirichlet problem in the sense of convex analysis) as $F^*(1) - F^*(F'(1))$, we observe that condition (1.9) is satisfied if and only if the conjugate function F^* explodes when approaching the upper boundary of its domain $\{s^* \in \mathbb{R}^+ : F^*(s^*) < \infty\}$, which in the setting (1.10) is given by 1.

Let us note that the result of Theorem 1.1 could in fact be extended to less regular settings. One possibility is to consider convex functions F which are of class \mathcal{C}^2 only for large values. Since precisely only large gradients need to be avoided (uniformly) for the solutions to the approximative problems, such an asymptotic condition is in general sufficient (see *e.g.* [2], Thm. 1.2 for a related result). Moreover, if one is interested only in the existence of solutions in the space $u_0 + W_0^{1,1}(\Omega)$ (and not necessarily Lipschitz), one might work on domains which are only piece-wise of class $\mathcal{C}^{1,1}$ and for more general (not Lipschitz) boundary values u_0 .

We conclude the introduction with some comments on the notation used throughout the paper. For a set S in \mathbb{R}^d we write ∂S for its topological boundary and \bar{S} for its closure. Furthermore, for points in \mathbb{R}^d we use bold letters (like \mathbf{x} and \mathbf{x}_0), and the open ball in \mathbb{R}^d with center \mathbf{x}_0 and radius r is denoted by $B_r(\mathbf{x}_0)$. Concerning function spaces, we denote by \mathcal{C}^k the space of continuous functions with continuous derivatives up to order $k \in \mathbb{N}_0$ and by $\mathcal{C}^{k,\alpha}$ the related Hölder spaces, with Hölder exponent $\alpha \in (0, 1]$. We further work with the standard Lebesgue spaces L^p and Sobolev spaces $W^{1,p}$, for $p \in [1, \infty]$, and we abbreviate the respective norms by $\|\cdot\|_p$ and $\|\cdot\|_{1,p}$, when the domain of reference is clear from the context.

2. NON-EXISTENCE OF LIPSCHITZ MINIMIZERS

We first show the necessity of assumption (1.9) for the existence of Lipschitz minimizers to any regular boundary value problem to (1.1). In what follows, we provide a simple example of a regular domain Ω and of regular boundary values u_0 for which no Lipschitz solution (and in fact also no $W^{1,1}$ solution) to (1.1) exists. The construction is motivated by a well-known counterexample due to Finn [10] for the minimal surface equation

with $a_2(s) = (1 + s^2)^{-\frac{1}{2}}$, and we shall also here work on the annulus $\Omega := B_2(\mathbf{0}) \setminus B_1(\mathbf{0})$ and with boundary values u_0 which are constant on every connected component of $\partial\Omega$, that is, we consider

$$\begin{aligned} -\operatorname{div} \left(F'(|\nabla u|) \frac{\nabla u}{|\nabla u|} \right) &= 0 && \text{in } B_2(\mathbf{0}) \setminus B_1(\mathbf{0}), \\ u &= 0 && \text{on } \partial B_1(\mathbf{0}), \\ u &= M && \text{on } \partial B_2(\mathbf{0}), \end{aligned} \quad (2.1)$$

for some positive number $M \in \mathbb{R}^+$ (to be specified later). Moreover, we take $F \in \mathcal{C}^2(\mathbb{R}^+)$ as in Theorem 1.1 satisfying

$$\int_1^\infty t F''(t) dt = C_0 \quad (2.2)$$

for some positive constant C_0 and, without loss of generality, also the normalization assumption (1.10). Thanks to the strict convexity of F , the monotonicity of F' and the radial symmetry of both the domain and the prescribed boundary values, the Lipschitz (or even $W^{1,1}$) solution to (2.1), if it exists, is radially symmetric and can consequently be written as $u(\mathbf{x}) = U(|\mathbf{x}|)$ for all $\mathbf{x} \in \Omega$ and some (Lipschitz) function $U: [1, 2] \rightarrow \mathbb{R}$ with $U(1) = 0$ and $U(2) = M$. Thus, we also have $\nabla u(\mathbf{x}) = U'(|\mathbf{x}|) \frac{\mathbf{x}}{|\mathbf{x}|}$ for almost all $\mathbf{x} \in \Omega$. In order to find a representation formula of U , we take an arbitrary function $\Phi \in \mathcal{D}([1, 2])$ and extend it radially to a function $\varphi \in \mathcal{D}(\Omega)$ by setting $\varphi(\mathbf{x}) := \Phi(|\mathbf{x}|)$. Then we test the weak formulation of (2.1) with φ and find, by $\nabla \varphi(\mathbf{x}) = \Phi'(|\mathbf{x}|) \frac{\mathbf{x}}{|\mathbf{x}|}$, the transformation to polar coordinates and the radial symmetry of both functions U and Φ , the identity

$$\begin{aligned} 0 &= \int_\Omega F'(|\nabla u(\mathbf{x})|) \frac{\nabla u(\mathbf{x})}{|\nabla u(\mathbf{x})|} \cdot \nabla \varphi(\mathbf{x}) d\mathbf{x} \\ &= \int_\Omega F'(|U'(|\mathbf{x}|)|) \frac{U'(|\mathbf{x}|)}{|U'(|\mathbf{x}|)|} \Phi'(|\mathbf{x}|) d\mathbf{x} \\ &= d\omega_d \int_1^2 r^{d-1} F'(|U'(r)|) \frac{U'(r)}{|U'(r)|} \Phi'(r) dr, \end{aligned}$$

where ω_d denotes the Lebesgue measure of the unit ball in \mathbb{R}^d (and hence, $d\omega_d$ is the $(d-1)$ -dimensional Hausdorff measure of the unit sphere in \mathbb{R}^d). Since $\Phi \in \mathcal{D}([1, 2])$ was arbitrary, we deduce in a first step

$$F'(|U'(r)|) \frac{U'(r)}{|U'(r)|} = \frac{c}{r^{d-1}}$$

for some constant c and all $r \in [1, 2]$. By assumptions on F , we next observe that no sign change of U' may occur, hence, U' is positive everywhere in $[1, 2]$ and we have indeed

$$1 > F'(U'(r)) = \frac{c}{r^{d-1}},$$

hence also $c \in (0, 1)$. After inverting the last identity we can integrate (keeping in mind the boundary condition $U(1) = 0$) and find the desired representation formula

$$U(r) = \int_1^r (F')^{-1} \left(\frac{c}{s^{d-1}} \right) ds$$

for all $r \in [1, 2]$. In turn, with the substitution $c/s^{d-1} = z$, that is, $s = c^{\frac{1}{d-1}} z^{\frac{1}{1-d}}$, we obtain the following upper bound on U :

$$\begin{aligned} U(r) &= \int_c^{c/r^{d-1}} \frac{c^{\frac{1}{d-1}}}{1-d} z^{\frac{d}{1-d}} (F')^{-1}(z) dz \\ &= \frac{c^{\frac{1}{d-1}}}{d-1} \int_{c/r^{d-1}}^c z^{\frac{d}{1-d}} (F')^{-1}(z) dz \leq \frac{2^d}{c(d-1)} \int_{c/r^{d-1}}^c (F')^{-1}(z) dz. \end{aligned}$$

This provides indeed a nontrivial upper bound, as can be seen by a case distinction between small and large values of c , in order to estimate the integral appearing on the right-hand side. In fact, in the case $0 < c \leq F'(1) < 1$, we find by monotonicity of $(F')^{-1}$

$$\int_{c/r^{d-1}}^c (F')^{-1}(z) \, dz \leq c(1 - r^{1-d}) < c,$$

while in the opposite case $0 < F'(1) < c < 1$ the change of variables with $z = F'(t)$ combined with (2.2) yields

$$\int_{c/r^{d-1}}^c (F')^{-1}(z) \, dz = \int_{(F')^{-1}(c/r^{d-1})}^{(F')^{-1}(c)} tF''(t) \, dt \leq \int_0^\infty tF''(t) \, dt \leq F'(1) + C_0.$$

In conclusion, for all $r \in [1, 2]$, we have derived the explicit upper bound

$$U(r) \leq \frac{2^d}{d-1} \left(1 + \frac{C_0}{F'(1)} \right),$$

which is a contradiction to $U(2) = M$ for $M \in \mathbb{R}^+$ sufficiently large. Thus, we have just proven that if (1.9) does not hold, then we can find a smooth domain Ω and smooth boundary values u_0 such that all assumptions of Theorem (1.1) are satisfied, and such that no Lipschitz solution to problem (1.1) exists.

Remark 2.1. More subtle non-existence results can be obtained, similarly as in the case of the minimal surface equation. In particular, the class of regular domains which allow for non-existence results can be investigated. Based on the previous construction and a comparison principle, one can in fact show that for every non-pseudoconvex regular domain Ω there exist smooth prescribed boundary values u_0 such that no Lipschitz solution to the Dirichlet problem (1.1) exists, cp. [10].

3. AUXILIARY LEMMATA

In order to proceed to the proof of the second implication of Theorem 1.1, we first derive some auxiliary algebraic inequalities.

Lemma 3.1. *Let $F \in \mathcal{C}^2(\mathbb{R}^+)$ be a strictly convex function with $\lim_{s \rightarrow 0} F'(s) = 0$ which satisfies (1.8), (1.9) and (1.10), and let $a \in \mathcal{C}^1(\mathbb{R}^+)$ be given by (1.3). Then, there hold*

$$C_1 s - C_2 \leq sF'(s) \leq s \quad \text{for every } s > 0, \quad (3.1)$$

$$\lim_{s \rightarrow \infty} sF''(s) = 0, \quad (3.2)$$

$$\lim_{s \rightarrow \infty} s^2 a'(s) = -1. \quad (3.3)$$

Proof. We start by observing some simple consequences of the strict convexity of F . As F' is monotonically increasing, assumptions (1.8) and (1.10) give

$$C_1 s - C_2 \leq F(s) = \int_0^s F'(r) \, dr \leq \int_0^s F'(s) \, dr = sF'(s),$$

which is the first inequality in (3.1). We further observe that the assumptions $\lim_{s \rightarrow 0} F'(s) = 0$ and $\lim_{s \rightarrow \infty} F'(s) = 1$ yield immediately $0 < F'(s) < 1$ for all $s \in (0, \infty)$, which implies the second inequality in (3.1), and moreover, $F'' \in L^1(0, \infty)$ holds. Due to the integrability of F'' , we next deduce the identity

$$a(s)s = F'(s) = 1 - \int_s^\infty F''(t) \, dt, \quad (3.4)$$

and since by the bounded oscillation assumption in (1.8) we have in particular

$$F''(s)s \leq C_2 \int_s^{2s} F''(t) dt$$

for all $s > 0$, also the claim (3.2) follows. Finally, differentiating (3.4) we find

$$a'(s)s + a(s) = F''(s), \quad (3.5)$$

and therefore, thanks to (3.2) and (1.10), we obtain

$$\lim_{s \rightarrow \infty} s^2 a'(s) = \lim_{s \rightarrow \infty} (sF''(s) - F'(s)) = -1,$$

which is the last claim (3.3). \square

Secondly, we define the integrand for a comparison functional, which will be used later for the construction of appropriate barrier functions. To this end, we will essentially decrease the convexity for large values of the original integrand F (note that the properties of the integrand for large values are the most crucial ones), which from a heuristic point of view will make it harder to construct solutions (cp. the calculations for annular domains in Sect. 2). However, it turns out that as long as the fundamental condition (1.9) for the new integrand is satisfied, this construction is still possible, and it is precisely the weaker convexity for large values which will allow for the verification of the barrier condition.

Lemma 3.2. *Let $F \in \mathcal{C}^2(\mathbb{R}^+)$ be a strictly convex function with $\lim_{s \rightarrow 0} F'(s) = 0$ which satisfies (1.8), (1.9) and (1.10). Then there exists a strictly positive, decreasing function $g \in \mathcal{C}(\mathbb{R}_0^+)$ with $\lim_{s \rightarrow \infty} g(s) = 0$ such that*

$$\int_0^\infty tF''(t)g(t) dt = \infty. \quad (3.6)$$

Moreover, the function F_g defined via

$$F_g(s) := \int_0^s \left(1 - \int_r^\infty F''(t)g(t) dt \right) dr \quad (3.7)$$

belongs to $\mathcal{C}^2(\mathbb{R}^+)$, is strictly convex and satisfies $\lim_{s \rightarrow 0} F'_g(s) = 0$ and (1.8)–(1.10), with possibly different constants C_1 and C_2 .

Proof. We start by defining

$$\tilde{g}(s) := \frac{1}{1 + \int_0^s tF''(t) dt}, \quad (3.8)$$

and we note that \tilde{g} is a continuous, strictly positive, strictly decreasing function on \mathbb{R}_0^+ , which is bounded from above by 1 and which fulfills $\tilde{g}(s) \rightarrow 0$ as $s \rightarrow \infty$. Next, we define a constant

$$A := \int_0^\infty F''(t)\tilde{g}(t) dt,$$

and due to the properties of F (more precisely, $\lim_{s \rightarrow 0} F'(s) = 0$ and $\lim_{s \rightarrow \infty} F'(s) = 1$) combined with $0 < \tilde{g}(s) \leq 1$ for $s \in \mathbb{R}^+$, we observe $A \in (0, 1]$. Finally, we define

$$g(s) := \frac{\tilde{g}(s)}{A}, \quad (3.9)$$

which again is strictly positive, decreasing and with $\lim_{s \rightarrow \infty} g(s) = 0$. We now show that all statements of the lemma are indeed fulfilled with such a choice of g . Let us start with (3.6). By the definition of g it directly follows that for any $s \geq 0$ we have

$$\begin{aligned} \int_0^s tF''(t)g(t) \, dt &= \frac{1}{A} \int_0^s \frac{tF''(t)}{1 + \int_0^t rF''(r) \, dr} \, dt \\ &= \frac{1}{A} \int_0^s \frac{d}{dt} \ln \left(1 + \int_0^t rF''(r) \, dr \right) \, dt \\ &= \frac{1}{A} \ln \left(1 + \int_0^s rF''(r) \, dr \right). \end{aligned}$$

Since F'' and g are positive and due to the assumption (1.9), we find

$$\begin{aligned} \int_0^\infty tF''(t)g(t) \, dt &= \lim_{s \rightarrow \infty} \int_0^s tF''(t)g(t) \, dt \\ &= \frac{1}{A} \lim_{s \rightarrow \infty} \ln \left(1 + \int_0^s rF''(r) \, dr \right) = \infty. \end{aligned}$$

Hence, (3.6) holds. Next, we show the properties of the function F_g . By its definition (3.7) we immediately obtain $F_g(0) = 0$. Moreover, we calculate its derivatives

$$F'_g(s) = 1 - \int_s^\infty F''(t)g(t) \, dt, \quad F''_g(s) = F''(s)g(s) > 0, \quad (3.10)$$

which shows the strict convexity and, via (3.6), the validity of (1.9) for F_g . Since by the definition of g we have

$$\int_0^\infty F''(t)g(t) \, dt = 1,$$

we find $\lim_{s \rightarrow 0} F'_g(s) = 0$ and $\lim_{s \rightarrow \infty} F'_g(s) = 1$, thus also (1.10) is satisfied for F_g . Finally, concerning (1.8), we note that the linear growth assumption follows immediately from the properties of F'_g , while the oscillation assumption of F carries directly over to F_g (with constant $C_2 + 4C_2^2$ instead of C_2). This concludes the proof of the lemma. \square

4. PROOF OF THEOREM 1.1

We now prove the second (and main) implication of Theorem 1.1, namely that assumption (1.9) implies the general solvability of the Dirichlet problem (1.1). To this end, we perform an approximation procedure and introduce suitable approximation problems to (1.1), and we will then prove uniform estimates for their minimizers. This is indeed sufficient to recover the claim with the passage to the limit since the strict convexity of our functional implies uniqueness of minimizers (if a minimizer exists in the desired Dirichlet class at all). In order to follow this strategy, we define, for arbitrary $\varepsilon > 0$, the approximate functionals

$$w \mapsto \frac{\varepsilon}{2} \int_\Omega |\nabla w|^2 \, dx + \int_\Omega F(|\nabla w|) \, dx$$

and look for minimizers u_ε in the Dirichlet class $u_0 + W_0^{1,2}(\Omega)$, which is equivalent to looking for weak solutions to the following approximate Dirichlet problem to (1.1)

$$\begin{aligned} -\varepsilon \Delta u_\varepsilon - \operatorname{div} (a(|\nabla u_\varepsilon|) \nabla u_\varepsilon) &= 0 & \text{in } \Omega, \\ u_\varepsilon &= u_0 & \text{on } \partial\Omega. \end{aligned} \quad (4.1)$$

Note that for the rest of the paper, we suppose, without explicit mentioning, that the function F satisfies (1.8), (1.9) and (1.10), and that a is related to F via (1.3).

Due to the application of the direct method of the calculus of variations to the approximate functionals (or the theory of monotone operators to the approximate Dirichlet problems), we obtain the existence of a unique solution $u_\varepsilon \in u_0 + W_0^{1,2}(\Omega)$. In addition, in view of the regularity of the prescribed boundary values, we observe that we have $u_\varepsilon \in C^{1,\alpha}(\bar{\Omega})$ for some $\alpha > 0$, see *e.g.* [12], and by difference quotient techniques we also have $u_\varepsilon \in W_{\text{loc}}^{2,2}(\Omega)$. Our main goal is to show that the following uniform estimate holds

$$\|\nabla u_\varepsilon\|_\infty \leq C(\Omega, F, u_0) \quad (4.2)$$

with some constant $C(\Omega, F, u_0)$ being independent of ε . Indeed, having (4.2) in hands, we first find a subsequence converging weakly-* to a function u in $u_0 + W_0^{1,\infty}(\Omega)$. Then, when passing to the limit $\varepsilon \rightarrow 0$ in the approximate functionals (*via* lower semicontinuity) or in the approximate Dirichlet problems (4.1) (*via* the theory of monotone operators), the limit $u \in u_0 + W_0^{1,\infty}(\Omega)$ turns out to be the desired solution. As it is actually Lipschitz regular, Theorem 1.1 is therefore proven, provided that we can show that the crucial uniform estimate (4.2) holds.

4.1. Reduction to the boundary estimates

We first show that, in order to establish (4.2), it is actually sufficient to control the normal derivatives of the solutions u_ε to the approximate problems (4.1) uniformly on the boundary $\partial\Omega$. To this end, we start by deriving some standard uniform estimates and denote by C an universal constant depending only on F , u_0 and Ω , but not on ε . For simplicity of notation, we shall drop from now on the index ε and write u instead of u_ε . Testing the weak formulation to (4.1) with the function $u - u_0 \in W_0^{1,2}(\Omega)$, keeping in mind relation (1.3) and applying Hölder's inequality, we obtain

$$\varepsilon \|\nabla u\|_2^2 + \int_\Omega F'(|\nabla u|) |\nabla u| \, dx \leq \varepsilon \|\nabla u\|_2 \|\nabla u_0\|_2 + \int_\Omega F'(|\nabla u|) |\nabla u_0| \, dx.$$

Hence, using Young's inequality and (3.1), we deduce

$$\varepsilon \|\nabla u\|_2^2 + \|\nabla u\|_1 \leq C. \quad (4.3)$$

Similarly, testing the weak formulation to (4.1) with the functions $(u \mp \|u_0\|_\infty)_\pm$ (note that these functions are admissible since $(u \mp \|u_0\|_\infty)_\pm = 0$ holds on $\partial\Omega$), we get

$$\varepsilon \int_\Omega |\nabla(u \mp \|u_0\|_\infty)_\pm|^2 \, dx + \int_\Omega a(|\nabla u|) |\nabla(u \mp \|u_0\|_\infty)_\pm|^2 \, dx = 0.$$

Thus, it follows that

$$\|u\|_\infty \leq \|u_0\|_\infty \leq C. \quad (4.4)$$

To proceed further, we identify the equation for $|\nabla u|$, which is possible due to the *a priori* (non-uniform) regularity of u . Applying $\frac{\partial}{\partial x_k} =: D_k$ to (4.1), multiplying the result by $D_k u$ and summing over $k = 1, \dots, d$, we obtain

$$\begin{aligned} 0 &= -\varepsilon \sum_{k=1}^d D_k u \Delta D_k u - \sum_{k,i=1}^d D_k u D_i D_k \left(F'(|\nabla u|) \frac{D_i u}{|\nabla u|} \right) \\ &= -\frac{\varepsilon}{2} \Delta |\nabla u|^2 + \varepsilon |\nabla^2 u|^2 - \sum_{k,i=1}^d D_i \left(D_k \left(F'(|\nabla u|) \frac{D_i u}{|\nabla u|} \right) D_k u \right) \\ &\quad + \sum_{k,i=1}^d D_{ik} u D_k \left(F'(|\nabla u|) \frac{D_i u}{|\nabla u|} \right) \\ &= -\frac{\varepsilon}{2} \Delta |\nabla u|^2 - \sum_{k,i=1}^d D_i (A_{ik}(\nabla u) D_k |\nabla u|) + \varepsilon |\nabla^2 u|^2 + F''(|\nabla u|) |\nabla |\nabla u||^2 + F'(|\nabla u|) \frac{|\nabla^2 u|^2 - |\nabla |\nabla u||^2}{|\nabla u|}, \end{aligned}$$

where

$$A_{ik}(\nabla u) := \left(|\nabla u| F''(|\nabla u|) \frac{D_i u D_k u}{|\nabla u|^2} + F'(|\nabla u|) \delta_{ik} - F'(|\nabla u|) \frac{D_i u D_k u}{|\nabla u|^2} \right).$$

Consequently,

$$-\frac{\varepsilon}{2} \Delta |\nabla u|^2 - \sum_{k,i=1}^d D_i (A_{ik}(\nabla u) D_k |\nabla u|) \leq 0.$$

Since A is positively definite, we see that $|\nabla u|^2$ is a sub-solution to a linear elliptic equation and therefore satisfies the minimum principle, *i.e.*,

$$\|\nabla u\|_{L^\infty(\Omega)} \leq \|\nabla u\|_{L^\infty(\partial\Omega)}.$$

In addition, since $u = u_0$ on $\partial\Omega$, this implies

$$\|\nabla u\|_{L^\infty(\Omega)} \leq \|\nabla u_0\|_{L^\infty(\Omega)} + \left\| \frac{\partial u}{\partial \mathbf{n}} \right\|_{L^\infty(\partial\Omega)},$$

where $\frac{\partial u}{\partial \mathbf{n}}$ denotes the normal derivative of u on $\partial\Omega$. Thus, in order to check (4.2) it remains to show that

$$\left\| \frac{\partial u}{\partial \mathbf{n}} \right\|_{L^\infty(\partial\Omega)} \leq C(\Omega, F, u_0). \quad (4.5)$$

The rest of the paper is devoted to the proof of (4.5), which will be shown *via* the barrier function technique.

4.2. Prototype barrier function

From now on, we fix the functions g and F_g according to Lemma 3.2 and define

$$a_g(s) := \frac{F'_g(s)}{s} \quad \text{for } s \in \mathbb{R}^+. \quad (4.6)$$

Clearly, all statements of Lemma 3.1 hold also for F_g and a_g with possibly different constants $C_1, C_2 > 0$. Moreover, F'_g is a strictly monotonically increasing mapping from $[0, \infty)$ to $[0, 1)$ with continuous inverse. With the help of F'_g we now define our prototype barrier function.

Let $r_0 > 0$ and $\delta \in (0, 1)$ be arbitrary. We set for all $r \geq r_0$

$$b_{r_0}^\delta(r) := (F'_g)^{-1} \left(\frac{(1-\delta)^{d-1} r_0^{d-1}}{r^{d-1}} \right). \quad (4.7)$$

It can be easily seen that $b_{r_0}^\delta \in C^1[r_0, \infty)$ is a non-negative decreasing function. Finally, for all $\mathbf{x} \in \mathbb{R}^d \setminus B_{r_0}(\mathbf{0})$, we define

$$\omega_{r_0}^\delta(\mathbf{x}) := \int_{r_0}^{|\mathbf{x}|} b_{r_0}^\delta(r) \, dr. \quad (4.8)$$

By construction, $\omega_{r_0}^\delta$ is a minimizer of the functional with integrand F_g and equivalently a solution to the associated Dirichlet problem on the set $\mathbb{R}^d \setminus \overline{B_{r_0}(\mathbf{0})}$, but moreover, it also turns out to be super-harmonic on a subset of it.

Lemma 4.1. *For every $r_0 > 0$ and $\delta \in (0, 1)$ the function $\omega_{r_0}^\delta$ defined in (4.8) satisfies*

$$\begin{aligned} -\operatorname{div} (a_g(|\nabla \omega_{r_0}^\delta|) \nabla \omega_{r_0}^\delta) &= 0 & \text{in } \mathbb{R}^d \setminus \overline{B_{r_0}(\mathbf{0})}, \\ \omega_{r_0}^\delta &= 0 & \text{on } \partial B_{r_0}(\mathbf{0}). \end{aligned} \quad (4.9)$$

Furthermore, there holds

$$-\Delta \omega_{r_0}^\delta(\mathbf{x}) \geq 0 \quad \text{for all } \mathbf{x} \in \mathbb{R}^d \setminus \overline{B_{r_0}(\mathbf{0})} \quad \text{such that} \quad a'_g(b_{r_0}^\delta(|\mathbf{x}|)) \leq 0. \quad (4.10)$$

Proof. Using the definition of $\omega_{r_0}^\delta$, we immediately see that $\omega_{r_0}^\delta$ vanishes on $\partial B_{r_0}(\mathbf{0})$, and we further observe

$$\nabla \omega_{r_0}^\delta(\mathbf{x}) = b_{r_0}^\delta(|\mathbf{x}|) \frac{\mathbf{x}}{|\mathbf{x}|} \quad \text{and} \quad |\nabla \omega_{r_0}^\delta(\mathbf{x})| = b_{r_0}^\delta(|\mathbf{x}|). \quad (4.11)$$

Via the definition of $b_{r_0}^\delta$, we thus have

$$F'_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|) = F'_g(b_{r_0}^\delta(|\mathbf{x}|)) = \frac{(1-\delta)^{d-1} r_0^{d-1}}{|\mathbf{x}|^{d-1}}.$$

Consequently, for all $|\mathbf{x}| > r_0$ there holds

$$\begin{aligned} \operatorname{div} (a_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|) \nabla \omega_{r_0}^\delta(\mathbf{x})) &= \operatorname{div} \left(F'_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|) \frac{\nabla \omega_{r_0}^\delta(\mathbf{x})}{|\nabla \omega_{r_0}^\delta(\mathbf{x})|} \right) \\ &= (1-\delta)^{d-1} r_0^{d-1} \operatorname{div} \left(\frac{\mathbf{x}}{|\mathbf{x}|^d} \right) = 0 \end{aligned} \quad (4.12)$$

and the solution property (4.9) follows. Finally, we check the super-harmonicity property of $\omega_{r_0}^\delta$. In view of (4.9) and (4.11) we get

$$\begin{aligned} 0 &= -\operatorname{div} (a_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|) \nabla \omega_{r_0}^\delta(\mathbf{x})) \\ &= -a_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|) \Delta \omega_{r_0}^\delta(\mathbf{x}) - \nabla a_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|) \cdot \nabla \omega_{r_0}^\delta(\mathbf{x}) \\ &= -a_g(b_{r_0}^\delta(|\mathbf{x}|)) \Delta \omega_{r_0}^\delta(\mathbf{x}) - a'_g(b_{r_0}^\delta(|\mathbf{x}|)) b_{r_0}^\delta(|\mathbf{x}|) (b_{r_0}^\delta(|\mathbf{x}|))'(|\mathbf{x}|). \end{aligned}$$

Therefore, since the functions a_g and $b_{r_0}^\delta$ are positive and $b_{r_0}^\delta$ is monotonically decreasing, also the second claim (4.10) follows. \square

Thus, $\omega_{r_0}^\delta$ is a good prototype super-solution to (4.1) on a certain set. However, due to the possibly non-constant prescribed boundary values u_0 , it must be corrected, which will be done in the next step.

4.3. True barrier function

Here, we correct $\omega_{r_0}^\delta$ via an affine function such that it will finally give us the desired super-solution property to (4.1). For this purpose, let $\mathbf{k} \in \mathbb{R}^d$, $c \in \mathbb{R}$, $r_0 > 0$ and $\delta \in (0, 1)$ be arbitrary. For all $\mathbf{x} \in \mathbb{R}^d \setminus B_{r_0}(\mathbf{0})$, we define

$$v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) := \omega_{r_0}^\delta(\mathbf{x}) + \mathbf{k} \cdot \mathbf{x} + c. \quad (4.13)$$

The key properties of the function $v_{\mathbf{k},c}^{\delta,r_0}$ are formulated in the following lemma.

Lemma 4.2. *For every $K > 0$ there exists a number $M > 0$ depending only on F and K such that for all $\mathbf{k} \in B_K(\mathbf{0})$, all $c \in \mathbb{R}$, all $\delta \in (0, 1)$ and all $r_0 > 0$ the function $v_{\mathbf{k},c}^{\delta,r_0}$ defined in (4.13) satisfies the inequalities*

$$\begin{aligned} -\operatorname{div} (a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})) &\geq 0, \\ -\Delta v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) &\geq 0 \end{aligned} \quad (4.14)$$

for all $\mathbf{x} \in \mathbb{R}^d \setminus B_{r_0}(\mathbf{0})$ fulfilling $b_{r_0}^\delta(|\mathbf{x}|) \geq M$ with $b_{r_0}^\delta$ given by (4.7).

Proof. First, it is evident that for all $\mathbf{x} \in \mathbb{R}^d \setminus B_{r_0}(\mathbf{0})$

$$\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) = \nabla \omega_{r_0}^\delta(\mathbf{x}) + \mathbf{k} = b_{r_0}^\delta(|\mathbf{x}|) \frac{\mathbf{x}}{|\mathbf{x}|} + \mathbf{k}. \quad (4.15)$$

Consequently, a direct computation leads to

$$\begin{aligned} |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|^2 &= (b_{r_0}^\delta)^2(|\mathbf{x}|) + |\mathbf{k}|^2 + 2b_{r_0}^\delta(|\mathbf{x}|) \frac{\mathbf{k} \cdot \mathbf{x}}{|\mathbf{x}|}, \\ \nabla |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})| &= \frac{b_{r_0}^\delta(|\mathbf{x}|)(b_{r_0}^\delta)'(|\mathbf{x}|) \frac{\mathbf{x}}{|\mathbf{x}|} + (b_{r_0}^\delta)'(|\mathbf{x}|) \frac{\mathbf{x}}{|\mathbf{x}|} \frac{\mathbf{k} \cdot \mathbf{x}}{|\mathbf{x}|} + b_{r_0}^\delta(|\mathbf{x}|) \left(\frac{\mathbf{k}}{|\mathbf{x}|} - \frac{(\mathbf{k} \cdot \mathbf{x})\mathbf{x}}{|\mathbf{x}|^3} \right)}{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|}. \end{aligned}$$

Hence, using these identities, we obtain the following auxiliary results that will be used later

$$\nabla |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})| \cdot \frac{\mathbf{x}}{|\mathbf{x}|} = (b_{r_0}^\delta)'(|\mathbf{x}|) \frac{b_{r_0}^\delta(|\mathbf{x}|) + \frac{\mathbf{k} \cdot \mathbf{x}}{|\mathbf{x}|}}{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|} \quad (4.16)$$

and

$$\nabla |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})| \cdot \mathbf{k} = \frac{b_{r_0}^\delta(|\mathbf{x}|)(b_{r_0}^\delta)'(|\mathbf{x}|) \frac{\mathbf{x} \cdot \mathbf{k}}{|\mathbf{x}|} + (b_{r_0}^\delta)'(|\mathbf{x}|) \frac{(\mathbf{k} \cdot \mathbf{x})^2}{|\mathbf{x}|^2} + b_{r_0}^\delta(|\mathbf{x}|) \left(\frac{|\mathbf{k}|^2}{|\mathbf{x}|} - \frac{(\mathbf{k} \cdot \mathbf{x})^2}{|\mathbf{x}|^3} \right)}{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|}. \quad (4.17)$$

Let us now evaluate the super-solution and super-harmonicity properties. To this end, we introduce the abbreviation

$$\begin{aligned} L(\mathbf{x}) &:= -\operatorname{div} (a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})) \\ &= -\nabla a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \cdot \nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) - a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \operatorname{div} (\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})) =: L_1(\mathbf{x}) + L_2(\mathbf{x}). \end{aligned}$$

Employing (4.15), (4.16) and (4.17), we first calculate

$$\begin{aligned} L_1(\mathbf{x}) &= -a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \nabla |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})| \cdot \nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) \\ &= -a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \nabla |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})| \cdot \left[b_{r_0}^\delta(|\mathbf{x}|) \frac{\mathbf{x}}{|\mathbf{x}|} + \mathbf{k} \right] \\ &= -\frac{a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|)}{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|} \left[b_{r_0}^\delta(|\mathbf{x}|)(b_{r_0}^\delta)'(|\mathbf{x}|) \left(b_{r_0}^\delta(|\mathbf{x}|) + \frac{\mathbf{k} \cdot \mathbf{x}}{|\mathbf{x}|} \right) \right. \\ &\quad \left. + b_{r_0}^\delta(|\mathbf{x}|)(b_{r_0}^\delta)'(|\mathbf{x}|) \frac{\mathbf{x} \cdot \mathbf{k}}{|\mathbf{x}|} + (b_{r_0}^\delta)'(|\mathbf{x}|) \frac{(\mathbf{k} \cdot \mathbf{x})^2}{|\mathbf{x}|^2} + b_{r_0}^\delta(|\mathbf{x}|) \left(\frac{|\mathbf{k}|^2}{|\mathbf{x}|} - \frac{(\mathbf{k} \cdot \mathbf{x})^2}{|\mathbf{x}|^3} \right) \right] \\ &= \frac{a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|)}{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|} (b_{r_0}^\delta)'(|\mathbf{x}|) \left(|\mathbf{x}| - \frac{b_{r_0}^\delta(|\mathbf{x}|)}{(b_{r_0}^\delta)'(|\mathbf{x}|)} \right) \left(\frac{|\mathbf{k}|^2}{|\mathbf{x}|} - \frac{(\mathbf{k} \cdot \mathbf{x})^2}{|\mathbf{x}|^3} \right) \\ &\quad - a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) (b_{r_0}^\delta)'(|\mathbf{x}|) |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|. \end{aligned}$$

Next, taking into account once again (4.15), the relation (4.11) and the fact that $\omega_{r_0}^\delta$ solves equation (4.9), we find

$$\begin{aligned} L_2(\mathbf{x}) &= -a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \operatorname{div} \left(\frac{a_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|) \nabla \omega_{r_0}^\delta(\mathbf{x})}{a_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|)} \right) \\ &= \frac{a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) a_g'(|\nabla \omega_{r_0}^\delta(\mathbf{x})|)}{a_g(|\nabla \omega_{r_0}^\delta(\mathbf{x})|)} \nabla |\nabla \omega_{r_0}^\delta(\mathbf{x})| \cdot \nabla \omega_{r_0}^\delta(\mathbf{x}) \\ &= \frac{a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) a_g'(b_{r_0}^\delta(|\mathbf{x}|))}{a_g(b_{r_0}^\delta(|\mathbf{x}|))} (b_{r_0}^\delta)'(|\mathbf{x}|) b_{r_0}^\delta(|\mathbf{x}|). \end{aligned}$$

In conclusion, after a simple algebraic manipulation, we have

$$\begin{aligned} L(\mathbf{x}) &= \frac{a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|)}{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|} (b_{r_0}^\delta)'(|\mathbf{x}|) \left(|\mathbf{x}| - \frac{b_{r_0}^\delta(|\mathbf{x}|)}{(b_{r_0}^\delta)'(|\mathbf{x}|)} \right) \left(\frac{|\mathbf{k}|^2}{|\mathbf{x}|} - \frac{(\mathbf{k} \cdot \mathbf{x})^2}{|\mathbf{x}|^3} \right) \\ &\quad - a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) (b_{r_0}^\delta)'(|\mathbf{x}|) \left(\frac{a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|}{a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|)} - \frac{a'_g(b_{r_0}^\delta(|\mathbf{x}|))}{a_g(b_{r_0}^\delta(|\mathbf{x}|))} b_{r_0}^\delta(|\mathbf{x}|) \right) \\ &=: \tilde{L}_1(\mathbf{x}) + \tilde{L}_2(\mathbf{x}). \end{aligned}$$

We now focus on estimating the resulting terms and we will show that both are non-negative in a suitably chosen set. To this end, we first relate $b_{r_0}^\delta(|\mathbf{x}|)$ and $|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|$ and provide some basic estimates, for sufficiently large values of $b_{r_0}^\delta(|\mathbf{x}|)$. Since $|\mathbf{k}| \leq K$, we deduce from (4.15) that for $M_1 := 2K > 0$ there holds

$$b_{r_0}^\delta(|\mathbf{x}|) \geq M_1 \implies b_{r_0}^\delta(|\mathbf{x}|) \leq 2|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})| \leq 4b_{r_0}^\delta(|\mathbf{x}|). \quad (4.18)$$

In turn, relying on (3.3) (for both functions a and a_g), we find a constant $M_2 \geq M_1$ depending only on F , g and K such that

$$\begin{aligned} b_{r_0}^\delta(|\mathbf{x}|) \geq M_2 &\implies \begin{cases} (b_{r_0}^\delta)^2(|\mathbf{x}|) a'_g(b_{r_0}^\delta(|\mathbf{x}|)) \leq -\frac{1}{2} \\ |\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|^2 a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \leq -\frac{1}{2} \end{cases} \\ &\implies a'_g(b_{r_0}^\delta(|\mathbf{x}|)) < 0 \quad \text{and} \quad a'(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) < 0. \end{aligned} \quad (4.19)$$

This implication now allows us to deduce the positivity of $\tilde{L}_1(\mathbf{x})$ and the super-harmonicity of $v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})$ (thus, the second claim of the lemma), provided that $b_{r_0}^\delta(|\mathbf{x}|) \geq M_2$ holds. In fact, since $b_{r_0}^\delta$ is a non-negative decreasing function and by using the Cauchy-Schwarz inequality, we see that the first and second expression in large brackets in the definition of $\tilde{L}_1(\mathbf{x})$ are non-negative. Thus, in view of (4.19) and once again the monotonicity of $b_{r_0}^\delta$, the sign of $\tilde{L}_1(\mathbf{x})$ is non-negative. Secondly, (4.15) yields $\Delta v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) = \Delta \omega_{r_0}^\delta(\mathbf{x})$, thus the super-harmonicity of $v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})$ follows from (4.10) and (4.19). In conclusion, we have the implication

$$b_{r_0}^\delta(|\mathbf{x}|) \geq M_2 \implies -\Delta v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) \geq 0 \quad \text{and} \quad \tilde{L}_1(\mathbf{x}) \geq 0. \quad (4.20)$$

Finally, we discuss the sign of $\tilde{L}_2(\mathbf{x})$. Using (3.5), (3.4), (3.10) and (4.6), we evaluate

$$\begin{aligned} \frac{a'(s)s}{a(s)} &= \frac{F''(s)}{a(s)} - 1 = \frac{sF''(s)}{1 - \int_s^\infty F''(t) dt} - 1, \\ \frac{a'_g(s)s}{a_g(s)} &= \frac{sF_g''(s)}{F_g'(s)} - 1 = \frac{s g(s)F''(s)}{1 - \int_s^\infty g(t)F''(t) dt} - 1. \end{aligned}$$

In this way the expression for $\tilde{L}_2(\mathbf{x})$ reduces to

$$\tilde{L}_2(\mathbf{x}) = -a(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) (b_{r_0}^\delta)'(|\mathbf{x}|) \times \left(\frac{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})| F''(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|)}{1 - \int_{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|}^\infty F''(t) dt} - \frac{b_{r_0}^\delta(|\mathbf{x}|) g(b_{r_0}^\delta(|\mathbf{x}|)) F''(b_{r_0}^\delta(|\mathbf{x}|))}{1 - \int_{b_{r_0}^\delta(|\mathbf{x}|)}^\infty g(t) F''(t) dt} \right). \quad (4.21)$$

For $b_{r_0}^\delta(|\mathbf{x}|) \geq M_2$, we then find, employing (4.18), the oscillation assumption (1.8) on F and the fact that g is a positive and monotonically decreasing function, the inequalities

$$|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})| F''(|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|) \geq \frac{1}{2C_2} b_{r_0}^\delta(|\mathbf{x}|) F''(b_{r_0}^\delta(|\mathbf{x}|))$$

and

$$\int_{|\nabla v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})|}^{\infty} F''(t) dt \geq \frac{1}{2C_2 g(b_{r_0}^{\delta}(|\mathbf{x}|))} \int_{b_{r_0}^{\delta}(|\mathbf{x}|)}^{\infty} g(t) F''(t) dt.$$

At this stage, we select a number $M \geq M_2$ depending only on F , g and K such that $2C_2 g(M) \leq 1$ holds which is possible because of $g(t) \rightarrow 0$ as $t \rightarrow \infty$, according to Lemma 3.2. With this choice and the integrability of F'' over \mathbb{R}^+ (with integral equal to 1), we see easily that the expression in large brackets on the right-hand side of (4.21) is non-negative, whenever $b_{r_0}^{\delta}(|\mathbf{x}|) \geq M$ holds. Consequently, using also the facts that $b_{r_0}^{\delta}$ is monotonically decreasing and that a is non-negative (thanks to (1.3) and the non-negativity of F'), we arrive at the implication

$$b_{r_0}^{\delta}(|\mathbf{x}|) \geq M \implies \tilde{L}_2(\mathbf{x}) \geq 0.$$

Combined with (4.20), we finally conclude that $L(\mathbf{x}) \geq 0$ holds for all $\mathbf{x} \in \mathbb{R}^d \setminus B_{r_0}(\mathbf{0})$ with $b_{r_0}^{\delta}(|\mathbf{x}|) \geq M$, and the proof of the lemma is complete. \square

4.4. Estimates of the normal derivatives

Once the true barrier function from Lemma 4.2 is at our disposal, we can return to study the normal derivative, with the aim to prove an estimate of the form (4.5).

The strategy of proof is as follows: by adjusting the true barrier function from Lemma 4.2 to our needs, we construct in a first step a (*local upper*) *barrier function* relative to the Dirichlet problem (1.1) for an arbitrary given boundary point \mathbf{x}_0 . This means that we specify a relative neighborhood $U(\mathbf{x}_0)$ of \mathbf{x}_0 in Ω and a local Lipschitz continuous function v (which will be an affine perturbation of the function from Lemma 4.2) defined on $U(\mathbf{x}_0)$ such that

- (i) v is a super-solution of the equation in $U(\mathbf{x}_0)$, i.e., $-\operatorname{div}(a(|\nabla v|)\nabla v) \geq 0$ in $U(\mathbf{x}_0)$,
- (ii) v lies above the solution u on $\partial U(\mathbf{x}_0)$ and coincides with it in \mathbf{x}_0 , i.e. $v \geq u$ on $\partial U(\mathbf{x}_0)$ and $v(\mathbf{x}_0) = u(\mathbf{x}_0)$.

Via a comparison principle applied to the solution u and the super-solution v , we can finally estimate the normal derivative of u at \mathbf{x}_0 by the sup-norm of the derivative of the barrier function v (which in turn is bounded in terms of the data) and arrive at the assertion (4.5).

Now we start with the rigorous derivation of the estimates for the normal derivative. Since Ω is by assumption of class \mathcal{C}^1 and satisfies an exterior ball condition, we find positive constants r_0 , L , L_d and N depending only on Ω such that we can suppose that an arbitrary boundary point $\mathbf{x}_0 \in \partial\Omega$ is given, after an orthogonal transformation, by $\mathbf{x}_0 = (\mathbf{0}, -r_0)$ (we use the notation $\mathbf{x} = (\mathbf{x}', x_d)$) and that we have the inclusions

$$\begin{aligned} \Gamma &:= \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}'| < L, f(\mathbf{x}') = x_d\} \\ &\subset \partial\Omega, \end{aligned}$$


$$\begin{aligned} \Omega_+ &:= \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}'| < L, f(\mathbf{x}') - L_d < x_d < f(\mathbf{x}')\} \\ &\subset \Omega, \end{aligned}$$

$$\begin{aligned} \Omega_- &:= \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}'| < L, f(\mathbf{x}') < x_d < f(\mathbf{x}') + L_d\} \\ &\subset \mathbb{R}^d \setminus \Omega, \end{aligned}$$

with a function $f \in \mathcal{C}^1(-L, L)^{d-1}$ fulfilling $\|f\|_{1,\infty} \leq N$, $f(\mathbf{0}') = -r_0$ and $D_i f(\mathbf{0}') = 0$ for all $i = 1, \dots, d-1$.

In addition, we may suppose that r_0 is so small that $B_{r_0}(\mathbf{0}) \subset \Omega_-$ holds and that

$$M^*(|\mathbf{x}| - r_0) \geq |\mathbf{x} - \mathbf{x}_0|^2 \tag{4.22}$$



is satisfied for all $\mathbf{x} \in \Gamma$, for some constant M^* depending only on Ω and r_0 (this can for example be seen easily if also $B_{2r_0}(\mathbf{0}', r_0) \subset \Omega_-$ holds).

In this setting, for an arbitrary $\delta \in (0, 1)$ (to be specified later on) we can work with the functions $b_{r_0}^\delta$ and $\omega_{r_0}^\delta$ introduced in (4.7) and (4.8), and with the function $v_{\mathbf{k},c}^{\delta,r_0}$ from (4.13) for the specific choices $\mathbf{k} := \nabla u_0(\mathbf{x}_0)$ and $c = u_0(\mathbf{x}_0) - \nabla u_0(\mathbf{x}_0) \cdot \mathbf{x}_0$, that is, with

$$v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) = \omega_{r_0}^\delta(\mathbf{x}) + \nabla u_0(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) + u_0(\mathbf{x}_0). \quad (4.23)$$

Note that $v_{\mathbf{k},c}^{\delta,r_0}$ is well defined outside the ball $B_{r_0}(\mathbf{0})$ and so it is well-defined also in Ω_+ . In addition, it is clear that $|\mathbf{k}| \leq \|\nabla u_0\|_\infty$ holds, hence, we can choose $K := \|\nabla u_0\|_\infty$ and fix the number M (depending only on F and this K) according to Lemma 4.2. Furthermore, since $(F'_g)^{-1}$ maps $[0, 1]$ to $[0, \infty)$ and is monotonically increasing, we can fix a number $\delta_{\max} \in (0, 1/2)$ such that

$$(F'_g)^{-1}(s) \geq \max\{M, M^*\|u_0\|_{1,\infty}\} \quad \text{for all } s \in [(1 - 2\delta_{\max})^{d-1}, 1]. \quad (4.24)$$

From now on, we will consider arbitrary $\delta \in (0, \delta_{\max})$. Then, from (4.7) and (4.24) it follows that

$$r_0 < |\mathbf{x}| \leq \frac{(1 - \delta_{\max})r_0}{1 - 2\delta_{\max}} =: r_{\max} \implies b_{r_0}^\delta(|\mathbf{x}|) \geq M. \quad (4.25)$$

Consequently, using Lemma 4.2, we see that $v_{\mathbf{k},c}^{\delta,r_0}$ is a super-solution to (4.1) in the set $(B_{r_{\max}} \setminus B_{r_0}) \cap \Omega_+$, which is the first crucial property of an upper barrier.

Next, we want to identify a part of Γ on which $v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) \geq u(\mathbf{x}) = u_0(\mathbf{x})$ holds, that is, where

$$\omega_{r_0}^\delta(\mathbf{x}) + \nabla u_0(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) + u_0(\mathbf{x}_0) - u_0(\mathbf{x}) \geq 0. \quad (4.26)$$

From Taylor expansion of u_0 and the $\mathcal{C}^{1,1}$ -regularity assumption on u_0 we know that

$$|u_0(\mathbf{x}) - u_0(\mathbf{x}_0) - \nabla u_0(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0)| \leq \|u_0\|_{1,\infty} |\mathbf{x} - \mathbf{x}_0|^2,$$

so to verify (4.26) it is enough to check where

$$\omega_{r_0}^\delta(\mathbf{x}) - \|u_0\|_{1,\infty} |\mathbf{x} - \mathbf{x}_0|^2 \geq 0 \quad (4.27)$$

holds. Using the definitions of $b_{r_0}^\delta$ in (4.7) and of $\omega_{r_0}^\delta$ in (4.8), combined with the fact that $(F'_g)^{-1}$ is monotonically increasing, we have for all $\mathbf{x} \in \Gamma$

$$\omega_{r_0}^\delta(\mathbf{x}) \geq (|\mathbf{x}| - r_0)(F'_g)^{-1}\left(\frac{(1 - \delta)^{d-1}r_0^{d-1}}{|\mathbf{x}|^{d-1}}\right).$$

Consequently, in order to guarantee (4.27) and thus (4.26) it is sufficient, in view of (4.22), to have


$$(F'_g)^{-1}\left(\frac{(1 - \delta)^{d-1}r_0^{d-1}}{|\mathbf{x}|^{d-1}}\right) \geq M^*\|u_0\|_{1,\infty},$$

which is indeed true for all \mathbf{x} with $r_0 \leq |\mathbf{x}| \leq r_{\max}$, by the choices of the parameter δ_{\max} in (4.24) and of the radius r_{\max} in (4.25). Thus, we have verified

$$u(\mathbf{x}) \leq v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) \quad \text{for all } \mathbf{x} \in \Gamma \text{ with } r_0 \leq |\mathbf{x}| \leq r_{\max}. \quad (4.28)$$

Finally, in order to complete the second property of the barrier function, namely that it lies above the solution on all of the boundary of a relative neighborhood of \mathbf{x}_0 , we still need to take care of the values of $v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})$

inside of Ω (but close to \mathbf{x}_0). This shall now be accomplished by suitable choices of a local neighborhood and of $\delta \in (0, \delta_{\max})$.



First, since r_0 and r_{\max} are already fixed (in dependence on Ω , F and u_0), we can zoom in the neighborhood of \mathbf{x}_0 and find two constants L^* and $L_d^* \leq L_d$ sufficiently small (depending again only on the data) such that

$$\begin{aligned} \Gamma^* &:= \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}'| < L^*, f(\mathbf{x}') = x_d\} \\ &\subset \partial\Omega, \end{aligned}$$

$$\begin{aligned} \Omega_+^* &:= \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}'| < L^*, f(\mathbf{x}') - L_d^* < x_d < f(\mathbf{x}')\} \\ &\subset \Omega \cap (B_{r_{\max}} \setminus B_{r_0}), \end{aligned}$$

$$\begin{aligned} \Omega_-^* &:= \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}'| < L^*, f(\mathbf{x}') < x_d < f(\mathbf{x}') + L_d\} \\ &\subset \mathbb{R}^d \setminus \Omega. \end{aligned}$$

By these choices, due to (4.25) and (4.28), $v_{\mathbf{k},c}^{\delta,r_0}$ is a super-solution to (4.1) in the relative neighborhood Ω_+^* of \mathbf{x}_0 and satisfies $v_{\mathbf{k},c}^{\delta,r_0} \geq u$ on Γ^* , for all $\delta < \delta_{\max}$. Our goal is to show $v_{\mathbf{k},c}^{\delta,r_0} \geq u$ on the remaining part of $\partial\Omega_+^*$, and this is indeed the point, where we shall use the assumption (1.9).

In view of the choices of r_0 and Ω_+^* there exists $\eta > 0$ independent of δ such that for all $\mathbf{x} \in \partial\Omega_+^* \setminus \Gamma^*$ there holds

$$|\mathbf{x}| \geq r_0 + \eta.$$

Hence, from the definition of $v_{\mathbf{k},c}^{\delta,r_0}$ in (4.23) and of $\omega_{r_0}^\delta$ in (4.8) we find

$$v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) \geq \int_{r_0}^{r_0+\eta} b_{r_0}^\delta(r) \, dr - C^*(\Omega) \|u_0\|_{1,\infty}. \quad (4.29)$$

On the other hand, we know $\|u\|_\infty \leq \|u_0\|_\infty$ from (4.4). Therefore, in order to show that $v_{\mathbf{k},c}^{\delta,r_0} \geq u$ on $\partial\Omega_+^*$, it is enough to verify that we can choose $\delta \in (0, \delta_{\max})$ in such way that

$$\int_{r_0}^{r_0+\eta} b_{r_0}^\delta(r) \, dr \geq C^*(\Omega) \|u_0\|_{1,\infty} + \|u_0\|_\infty. \quad (4.30)$$

Using the definition of $b_{r_0}^\delta$ in (4.7) and the substitution formula, we deduce that

$$\begin{aligned} \int_{r_0}^{r_0+\eta} b_{r_0}^\delta(r) \, dr &= \int_{r_0}^{r_0+\eta} (F_g')^{-1} \left(\frac{(1-\delta)^{d-1} r_0^{d-1}}{r^{d-1}} \right) \, dr \\ &= \frac{(1-\delta)r_0}{d-1} \int_{\frac{(1-\delta)^{d-1} r_0^{d-1}}{(r_0+\eta)^{d-1}}}^{(1-\delta)^{d-1}} (F_g')^{-1}(s) s^{-\frac{d}{d-1}} \, ds. \end{aligned}$$

If we now introduce

$$\alpha := \min \left\{ \frac{r_0}{d-1}, 1 - \frac{r_0^{d-1}}{(r_0+\eta)^{d-1}} \right\}$$

(depending only on Ω , d , F and \mathbf{u}_0 , as r_0 and η are already fixed), the above integral can be estimated by

$$\int_{r_0}^{r_0+\eta} b_{r_0}^\delta(r) \, dr \geq \alpha \int_{1-\alpha}^{(1-\delta)^{d-1}} (F_g')^{-1}(s) \, ds = \alpha \int_{(F_g')^{-1}(1-\alpha)}^{(F_g')^{-1}((1-\delta)^{d-1})} t F_g''(t) \, dt$$

(note that the integral on the right-hand side is negative whenever $1 - \alpha > (1 - \delta)^{d-1}$ holds, hence, the previous inequality is trivially satisfied in this case). Thanks to Lemma 3.2 (more precisely, by (1.10) and (1.9) for F_g), we know first that $(F'_g(s))^{-1} \rightarrow \infty$ as $s \rightarrow 1$ and secondly that

$$\int_1^\infty t F_g''(t) dt = \infty.$$

Therefore, we can fix $\delta \in (0, \delta_{\max})$ (depending only on Ω , d , F and u_0) such that

$$\int_{r_0}^{r_0+\eta} b_{r_0}^\delta(r) dr \geq \alpha \int_{(F'_g)^{-1}(1-\alpha)}^{(F'_g)^{-1}((1-\delta)^{d-1})} t F_g''(t) dt \geq C^*(\Omega) \|u_0\|_{1,\infty} + \|u_0\|_\infty$$

holds, which yields the desired inequality (4.30). Therefore, we have proved

$$v_{\mathbf{k},c}^{\delta,r_0} \geq u \quad \text{on } \partial\Omega_+^* \quad (4.31)$$

and have finished the construction of the barrier function on the relative neighborhood Ω_+^* of \mathbf{x}_0 .

It now remains to establish the bound (4.5) for the normal derivative of u , locally at \mathbf{x}_0 , using the properties of the barrier function $v_{\mathbf{k},c}^{\delta,r_0}$. For this purpose, we recall that $v_{\mathbf{k},c}^{\delta,r_0}$ is a super-solution to (4.1) and that u is a solution to (4.1) in Ω_+^* . Thus, in view of (4.31) we obtain from the comparison principle for elliptic equations in divergence form

$$v_{\mathbf{k},c}^{\delta,r_0} \geq u \quad \text{in } \Omega_+^*.$$

This now allows to estimate the normal derivative. Indeed, for any $0 < h \leq L_d^*$ we have $\mathbf{x} := (\mathbf{0}', -r_0 - h) \in \Omega_+^*$, and therefore, taking into account also $v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}_0) = u(\mathbf{x}_0)$, we find the following estimate

$$\begin{aligned} \frac{u(\mathbf{x}_0) - u(\mathbf{x})}{h} &= \frac{v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}_0) - v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})}{h} + \frac{v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}) - u(\mathbf{x})}{h} \\ &\geq \frac{v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x}_0) - v_{\mathbf{k},c}^{\delta,r_0}(\mathbf{x})}{h} \geq -\|v_{\mathbf{k},c}^{\delta,r_0}\|_{1,\infty} \geq -C(\Omega, F, u_0). \end{aligned}$$

Thus, recalling that the outer unit normal to $\partial\Omega$ in \mathbf{x}_0 is given by e_n , we obtain in the passage $h \rightarrow 0_+$ the lower bound

$$\frac{\partial u(\mathbf{x}_0)}{\partial \mathbf{n}} \geq -C(\Omega, F, u_0).$$

Repeating the whole procedure with $\omega_{r_0}^\delta$ replaced by $-\omega_{r_0}^\delta$, we get the opposite inequality

$$\frac{\partial u(\mathbf{x}_0)}{\partial \mathbf{n}} \leq C(\Omega, F, u_0).$$

The latter two inequalities imply (4.5), which in turn, due to the reasoning in Section 4.1, provides

$$\|\nabla u\|_\infty \leq C(\Omega, F, u_0)$$

(which, as the index ε was dropped, is precisely the uniform bound (4.2)). This finishes the proof of the theorem. \square

REFERENCES

- [1] L. Beck and T. Schmidt, On the Dirichlet problem for variational integrals in *BV*. *J. Reine Angew. Math.* **674** (2013) 113–194.
- [2] L. Beck and T. Schmidt, Interior gradient regularity for BV-minimizers of singular variational problems. *Nonl. Anal.* **120** (2015) 86–106.

- [3] S. Bernstein, Sur les équations du calcul des variations. *Ann. Sci. École Norm. Sup.* **29** (1912) 431–485.
- [4] M. Bildhauer, A priori gradient estimates for bounded generalized solutions of a class of variational problems with linear growth. *J. Convex Anal.* **9** (2002) 117–137.
- [5] M. Bildhauer, Convex variational problems. Linear, nearly linear and anisotropic growth conditions. Vol. 1818 of *Lect. Notes Math.* Berlin, Springer (2003).
- [6] M. Bildhauer, Two dimensional variational problems with linear growth. *Manuscripta Math.* **110** (2003) 325–342.
- [7] M. Bildhauer and M. Fuchs, On a class of variational integrals with linear growth satisfying the condition of μ -ellipticity. *Rend. Mat. Appl., VII. Ser.* **22** (2002) 249–274.
- [8] M. Bulíček, J. Málek, K.R. Rajagopal and J.R. Walton, Existence of solutions for the anti-plane stress for a new class of “strain-limiting” elastic bodies. *Calc. Var. Partial Differ. Equ.* **54** (2015) 2115–2147.
- [9] J. Dalphin, Some characterizations of a uniform ball property. Congrès SMAI 2013. *ESAIM: PROCs.* **45** (2014) 437–446.
- [10] R. Finn, Remarks relevant to minimal surfaces, and to surfaces of prescribed mean curvature. *J. Anal. Math.* **14** (1965) 139–160.
- [11] M. Fuchs and G. Mingione, Full $C^{1,\alpha}$ -regularity for free and constrained local minimizers of elliptic variational integrals with nearly linear growth. *Manuscripta Math.* **102** (2000) 227–250.
- [12] M. Giaquinta and E. Giusti, Global $C^{1,\alpha}$ -regularity for second order quasilinear elliptic equations in divergence form. *J. Reine Angew. Math.* **351** (1984) 55–65.
- [13] M. Giaquinta, G. Modica and J. Souček, Functionals with linear growth in the calculus of variations I, II. *Comment. Math. Univ. Carol.* **20** (1979) 143–156, 157–172.
- [14] E. Giusti, Minimal surfaces and functions of bounded variation. Birkhäuser, Basel (1984).
- [15] H. Lebesgue, *Intégrale, longueur, aire*. Thèse (1902).
- [16] J. Leray, Discussion d’un problème de Dirichlet. *J. Math. Pures Appl.* **18** (1939) 249–284.
- [17] P. Marcellini and G. Papi, Nonlinear elliptic systems with general growth. *J. Differ. Equ.* **221** (2006) 412–443.
- [18] G. Mingione and F. Siepe, Full $C^{1,\alpha}$ -regularity for minimizers of integral functionals with $L \log L$ -growth. *Z. Anal. Anwend.* **18** (1999) 1083–1100.
- [19] M. Miranda, Un principio di massimo forte per le frontiere minimali e una sua applicazione alla risoluzione del problema al contorno per l’equazione delle superfici di area minima. *Rend. Sem. Mat. Univ. Padova* **45** (1971) 355–366.
- [20] Y.G. Reshetnyak, Weak convergence of completely additive vector functions on a set. *Sib. Math. J.* **9** (1968) 1039–1045.
- [21] J. Serrin, On the definition and properties of certain variational integrals. *Trans. Am. Math. Soc.* **101** (1961) 139–167.
- [22] J. Serrin, The problem of Dirichlet for quasilinear elliptic differential equations with many independent variables. *Philos. Trans. R. Soc. Lond. Ser. A* **264** (1969) 413–496.