

Invariance entropy, quasi-stationary measures and control sets

Fritz Colonius

Angaben zur Veröffentlichung / Publication details:

Colonius, Fritz. 2018. "Invariance entropy, quasi-stationary measures and control sets."
Discrete and Continuous Dynamical Systems - A 38 (4): 2093–123.
<https://doi.org/10.3934/dcds.2018086>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



INVARIANCE ENTROPY, QUASI-STATIONARY MEASURES AND CONTROL SETS

FRITZ COLONIUS

Institut für Mathematik
Universität Augsburg
Universitätsstrasse 14
86159 Augsburg, Germany

ABSTRACT. For control systems in discrete time, this paper discusses measure-theoretic invariance entropy for a subset Q of the state space with respect to a quasi-stationary measure obtained by endowing the control range with a probability measure. The main results show that this entropy is invariant under measurable transformations and that it is already determined by certain subsets of Q which are characterized by controllability properties.

1. Introduction. Metric invariance entropy provides a measure-theoretic analogue of the topological notion of (feedback) invariance entropy $h_{\text{inv}}(Q)$ of deterministic control systems, cf. Nair, Evans, Mareels and Moran [19] and Kawan [17]. The present paper discusses metric invariance entropy and its relations to controllability properties. We consider control systems in discrete time of the form

$$x_{k+1} = f(x_k, u_k), k \in \mathbb{N} = \{0, 1, \dots\}, \quad (1)$$

where $f : M \times \Omega \rightarrow M$ is continuous and M and Ω are metric spaces.

For an initial value $x_0 \in M$ at time $k = 0$ and control $u = (u_k)_{k \geq 0} \in \mathcal{U} := \Omega^{\mathbb{N}}$ we denote the solutions by $x_k = \varphi(k, x_0, u)$, $k \in \mathbb{N}$. The notion of invariance entropy $h_{\text{inv}}(Q)$ describes the average data rate needed to keep the system in a given subset Q of M (forward in time). It is constructed with some analogy to topological entropy of dynamical systems. A major difference of entropy in a control context to entropy for dynamical systems (cf. Walters [25] or Viana and Oliveira [24]) is that the minimal required entropy for the considered control task is of interest instead of the “total” entropy generated by the dynamical system, hence the infimum over open covers or partitions is taken instead of the supremum.

The present paper discusses notions of metric invariance entropy modifying and extending the analysis in Colonius [6, 7]. A probability measure on the space Ω of control values is fixed. Then an associated quasi-stationary measure η for Q is considered and an entropy notion is constructed that takes into account information on feedbacks. A significant relaxation compared to topological invariance entropy is that only invariance with η -probability one is required (this was not done in [6, 7]).

2010 *Mathematics Subject Classification.* Primary: 93C41, 37A35; Secondary: 94A17.

Key words and phrases. Invariance entropy, quasi-stationary measures, control sets, coder-controllers, transitivity set.

Research supported by DFG grant 124/19-2.

The main results are Theorem 2.6 showing that the invariance entropy does not decrease under semi-conjugacy, Theorem 2.8 showing that the topological invariance entropy is an upper bound for the metric invariance entropy and Theorems 5.4 and 5.5 providing conditions under which the metric invariance entropy is already determined by certain maximal subsets of approximate controllability within the interior of Q (i.e., invariant W -control sets with $W := \text{int}Q$).

A general reference to quasi-stationary measures is the monograph Collett, Martinez and San Martin [5]; the survey Méléard and Villemonais [18] presents, in particular, applications to population dynamics where quasi-stationary distributions correspond to plateaus of mortality rates. Intuitively speaking, quasi-stationarity measures may exist when exit from Q occurs with probability one for time tending to infinity, while in finite time a quasi-stationary behavior develops. A bibliography for quasi-stationary measures with more than 400 entries is due to Pollett [21].

For controllability properties in discrete time, the results by Jakubczyk and Sontag [14] on reachability are fundamental. Related results, in particular on control sets, are due to Albertini and Sontag [2, 1], Sontag and Wirth [23] and Wirth [27], Patrão and San Martin [20] and Colonius, Homburg and Kliemann [8]. The case of W -control sets has only been discussed in the continuous-time case, cf. Colonius and Lettau [9]. Although many properties of control sets and W -control sets in discrete time are analogous to those in continuous time, some additional difficulties occur. In particular, in the proofs one has to replace the interior of a control set by its transitivity set or its closely related core.

System (1) together with the measure ν on Ω generates a random dynamical system. Metric and topological entropy of such systems have been intensely studied, see, e.g., Bogenschütz [4]. If the control range Ω in (1) is a finite set, say $\Omega = \{1, \dots, p\}$, then the associated right hand sides $g_i = f(\cdot, i)$, $i \in \Omega$, generate a semigroup of continuous maps acting on M . The metric and topological entropy theory of finitely generated semigroups acting on compact metric spaces has recently found interest, see, e.g., Rodrigues and Varandas [22].

The contents of this paper is as follows: In Section 2 definitions of metric invariance entropy are presented and discussed. In particular, the behavior under measurable transformations is characterized and it is shown that the topological invariance entropy is an upper bound for metric invariance entropy. Section 3 relates the metric invariance entropy to properties of coder-controllers rendering Q invariant. Section 4 presents conditions ensuring that this entropy is already determined on a subset K which is invariant in Q , i.e., a set which cannot be left by the system without leaving Q . In Section 5, invariant W -control sets are introduced and their properties are analyzed. The union of their closures yields a set K satisfying the conditions derived earlier guaranteeing that the metric invariance entropy of Q coincides with the metric invariance entropy of K . Examples 2.9 and 5.7 illustrate some of the concepts in simple situations.

Notation. Given a probability measure μ we say that a property holds for μ -a.a. (almost all) points if it is valid outside a set of μ -measure zero.

2. Definition of metric invariance entropy. In this section, we present definitions of metric invariance entropy and discuss their motivation. First we recall entropy of dynamical systems which also serves to introduce some notation.

Let μ be a probability measure on a space X endowed with a σ -algebra \mathfrak{F} . For every finite partition $\mathcal{P} = \{P_1, \dots, P_n\}$ of X into measurable sets the entropy is

defined as $H_\mu(\mathcal{P}) = -\sum_i \phi(\mu(P_i))$, where $\phi(x) = x \log x, x \in (0, 1]$ with $\phi(0) = 0$. The entropy specifies the expected information gained from the outcomes in \mathcal{P} of an experiment, or the amount of uncertainty removed upon learning the \mathcal{P} -address of a randomly chosen point. For a dynamical system generated by a continuous map T on a compact metric space X one considers an invariant measure on the Borel σ -algebra $\mathcal{B}(X)$, i.e., $\mu(T^{-1}E) = \mu(E)$ for all $E \in \mathcal{B}(X)$. For a finite partition \mathcal{P} of X and $j \in \mathbb{N}$ one finds with $T^{-j}\mathcal{P} := \{T^{-j}P | P \in \mathcal{P}\}$ that

$$\mathcal{P}_n := \bigvee_{j=0}^{n-1} T^{-j}\mathcal{P} = \mathcal{P} \vee T^{-1}\mathcal{P} \vee \dots \vee T^{-(n-1)}\mathcal{P}$$

again is a finite partition of X (for two collections \mathfrak{A} and \mathfrak{B} of sets the join is $\mathfrak{A} \vee \mathfrak{B} = \{A \cap B | A \in \mathfrak{A} \text{ and } B \in \mathfrak{B}\}$). The entropy of T with respect to the partition \mathcal{P} is $h_\mu(T, \mathcal{P}) := \lim_{n \rightarrow \infty} \frac{1}{n} H_\mu(\mathcal{P}_n)$. Using conditional entropy, one can also write

$$H_\mu(\mathcal{P}_n) = \sum_{i=0}^{n-1} H_\mu(\mathcal{P}_{i+1} | \mathcal{P}_i). \quad (2)$$

The metric entropy of T is $h_\mu(T) := \sup_{\mathcal{P}} h_\mu(T, \mathcal{P})$, where the supremum is taken over all finite partitions \mathcal{P} of X , i.e., it is the total information generated by the dynamical system generated by T .

This concept has to be modified when we want to determine the minimal information that is needed to make a subset Q of the state space of a control system (1) invariant under feedbacks. We suppose that a closed set $Q \subset M$ is given and fix a probability measure ν on the Borel σ -algebra $\mathcal{B}(\Omega)$ of the control range Ω . Let $p(x, A) = \nu\{\omega \in \Omega | f(x, \omega) \in A\}$, $x \in M$, $A \subset M$, be the associated Markov transition probabilities. A quasi-stationary measure with respect to Q of M is a probability measure η on $\mathcal{B}(M)$ such that for some $\rho \in (0, 1]$

$$\rho \eta(A) = \int_Q p(x, A) \eta(dx) \text{ for all } A \in \mathcal{B}(Q). \quad (3)$$

The measure η is stationary if and only if $\rho = 1$. With $A = Q$ one obtains $\rho = \int_Q p(x, Q) \eta(dx)$ and the support $\text{supp} \eta$ is contained in Q . Results on the existence of quasi-stationary measures are given, e.g., in Collett, Martinez and San Martin [5, Proposition 2.10 and Theorem 2.11] and Colonius [6, Theorem 2.9].

With the shift $\theta : \mathcal{U} \rightarrow \mathcal{U}, (u_k)_{k \geq 0} \mapsto (u_{k+1})_{k \geq 0}$, control system (1) can equivalently be described by the continuous skew product map

$$S : \mathcal{U} \times M \rightarrow \mathcal{U} \times M, (u, x) \mapsto (\theta u, f(x, u_0)), \quad (4)$$

where $\mathcal{U} = \Omega^{\mathbb{N}}$ is endowed with the product topology. Then $S^k(u, x) = (\theta^k u, \varphi(k, x, u))$. A conditionally invariant measure μ for the map S with respect to $Q \subset M$ is a probability measure on the Borel σ -algebra of $\mathcal{U} \times M$ such that $0 < \rho := \mu(S^{-1}(\mathcal{U} \times Q) \cap (\mathcal{U} \times Q)) \leq 1$ and

$$\rho \mu(B) = \mu(S^{-1}B \cap (\mathcal{U} \times Q)) \text{ for all } B \in \mathcal{B}(\mathcal{U} \times M). \quad (5)$$

We write $S_Q := S|_{\mathcal{U} \times Q} : \mathcal{U} \times Q \rightarrow \mathcal{U} \times M$ for the restriction. Then the condition in (5) can be written as $\rho \mu(B) = \mu(S_Q^{-1}B)$. For $k \in \mathbb{N}$ the measure μ is conditionally invariant for S_Q^k with constant ρ^k and, in particular, $\rho^{-k} \mu$ is a probability measure on $S_Q^{-k}(\mathcal{U} \times Q)$.

If η is a quasi-stationary measure, one finds that with the product measure $\nu^{\mathbb{N}}$ on $\mathcal{U} = \Omega^{\mathbb{N}}$ the measure $\mu = \nu^{\mathbb{N}} \times \eta$ is a probability measure on the product space

$\mathcal{U} \times M$ satisfying (5), cf. [6, Proposition 2.8]. In the present paper only conditionally invariant measures of this form are considered and ν will be fixed (cf., e.g., Demers and Young [12], Demers [11] for results on general conditionally invariant measures). Often we will suppress the dependence on ν and only indicate the dependence on the quasi-stationary measure η .

Next we construct certain partitions for subsets of $\mathcal{U} \times Q$ whose entropy with respect to $\mu = \nu^{\mathbb{N}} \times \eta$ will be used to define metric invariance entropy.

Definition 2.1. For a closed subset $Q \subset M$ an invariant (Q, η) -partition $\mathcal{C}_\tau = \mathcal{C}_\tau(\mathcal{P}, F)$ is given by $\tau \in \mathbb{N}$, a finite partition \mathcal{P} of Q into Borel measurable sets and a map $F : \mathcal{P} \rightarrow \Omega^\tau$ assigning to each set P in \mathcal{P} a control function such that

$$\varphi(k, x, F(P)) \in Q \text{ for } k \in \{1, \dots, \tau\} \text{ and } \eta\text{-a.a. } x \in P. \quad (6)$$

When no misunderstanding can occur, we just talk about invariant Q -partitions or just invariant partitions. Clearly, condition (6) means that

$$\eta\{x \in P \mid \varphi(k, x, F(P)) \in Q \text{ for } k = 1, \dots, \tau\} = \eta(P).$$

Fix an invariant (Q, η) -partition $\mathcal{C}_\tau = \mathcal{C}_\tau(\mathcal{P}, F)$ with $\mathcal{P} = \{P_1, \dots, P_q\}$. Abbreviate $F_i := F(P_i) \in \Omega^\tau$, $i = 1, \dots, q$, and define for every word $a := [a_0, a_1, \dots, a_{n-1}]$, $n \in \mathbb{N}$, with $a_j \in \{1, \dots, q\}$ a control function u_a on $\{0, \dots, n\tau - 1\}$ by applying these feedback maps one after the other: for $i = 0, \dots, n-1$ and $k = 0, \dots, \tau-1$

$$(u_a)_{i\tau+k} := (F_{a_i})_k. \quad (7)$$

We also write $u_a := (F_{a_0}, F_{a_1}, \dots, F_{a_{n-1}})$. A word a is called (η, \mathcal{C}_τ) -admissible if

$$\eta\{x \in Q \mid \varphi(i\tau, x, u_a) \in P_{a_i} \text{ for } i = 0, 1, \dots, n-1\} > 0. \quad (8)$$

Note that for η -a.a. x it follows that $\varphi(k, x, u_a) \in Q$ for $k = 0, \dots, n\tau$ if $\varphi(i\tau, x, u_a) \in P_{a_i}$ for $i = 0, \dots, n-1$. If η and \mathcal{C}_τ are clear from the context, we just say that a is admissible. The admissible words describe the sequences of partition elements under the feedbacks associated with \mathcal{C}_τ which are followed with positive probability.

For $P \in \mathcal{P}$ we define

$$A(P, \eta) = \{u \in \mathcal{U} \mid \varphi(k, x, u) \in Q \text{ for } k = 1, \dots, \tau \text{ and } \eta\text{-a.a. } x \in P\} \times P \quad (9)$$

and

$$\mathfrak{A}(\mathcal{C}_\tau, \eta) = \{A(P, \eta) \mid P \in \mathcal{P}\} \text{ with union } \mathcal{A}(\mathcal{C}_\tau, \eta) = \bigcup_{P \in \mathcal{P}} A(P, \eta).$$

Here and in the following the dependence on \mathcal{C}_τ (actually, these sets only depend on (\mathcal{P}, τ)) or η is omitted, if it is clear from the context. The controls u in (9) can be considered as constant parts of feedbacks keeping η -almost all $x \in P$ in Q up to time τ .

Lemma 2.2. *The sets $A(P, \eta)$ defined in (9) are Borel measurable, hence \mathfrak{A} is a measurable partition of \mathcal{A} which, in general, is a proper subset of $\mathcal{U} \times Q$.*

Proof. Clearly the sets $A(P, \eta)$ are pairwise disjoint, hence it only remains to show measurability. We only prove this for the case $\tau = 1$, where it suffices to show that

$$\{u \in \Omega \mid f(x, u) \in Q \text{ for } \eta\text{-a.a. } x \in P\} = \{u \in \Omega \mid \eta(f(\cdot, u)^{-1}Q \cap P) \geq \eta(P)\}$$

is measurable. First we claim that for compact $K \subset P$ and $\delta > 0$ the set

$$\{u \in \Omega \mid \eta(f(\cdot, u)^{-1}Q \cap K) \geq \eta(P) - \delta\} \quad (10)$$

is closed. In fact, if this set is nonvoid, let $u_n \in \Omega, u_n \rightarrow u \in \Omega$ with

$$\eta(f(\cdot, u_n)^{-1}Q \cap K) \geq \eta(P) - \delta.$$

The sets B_m defined by

$$B_m := \bigcup_{i=m}^{\infty} f(\cdot, u_i)^{-1}Q \cap K, B := \bigcap_{m=1}^{\infty} B_m,$$

are decreasing and $\eta(B) = \lim_{m \rightarrow \infty} \eta(B_m) \geq \eta(P) - \delta$. Furthermore, suppose that a subsequence of $y_i \in f(\cdot, u_i)^{-1}Q \cap K, i \in \mathbb{N}$, converges to y . Since $f^{-1}(Q, \Omega)$ is closed, it follows that $y \in f(\cdot, u)^{-1}Q \cap K$. This shows that $B \subset f(\cdot, u)^{-1}Q \cap K$ and hence closedness of the set in (10) follows from

$$\eta(f(\cdot, u)^{-1}Q \cap K) \geq \eta(B) \geq \eta(P) - \delta.$$

Since Ω is a compact metric space, regularity of the probability measure η implies that there are compact sets $K_n \subset P$ with $\eta(P \setminus K_n) \leq \frac{1}{n}$ (cf. Viana and Oliveira [24, Proposition A.3.2]). Hence

$$\{u \in \Omega \mid \eta(f(\cdot, u)^{-1}Q \cap P) \geq \eta(P)\} \subset \{u \in \Omega \mid \eta(f(\cdot, u)^{-1}Q \cap K_n) \geq \eta(P) - 1/n\},$$

and it follows that

$$\{u \in \Omega \mid \eta(f(\cdot, u)^{-1}Q \cap P) \geq \eta(P)\} = \bigcap_{n=1}^{\infty} \left\{u \in \Omega \mid \eta(f(\cdot, u)^{-1}Q \cap K_n) \geq \eta(P) - \frac{1}{n}\right\}.$$

Thus the set on the left hand side is measurable as countable intersection of closed sets. \square

A sequence (A_0, \dots, A_{n-1}) of sets in \mathfrak{A} is called \mathcal{C}_τ -admissible (or a \mathcal{C}_τ -itinerary), if there is an admissible word $a = [a_0, \dots, a_{n-1}]$ of length n with $A_i = A(P_{a_i}) \in \mathfrak{A}$ for all i . Then also the set

$$D_a = A_0 \cap S^{-\tau}A_1 \cap \dots \cap S^{-(n-1)\tau}A_{n-1} \in \bigvee_{i=0}^{n-1} S_Q^{-i\tau} \mathfrak{A} \quad (11)$$

is called admissible. Only the sets D_a with $\mu(D_a) > 0$ will be relevant (as usual, if $\mu(D_a) = 0$, this set is simply omitted in the following).

Note that μ -a.a. $(u, x) \in D_a$ satisfy $\varphi(k, x, u) \in Q$ for $k = 0, \dots, n\tau$. The collection of all sets D_a is

$$\mathfrak{A}_n := \left\{ D_a \in \bigvee_{i=0}^{n-1} S_Q^{-i\tau} \mathfrak{A} \mid a \text{ admissible} \right\}, \mathcal{A}_n := \bigcup_{a \text{ admissible}} D_a \subset S_Q^{-(n-1)\tau}(\mathcal{U} \times Q). \quad (12)$$

Observe that $\mathfrak{A}_1 = \mathfrak{A}$ and that \mathfrak{A}_n is a measurable partition of \mathcal{A}_n and, for convenience, we set $\mathfrak{A}_0 = \mathcal{U} \times Q$. Note that the inclusion $\mathcal{A}_{n+1} \subset \mathcal{A}_n$ holds for all $n \in \mathbb{N}$, and that, in general, it is proper.

Remark 1. For invariant (Q, η) -partitions the inclusion

$$\mathfrak{A}_{n+m} \subset \mathfrak{A}_n \vee S^{-n\tau} \mathfrak{A}_m, n, m \in \mathbb{N}, \quad (13)$$

does not hold, in general (in contrast to Colonius [6, Lemma 3.3] where only existence of a trajectory following a sequence of partition elements is required). The

problem is that for an (η, \mathcal{C}_τ) -admissible word $a = [a_0, \dots, a_{n-1}, a_n, \dots, a_{n+m-1}]$ the word $[a_n, \dots, a_{n+m-1}]$ need not be (η, \mathcal{C}_τ) -admissible: Certainly, the inequality

$$\eta \{y \in Q \mid \varphi(i\tau, y, u_{[a_n, \dots, a_{n+m-1}]}) \in P_{a_{n+i}} \text{ for } i = 0, \dots, m-1\} \\ \geq \eta \{\varphi(n\tau, x, u_a) \in Q \mid x \in Q, \varphi(i\tau, x, u_a) \in P_{a_i} \text{ for } i = 0, \dots, n+m-1\}$$

holds, but the term on the right hand side need not be positive. This could be guaranteed by changing the definition of (η, \mathcal{C}_τ) -admissible words $a = [a_0, \dots, a_{n-1}]$ to: For all $j = 0, 1, \dots, n-1$

$$\eta \{y \in Q \mid \varphi(i\tau, y, u_a) \in P_{a_i} \text{ for } i = j, j+1, \dots, n-1\} > 0.$$

We do not adapt this definition since the inclusion (13) is not needed below.

The direct way to define a notion of metric invariance entropy is to consider the entropy of the partitions \mathfrak{A}_n of the sets \mathcal{A}_n . An alternative is to consider the additional information in every step. We start with the first choice.

We consider the entropy $H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau))$ of $\mathfrak{A}_n(\mathcal{C}_\tau)$ in $\mathcal{A}_n(\mathcal{C}_\tau) \subset S_Q^{-(n-1)\tau}(\mathcal{U} \times Q)$ with respect to the probability measure $\rho^{-(n-1)\tau}\mu$ and then take the average of the required information as time tends to ∞ to get the invariance μ -entropy of \mathcal{C}_τ ,

$$h_\mu(\mathcal{C}_\tau, Q) := \limsup_{n \rightarrow \infty} \frac{1}{n\tau} H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau)).$$

Definition 2.3. Let η be a quasi-stationary measure on a closed set Q for a measure ν on Ω and set $\mu = \nu^{\mathbb{N}} \times \eta$. The invariance entropy for control system (1) is

$$h_\mu(Q) := \limsup_{\tau \rightarrow \infty} \inf_{\mathcal{C}_\tau} h_\mu(\mathcal{C}_\tau, Q), \quad (14)$$

where for fixed $\tau \in \mathbb{N}$ the infimum is taken over all invariant (Q, η) -partitions $\mathcal{C}_\tau = \mathcal{C}_\tau(\mathcal{P}, F)$. If no invariant (Q, η) -partition \mathcal{C}_τ exists, we set $h_\mu(Q) := \infty$.

The following remarks comment on this definition.

Remark 2. An objection to the consideration of $H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n)$ might be that \mathfrak{A}_n is not a partition of $S_Q^{-(n-1)\tau}(\mathcal{U} \times Q)$, while $\rho^{-(n-1)\tau}\mu$ is a probability measure on this space. However, one may add to the collection $\mathfrak{A}_n(\mathcal{C}_\tau)$ the complement

$$Z_n := \left(S_Q^{-(n-1)\tau}(\mathcal{U} \times Q) \right) \setminus \mathcal{A}_n(\mathcal{C}_\tau).$$

Thus one obtains a partition $\mathfrak{A}_n \cup \{Z_n\}$ of $S_Q^{-(n-1)\tau}(\mathcal{U} \times Q)$ with entropy

$$H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau) \cup \{Z_n\}) = H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau)) - \phi\left(\rho^{-(n-1)\tau}\mu(Z_n)\right).$$

For each $n \in \mathbb{N}$ the second summand is bounded by $1/e = \max_{x \in [0,1]}(-\phi(x))$ and hence

$$h_\mu(\mathcal{C}_\tau) = \limsup_{n \rightarrow \infty} \frac{1}{n\tau} H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau)) = \limsup_{n \rightarrow \infty} \frac{1}{n\tau} H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau) \cup \{Z_n\}).$$

This shows that $h_\mu(\mathcal{C}_\tau)$ is given by the entropy of bona fide partitions.

Remark 3. Definition (14) ensures that $\tau \rightarrow \infty$. This will be needed in the proof of Theorem 4.3(ii). Instead of the limit superior for $n \rightarrow \infty$ and $\tau \rightarrow \infty$ one also might consider the limit inferior. However, the limit superior is advantageous in Theorem 4.3(iii). For topological invariance entropy, one takes instead an infimum over all invariant open covers (where the partition \mathcal{P} is replaced by an open cover of Q). Then it follows that it suffices to take the limit for $\tau \rightarrow \infty$, cf. Kawan [17, Theorem 2.3 and its proof].

Remark 4. If the sets in invariant (Q, η) -partitions \mathcal{C}_τ and \mathcal{C}'_τ coincide modulo η -null sets, the entropies $H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau))$ and $H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}'_\tau))$, $n \in \mathbb{N}$, coincide. Hence it suffices to specify a partition of Q outside of a set of η -measure zero.

Remark 5. For a stationary measure η the trivial partition of Q yields an invariant partition. In fact, for every $\tau > 0$ there is an invariant (Q, η) -partition $\mathcal{C}_\tau = \mathcal{C}_\tau(\{Q\}, F)$, where the control $F(Q) \in \mathcal{U}$ can be chosen arbitrarily in a set of full $\nu^{\mathbb{N}}$ -measure in \mathcal{U} . Thus the associated metric invariance entropy vanishes. This is seen as follows: Assume, contrary to the assertion, that there is a set $\mathcal{U}_0 \subset \mathcal{U}$ with $\nu^{\mathbb{N}}(\mathcal{U}_0) > 0$ such that for every $u \in \mathcal{U}_0$

$$\eta\{x \in Q \mid \varphi(k, x, u) \in Q \text{ for } k = 1, \dots, \tau\} < \eta(Q) = 1. \quad (15)$$

We may assume that there is $k \in \{1, \dots, \tau\}$ such that for every $u \in \mathcal{U}_0$

$$\eta\{x \in Q \mid \varphi(k, x, u) \in Q\} < 1.$$

By invariance of $\nu^{\mathbb{N}} \times \eta$ Fubini's theorem yields the contradiction

$$\begin{aligned} 1 &= \int_{\mathcal{U} \times Q} S_Q^{-k}(\mathcal{U} \times Q)(\nu^{\mathbb{N}} \times \eta)(du, dx) = \int_{\mathcal{U}} \int_Q \chi_{\{\{u, x\} \mid \varphi(k, x, u) \in Q\}} \eta(dx) \nu^{\mathbb{N}}(du) \\ &< \int_{\mathcal{U}_0} \eta(Q) \nu^{\mathbb{N}}(du) + \int_{\mathcal{U} \setminus \mathcal{U}_0} \eta(Q) \nu^{\mathbb{N}}(du) = \int_{\mathcal{U}} \eta(Q) \nu^{\mathbb{N}}(du) = 1. \end{aligned}$$

Remark 6. If there exists an invariant partition, then invariant partitions with arbitrarily large time step τ exist. It suffices to see that for every invariant partition $\mathcal{C}_\tau = \mathcal{C}_\tau(\mathcal{P}, F)$ there exists an invariant partition $\mathcal{C}_{2\tau} = \mathcal{C}_{2\tau}(\mathcal{P}^2, F^2)$. In fact, for $P_i, P_j \in \mathcal{P}$ let

$$P_{ij} := \{x \in P_i \mid \varphi(\tau, x, F(P_i)) \in P_j\} = P_i \cap \varphi(\tau, \cdot, F(P_i))^{-1} P_j.$$

This yields a partition of Q given by $\mathcal{P}^2 := \{P_{ij} \mid P_i, P_j \in \mathcal{P}\}$. Define feedbacks $F^2 : \mathcal{P}^2 \rightarrow \Omega^{2\tau}$ by

$$F^2(P_{ij})(r) := \begin{cases} F(P_i)(r) & \text{for } r = 0, \dots, \tau - 1 \\ F(P_j)(r - \tau) & \text{for } r = \tau, \dots, 2\tau - 1 \end{cases}.$$

Then $\mathcal{C}_{2\tau} = (\mathcal{P}^2, F^2)$ is an invariant partition.

An alternative concept of metric invariance entropy can be based on the additional information gained in every time step (this was proposed in Colonius [6] and is slightly reformulated below). The following construction has to take into account that the space \mathcal{A}_n that is partitioned decreases in every time step.

Let an invariant partition \mathcal{C}_τ be given. Then the partition $\mathfrak{A}_n = \mathfrak{A}_n(\mathcal{C}_\tau)$ of \mathcal{A}_n induces a partition of \mathcal{A}_{n+1} : For $D \in \mathfrak{A}_n$ let

$$\mathfrak{A}_{n+1}(D) = \{E \in \mathfrak{A}_{n+1} \mid E \cap D \neq \emptyset\}, \mathcal{A}_{n+1}(D) = \bigcup_{E \in \mathfrak{A}_{n+1}(D)} E. \quad (16)$$

Since $E \cap D \neq \emptyset$ implies $E \subset D$ and the sets $\mathcal{A}_{n+1}(D)$ are mutually disjoint, one obtains an induced partition

$$\mathfrak{A}_n^{n+1} := \{\mathcal{A}_{n+1}(D) \mid D \in \mathfrak{A}_n\} \text{ of } \mathcal{A}_{n+1} = \bigcup_{D \in \mathfrak{A}_n} \mathcal{A}_{n+1}(D). \quad (17)$$

Clearly, \mathfrak{A}_{n+1} is a refinement of \mathfrak{A}_n^{n+1} . The information from \mathfrak{A}_n that is relevant for \mathfrak{A}_{n+1} comes from the partition \mathfrak{A}_n^{n+1} . Assuming that the information encoded in \mathfrak{A}_n^{n+1} is known at the time step n , the incremental information is the conditional entropy of \mathfrak{A}_{n+1} given \mathfrak{A}_n^{n+1} (with respect to $\rho^{-n\tau}\mu$). For every $n \in \mathbb{N}$

$$H_{\rho^{-n\tau}\mu}(\mathfrak{A}_{n+1}) = H_{\rho^{-n\tau}\mu}(\mathfrak{A}_n^{n+1}) + H_{\rho^{-n\tau}\mu}(\mathfrak{A}_{n+1} \mid \mathfrak{A}_n^{n+1}),$$

where the conditional entropy of \mathfrak{A}_{n+1} given \mathfrak{A}_n^{n+1} is

$$H_{\rho^{-n\tau}\mu}(\mathfrak{A}_{n+1} | \mathfrak{A}_n^{n+1}) = - \sum_{D \in \mathfrak{A}_n} \rho^{-n\tau} \mu(\mathcal{A}_{n+1}(D)) \sum_{E \in \mathfrak{A}_{n+1}} \phi \left(\frac{\mu(D \cap E)}{\mu(\mathcal{A}_{n+1}(D))} \right). \quad (18)$$

(Observe that in the argument of ϕ one may multiply numerator and denominator by $\rho^{-n\tau}$.) Taking the average incremental information one arrives at the following notion.

Definition 2.4. Let η be a quasi-stationary measure on a closed set Q for a measure ν on Ω and set $\mu = \nu^{\mathbb{N}} \times \eta$. For an invariant (Q, η) -partition $\mathcal{C}_\tau = \mathcal{C}_\tau(\mathcal{P}, F)$ define the incremental invariance entropy of \mathcal{C}_τ by

$$h_\mu^{\text{inc}}(\mathcal{C}_\tau, Q) = \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \sum_{j=0}^{n-1} H_{\rho^{-j\tau}\mu}(\mathfrak{A}_{j+1}(\mathcal{C}_\tau) | \mathfrak{A}_j^{j+1}(\mathcal{C}_\tau)),$$

and define the incremental invariance entropy for control system (1) by

$$h_\mu^{\text{inc}}(Q) := \limsup_{\tau \rightarrow \infty} \inf_{\mathcal{C}_\tau} h_\mu^{\text{inc}}(\mathcal{C}_\tau, Q), \quad (19)$$

where the infimum is taken over all invariant (Q, η) -partitions $\mathcal{C}_\tau(\mathcal{P}, F)$. If no invariant (Q, η) -partition \mathcal{C}_τ exists, we set $h_\mu^{\text{inc}}(Q) := \infty$.

Regrettably, a formula analogous to (2) for dynamical systems is not available for invariance entropy of control systems. Hence the relation between the invariance entropy and the incremental invariance entropy remains unknown. The following proposition only describes a relation between the entropy of \mathfrak{A}_n and of the induced partition \mathfrak{A}_n^{n+1} .

Proposition 1. *There is $K \in \mathbb{N}$ such that for all $n \in \mathbb{N}$*

$$H_{\rho^{-n\tau}\mu}(\mathfrak{A}_n^{n+1}) \leq H_{\rho^{-n\tau}\mu}(\mathfrak{A}_n) + K/e.$$

Proof. There is $K \in \mathbb{N}$ such that for every probability measure m there are at most K mutually disjoint sets A_1, \dots, A_K with $m(A_i) > \rho^\tau/e$, since $\sum_{i=1}^K m(A_i) \leq 1$. In particular, for every $n \in \mathbb{N}$ there are at most K sets $D \in \mathfrak{A}_n$ such that $\rho^{-(n-1)\tau}\mu(D) > \rho^\tau/e$, since every $D \in \mathfrak{A}_n$ is contained in $S_Q^{-(n-1)\tau}(\mathcal{U} \times Q)$ and $\rho^{-(n-1)\tau}\mu$ is a probability measure on this set. This inequality is equivalent to $\rho^{-n\tau}\mu(D) > 1/e$. Let $\mathfrak{A}_n^{\text{big}}$ be the set of elements in \mathfrak{A}_n with $\rho^{-n\tau}\mu(D) > 1/e$. The other elements D in \mathfrak{A}_n satisfy

$$\rho^{-n\tau}\mu(\mathcal{A}_{n+1}(D)) \leq \rho^{-n\tau}\mu(D) \leq 1/e.$$

Since ϕ is monotonically decreasing on $[0, 1/e]$ it follows that $\phi(\rho^{-n\tau}\mu(\mathcal{A}_{n+1}(D))) \geq \phi(\rho^{-n\tau}\mu(D))$ and hence

$$H_{\rho^{-n\tau}\mu}(\mathfrak{A}_n^{n+1}) \leq H_{\rho^{-n\tau}\mu}(\mathfrak{A}_n) - K \min \phi \leq H_{\rho^{-n\tau}\mu}(\mathfrak{A}_n) + K/e.$$

□

Next we analyze the behavior of both notions of invariance entropy under measure preserving transformations (cf. Walters [25, §2.3]). For notational simplicity, we suppose that the control ranges and the measures on them coincide.

Definition 2.5. Consider two control systems of the form (1) on M_1 and M_2 , respectively, given by

$$x_{k+1} = f_1(x_k, u_k) \text{ and } y_{k+1} = f_2(y_k, u_k) \text{ with } (u_k) \in \mathcal{U} = \Omega^{\mathbb{N}}. \quad (20)$$

Let ν be a probability measure on Ω and suppose that η_1 and η_2 are corresponding quasi-stationary measures with respect to closed subsets $Q_1 \subset M_1$ and $Q_2 \subset M_2$, respectively. We say that (f_2, η_2) is semi-conjugate to (f_1, η_1) , if there are subsets $\hat{\Omega} \subset \Omega$ and $\hat{Q}_i \subset Q_i$ of full ν -measure and full η_i -measure, $i = 1, 2$, resp., with the following properties:

(i) there exists a measurable map $\pi : \hat{Q}_1 \rightarrow \hat{Q}_2$ such that π maps η_1 onto η_2 , i.e.,

$$(\pi_* \eta_1)(B) := \eta_1(\pi^{-1}B) = \eta_2(B) \text{ for all } B \subset \hat{Q}_2, \quad (21)$$

(ii) one has $f_i(x, \omega) \in \hat{Q}_i$ for all $(\omega, x) \in \hat{\Omega} \times \hat{Q}_i$, $i = 1, 2$, and

$$\pi(f_1(x, \omega)) = f_2(\pi x, \omega) \text{ for } \omega \in \hat{\Omega} \text{ and } x \in \hat{Q}_1. \quad (22)$$

The map π is called a semi-conjugacy from (f_1, η_1) to (f_2, η_2) . In terms of the solutions, condition (22) implies that for $\nu^{\mathbb{N}}$ -a.a. $u \in \Omega^{\mathbb{N}}$ and η_1 -a.a. $x \in Q_1$

$$\pi \varphi_1(k, x_0, u) = \varphi_2(k, \pi x_0, u) \text{ for all } k \in \mathbb{N}.$$

With the associated skew product maps $S_i(u, x) = (\theta u, f_i(x, u_0))$ one obtains $S_i(\hat{\Omega}^{\mathbb{N}} \times \hat{Q}_i) \subset \hat{\Omega}^{\mathbb{N}} \times \hat{Q}_i$, $i = 1, 2$, and for all $(u, x) \in \hat{\Omega}^{\mathbb{N}} \times \hat{Q}_1$

$$\begin{aligned} (\text{id}_{\mathcal{U}} \times \pi) \circ S_1(u, x) &= (\text{id}_{\mathcal{U}} \times \pi)(\theta u, f_1(x, u_0)) = (\theta u, f_2(\pi x, u_0)) \\ &= S_2 \circ (\text{id}_{\mathcal{U}} \times \pi)(u, x). \end{aligned} \quad (23)$$

If the map π is a bimeasurable bijection, we obtain an equivalence relation called conjugacy. A consequence of the following theorem is that the metric invariance entropies are invariant under conjugacies.

Theorem 2.6. *Suppose that for two control systems given by (20) there is a semi-conjugacy π from (f_1, η_1) to (f_2, η_2) . Then the constants ρ_i coincide and with $\mu_i = \nu^{\mathbb{N}} \times \eta_i$, $i = 1, 2$, the metric invariance entropy satisfies*

$$h_{\mu_1}(Q_1) \leq h_{\mu_2}(Q_2) \text{ and } h_{\mu_1}^{\text{inc}}(Q_1) \leq h_{\mu_2}^{\text{inc}}(Q_2).$$

Proof. First observe that $\eta_2(\hat{Q}_2) = \eta_1(\pi^{-1}\hat{Q}_2) = \eta_1(\hat{Q}_1)$ and

$$\mu_2 = \nu^{\mathbb{N}} \times \eta_2 = (\text{id}_{\mathcal{U}} \times \pi)_* (\nu^{\mathbb{N}} \times \eta_1) = (\text{id}_{\mathcal{U}} \times \pi)_* \mu_1.$$

Furthermore $\rho_1 = \rho_2$, since properties (21) and (23) imply

$$\begin{aligned} \rho_2 &= \rho \mu_2(\mathcal{U} \times \hat{Q}_2) = \mu_2\{(u, y) \in \mathcal{U} \times Q_2 \mid S_2(u, y) \in \mathcal{U} \times \hat{Q}_2\} \\ &= \mu_2\{(u, y) \in \hat{\Omega}^{\mathbb{N}} \times \hat{Q}_2 \mid S_2(u, y) \in \mathcal{U} \times \hat{Q}_2\} \\ &= \mu_1(\text{id}_{\mathcal{U}} \times \pi)^{-1}\{(u, y) \in \hat{\Omega}^{\mathbb{N}} \times \hat{Q}_2 \mid S_2(u, y) \in \mathcal{U} \times \hat{Q}_2\} \\ &= \mu_1\{(u, x) \in \hat{\Omega}^{\mathbb{N}} \times \hat{Q}_1 \mid S_2 \circ (\text{id}_{\mathcal{U}} \times \pi)(u, x) \in \mathcal{U} \times \hat{Q}_2\} \\ &= \mu_1\{(u, x) \in \hat{\Omega}^{\mathbb{N}} \times \hat{Q}_1 \mid (\text{id}_{\mathcal{U}} \times \pi) \circ S_1(u, x) \in \mathcal{U} \times \hat{Q}_2\} \\ &= \mu_1\{(u, x) \in \hat{\Omega}^{\mathbb{N}} \times \hat{Q}_1 \mid S_1(u, x) \in \mathcal{U} \times \hat{Q}_1\} \\ &= \mu_1\{(u, x) \in \mathcal{U} \times Q_1 \mid S_1(u, x) \in \mathcal{U} \times \hat{Q}_1\} \\ &= \rho_1 \mu_1(\mathcal{U} \times \hat{Q}_1) = \rho_1. \end{aligned}$$

Let $\mathcal{C}_{2,\tau} = \mathcal{C}_{2,\tau}(\mathcal{P}_2, F)$ be an invariant (Q_2, η_2) -partition. Then it follows that $\pi^{-1}\mathcal{P}_2 = \{\pi^{-1}P \mid P \in \mathcal{P}_2\}$ is a measurable partition of $Q_1 = \pi^{-1}Q_2$ modulo η_1 -null sets

and we may assume that $\pi^{-1}P \subset \hat{Q}_1$ for all P . For $P \in \mathcal{P}_2$ it follows that for η_2 -a.a. $x \in P$ one has $\varphi_2(k, x, F(P)) \in Q_2$ for all $k \in \{1, \dots, \tau\}$ if and only if for η_1 -a.a. $y \in \pi^{-1}P$ one has $y = \pi^{-1}x$ for some $x \in P$ and

$$\varphi_1(k, y, F(P)) \in \pi^{-1}\varphi_2(k, x, F(P)) \in \pi^{-1}Q_2 = Q_1 \text{ for all } k \in \{1, \dots, \tau\}$$

(note that the preimage under π of an η_2 -null set is an η_1 -null set). By Remark 4 it follows that $\mathcal{C}_{1,\tau} = \mathcal{C}_{1,\tau}(\pi^{-1}\mathcal{P}_2, F)$ with $F(\pi^{-1}P) := F(P)$, $\pi^{-1}P \in \pi^{-1}\mathcal{P}_1$, is an invariant (Q_1, η_1) -partition. Then the preimage of the collection $\mathfrak{A}(\mathcal{C}_{2,\tau})$ of $\mathcal{U} \times Q_1$ equals the collection

$$\mathfrak{A}(\mathcal{C}_{1,\tau}) = \{(\text{id}_{\mathcal{U}} \times \pi)^{-1}A \mid A \in \mathfrak{A}(\mathcal{C}_{2,\tau})\}.$$

Let a be a (Q_2, η_2) -admissible word. Then

$$\begin{aligned} 0 &< \eta_2 \{y \in Q_2 \mid \varphi_2(i\tau, y, u_a) \in P_{a_i} \text{ for } i = 0, 1, \dots, n-1\} \\ &= \eta_2 \{y \in \hat{Q}_2 \mid \varphi_2(i\tau, y, u_a) \in P_{a_i} \text{ for } i = 0, 1, \dots, n-1\} \\ &= \eta_1 \{\pi^{-1}y \in \hat{Q}_1 \mid \varphi_2(i\tau, y, u_a) \in P_{a_i} \text{ for } i = 0, 1, \dots, n-1\} \\ &= \eta_1 \{x \in \hat{Q}_1 \mid \varphi_2(i\tau, \pi x, u_a) \in P_{a_i} \text{ for } i = 0, 1, \dots, n-1\} \\ &= \eta_1 \{x \in \hat{Q}_1 \mid \pi\varphi_1(i\tau, x, u_a) \in P_{a_i} \text{ for } i = 0, 1, \dots, n-1\} \\ &= \eta_1 \{x \in \hat{Q}_1 \mid \varphi_1(i\tau, x, u_a) \in \pi^{-1}P_{a_i} \text{ for } i = 0, 1, \dots, n-1\}. \end{aligned}$$

It follows that a is also (Q_1, η_1) -admissible. The same arguments show that every (Q_1, η_1) -admissible word is also (Q_2, η_2) -admissible and hence for all $n \in \mathbb{N}$

$$(\text{id}_{\mathcal{U}} \times \pi)^{-1}\mathfrak{A}_n(\mathcal{C}_{2,\tau}) = \mathfrak{A}_n(\mathcal{C}_{1,\tau}).$$

One finds for the entropy

$$H_{\rho_1^{-(n-1)\tau} \mu_1}(\mathfrak{A}_n(\mathcal{C}_{1,\tau})) = H_{\rho_2^{-(n-1)\tau} \mu_2}(\mathfrak{A}_n(\mathcal{C}_{2,\tau})), n \in \mathbb{N},$$

and hence $h_{\mu_1}(\mathcal{C}_{1,\tau}) = h_{\mu_2}(\mathcal{C}_{2,\tau})$. Taking first the infimum over all invariant (Q_2, η_2) -partitions and then over all invariant (Q_1, η_1) -partitions one finds that $h_{\mu_1}(Q_1) \leq h_{\mu_2}(Q_2)$. These arguments also show that $h_{\mu_2}(Q_1) = \infty$ if $h_{\mu_1}(Q_2) = \infty$. For the incremental invariance entropy one similarly finds that for all $n \in \mathbb{N}$

$$(\text{id}_{\mathcal{U}} \times \pi)\mathfrak{A}_n^{n+1}(\mathcal{C}_{1,\tau}) = \mathfrak{A}_n^{n+1}(\mathcal{C}_{2,\tau}).$$

Then it follows that

$$H_{\rho^{-n\tau} \mu_1}(\mathfrak{A}_{n+1}(\mathcal{C}_{1,\tau}) \mid \mathfrak{A}_n^{n+1}(\mathcal{C}_{1,\tau})) = H_{\rho^{-n\tau} \mu_2}(\mathfrak{A}_{n+1}(\mathcal{C}_{2,\tau}) \mid \mathfrak{A}_n^{n+1}(\mathcal{C}_{2,\tau}))$$

and the inequality of the incremental invariance entropies is a consequence. \square

Remark 7. Observe that the inequalities for invariance entropies under semi-conjugacy are opposite to the inequalities for entropy of dynamical systems, cf. Viana and Oliveira [24, Exercise 9.1.5]. This is due to the fact that we construct invariant (Q_1, η_1) -partitions from invariant (Q_2, η_2) -partitions and then take the infimum (instead of the supremum) of partitions. Note also that for topological invariance entropy Kawan [17, Proposition 2.13] constructs from spanning sets of controls for Q_1 spanning sets for Q_2 . Then letting the time tend to infinity and taking the infimum over spanning sets one gets that the invariance entropy of Q_1 is greater than or equal to the invariance entropy of Q_2 .

To conclude this section we show that the metric invariance entropy is bounded above by the topological invariance entropy. As in Kawan [17, Definition 2.2 and Proposition 2.3(ii)] consider for system (1) a compact controlled invariant set $Q \subset M$, i.e., for every $x \in Q$ there is $\omega_x \in \Omega$ with $f(x, \omega_x) \in Q$. For $\tau \in \mathbb{N}$ a set $\mathcal{R} \subset \mathcal{U}$ is called (τ, Q) -spanning if for all $x \in Q$ there is $u \in \mathcal{R}$ with $\varphi(n, x, u) \in Q$ for all $n = 1, \dots, \tau$. Denote by $r_{inv}(\tau, Q)$ the minimal number of elements such a set can have (if no finite (τ, Q) -spanning set exists, $r_{inv}(\tau, Q) := \infty$). The topological invariance entropy is defined by

$$h_{inv}(Q) := \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \log r_{inv}(\tau, Q).$$

In order to relate this notion to metric invariance entropy, we use a characterization of topological invariance entropy by invariant partitions (the original definition due to Nair et al. [19] uses invariant open covers). Here a (topological) invariant partition $\mathcal{C}_\tau = \mathcal{C}(\mathcal{P}, \tau, F)$ is defined by $\tau \in \mathbb{N}$, a finite measurable partition $\mathcal{P} = \{P_1, \dots, P_q\}$ of Q and $F: \mathcal{P} \rightarrow \Omega^\tau$ such that

$$\varphi(k, P, F(P)) \subset Q \text{ for } k = 1, \dots, \tau.$$

Thus, in contrast to (Q, η) -invariant partitions (cf. Definition 2.1), it is required that every $x \in P$ remains in Q under the feedback $F(P)$. Then a word $a = [a_0, a_1, \dots, a_{n-1}]$ is called admissible if there exists a point $x \in Q$ with $\varphi(i\tau, x, u_a) \in P_{a_i}$ for $i = 0, 1, \dots, n-1$. Write $\#\mathcal{W}_n(\mathcal{C}_\tau)$ for the number of elements in the set $\mathcal{W}_n(\mathcal{C}_\tau)$ of all admissible words of length n and define the entropy of \mathcal{C}_τ by

$$h_{top}(\mathcal{C}_\tau) := \lim_{n \rightarrow \infty} \frac{\log \#\mathcal{W}_n(\mathcal{C}_\tau)}{n\tau} = \inf_{n \in \mathbb{N}} \frac{\log \#\mathcal{W}_n(\mathcal{C}_\tau)}{n\tau}.$$

A topological invariant partition of Q is also a (Q, η) -partition and an η -admissible word is also admissible in the topological sense. The following characterization of topological invariance entropy is given in Kawan [17, Theorem 2.3 and its proof].

Theorem 2.7. *For a compact and controlled invariant set Q it holds that*

$$h_{inv}(Q) = \inf_{\mathcal{C}_\tau} h(\mathcal{C}_\tau) = \lim_{\tau \rightarrow \infty} \inf_{\mathcal{C}_\tau} h(\mathcal{C}_\tau),$$

where the first infimum is taken over all invariant Q -partitions \mathcal{C}_τ and the second infimum is taken over all invariant Q -partitions \mathcal{C}_τ with fixed $\tau \in \mathbb{N}$.

The following theorem relates metric and topological invariance entropy.

Theorem 2.8. *Let Q be a compact and controlled invariant set Q . Then for every quasi-stationary measure η on Q the metric entropy with respect to $\mu = \nu^N \times \eta$ satisfies*

$$h_\mu(Q) \leq h_{inv}(Q).$$

Proof. Let η be a quasi-stationary measure on Q and fix a topological invariant partition $\mathcal{C}_\tau(\mathcal{P}, F)$. Then $\mathcal{C}_\tau(\mathcal{P}, F)$ is also a (Q, η) -invariant partition and $\#\mathfrak{A}_n(\mathcal{C}_\tau) \leq \#\mathcal{W}_n(\mathcal{C}_\tau)$ for every $n \in \mathbb{N}$, since an η -admissible word is also trivially admissible in the topological sense. Using

$$H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau)) \leq \log \#\mathfrak{A}_n(\mathcal{C}_\tau)$$

one finds that

$$h_\mu(\mathcal{C}_\tau, Q) = \limsup_{n \rightarrow \infty} \frac{1}{n\tau} H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau)) \leq \lim_{n \rightarrow \infty} \frac{\log \#\mathcal{W}_n(\mathcal{C}_\tau)}{n\tau} = h(\mathcal{C}_\tau).$$

This yields the assertion $h_\mu(Q) \leq h_{inv}(Q)$. \square

The following example illustrates the existence of quasi-stationary measures in a simple situation (it is a modification of Colonius, Homburg and Kliemann [8, Example 1]).

Example 2.9. Consider the family of control systems depending on a real parameter given by $f_\alpha : \mathbb{R}/\mathbb{Z} \times [-1, 1] \rightarrow \mathbb{R}/\mathbb{Z}$,

$$f_\alpha(x, \omega) = x + \sigma \cos(2\pi x) + A\omega + \alpha \pmod{1}. \quad (24)$$


Suppose that the amplitudes A and σ as well as α take on small positive values. Let a probability measure ν on $\Omega := [-1, 1]$ be given. One obtains Markov transition probabilities

$$p(x, B) = \nu\{\omega \in \Omega \mid f_\alpha(x, \omega) \in B\} \text{ for } x \in \mathbb{R}/\mathbb{Z} \text{ and } B \subset \mathbb{R}/\mathbb{Z}.$$

For $\alpha_0 = \sigma - A$ the extremal graph $f_{\alpha_0}(\cdot, 1)$ is tangent to the diagonal at a point c_0 . Now let $\alpha > \alpha_0$ and consider $Q = [0.2, 0.5]$. Colonius [6, Theorem 2.9] implies the existence of a quasi-stationary measure η for Q with $0 < \rho < 1$ if ν has a density with respect to Lebesgue measure and there is $\gamma > 0$ such that $p(x, Q) \geq \gamma > 0$ for all $x \in Q$. If we take the uniform distribution on $\Omega = [-1, 1]$, these conditions are satisfied. By [6, Proposition 2.4], the support of the corresponding conditionally invariant measure $\mu = \nu^N \times \eta$ is contained in

$$\{(u, x) \in \mathcal{U} \times Q \mid S_Q^{-n}(u, x) \cap (\mathcal{U} \times Q) \neq \emptyset \text{ for all } n \in \mathbb{N}\}.$$

Let $d(\alpha) < 0.5$ be given by the intersection of the lower sinusoidal curve $f_\alpha(\cdot, -1)$ (cf. the figure below) with the diagonal. Then points to the left of $[d(\alpha), 0.5]$ leave Q backwards in time, hence they cannot be in the support of η . Thus the quasi-stationary measure η has support contained in $[d(\alpha), 0.5]$. Observe that for the uniform distribution on Ω there is no stationary measure η with support in Q , hence there is no invariant measure μ of the form $\mu = \nu^N \times \eta$.



Extremal graphs for (24) and the set $[d(\alpha), 0.5]$ in $Q = [0.2, 0.5]$
(here $A = 0.05, \sigma = 0.1$ and $\alpha = 0.08$)

The Variational Principle for dynamical systems states that the supremum of the metric entropies coincides with the topological entropy. Certainly, an analogous result for invariance entropy would give considerable structural insight. Apart from this, however, it would only be of limited interest. The metric invariance entropy is introduced since it is smaller than the topological invariance entropy (see also the discussion of coder-controllers in Section 3). Instead of asking for measures μ maximizing the entropy one should instead look for measures minimizing h_μ over a class \mathcal{M}_{ad} of admissible measures, hence to determine measures μ_0 with minimal invariance entropy, i.e.,

$$h_{\mu_0}(Q) = \inf_{\mu \in \mathcal{M}_{ad}} h_\mu(Q).$$

This induces the question over which class \mathcal{M}_{ad} of measures one should minimize. The set of all measures $\mu = \nu^{\mathbb{N}} \times \eta$ where η is quasi-stationary with respect to ν is too big, since one would often obtain that the minimum is zero: This is illustrated by Example 2.9. There are many stationary measures S with support in Q : Take $\mu = (\delta_\omega)^{\mathbb{N}} \times \delta_{x(u)}$, where δ_ω and $\delta_{x(\omega)}$ are Dirac measures with $f(x(\omega), \omega) = x(\omega) \in Q$ and $h_\mu = 0$. The reason is that by the choice $\nu = \delta_\omega$ the invariance problem is already solved. Instead it seems reasonable to require at least that the support of ν coincides with the control range Ω , since otherwise we would already know that we do not need controls in $\Omega \setminus \text{supp } \nu$. If $\Omega \subset \mathbb{R}^m$ has positive Lebesgue measure $\lambda(\Omega)$ one might even restrict attention to the measures ν which are absolutely continuous with respect to λ . Concerning the support of the quasi-stationary measure η it is immediately clear that $Q \setminus \text{supp } \eta$ does not contribute to h_μ . Some further results will be given in Sections 4 and 5.

3. Relations to coder-controllers. This section defines coder-controllers associated with a quasi-stationary measure rendering Q invariant, and shows that their minimal entropy coincides with the entropy $h_\mu(Q)$.

A coder-controller (cf., e.g., Kawan [17, Section 2.5]) may be defined as a quadruple $\mathcal{H} = (S, \gamma, \delta, \tau)$ where $S = (S_k)_{k \in \mathbb{N}}$ denotes finite coding alphabets and the coder mapping $\gamma_k : M^{k+1} \rightarrow S_k$ associates to the present and past states the symbol $s_k \in S_k$. At time $k\tau$, the controller has $k+1$ symbols s_0, \dots, s_k available and generates a control $u_k \in \Omega^\tau$. The corresponding controller mapping is $\delta_k : S_0 \times \dots \times S_k \rightarrow \Omega^\tau$. Thus of interest are for $x_0 \in Q$ the sequences

$$x_{k+1} := \varphi(\tau, x_k, u_k), k \in \mathbb{N}, \quad (25)$$

with

$$u_k = \delta_k(\gamma_0(x_0), \gamma_1(x_0, x_1), \dots, \gamma_k(x_0, x_1, \dots, x_k)) \in \Omega^\tau \quad (26)$$

satisfying

$$\varphi(i, x_k, u_k) \in Q \text{ for all } i \in \{1, \dots, \tau\} \text{ and all } k \in \mathbb{N}. \quad (27)$$

In the following construction, we suppose that the coding alphabet S is independent of k . Hence

$$\gamma_k : M^{k+1} \rightarrow S \text{ and } \delta_k : S^{k+1} \rightarrow \Omega^\tau.$$

Each sequence of symbols in S^k defines a coding region in Q which is defined as the set of all initial states x_0 which force the coder to generate this sequence. More precisely, for $s = (s_0, \dots, s_{k-1}) \in S^k$ let

$$P_s := \{x_0 \in Q \mid \gamma_0(x_0) = s_0, \dots, \gamma_{k-1}(x_0, x_1, \dots, x_{k-1}) = s_{k-1}\},$$

where $x_j, j = 1, \dots, k-1$, are generated by (25), (26). Furthermore, let

$$u_s := (\delta_0(s_0), \delta_1(s_0, s_1), \dots, \delta_{k-1}(s_0, \dots, s_{k-1})) \in \Omega^{k\tau}.$$

Thus P_s consists of the points x_0 satisfying for $j = 0, \dots, k-1$

$$(x_0, \varphi(\tau, x_0, u_s), \dots, \varphi(j\tau, x_0, u_s)) \in \gamma_j^{-1}(s_j),$$

and one has $\varphi(i, x_0, u_s) \in Q$ for all $i = 0, \dots, k\tau$.

Again, suppose that a quasi-stationary measure η with $\rho = \int_Q p(x, Q)\eta(dx)$ corresponding to ν on Ω is fixed and let $\mu = \nu^{\mathbb{N}} \times \eta$. The set S_{ad}^k of (Q, η) -admissible words is the set of $s \in S^k$ with $\eta(P_s) > 0$.

For a (Q, η) -admissible word we consider all controls u such that the application of u yields the same word and renders Q invariant with probability 1. More explicitly, define for $s_0 \in S$

$$A(P_{s_0}) := \{u \in \mathcal{U} \mid \varphi(i, x, u) \in Q \text{ for } i = 1, \dots, \tau \text{ and } \eta\text{-a.a. } x \in P_{s_0}\} \times P_{s_0}$$

and for $s = (s_0, \dots, s_{k-1}) \in S_{ad}^k$

$$D_s := A(P_{s_0}) \cap S^{-\tau} A(P_{s_1}) \cap \dots \cap S^{-(k-1)\tau} A(P_{s_{k-1}}) \subset S_Q^{-(k-1)\tau} (\mathcal{U} \times Q).$$

Observe that for $(u, x_0) \in D_s$ one obtains $\varphi(j\tau, x_0, u) \in P_{s_j}$, $j = 0, \dots, k-1$. Define a measure λ_k on the (finite) set S_{ad}^k by

$$\lambda_k(s) := \rho^{-(k-1)\tau} \mu(D_s). \quad (28)$$

The considered coder maps $\gamma_k : M^{k+1} \rightarrow S$ will only be defined on a set of full η^{k+1} -measure and the considered controller maps δ_k will only be defined on a set of full λ_{k+1} -measure. The following coder-controllers render Q invariant.

Definition 3.1. A (Q, η) -coder-controller is a quadruple $\mathcal{H} = (S, \gamma, \delta, \tau)$ as above such that $\{P_s \mid s \in S\}$ forms a partition (modulo η -null sets) of Q . The entropy of \mathcal{H} is defined as

$$R(\mathcal{H}) = \limsup_{k \rightarrow \infty} \frac{1}{k\tau} H_{\lambda_k}(S_{ad}^k).$$

Remark 8. In general, $\lambda_k(S_{ad}^k) < 1$, hence the measures λ_k are not probability measures. Nevertheless, it makes sense to consider the associated entropy, cf. Remark 2 for a similar situation.

Remark 9. Kawan [17, p. 72 and p. 83] (see also Nair, Evans, Mareels and Moran [19]) defines coder-controllers with and without τ (i.e., $\tau = 1$) and the coder maps γ_k may or may not depend on the past symbols (in addition to the past states). Furthermore, it is usually assumed that the size of the set of symbols may vary with k . Using the same set of symbols for every k amounts to requiring that $\sup_{k \in \mathbb{N}} \#S_k < \infty$ where $\#S_k$ is the size of the set S_k of symbols at time $k\tau$. In the (topological) definition of data rates this might be taken into account by looking at the number of symbols actually used at time $k\tau$. In fact, $\sup_{k \in \mathbb{N}} \#S_k < \infty$ in the situations considered in [17, Theorem 2.1 and Theorem 2.4].

When one wants to relate coder-controllers to the invariance entropy in Definition 2.3, it may appear rather straightforward to identify the elements of an invariant partition with the set of symbols for a coder-controller and the feedbacks $F(P)$ with the controls generated by a coder-controller. This is the content of the following theorem.

Theorem 3.2. Consider system (1) and let η be a quasi-stationary measure for a closed set Q corresponding to ν on Ω and denote $\mu = \nu^{\mathbb{N}} \times \eta$. The μ -invariance entropy satisfies

$$h_\mu(Q) = \limsup_{\tau \rightarrow \infty} \inf_{\mathcal{H}} R(\mathcal{H}),$$

where the infimum is taken over all (Q, η) -coder-controllers $\mathcal{H} = (S, \gamma, \delta, \tau)$.

Proof. (i) First we show that for every invariant (Q, η) -partition $\mathcal{C}_\tau = \mathcal{C}_\tau(\mathcal{P}, F)$ with $\mathcal{P} = \{P_1, \dots, P_q\}$ there exists a (Q, η) -coder-controller $\mathcal{H} = (S, \gamma, \delta, \tau)$ rendering Q invariant such that $R(\mathcal{H}) = h_\mu(\mathcal{C}_\tau)$. In order to construct \mathcal{H} let $S := \{1, \dots, q\}$ and for $k \geq 1$ let S_{ad}^k be the set of \mathcal{C}_τ -admissible words a of length k . Denote by λ_k the measure on S_{ad}^k given by

$$\lambda_k(a) = \rho^{-(k-1)\tau} \mu(D_a), D_a \in \mathfrak{A}_k(\mathcal{C}_\tau). \quad (29)$$

It follows that

$$H_{\lambda_k}(S_{ad}^k) = H_{\rho^{-(k-1)\tau} \mu}(\mathfrak{A}_k(\mathcal{C}_\tau)).$$

The coders $\gamma_k : Q^{k+1} \rightarrow S$ are defined by

$$\gamma_k(x_0, \dots, x_k) := a_k \text{ if } x_k \in P_{a_k}.$$

Then the entropy associated to this coder is

$$\limsup_{k \rightarrow \infty} \frac{1}{k\tau} H_{\lambda_k}(S_{ad}^k) = \limsup_{k \rightarrow \infty} \frac{1}{k\tau} H_{\rho^{-(k-1)\tau} \mu}(\mathfrak{A}_k(\mathcal{C}_\tau)) = h_\mu(\mathcal{C}_\tau).$$

The controller is constructed as follows. Each set $D_a \in \mathfrak{A}_k(\mathcal{C}_\tau)$ is of the form

$$D_a = A(P_{a_0}) \cap \dots \cap S^{-(k-1)\tau} A(P_{a_{k-1}})$$

with $A(P_{a_i}) \in \mathfrak{A}(\mathcal{C}_\tau)$ for all i . Upon receiving the symbol a_{k-1} in addition to the previous symbols a_0, \dots, a_{k-2} the controller finds $a = (a_0, \dots, a_{k-1})$ which indexes an element of $\mathfrak{A}_k(\mathcal{C}_\tau)$. Then the controller is given by the maps $\delta_{k-1} : S_{ad}^k \rightarrow \Omega^\tau$,

$$\delta_{k-1}(a_0, \dots, a_{k-1}) := F(P_{a_{k-1}}).$$

By the definition this yields for the corresponding solution that $x_{(k-1)\tau+i} \in Q, i = 0, \dots, \tau$, and hence the constructed coder-controller renders Q invariant. Taking the infimum over all invariant (Q, η) -partitions \mathcal{C}_τ , then the infimum over all (Q, η) -coder-controllers $\mathcal{H} = (S, \gamma, \delta, \tau)$ and, finally, the limit superior for $\tau \rightarrow \infty$ one obtains

$$\limsup_{\tau \rightarrow \infty} \inf_{\mathcal{H}} R(\mathcal{H}) \leq h_\mu(Q).$$

(ii) For the converse inequality it suffices to show that for an arbitrary (Q, η) -coder-controller $\mathcal{H} = (S, \gamma, \delta, \tau)$ there is an invariant (Q, η) -partition \mathcal{C}_τ with $h_\mu(\mathcal{C}_\tau) = R(\mathcal{H})$. For every $s_0 \in S$ consider the set P_{s_0} of all points $x_0 \in Q$ such that $s_0 = \gamma_0(x_0)$. Since \mathcal{H} renders Q invariant it follows that for $u_0 = \delta_0(s) \in \Omega^\tau$ (this does not denote a Dirac measure!) one has $\varphi(i, x_0, u_0) \in Q$ for $i = 1, \dots, \tau$ and η -a.a. $x_0 \in P_{s_0}$. Since $\mathcal{P} = \{P_{s_0} | s_0 \in S\}$ is a partition modulo η -null sets of Q , one obtains that $\mathcal{C}_\tau = \mathcal{C}_\tau(\mathcal{P}, F)$ with $F(P_{s_0}) = u_0 = \delta_0(s_0), s_0 \in S$, is an invariant partition of Q . Then

$$A(P_{s_0}) = \{u \in \Omega^\tau | \varphi(i, x, u) \in Q \text{ for } i = 1, \dots, \tau \text{ and for } \eta\text{-a.a. } x \in P_{s_0}\} \times P_{s_0}$$

and $\mathfrak{A}(\mathcal{C}_\tau) = \{A(P_{s_0}) | s_0 \in S\}$. For $s = (s_0, \dots, s_{k-1}) \in S_{ad}^k$ we have

$$D_s = A(P_{s_0}) \cap S^{-\tau} A(P_{s_1}) \cap \dots \cap S^{-(k-1)\tau} A(P_{s_{k-1}}).$$

Then the probability measure λ_k on S_{ad}^k satisfies $\lambda_k(s) = \rho^{-(k-1)\tau} \mu(D_s)$ and hence

$$H_{\lambda_k}(S_{ad}^k) = H_{\rho^{-(k-1)\tau} \mu}(\mathfrak{A}_k(\mathcal{C}_\tau)).$$

Thus one obtains for the entropy

$$R(\mathcal{H}) = \limsup_{k \rightarrow \infty} \frac{1}{k\tau} H_{\lambda_k}(S_{ad}^k) = \limsup_{k \rightarrow \infty} \frac{1}{k\tau} H_{\rho^{-(k-1)\tau} \mu}(\mathfrak{A}_k(\mathcal{C}_\tau)) = h_\mu(\mathcal{C}_\tau). \quad \square$$

4. Invariance entropy and relative invariance. In this section and in Section 5, we analyze when the metric invariance entropy of Q is already determined on certain subsets of Q . The analysis is based on the following relative invariance property.

Definition 4.1. Consider for system (1) subsets $K \subset Q \subset M$. The set K is called invariant in Q , if $x \in K$ and $f(x, \omega) \notin K$ for some $\omega \in \Omega$ implies $f(x, \omega) \notin Q$.

Thus a solution $\varphi(\cdot, x, u)$ can leave a set K which is invariant in Q only if it also leaves Q .

For (measurable) subsets K which are invariant in Q and a quasi-stationary measure η on Q we define invariant (K, η) -partitions as in Definition 2.1 with Q replaced by K . Then one can define the invariance entropy $h_\mu(K)$ of K as in Definition 2.3, again replacing Q by K . In the following, objects associated with invariant (K, η) -partitions and invariant (Q, η) -partitions are denoted with a superscript K and Q , respectively.

First we determine relations between invariant partitions of K and Q . We call a map F on Q nonsingular with respect to η if $\eta(E) = 0$ implies $\eta(F^{-1}E) = 0$ for $E \subset Q$.

Lemma 4.2. Let K be a closed set which is invariant in a closed set $Q \subset M$.

(i) Then every invariant (Q, η) -partition $\mathcal{C}_\tau^Q = \mathcal{C}_\tau(\mathcal{P}^Q, F^Q)$ induces an invariant (K, η) -partition $\mathcal{C}_\tau^K = \mathcal{C}_\tau(\mathcal{P}^K, F^K)$ given by

$$\mathcal{P}^K := \{P \cap K \mid P \in \mathcal{P}^Q\} \text{ and } F^K(P \cap K) := F^Q(P).$$

(ii) Assume that there are a finite measurable cover of Q by sets V^1, \dots, V^N , control functions $v^1, \dots, v^N \in \mathcal{U}$ and times $\tau^1, \dots, \tau^N \in \mathbb{N}$ such that for all $j = 1, \dots, N$ and a.a. $x \in V^j$

$$\varphi(k, x, v^j) \in Q \text{ for } k = 1, \dots, \tau^j \text{ and } \varphi(\tau^j, x, v^j) \in K$$

and the maps $\varphi(\tau^j, \cdot, v^j)$ on Q are nonsingular with respect to η .

Then every invariant (K, η) -partition $\mathcal{C}_\tau^K = \mathcal{C}_\tau(\mathcal{P}^K, F^K)$ with $\tau \geq \bar{\tau} := \max_{j=1, \dots, N} \tau^j$ can be extended to an invariant (Q, η) -partition $\mathcal{C}_\tau^Q = \mathcal{C}_\tau(\mathcal{P}^Q, F^Q)$ such that $\#\mathcal{P}^Q \leq (1 + N)\#\mathcal{P}^K$, $\mathcal{P}^K \subset \mathcal{P}^Q$ and

$$F^Q(P) = F^K(P) \text{ if } P \in \mathcal{P}^K \text{ and } \varphi(\tau, P^Q, F^Q(P^Q)) \subset K \text{ for } P^Q \in \mathcal{P}^Q.$$

Proof. (i) Let $P \in \mathcal{P}^Q$. Since K is invariant in Q it follows from $P \cap K \subset K$ and $\varphi(k, x, F^Q(P)) \in Q$ for all $k = 0, \dots, \tau$ and η -a.a. $x \in P$ that

$$\varphi(k, x, F^K(P \cap K)) = \varphi(k, x, F^Q(P)) \in K \text{ for } \eta\text{-a.a. } x \in P.$$

Thus $\mathcal{C}_\tau(\mathcal{P}^K, F^K)$ is an invariant (K, η) -partition. (ii) Let $\mathcal{C}_\tau^K = \mathcal{C}_\tau(\mathcal{P}^K, F^K)$ be an invariant K -partition with $\tau \geq \bar{\tau}$. The cover of Q induces a finite partition \mathcal{P}_1 of $Q \setminus K$ such that for every $P_1^j \in \mathcal{P}_1$ the control $F_1(P_1^j) := (v_0^j, \dots, v_{\tau^j-1}^j) \in \Omega^{\tau^j}$ satisfies for a.a. $x \in P_1^j$ one has $\varphi(k, x, F_1(P_1^j)) \in Q$ for all $k = 0, \dots, \tau^j$ and $\varphi(\tau^j, x, F_1(P_1^j)) \in K$. In fact, we obtain a partition of $Q \setminus K$ by defining $P_1^1 := (Q \setminus K) \cap V^1$ and

$$P_1^j := [(Q \setminus K) \cap V^j] \setminus \bigcup_{i < j} P_1^i \text{ for } j > 1.$$

Then $P_1^j \subset V^j$ implying for a.a. $x \in P_1^j$ that $\varphi(k, x, F_1(P_1^j)) \in Q$ for $k = 1, \dots, \tau^j$ and $\varphi(\tau^j, x, F_1(P_1^j)) \in P^i \subset K$ for some $P^i \in \mathcal{P}^K$. By nonsingularity with respect

to η of $\varphi(\tau^j, \cdot, v^j) = \varphi(\tau^j, \cdot, F_1(P_1^j))$ it follows that for η -a.a. $x \in P_1^j$ there is $P^i \in \mathcal{P}^K$ such that for all $k = 0, \dots, \tau$

$$\varphi(k, \varphi(\tau^j, x, F_1(P_1^j)), F^K(P^i)) \in K.$$

Define an invariant Q -partition $\mathcal{C}_\tau^Q = \mathcal{C}_\tau(\mathcal{P}^Q, F^Q)$ in the following way: The partition consists of the sets in \mathcal{P}^K together with all (nonvoid) sets of the form

$$P^{i,j} := \left\{ x \in P_1^j \mid \varphi(\tau^j, x, F_1(P_1^j)) \in P^i \right\}$$

with feedbacks defined as follows: Let

$$F^Q(P^{i,j})_k := \begin{cases} v_k^j & \text{for } k = 0, \dots, \tau^j - 1 \\ u_{k-\tau^j}^i & \text{for } k = \tau^j, \dots, \tau - 1. \end{cases}$$

and for $P^i \in \mathcal{P}^K$ let

$$F^Q(P^i) := F^K(P^i) = (u_0^i, \dots, u_{\tau-1}^i).$$

This is well defined, since $\tau - \tau^j \geq 0$ (we use only the first part of $F^K(P^i)$) and hence $\mathcal{C}_\tau(\mathcal{P}^Q, F^Q)$ is an invariant Q -partition with $\#\mathcal{P}^Q \leq \#\mathcal{P}^K + (\#\mathcal{P}^K \cdot \#\mathcal{P}_1) \leq (1+N)\#\mathcal{P}^K$. \square

The following theorem shows when the invariance entropy of Q is already determined on an subset K that is invariant in Q .

Theorem 4.3. *Consider control system (1). Let K be a closed invariant subset in Q , fix a quasi-stationary measure η on Q for a probability measure ν on the control range Ω and let $\mu = \nu^{\mathbb{N}} \times \eta$.*

(i) *Then the invariance entropy of K is bounded above by the invariance entropy of Q , $h_\mu(K) \leq h_\mu(Q)$.*

(ii) *Suppose that there are a finite measurable cover of Q by sets V^1, \dots, V^N , control functions $v^1, \dots, v^N \in \mathcal{U}$ and times $\tau^1, \dots, \tau^N \in \mathbb{N}$ such that for all $j = 1, \dots, N$ and a.a. $x \in V^j$*

$$\varphi(k, x, v^j) \in Q \text{ for } k = 1, \dots, \tau^j \text{ and } \varphi(\tau^j, x, v^j) \in K,$$

and the maps $\varphi(\tau^j, \cdot, v^j)$ on Q are nonsingular with respect to η .

Then $h_\mu(Q) = h_\mu(K)$ follows.

(iii) *If the assumptions in (ii) are satisfied and K is the disjoint union of sets K_1, \dots, K_m which are closed and invariant in Q , then*

$$\max_i h_\mu(K_i) \leq h_\mu(Q) \leq h_\mu(K_1) + \dots + h_\mu(K_m).$$

Proof. (i) Let $\mathcal{C}_\tau^Q = \mathcal{C}_\tau(\mathcal{P}^Q, F^Q)$ be an invariant (Q, η) -partition and consider the induced invariant (K, η) -partition $\mathcal{C}_\tau^K = \mathcal{C}_\tau(\mathcal{P}^K, F^K)$ according to Lemma 4.2(i). We will show that $h_\mu(\mathcal{C}_\tau^K) \leq h_\mu(\mathcal{C}_\tau^Q)$. Then, taking first the infimum over all invariant Q -partitions \mathcal{C}_τ^Q and then over all invariant K -partitions \mathcal{C}_τ^K and, finally, the limit superior for $\tau \rightarrow \infty$, one concludes, as claimed, that $h_\mu(K) \leq h_\mu(Q)$. For $P \in \mathcal{P}^Q$ consider $v \in \mathcal{U}$ with $\varphi(i, x, v) \in Q$ for all $i = 1, \dots, \tau$ and η -a.a. $x \in P$. If x is even in $P \cap K$, then invariance of K in Q implies that $\varphi(i, x, u) \in K$ for all $i = 1, \dots, \tau$. It follows for all $P_K = P \cap K \in \mathcal{P}^K$ that

$$A(P \cap K, \mathcal{C}_\tau^K) = A(P, \mathcal{C}_\tau^Q) \cap (\mathcal{U} \times K)$$

showing that $\mathfrak{A}^K = \mathfrak{A}^Q \cap (\mathcal{U} \times K)$. For a \mathcal{C}_τ^K -admissible partition sequence corresponding to a word a abbreviate $A_{a_i}^K = A(P_{a_i} \cap K, \mathcal{C}_\tau^K)$ with $P_{a_i} \in \mathcal{P}$. Then

$(P_{a_0}, \dots, P_{a_{n-1}})$ is a \mathcal{C}_τ^Q -admissible partition sequence and hence a \mathcal{C}_τ^K -admissible word a is also \mathcal{C}_τ^Q -admissible. Consider

$$D_a^K = A_{a_0}^K \cap S^{-\tau} A_{a_1}^K \cap \dots \cap S^{-(n-1)\tau} A_{a_{n-1}}^K.$$

The corresponding \mathcal{C}_τ^Q -admissible set

$$D_a^Q = A_{a_0}^Q \cap S^{-\tau} A_{a_1}^Q \cap \dots \cap S^{-(n-1)\tau} A_{a_{n-1}}^Q$$

satisfies $D_a^Q \cap (\mathcal{U} \times K) = D_a^K$, since $(u, x) \in A_{a_0}^Q \cap (\mathcal{U} \times K)$ satisfies $(\theta^{i\tau} u, \varphi(i\tau, x, u)) \in A_{a_i}^K, i = 0, \dots, (n-1)\tau$. This shows that

$$\mathfrak{A}_n^K \subset \mathfrak{A}_n^Q \cap (\mathcal{U} \times K) \text{ for all } n, \quad (30)$$

where at the right hand side consists of the elements of \mathfrak{A}_n^Q intersected with $\mathcal{U} \times K$. In order to compute the entropy, first consider $D \in \mathfrak{A}_n^Q$ with $\rho^{-(n-1)\tau} \mu(D) \leq 1/e$. Then also $\rho^{-(n-1)\tau} \mu(D \cap (\mathcal{U} \times K)) \leq 1/e$ and it follows that

$$\phi\left(\rho^{-(n-1)\tau} \mu(D)\right) \leq \phi\left(\rho^{-(n-1)\tau} \mu(D \cap (\mathcal{U} \times K))\right), \quad (31)$$

since ϕ is monotonically decreasing on $[0, 1/e]$. For every $n \in \mathbb{N}$ there are at most three sets $D \in \mathfrak{A}_n^Q$ with $\rho^{-(n-1)\tau} \mu(D) \geq 1/e$, since they are disjoint and the sum of the measures of four mutually disjoint sets $D \subset A_n^Q \subset S_Q^{-(n-1)\tau}(\mathcal{U} \times Q)$ would be greater than or equal to $4/e > 1 = \rho^{-(n-1)\tau} \mu(S_Q^{-(n-1)\tau}(\mathcal{U} \times Q))$. Let

$$\mathfrak{A}_n^{Q, big} := \{D \in \mathfrak{A}_n^Q \mid \rho^{-(n-1)\tau} \mu(D) \geq 1/e\}.$$

Then $\#\mathfrak{A}_n^{Q, big} \leq 3$ and, using $\phi(x) \geq \phi(1/e) = -1/e, x \in [0, 1]$, it follows that

$$\sum_{D \in \mathfrak{A}_n^{Q, big}} \phi\left(\rho^{-(n-1)\tau} \mu(D \cap (\mathcal{U} \times K))\right) \geq -3/e.$$

We find, using $\phi(x) \leq 0$ and (31), (30),

$$\begin{aligned} & -H_{\rho^{-(n-1)\tau} \mu}(\mathfrak{A}_n^Q) \\ &= \sum_{D \in \mathfrak{A}_n^Q \setminus \mathfrak{A}_n^{Q, big}} \phi\left(\rho^{-(n-1)\tau} \mu(D)\right) + \sum_{D \in \mathfrak{A}_n^{Q, big}} \phi\left(\rho^{-(n-1)\tau} \mu(D)\right) \\ &\leq \sum_{D \in \mathfrak{A}_n^Q \setminus \mathfrak{A}_n^{Q, big}} \phi\left(\rho^{-(n-1)\tau} \mu(D \cap (\mathcal{U} \times K))\right) \\ &= \sum_{D \in \mathfrak{A}_n^Q} \phi\left(\rho^{-(n-1)\tau} \mu(D \cap (\mathcal{U} \times K))\right) - \sum_{D \in \mathfrak{A}_n^{Q, big}} \phi\left(\rho^{-(n-1)\tau} \mu(D \cap (\mathcal{U} \times K))\right) \\ &\leq \sum_{D^K \in \mathfrak{A}_n^K} \phi\left(\rho^{-(n-1)\tau} \mu(D^K)\right) + 3/e \\ &\leq -H_{\rho^{-(n-1)\tau} \mu}(\mathfrak{A}_n^K) + 3/e. \end{aligned}$$

This implies

$$\begin{aligned} h_\mu(\mathcal{C}_\tau^Q) &= \limsup_{n \rightarrow \infty} \frac{1}{n\tau} H_{\rho^{-(n-1)\tau} \mu}(\mathfrak{A}_n(\mathcal{C}_\tau^Q)) \geq \limsup_{n \rightarrow \infty} \frac{1}{n\tau} [H_{\rho^{-(n-1)\tau} \mu}(\mathfrak{A}_n(\mathcal{C}_\tau^K)) - 3/e] \\ &= h_\mu(\mathcal{C}_\tau^K). \end{aligned}$$

(ii) By Lemma 4.2 we can extend an invariant (K, η) -partition $\mathcal{C}_\tau^K = \mathcal{C}_\tau(\mathcal{P}^K, F^K)$ with $\tau > \bar{\tau} := \max_{j=1, \dots, N} \tau^j$ to an invariant (Q, η) -partition $\mathcal{C}_\tau^Q = \mathcal{C}_\tau(\mathcal{P}^Q, F^Q)$.

We claim that $h_\mu(\mathcal{C}_\tau^Q) \leq h_\mu(\mathcal{C}_\tau^K)$. Then assertion (ii) will follow, if we take first the infimum over all invariant (K, η) -partitions \mathcal{C}_τ^K , then the infimum over all invariant (Q, η) -partitions \mathcal{C}_τ^Q and, finally, let $\tau \rightarrow \infty$. Recall from Lemma 4.2 that $\mathcal{P}^K \subset \mathcal{P}^Q$, $F^K(P) = F^Q(P)$ for $P \in \mathcal{P}^K$ and $\varphi(\tau, x, F^Q(P)) \in K$ for η -a.a. $x \in P$ and all $P \in \mathcal{P}^Q$. It follows for all $P \in \mathcal{P}^K \subset \mathcal{P}^Q$ that

$$A^K(P) = A^Q(P) \text{ and hence } \mathfrak{A}^K := \mathfrak{A}(\mathcal{C}_\tau^K) \subset \mathfrak{A}^Q := \mathfrak{A}(\mathcal{C}_\tau^Q).$$

Let (P_0, \dots, P_{n-1}) be a \mathcal{C}_τ^Q -admissible partition sequence in \mathcal{P}^Q . Since K is invariant in Q , it follows that $P_i \subset K$ for all $i = 1, \dots, n-1$, and hence for every $k \geq \tau$ the controls are given by feedbacks F^K keeping the system in K . Hence (P_1, \dots, P_{n-1}) is a \mathcal{C}^K -admissible sequence in \mathcal{P}^K . Together, this implies for a \mathcal{C}_τ^Q -admissible word $a^{n+1} = [a_0, a_1, \dots, a_n]$ of length $n+1$ and for the elements

$$D_{a^{n+1}} = A_{a_0} \cap S^{-\tau} A_{a_1} \cap \dots \cap S^{-n\tau} A_{a_n} \in \mathfrak{A}_{n+1}^Q,$$

that $a^n := [a_1, \dots, a_n]$ is a \mathcal{C}^K -admissible word with

$$D_{a^n} = A_{a_1} \cap \dots \cap S^{-(n-1)\tau} A_{a_n} \in \mathfrak{A}_n^K.$$

Hence $D_{a^{n+1}} = A_{a_0} \cap S^{-\tau} D_{a^n}$ with $A_{a_0} \in \mathfrak{A}^Q$ and we find

$$\mu(D_{a^{n+1}}) = \mu(A_{a_0} \cap S^{-\tau} D_{a^n}) \leq \mu(S^{-\tau} D_{a^n}) = \rho^\tau \mu(D_{a^n}). \quad (32)$$

Define

$$\mathfrak{A}_n^{K,big} := \{D_{a^n} \in \mathfrak{A}_n^K \mid \rho^{-(n-1)\tau} \mu(D_{a^n}) \geq 1/e\},$$

$$\mathfrak{A}_{n+1}^{Q,big} := \{D_{a^{n+1}} = A_{a_0} \cap S^{-\tau} D_{a^n} \in \mathfrak{A}_{n+1}^Q \mid A_{a_0} \in \mathfrak{A}^Q \text{ and } D_{a^n} \in \mathfrak{A}_n^{K,big}\}.$$

Then, as above, $\#\mathfrak{A}_n^{K,big} \leq 3$. Furthermore, by Lemma 4.2, the number of elements of $\mathfrak{A}_{n+1}^{Q,big}$ is bounded, independently of n , by

$$\#\mathfrak{A}_{n+1}^{Q,big} \leq 3 \cdot \#\mathfrak{A}^Q \leq 3(N+1) \cdot \#\mathcal{P}^K.$$

For $D_{a^{n+1}} = A_{a_0} \cap S^{-\tau} D_{a^n} \in \mathfrak{A}_{n+1}^Q$ with $D_{a^n} \in \mathfrak{A}_n^K \setminus \mathfrak{A}_n^{K,big}$, it follows from (32) that

$$\rho^{-n\tau} \mu(D_{a^{n+1}}) \leq \rho^{-n\tau} \rho^\tau \mu(D_{a^n}) = \rho^{-(n-1)\tau} \mu(D_{a^n}) \leq 1/e,$$

and hence, using monotonicity of ϕ on $[0, 1/e]$, for all $D_{a^{n+1}} \in \mathfrak{A}_{n+1}^Q \setminus \mathfrak{A}_{n+1}^{Q,big}$

$$\phi(\rho^{-n\tau} \mu(D_{a^{n+1}})) \geq \phi(\rho^{-(n-1)\tau} \mu(D_{a^n})).$$

It follows that

$$\begin{aligned} H_{\rho^{-n\tau}\mu}(\mathfrak{A}_{n+1}^Q) &= - \sum_{D \in \mathfrak{A}_{n+1}^Q} \phi(\rho^{-n\tau} \mu(D)) \\ &\leq - \sum_{D \in \mathfrak{A}_{n+1}^Q \setminus \mathfrak{A}_{n+1}^{Q,big}} \phi(\rho^{-n\tau} \mu(D)) + 3(N+1) \cdot \#\mathcal{P}^K/e \\ &\leq - \sum_{D \in \mathfrak{A}_n^K \setminus \mathfrak{A}_n^{K,big}} \phi(\rho^{-(n-1)\tau} \mu(D)) + 3(N+1) \cdot \#\mathcal{P}^K/e \\ &\leq - \sum_{D \in \mathfrak{A}_n^K} \phi(\rho^{-(n-1)\tau} \mu(D)) + 3(N+1) \cdot \#\mathcal{P}^K/e \\ &= H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n^K) + 3(N+1) \cdot \#\mathcal{P}^K/e. \end{aligned}$$

We conclude that, as claimed,

$$\begin{aligned} h_\mu(\mathcal{C}_\tau^Q) &= \limsup_{n \rightarrow \infty} \frac{1}{n\tau} H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau^Q)) \\ &= \limsup_{n \rightarrow \infty} \frac{1}{n\tau} H_{\rho^{-n\tau}\mu}(\mathfrak{A}_{n+1}(\mathcal{C}_\tau^Q)) \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{n\tau} [H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau^K)) + 3(N+1) \cdot \#\mathcal{P}^K/e] = h_\mu(\mathcal{C}_\tau^K). \end{aligned}$$

(iii) The first assertion follows by (i) using that each K_i is invariant in Q . For the second inequality use that by (ii) $h_\mu(Q) = h_\mu(K)$. Take invariant partitions $\mathcal{C}_\tau^{K_i} = \mathcal{C}_\tau(\mathcal{P}^i, F^i)$ of K_i for each i . Then $\mathcal{C}_\tau^K = \mathcal{C}_\tau(\mathcal{P}^K, F^K)$ defined by $\mathcal{P}^K = \bigcup \mathcal{P}^i, F_{\mathcal{P}^i}^K = F_i, i = 1, \dots, m$, forms an invariant partition of K and one finds

$$\mathfrak{A}_n^K(\mathcal{C}_\tau) = \bigcup_{i=1}^m \mathfrak{A}_n^{K_i}(\mathcal{C}_\tau) \text{ and } H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau^K)) = \sum_{i=1}^m H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau^{K_i})).$$

This implies that

$$\begin{aligned} h_\mu(\mathcal{C}_\tau^K) &\leq \limsup_{n \rightarrow \infty} \frac{1}{n\tau} H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau^K)) \\ &= \limsup_{n \rightarrow \infty} \frac{1}{n\tau} \sum_{i=1}^m H_{\rho^{-(n-1)\tau}\mu}(\mathfrak{A}_n(\mathcal{C}_\tau^{K_i})) \leq \sum_{i=1}^m h_\mu(\mathfrak{A}_n(\mathcal{C}_\tau^{K_i})). \end{aligned}$$

Now take the infimum over all invariant partitions $\mathcal{C}_\tau^{K_j}$ of the K_j , then over all invariant partitions of K and, finally, the limit superior for $\tau \rightarrow \infty$. \square

Remark 10. If the assumptions of Theorem 4.3(ii) are satisfied, one might conjecture that the support of any quasi-stationary measure is contained in K . However, this cannot be expected as seen from the description of all quasi-stationary measures for finite state spaces given in van Doorn and Pollett [13, Theorem 4.2]. A simple example is given in Benaïm, Cloez and Panloup [3, Example 3.5].

For a quasi-stationary measure η and a closed invariant subset Q_1 in Q with $\eta(Q_1) > 0$ let $\mu_{Q_1} := \nu^{\mathbb{N}} \times \eta_{Q_1}$, where $\eta_{Q_1}(\cdot) = \eta(\cdot \cap Q_1)/\eta(Q_1)$ is the conditional measure on Q_1 .

Corollary 1. Consider control system (1) and fix a quasi-stationary measure η on Q for a probability measure ν on the control range Ω and let $\mu = \nu^{\mathbb{N}} \times \eta$. Suppose that Q is the disjoint union of closed pairwise disjoint sets $Q_i, i = 1, \dots, m$, which are invariant in Q with $\eta(Q_i) > 0$ for all i . Then all η_{Q_i} are quasi-stationary for Q_i with the same constant as η and

$$\max_{i=1, \dots, m} \{\eta(Q_i) h_{\mu_{Q_i}}(Q_1)\} \leq h_\mu(Q) \leq \eta(Q_1) h_{\mu_{Q_1}}(Q_1) + \dots + \eta(Q_m) h_{\mu_{Q_m}}(Q_m). \quad (33)$$

Proof. Abbreviate $\eta_i := \eta_{Q_i}$ for all i . One has the convex combination $\eta = \eta(Q_1)\eta_1 + \dots + \eta(Q_m)\eta_m$ implying for all $k \in \mathbb{N}$

$$\rho^{-k}\eta = \eta(Q_1)\rho^{-k}\eta_1 + \dots + \eta(Q_m)\rho^{-k}\eta_m(Q_m).$$

For $A \subset Q_i$, invariance of $Q_j, j \neq i$, in Q implies

$$\rho\eta_i(A) = \rho \frac{\eta(A)}{\eta(Q_i)} = \frac{1}{\eta(Q_i)} \int_Q p(x, A) \eta(dx) = \frac{1}{\eta(Q_i)} \int_{Q_i} p(x, A) \eta(dx)$$

$$= \int_{Q_i} p(x, A) \eta_i(dx).$$

Hence η_i is quasi-stationary for Q_i with the same constant ρ as η . Then one easily sees that $h_\mu(Q_i) = \eta(Q_i)h_{\mu_i}(Q_i)$. Now the first inequality in (33) follows from Theorem 4.3(i) and the second inequality follows from the arguments in the proof of 4.3(iii) if one notes that here $K = Q$. \square

For the incremental invariance entropy we can only show the analog of Theorem 4.3(i). Here the arguments are a bit more involved.

Theorem 4.4. *Consider control system (1). Let K be a closed invariant subset in Q , fix a quasi-stationary measure η on Q for a probability measure ν on the control range Ω and let $\mu = \nu^\Omega \times \eta$.*

Then the incremental invariance entropy of K is bounded above by the incremental invariance entropy of Q , $h_\mu^{\text{inc}}(K) \leq h_\mu^{\text{inc}}(Q)$.

Proof. Let $\mathcal{C}_\tau^Q = \mathcal{C}_\tau(\mathcal{P}^Q, F^Q)$ be an invariant (Q, η) -partition and consider the induced invariant (K, η) -partition $\mathcal{C}_\tau^K = \mathcal{C}_\tau(\mathcal{P}^K, F^K)$ according to Lemma 4.2. We will show that $h_\mu^{\text{inc}}(\mathcal{C}_\tau^K) \leq h_\mu^{\text{inc}}(\mathcal{C}_\tau^Q)$. Then the assertion will follow. The arguments used to prove formula (30) also show that for a set $D \cap (\mathcal{U} \times K) \in \mathfrak{A}_n^K = \mathfrak{A}_n(\mathcal{C}_\tau^K)$ one obtains (cf. (16))

$$\mathfrak{A}_{n+1}^K(D \cap (\mathcal{U} \times K)) \subset \mathfrak{A}_{n+1}^Q(D) \cap (\mathcal{U} \times K),$$

and hence the unions satisfy

$$\mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K)) \subset \mathcal{A}_{n+1}^Q(D) \cap (\mathcal{U} \times K) \subset \mathcal{A}_{n+1}^Q(D).$$

Next we consider the conditional entropy (cf. (18)),

$$\begin{aligned} & -H_{\rho^{-n}\tau_\mu}(\mathfrak{A}_{n+1}^K | \mathfrak{A}_n^{K, n+1}) \\ &= \rho^{-n\tau} \sum_{D \cap (\mathcal{U} \times K) \in \mathfrak{A}_n^{K, n+1}} \mu(\mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K))) \sum_{E \in \mathfrak{A}_{n+1}^K} \phi \left(\frac{\mu(D \cap (\mathcal{U} \times K) \cap E)}{\mu(\mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K)))} \right). \end{aligned} \quad (34)$$

Fix an element $D \cap (\mathcal{U} \times K) \in \mathfrak{A}_n^K$ and let

$$\begin{aligned} \alpha_1 &:= \mu(\mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K))), \quad \alpha_2 := \mu(\mathcal{A}_{n+1}^Q(D) \setminus \mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K))), \\ \alpha &:= \alpha_1 + \alpha_2 = \mu(\mathcal{A}_{n+1}^Q(D)). \end{aligned}$$

Observe that convexity of ϕ implies

$$\phi \left(\frac{\alpha_1}{\alpha} \mu_1 + \frac{\alpha_2}{\alpha} \mu_2 \right) \leq \frac{\alpha_1}{\alpha} \phi(\mu_1) + \frac{\alpha_2}{\alpha} \phi(\mu_2) \text{ for } \mu_1, \mu_2 \in [0, 1].$$

This, together with $\phi(x) \leq 0$, shows that for $E \in \mathfrak{A}_{n+1}^K$

$$\begin{aligned} & \frac{\alpha_1}{\alpha} \phi \left(\frac{\mu(D \cap (\mathcal{U} \times K) \cap E)}{\mu(\mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K)))} \right) \\ & \geq \frac{\alpha_1}{\alpha} \phi \left(\frac{\mu(D \cap (\mathcal{U} \times K) \cap E)}{\mu(\mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K)))} \right) + \frac{\alpha_2}{\alpha} \phi \left(\frac{\mu(D \cap (\mathcal{U} \times (Q \setminus K)) \cap E)}{\mu(\mathcal{A}_{n+1}^Q(D) \setminus \mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K)))} \right) \\ & \geq \phi \left(\frac{\alpha_1}{\alpha} \frac{\mu(D \cap (\mathcal{U} \times K) \cap E)}{\mu(\mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K)))} + \frac{\alpha_2}{\alpha} \frac{\mu(D \cap (\mathcal{U} \times (Q \setminus K)) \cap E)}{\mu(\mathcal{A}_{n+1}^Q(D) \setminus \mathcal{A}_{n+1}^K(D \cap (\mathcal{U} \times K)))} \right) \end{aligned}$$

$$= \phi \left(\frac{\mu(D \cap (\mathcal{U} \times K) \cap E)}{\mu(\mathcal{A}_{n+1}^Q(D))} + \frac{\mu(D \cap (\mathcal{U} \times (Q \setminus K)) \cap E)}{\mu(\mathcal{A}_{n+1}^Q(D))} \right) = \phi \left(\frac{\mu(D \cap E)}{\mu(\mathcal{A}_{n+1}^Q(D))} \right).$$

With (34) it follows that

$$-H_{\rho^{-n\tau}\mu}(\mathfrak{A}_{n+1}^K | \mathfrak{A}_n^K) \geq \rho^{-n\tau} \sum_{D \cap (\mathcal{U} \times K) \in \mathfrak{A}_n^K} \mu(\mathcal{A}_{n+1}^Q(D)) \sum_{E \in \mathfrak{A}_{n+1}^Q} \phi \left(\frac{\mu(D \cap E)}{\mu(\mathcal{A}_{n+1}^Q(D))} \right).$$

We may add further negative summands (corresponding to elements $D \in \mathfrak{A}_n^Q \setminus \mathfrak{A}_n^K$ and $E \in \mathfrak{A}_{n+1}^Q \setminus \mathfrak{A}_{n+1}^K$) and estimate this by

$$\geq \rho^{-n\tau} \sum_{D \in \mathfrak{A}_n^Q} \mu(\mathcal{A}_{n+1}^Q(D)) \sum_{E \in \mathfrak{A}_{n+1}^Q} \phi \left(\frac{\mu(D \cap E)}{\mu(\mathcal{A}_{n+1}^Q(D))} \right) = -H_{\rho^{-n\tau}\mu}(\mathfrak{A}_{n+1}^Q | \mathfrak{A}_n^{Q,n+1}).$$

Next take the sum over the conditional entropies as specified in Definition 2.4, divide by n and take the limit for $n \rightarrow \infty$. This shows, as claimed, that $h_\mu^{inc}(\mathcal{C}_\tau^K) \leq h_\mu^{inc}(\mathcal{C}_\tau^Q)$. Now take the infimum over all invariant (Q, η) -partitions \mathcal{C}_τ^Q , the infimum over all invariant (K, η) -partitions \mathcal{C}_τ^K and, finally, let $\tau \rightarrow \infty$. This concludes the proof. \square

5. Invariant W -control sets. In this section, we will describe certain subsets of complete approximate controllability within a subset of the state space called control sets. They will yield a relatively invariant set K as considered in Theorems 4.3 and Theorem 4.4.

For systems in discrete time, subsets of the state space where approximate or exact controllability holds, have been analyzed in diverse settings. Relevant contributions are due, in particular, to Albertini and Sontag [1, 2], Sontag and Wirth [23], Wirth [26, 27] as well as Patrão and San Martin [20] (there are subtle differences in the definitions).

We recall the following notions and facts from the abstract framework in [20], slightly modified for our purposes. A local semigroup \mathcal{S} on a topological space X is a family of continuous maps $\phi : \text{dom}\phi \rightarrow X$ with open domain $\text{dom}\phi \subset X$ such that for all $\phi, \psi \in \mathcal{S}$ with $\psi^{-1}(\text{dom}\phi) \neq \emptyset$ it follows that $\phi \circ \psi : \psi^{-1}(\text{dom}\phi) \rightarrow X$ also is in \mathcal{S} .

For $x \in X$ the orbit is $\mathcal{S}x = \{\phi(x) | \phi \in \mathcal{S} \text{ and } x \in \text{dom}\phi\}$ and the backward orbit is

$$\mathcal{S}^*x = \{y \in X | \exists \phi \in \mathcal{S} : \phi(y) = x\} = \bigcup_{\phi \in \mathcal{S}} \phi^{-1}(x). \quad (35)$$

A local semigroup \mathcal{S} is called accessible if $\text{int}(\mathcal{S}x) \neq \emptyset$ and $\text{int}(\mathcal{S}^*x) \neq \emptyset$ for all $x \in X$.

Definition 5.1. A control set for a local semigroup \mathcal{S} on X is a nonvoid subset $D \subset X$ such that (i) $y \in \text{cl}(\mathcal{S}x)$ for all $x, y \in D$ (ii) for every $x \in D$ there are $\phi_n \in \mathcal{S}, n \in \mathbb{N}$, such that $\phi_n \circ \dots \circ \phi_1(x) \in D$ for all $n \in \mathbb{N}$ and (iii) the set D is maximal with this property.

The transitivity set D_0 of D is the set of all elements $x \in D$ such that $x \in \text{int}(\mathcal{S}^*x)$.

Remark 11. Patrão and San Martin [20] define control sets in the following slightly different way: A control set for \mathcal{S} is a subset $D \subset X$ such that $y \in \text{cl}(\mathcal{S}x)$ for all $x, y \in D$, the set D is maximal with this property, and there is $x \in D$ with

$x \in \text{int}(S^*x)$. The latter condition means that the transitivity set is nonvoid. Hence a control set as defined above with nonvoid transitivity set is a control set in the sense of [20]. Conversely, for an accessible semigroup Patrão and San Martin show that a control set D in their sense has the following properties:

The transitivity set D_0 is open and dense in D and invariant in D , i.e., $SD_0 \cap D \subset D_0$ (cf. [20, Proposition 4.10]) and, by [20, Proposition 4.8],

$$D = \text{cl}(Sx) \cap S^*x \text{ for } x \in D_0. \quad (36)$$

It follows that property (ii) in Definition 5.1 is satisfied. In fact, this is clear for $x \in D_0$ and for an arbitrary point x in D there is $\phi \in S$ with $\phi(x) \in D_0$, since D_0 is open. Thus for an accessible semigroup the control sets as defined above with nonvoid transitivity set coincide with the control sets in the sense of [20].

The following result is Patrão and San Martin [20, Proposition 4.15].

Proposition 2. *Let D be a control set with nonvoid transitivity set D_0 for an accessible local semigroup S . Then the transitivity set D_0 is open and dense in D and the following statements are equivalent:*

- (i) $\text{cl}(Sx) \subset \text{cl}D$ for all $x \in D$.
- (ii) D is closed and S -invariant, i.e., $Sx \in D$ for all $x \in D$.
- (iii) $\text{cl}D$ is S -invariant.

Next we use these concepts in our context. Again we consider control system (1), but now we will restrict the state space to an open subset of the state space M . For $\omega \in \Omega$ let $f_\omega := f(\cdot, \omega) : M \rightarrow M$. Then the solutions $\varphi(k, x, u)$, $u = (\omega_i)$ can be written in the form

$$\varphi(k, x, u) = f_{\omega_{k-1}} \circ \cdots \circ f_{\omega_0}(x).$$

Let W be an open nonvoid subset of the state space M . The maps f_ω , $\omega \in \Omega$, generate the following family of continuous maps on W :

$$f_\omega^0(x) := f_\omega(x) : \text{dom} f_\omega^0 \rightarrow W, \text{dom} f_\omega^0 := \{x \in W \mid f(x, \omega) \in W\}$$

and for $k \geq 1$ and $u = (\omega_0, \dots, \omega_{k-1}) \in \Omega^k$

$$f_u^k(x) := f_{\omega_{k-1}} \circ \cdots \circ f_{\omega_0}(x) : \text{dom} f_u^k \rightarrow W,$$

$$\text{dom} f_u^k := \{x \in W \mid f_{\omega_{i-1}} \circ \cdots \circ f_{\omega_0}(x) \in W \text{ for } i = 1, \dots, k-1\}.$$

Observe that the domain of f_u^k is open and the maps f_ω are continuous. Hence it follows that

$$S := \{f_u^k \mid k \in \mathbb{N}, u \in \mathcal{U}\} \quad (37)$$

forms a local semigroup on $X := W$.

For $x \in W$, $u \in \mathcal{U}$ and $k \in \mathbb{N}$ the corresponding solution of (1) in W is denoted by $\varphi_W(k, x, u)$, if the solution of (1) satisfies $\varphi(i, x, u) \in W$ for $i = 1, \dots, k$. Thus $\varphi_W(k, x, u) = f_u^k(x)$.

Definition 5.2. For $x \in M$ the W -reachable set $\mathbf{R}^W(x)$ and the W -controllable set $\mathbf{C}^W(x)$, resp., are

$$\mathbf{R}^W(x) := \{y \in W \mid \exists k \geq 1 \exists u \in \mathcal{U} : y = \varphi_W(k, x, u)\},$$

$$\mathbf{C}^W(x) := \{y \in W \mid \exists u \in \mathcal{U} \exists k \geq 1 : \varphi_W(k, y, u) = x\}.$$

Note that in the language of local semigroups one has $\mathbf{R}^W(x) = Sx$ and $\mathbf{C}^W(x) = S^*x$. Next we specify maximal subsets of complete approximate controllability within W .

Definition 5.3. For system (1) a subset $D \subset W$ is called a W -control set if (i) $D \subset \text{cl}_W \mathbf{R}^W(x)$ for all $x \in D$, (ii) for every $x \in D$ there is $u \in \mathcal{U}$ with $\varphi_W(k, x, u) \in D, k \in \mathbb{N}$, and (iii) D is a maximal set with properties (i) and (ii). A W -control set D is called an invariant W -control set if $\mathbf{R}^W(x) \subset \text{cl}_W D$ for all $x \in D$.

Here the closures are taken with respect to W , and for an invariant W -control set $\text{cl}_W \mathbf{R}^W(x) = \text{cl}_W D$ for all $x \in D$. If $W = M$, we omit the index W and just speak of control sets and invariant control sets (if they have nonvoid interior they actually coincide with the control sets and invariant control sets, respectively, considered in Colonius, Homburg, Kliemann [8]). The transitivity set D_0 of a W -control set D is the set of all $x \in D$ with $x \in \text{int} S^* x = \text{int} C^W(x)$.

It is immediate that the W -control sets coincide with the control sets for the local semigroup \mathcal{S} on $X = W$ defined in (37). Accessibility of the considered local semigroup \mathcal{S} means that

$$\text{int} \mathbf{R}^W(x) \neq \emptyset \text{ and } \text{int} C^W(x) \neq \emptyset \text{ for all } x \in W. \quad (38)$$

This is certainly valid if

$$\text{int} f(x, \Omega) \cap W \neq \emptyset \text{ and } \text{int} \{y \in W \mid \exists \omega \in \Omega : f(y, \omega) = x\} \neq \emptyset \text{ for all } x \in W. \quad (39)$$

If accessibility holds, Proposition 2 shows that a W -control set D with nonvoid transitivity set D_0 is an invariant W -control set if and only if $\text{cl}_W D = D$ and $\mathbf{R}^W(x) \subset D$ for all $x \in D$ if and only if $\mathbf{R}^W(x) \subset \text{cl}_W D$ for all $x \in \text{cl}_W D$. In particular, an invariant W -control set is an invariant set in W .

It is also of interest to know when closedness in W of a W -control set already implies that it is an invariant W -control set. The proof of the following proposition is adapted from Wirth [26, Proposition 4.1.4].

Proposition 3. Suppose that control system (1) satisfies accessibility condition (38).

(i) Then every invariant W -control set D is closed in W and has nonvoid interior.

(ii) If for every $x \in W$ the set $\text{cl}_W [f(x, \Omega) \cap W]$ is path connected, then a W -control set D which is closed in W and has nonvoid interior is an invariant W -control set.

Proof. (i) By the previous remarks the set D is closed and has nonvoid interior since $\emptyset \neq \text{int} \mathbf{R}^W(y) \subset D$. (ii) If $D = W$, there is nothing to prove. Otherwise we have to show for every $x \in D$ that $\text{cl}_W \mathbf{R}^W(x) \subset \text{cl}_W D$ or, equivalently, that $\mathbf{R}^W(x) \subset D$, since D is closed in W . For every $y \in D$ there are $k \in \mathbb{N}$ and a control u such that $\varphi_W(k, y, u) \in \text{int} D$. By continuous dependence on initial values there exists an open neighborhood $V(y)$ of y with $\varphi_W(k, V(y), u) \subset \text{int} D$. Taking the union of all $V(y)$ one finds an open set $V \supset D$ such that for every $z \in V$ the intersection $\mathbf{R}^W(z) \cap \text{int} D \neq \emptyset$ and therefore $D \subset \text{cl}_W \mathbf{R}^W(z)$. Assume now, contrary to the assertion, that there exist $x \in D$ and a control value ω such that $f(x, \omega) \in W \setminus D$. As $D \subset \text{cl}_W \mathbf{R}^W(x)$ there exists $y \in \mathbf{R}^W(x) \cap D$ and hence there is $\omega \in \Omega$ with $f(x, \omega) \in D$, by maximality of W -control sets. We have shown that $\text{cl}_W [f(x, \Omega) \cap D] \neq \emptyset$ and $\text{cl}_W [f(x, \Omega) \cap W] \not\subset D$. Since $\text{cl}_W [f(x, \Omega) \cap W]$ is path connected by assumption, it follows that there exists $z \in \text{cl}_W [f(x, \Omega) \cap W] \cap (V \setminus D)$. By continuity, this implies that $z \in \text{cl}_W \mathbf{R}^W(y)$ for all $y \in D$ and, by construction of V , one has $D \subset \text{cl}_W \mathbf{R}^W(z)$ and thus $z \in D$ by the maximality property of control sets. This is a contradiction. \square

The following result constructs a set K which is invariant in Q and satisfies the assumptions of Theorem 4.3, hence it determines the invariance entropy.

Theorem 5.4. *Suppose that control system (1) satisfies accessibility condition (38) and let $Q \subset M$ be compact and equal to the closure of its interior $W := \text{int}Q$. Furthermore, assume*

(i) *there are only finitely many invariant W -control sets D_1, \dots, D_ℓ and their transitivity sets are nonvoid.*

(ii) *For every $x \in Q$ there is an invariant W -control set $D_i \subset \text{cl}\mathbf{R}^W(x)$.*

(iii) *Let $K := \bigcup_{i=1}^\ell \text{cl}D_i$, suppose that this union is disjoint and that $f(K, \Omega) \cap (\partial Q \setminus K) = \emptyset$.*

Then every set $\text{cl}D_i$ and hence K is invariant in Q .

If η is a quasi-stationary measure for ν on Ω and the maps $f(\cdot, \omega), \omega \in \Omega$, are nonsingular with respect to η , then the invariance entropies with respect to $\mu = \nu^N \times \eta$ of K and Q coincide with

$$\max_{i \in \{1, \dots, \ell\}} h_\mu(\text{cl}D_i) \leq h_\mu(K) = h_\mu(Q) \leq \sum_{i=1}^\ell h_\mu(\text{cl}D_i). \quad (40)$$

Proof. The assumption implies that the boundaries of Q and W coincide, $\partial Q = \partial W$. First we show that for every invariant W -control set D its closure $\text{cl}D$ is invariant in the set Q . This also implies that K is invariant in Q . By Proposition 2 D is closed in W and hence $\text{cl}D = D \cup (\partial D \cap \partial Q)$. Suppose, contrary to the assertion, that there are $x \in \text{cl}D$ and $\omega \in \Omega$ with $f(x, \omega) \in Q \setminus \text{cl}D$. If $x \in D$ then invariance in W of $D = \text{cl}_W D$ implies that $\emptyset = f(x, \Omega) \cap (W \setminus D) = f(x, \Omega) \cap (W \setminus \text{cl}D)$. Since $\partial Q = \partial W$ it follows that $f(x, \omega) \in \partial Q \setminus \text{cl}D$. But assumption (iii) excludes this case. It remains to discuss the case $x \in \partial D \cap \partial Q$. Then either $f(x, \omega) \in W \setminus D$ or $f(x, \omega) \in \partial Q \setminus D$. In the first case, Proposition 2 implies $D = \text{cl}(D_0)$ and hence continuity of f implies that there is $y \in D_0$ with $f(y, \omega) \in W \setminus D$. This is excluded since D is invariant in W . The second case $f(x, \omega) \in \partial Q \setminus D$ is excluded by assumption (iii) and it follows that $\text{cl}D$ is invariant in Q . Assumptions (i) and (ii) show that for every $x \in Q$ there are an invariant W -control set D_i , a natural number k_0 and a control u such that $\varphi(k_0, x, u)$ is in the transitivity set D_0 of D_i . Since D_0 is open, continuity with respect to x and compactness of Q imply that the assumptions of Theorem 4.3(ii) are satisfied and it follows that $h_\mu(Q) = h_\mu(K)$. The inequalities in (40) follow by invariance of $\text{cl}D_i$ in Q from Theorem 4.3(iii). \square

Next we discuss when the assumptions of Theorem 5.4 are satisfied. The following theorem characterizes the existence of finitely many invariant W -control sets.

Theorem 5.5. *Consider a control system of the form (1) satisfying accessibility condition (38).*

(i) *Let $x \in Q$ and assume that there exists a compact set $F \subset W$ such that for all $y \in \mathbf{R}^W(x)$ one has $\text{cl}\mathbf{R}^W(y) \cap F \neq \emptyset$. Then there exists an invariant W -control set $D \subset \text{cl}_W \mathbf{R}^W(x)$.*

(ii) *If there is a compact set $F \subset W$ such that $\text{cl}\mathbf{R}^W(x) \cap F \neq \emptyset$ for all $x \in Q$, then for every $x \in Q$ there is an invariant W -control set D with $D \subset \text{cl}\mathbf{R}^W(x)$ and there are only finitely many invariant W -control sets D_1, \dots, D_ℓ .*

(iii) *Conversely, suppose that for every $x \in Q$ there is an invariant W -control set D with $D \subset \text{cl}\mathbf{R}^W(x)$ and there are only finitely many invariant W -control sets D_1, \dots, D_ℓ and they all have nonvoid transitivity set. Then there is a compact set $F \subset W$ such that $\mathbf{R}^W(x) \cap F \neq \emptyset$ for all $x \in Q$.*

Proof. (i) For $y \in \mathbf{R}^W(x)$ let $F(y) := \text{cl}\mathbf{R}^W(y) \cap F$. Consider the family of nonvoid and compact subsets of W given by $\mathcal{F} = \{F(y) \mid y \in F(x)\}$. Then \mathcal{F} is ordered via

$$F(y) \preceq F(z) \text{ if } z \in \text{cl}\mathbf{R}^W(y).$$

Every linearly ordered subset $\{F(y_i), i \in I\}$ has an upper bound

$$F(y) = \bigcap_{i \in I} F(y_i) \text{ for some } y \in \bigcap_{i \in I} F(y_i),$$

since the intersection of decreasing compact subsets of the compact set F is nonvoid. Thus Zorn's lemma implies that the family \mathcal{F} has a maximal element $F(y)$. Now the set

$$D := \text{cl}_W \mathbf{R}^W(y)$$

is an invariant W -control set. Note first that by condition (38) the set D has nonvoid interior. Every $z \in D$ is in $\text{cl}_W \mathbf{R}^W(y)$ and, conversely, $y \in \text{cl}_W \mathbf{R}^W(z)$ for every $z \in D$ since otherwise $y \notin F(z) = \text{cl}\mathbf{R}^W(z) \cap F \subset \text{cl}_W \mathbf{R}^W(y) \cap F = F(y)$, hence $F(y) \preceq F(z)$ and $F(y) \neq F(z)$ contradicting the maximality of $F(y)$. Continuity implies that for all $z_1, z_2 \in \text{cl}_W \mathbf{R}^W(y)$ one has $z_2 \in \text{cl}_W \mathbf{R}^W(z_1)$, hence $D = \text{cl}_W \mathbf{R}^W(y)$ is a W -control set. It is an invariant D -control set since for $z \in D = \text{cl}_W \mathbf{R}^W(y)$ continuity implies $\mathbf{R}^W(z) \subset \text{cl}_W \mathbf{R}^W(y) = D$. (ii) Let $x \in Q$. Then, by (i), one finds an invariant W -control set in $\text{cl}\mathbf{R}^W(x)$. Suppose that there are countably many pairwise different invariant W -control sets $D_n, n \in \mathbb{N}$. Thus $\text{cl}\mathbf{R}^W(y) = \text{cl}D_n \subset \text{cl}_W D_n \cup \partial Q$ for all $y \in D_n$. Since D_n is closed in $W = \text{int}Q$ and $F \subset W$, the property $\text{cl}\mathbf{R}^W(y) \cap F \neq \emptyset$ implies that $D_n \cap F \neq \emptyset$. There are points $y_n \in D_n \cap F$ converging to some point $y \in F$. By part (i) one finds an invariant W -control set D contained in $\text{cl}_W \mathbf{R}^W(y)$, and hence there is a point z in the intersection of $\mathbf{R}^W(y)$ and the interior of D . Now continuity implies that for every n large enough there is a point z_n in D_n with $\mathbf{R}^W(z_n) \cap \text{int}D \neq \emptyset$. This contradicts invariance in W of the W -control sets D_n . (iii) Choose for each of the finitely many invariant W -control sets $D_i, i = 1, \dots, \ell$, a point $x_i \in D_i$ and define $F := \{x_1, \dots, x_\ell\}$. Let $x \in Q$. By assumption, there is an invariant W -control set D_i contained in $\text{cl}\mathbf{R}^W(x)$. Hence there is a point in the intersection of $\mathbf{R}^W(x)$ and the transitivity set of D_i , thus (36) implies that $x_i \in \mathbf{R}^W(x)$ and the assertion follows. \square

It remains to discuss when an invariant W -control set has a nonvoid transitivity set. Obviously, this holds if for the local semigroup \mathcal{S} one has that $\mathcal{S}x$ and \mathcal{S}^*x are open, cf. San Martin and Patrão [20, Corollary 5.4] for a situation where this occurs. Instead of this strong assumption, we will require smoothness of f and use Sard's Theorem as well as some arguments from Wirth [27].

Consider a control system of the form

$$x_{k+1} = f(x_k, u_k), k \in \mathbb{N} = \{0, 1, \dots\}, \quad (41)$$

under the following assumptions: The state space M is a C^∞ -manifold of dimension d endowed with a corresponding metric. The set of control values $\Omega \subset \mathbb{R}^m$ satisfies $\Omega \subset \text{cl}(\text{int}\Omega)$. Let $\tilde{\Omega}$ be an open set containing $\text{cl}\Omega$. The map $f : M \times \tilde{\Omega} \rightarrow M$ is a C^∞ -map and $W \subset M$ is a nonvoid open subset.

We define for $k \geq 1$ a C^∞ -map

$$F_k : W \times \text{int}\Omega^k \rightarrow W, F_k(x, u) := \varphi_W(k, x, u).$$

The domain of F_k is an open subset of $W \times \Omega^k$.

A pair $(x, u) \in W \times \text{int}\Omega^k$ is called regular, if $\text{rank} \frac{\partial F_k}{\partial u}(x, u) = d$ (clearly, this implies $mk \geq d$). For $x \in M$ and $k \in \mathbb{N}$ the regular W -reachability set and the regular W -controllability set, resp., are

$$\begin{aligned}\hat{\mathbf{R}}_k^W(x) &:= \{y \in W \mid \exists u \in \text{int}\Omega^k : y = \varphi_W(k, x, u) \text{ and } (x, u) \text{ is regular}\}, \\ \hat{\mathbf{C}}_k^W(x) &:= \{y \in W \mid \exists u \in \text{int}\Omega^k : \varphi_W(k, y, u) = x \text{ and } (y, u) \text{ is regular}\},\end{aligned}$$

and the regular W -reachability set $\hat{\mathbf{R}}^W(x)$ and W -controllability set $\hat{\mathbf{C}}^W(x)$ are given by the respective union over all $k \in \mathbb{N}$. It is not difficult to see that $\hat{\mathbf{R}}^W(x)$ and $\hat{\mathbf{C}}^W(x)$ are open for every x (cf. Wirth [27, Lemma 8]). In the notation of the local semigroup \mathcal{S} defined in (37) one has

$$\hat{\mathbf{C}}^W(x) \subset \text{int}\mathbf{C}^W(x) = \text{int}\mathcal{S}^*x.$$

In order to show that the transitivity set of a control set is nonvoid, we start with the following observations. If $\hat{\mathbf{R}}_{k_0}^W(x) \neq \emptyset$ it follows that $\hat{\mathbf{R}}_k^W(x) \neq \emptyset$ for all $k > k_0$. Accessibility condition (38) implies for all $x \in W$ that there is $k_0 \in \mathbb{N}$ such that for all $k \geq k_0$ one has $\text{int}\mathbf{R}_k^W(x) \neq \emptyset$ and

$$\mathbf{R}_k^W(x) \subset \text{cl}\{y = \varphi_W(k, x, u) \in \text{int}\mathbf{R}_k^W(x) \mid u \in \text{int}\Omega^k\}.$$

Sard's Theorem (cf., e.g., Katok and Hasselblatt [15, Theorem A.3.13]) implies that the set of points $\varphi_W(k, x, u) \in \mathbf{R}_k^W(x)$ such that (x, u) is not regular has Lebesgue measure zero.

The following proposition presents conditions which imply that the transitivity set of a control set is nonvoid.

Proposition 4. *Consider system (41) and assume that accessibility condition (38) holds. Then for every W -control set $D \subset W$ with nonvoid interior the transitivity set D_0 is nonvoid.*

Proof. Let $x \in \text{int}D$ and consider an open neighborhood $V_1 \subset D$ of x . There is $k_0 \in \mathbb{N}$ such that the reachable set $\mathbf{R}_k^W(x)$ at time k has nonvoid interior for all $k \geq k_0$. There are $k \geq k_0$ and $\varphi_W(k, x, u) \in \mathbf{R}_k^W(x) \cap \text{int}D$, hence we may assume that there is $y := \varphi_W(k, x, u) \in \text{int}\mathbf{R}_k^W(x) \cap \text{int}D$. Then, by Sard's Theorem, it follows that there is a point $y = \varphi_W(k, x, u) \in \text{int}D$ with regular (x, u) , i.e., $y \in \text{int}D \cap \hat{\mathbf{R}}_k^W(x)$. Then $x \in \hat{\mathbf{C}}_k(y) \subset \text{int}\mathbf{C}(y)$. Let $V \subset \text{int}\mathbf{C}(y)$ be a neighborhood of x . Then $x \in D \subset \text{cl}\mathbf{R}^W(y)$, hence there is $z \in V \cap \mathbf{R}^W(y) \subset D$ and thus $y \in \mathbf{C}(z)$. By construction, the point $z \in D$ satisfies $z \in \text{int}\mathbf{C}(y) \subset \text{int}\mathbf{C}(z)$, hence it is in the transitivity set of D . \square

Remark 12. Wirth [27] defines (for $W = M$) the regular core of a control set D denoted by $\text{core}(D)$ as the set of points in D for which the regular reachability and controllability sets intersect D and shows, under real analyticity assumptions, that $\text{core}(D)$ is open and dense in $\text{cl}D$. Thus $\text{core}(D) \subset D_0$. This generalizes earlier results by Albertini and Sontag [1, Section 3] and [2, Section 7] (again for $W = M$). They define the core of a control set as

$$\{x \in \text{int}D \mid \text{int}\mathbf{R}(x) \cap D \neq \emptyset \text{ and } \text{int}\mathbf{C}(x) \cap D \neq \emptyset\}, \quad (42)$$

and assume, in particular, that for every $\omega \in \Omega$ the map $f_\omega := f(\cdot, \omega)$ is a global diffeomorphism on M and that for all $x \in M$ the set of points which can be reached by finite compositions of maps of the form f_ω and f_ω^{-1} applied to x coincides with M . They show that the core is open and that it is dense in D . Again, it is clear that the transitivity set of D is contained in the core as defined in (42).

The results above give conditions which imply that the metric invariance entropy for a quasi-stationary measure is determined by the invariant W -control sets. It may be of interest to analyze the relations between the supports of quasi-stationary measures and W -control sets. For ergodic stationary measures η , Colonius, Homburg and Kliemann [8, Lemma 5] shows (for certain random diffeomorphisms) that the support of η coincides with an invariant control set. The following proposition gives a result in that direction.

In the setting of (3) define the k -step transition function $p_k^Q(x, A)$, $x \in Q$, $A \subset Q$,

$$p_1^Q(x, A) := p(x, A), p_k^Q(x, A) := \int_Q p_{k-1}^Q(y, A) p(x, dy), k > 1.$$

Then (cf. Colonius [6, Remark 2.8]) it follows for all $k \geq 1$ and all $A \subset Q$ that

$$\rho^k \eta(A) = \int_Q p_k^Q(z, A) \eta(dz). \quad (43)$$

Proposition 5. *Consider control system (1) and let $Q \subset M$ be equal to the closure of its interior $W := \text{int}Q$. Consider an invariant W -control set D with nonvoid transitivity set D_0 and let η be a quasi-stationary measure for ν on Ω . Assume that for every $x \in W$, $k \geq 1$ and $y \in \mathbf{R}_k^W(x)$ every neighborhood $V(y)$ satisfies $p_k(x, V(y)) > 0$.*

Then every quasi-stationary measure η with $\text{supp}\eta \cap D \neq \emptyset$ satisfies $D \subset \text{supp}\eta$.

Proof. Suppose, contrary to the assertion, that there is $y \in D \setminus \text{supp}\eta$. Since $W \setminus \text{supp}\eta$ is open in W there is a neighborhood $V(y)$ of y in W such that $V(y) \cap \text{supp}\eta = \emptyset$. By Proposition 2 the transitivity set D_0 is dense in D , hence we may take $y \in D_0$. By assumption, there is $x \in \text{supp}\eta \cap D$. Thus $x \in \mathbf{R}_k^W(y)$ for some $k \geq 1$. By continuity, there is a neighborhood $V(x)$ such that $\mathbf{R}_k^W(z) \cap V(y) \neq \emptyset$ for all $z \in V(x)$ and, by the definition of the support, one has $\eta(V(x)) > 0$. The assumption guarantees that $p_k^Q(z, V(y)) > 0$ for all $z \in V(x)$. This contradicts the quasi-stationarity property (43), since $\eta(D \setminus \text{supp}\eta) = 0$, while

$$\int_Q p_k^Q(z, D \setminus \text{supp}\eta) \eta(dz) \geq \int_{V(x)} p_k^Q(z, V(y)) \eta(dz) > 0.$$

□

The results above show that the metric invariance entropy of a subset Q of the state space is already determined on a subset K that can be characterized using controllability properties. For the topological invariance entropy of systems in continuous time, an analogous result has been shown in Colonius and Lettau [9, Theorem 5.2].

Finally, we present two examples illustrating W -control sets and their relation to invariance entropy. First we take a closer look at Example 2.9 in order to discuss W -control sets in a simple situation.

Example 5.6. Recall that $f_\alpha : \mathbb{R}/\mathbb{Z} \times [-1, 1] \rightarrow \mathbb{R}/\mathbb{Z}$ is given by


$$f_\alpha(x, \omega) = x + \sigma \cos(2\pi x) + A\omega + \alpha \mod 1.$$

For $\alpha \leq \alpha_0$ there is an invariant control set $\hat{D}^\alpha = [d(\alpha), c(\alpha)] \neq \mathbb{R}/\mathbb{Z}$ that varies continuously with α . For $\alpha > \alpha_0$ the only control set is the invariant control set $\hat{D}^\alpha = \mathbb{R}/\mathbb{Z}$.

Now consider $\alpha > \alpha_0$ and $W = (0.2, 0.5)$. There is a unique invariant W -control set D^α , which has the form $D^\alpha = [d(\alpha), 0.5)$. Then one easily sees that the

invariant W -control sets D^α are closed in W and their transitivity sets are nonvoid. Furthermore, the closure $\text{cl}D^\alpha = [d(\alpha), 0.5]$ is invariant in $Q := \text{cl}W = [0, 2, 0.5]$.

For the uniform distribution ν on $\Omega := [-1, 1]$, [8, Theorem 3] implies that for all $\alpha > 0$ there is a unique stationary measure $\hat{\eta}^\alpha$ satisfying $\eta(B) = \int_{\mathbb{R}/\mathbb{Z}} p(x, B) \eta(dx)$ for all $B \subset \mathbb{R}/\mathbb{Z}$. It has support equal to the invariant control set \hat{D}^α . For $\alpha > \alpha_0$ Theorem 5.4 shows that for every quasi-stationary measure η^α of $Q = \text{cl}W$ the invariance entropy of Q coincides with the invariance entropy of the closure $\text{cl}D^\alpha = [d(\alpha), 0.5]$ of the invariant W -control set D^α as already seen in Example 2.9. From Proposition 5 we obtain the additional information that the quasi-stationary measure η_α has support equal to $\text{cl}D^\alpha = [d(\alpha), 0.5]$.



Extremal graphs for (44) and the W -control sets
 $D_1(\alpha) = [a(\alpha), b(\alpha))$ and $D_2(\alpha) = [d(\alpha), 0.7]$ in $Q = [0.1, 0.7]$
 (here $A = 0.05, \sigma = 0.1$ and $\alpha = 0.08$)

We modify this example, so that in addition to an invariant W -control set D_2^α there is there is a second W -control set D_1^α (to the left of D_2^α) which is not invariant.

Example 5.7. Define $f_\alpha : \mathbb{R}/\mathbb{Z} \times [-1, 1] \rightarrow \mathbb{R}/\mathbb{Z}$ by

$$f_\alpha(x, \omega) = x + \sigma \cos(4\pi x) + A\omega + \alpha \pmod{1}. \quad (44)$$

This is illustrated in the figure above. Here $Q = [0, 1, 0.7]$ and with $W = (0.1, 0.7)$ the invariant W -control set is $D_2^\alpha = [d(\alpha), 0.7]$ (to the right). The W -control set $D_1^\alpha = [a(\alpha), b(\alpha))$ (to the left) is not invariant in W , since exit to the right is possible. Invoking Theorem 5.4 one sees that for every quasi-stationary measure η^α of $Q = \text{cl}W$ (such that $f_\alpha(\cdot, \omega), \omega \in [-1, 1]$, are nonsingular) the invariance entropy for μ^α of Q coincides with the invariance entropy of $\text{cl}D_2^\alpha$. As above, for the uniform distribution ν on Ω there is no stationary measure with support contained in Q .

The many open problems in this area include the following. When is the support of a quasi-stationary measure contained in the (closure of) the union of the invariant W -control sets? In this situation, Theorems 4.3 and 4.4 would hold trivially, since, naturally, the metric invariance entropy is determined on the support of the quasi-stationary measure (cf. also Remark 10). Theorem 5.4 reduces the analysis of the

invariance entropy from arbitrary closed sets Q to invariant W -control sets. Hence their measure theoretic invariance entropy is of particular interest. For control sets, the topological invariance entropy has been characterized in Kawan [16] and da Silva and Kawan [10] using hyperbolicity and Lyapunov exponents.

Acknowledgments. I would like to thank two anonymous reviewers for their helpful comments.

REFERENCES

- [1] F. Albertini and E. D. Sontag, Some connections between chaotic dynamical systems and control systems, in *Proc. European Control Conference, Grenoble*, 1991, 158–163.
- [2] ———, Discrete-time transitivity and accessibility: Analytic systems, *SIAM J. Control Optim.*, **31** (1993), 1599–1622.
- [3] M. Benaïm, B. Cloez and F. Panloup, *Stochastic Approximation of Quasi-Stationary Distributions on Compact Spaces and Applications*, arXiv:1606.06477v2 [math.PR] 6 Dec 2016.
- [4] T. Bogenschütz, Entropy, pressure, and a variational principle for random dynamical systems, *Random and Computational Dynamics*, **1** (1992/93), 99–116.
- [5] P. Collett, S. Martinez and J. San Martín, *Quasi-Stationary Distributions: Markov Chains, Diffusions, and Dynamical Systems*, Springer-Verlag, Berlin, 2013.
- [6] F. Colonius, Metric invariance entropy and conditionally invariant measures, *Ergodic Theory and Dynamical Systems*, (2016). First published online: 20 October 2016.
- [7] ———, *Metric Invariance Entropy and Relatively Invariant Control Sets*, in Proceedings of the 55th IEEE Conference on Decision and Control (Las Vegas, December 12–14, 2016), 2016.
- [8] F. Colonius, J.-A. Homburg and W. Kliemann, Near invariance and local transience for random diffeomorphisms, *J. Difference Equations and Applications*, **16** (2010), 127–141.
- [9] F. Colonius and R. Lettau, Relative controllability properties, *IMA Journal of Mathematical Control and Information*, **33** (2016), 701–722.
- [10] A. da Silva and C. Kawan, Invariance entropy of hyperbolic control sets, *Discrete Cont. Dyn. Syst. A*, **36** (2016), 97–136.
- [11] M. F. Demers, *Introductory Lectures on Open Systems*, given as part of the LMS-CMI Research School at Loughborough University, April 13–17, 2015.
- [12] M. F. Demers and L.-S. Young, Escape rates and conditionally invariant measures, *Nonlinearity*, **19** (2006), 377–397.
- [13] E.A. van Doorn and P. Pollett, Quasi-stationary distributions for reducible absorbing Markov chains in discrete time, *Markov Processes and Related Fields*, **15** (2009), 191–204.
- [14] B. Jakubczyk and E. D. Sontag, Controllability of nonlinear discrete time systems: A Lie algebraic approach, *SIAM J. Control Optim.*, **28** (1990), 1–33.
- [15] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems*, Cambridge University Press, 1995.
- [16] C. Kawan, Invariance entropy of control sets, *SIAM J. Control Optim.*, **49** (2011), 732–751.
- [17] ———, *Invariance Entropy for Deterministic Control Systems. An Introduction*, vol. 2089 of Lecture Notes in Mathematics, Springer-Verlag, 2013.
- [18] S. Méléard and D. Villemonais, Quasi-stationary distributions and population processes, *Probability Surveys*, **9** (2012), 340–410.
- [19] G. Nair, R. J. Evans, I. Mareels and W. Moran, Topological feedback entropy and nonlinear stabilization, *IEEE Trans. Aut. Control*, **49** (2004), 1585–1597.
- [20] M. Patrão and L. San Martín, Semiflows on topological spaces: Chain transitivity and semigroups, *J. Dyn. Diff. Equations*, **19** (2007), 155–180.
- [21] P. Pollett, *Quasi-stationary distributions: a bibliography*. <http://www.maths.uq.edu.au/~pkp/papers/qsds/qsds.pdf>, 2015.
- [22] F. Rodrigues and P. Varandas, Specification and thermodynamical properties of semigroup actions, *Journal Math. Phys.*, **57** (2016), 052704, 27pp.
- [23] E. Sontag and F. Wirth, Remarks on universal nonsingular controls for discrete-time systems, *Sys. Control Lett.*, **33** (1998), 81–88.
- [24] M. Viana and K. Oliveira, *Foundations of Ergodic Theory*, Cambridge University Press, 2016.
- [25] P. Walters, *An Introduction to Ergodic Theory*, Springer-Verlag, 1982.

- [26] F. Wirth, *Robust Stability of Discrete-Time Systems under Time-Varying Perturbations*, PhD thesis, Fachbereich Mathematik/Informatik, Universität Bremen, 1995.
- [27] F. Wirth, Dynamics and controllability of nonlinear discrete-time control systems, *IFAC Proceedings Volumes*, **31** (1998), 267–272.

E-mail address: fritz.colonius@math.uni-augsburg.de