

A C0 interior penalty discontinuous Galerkin method for fourth order total variation flow, I: Derivation of the method and numerical results

Chandi Bhandari, Ronald H. W. Hoppe, Rahul Kumar

Angaben zur Veröffentlichung / Publication details:

Bhandari, Chandi, Ronald H. W. Hoppe, and Rahul Kumar. 2019. "A C0 interior penalty discontinuous Galerkin method for fourth order total variation flow, I: Derivation of the method and numerical results." *Numerical Methods for Partial Differential Equations* 35 (4): 1458–76. <https://doi.org/10.1002/num.22359>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



A C^0 interior penalty discontinuous Galerkin method for fourth-order total variation flow. I: Derivation of the method and numerical results

Chandi Bhandari¹ Ronald H.W. Hoppe^{1,2} Rahul Kumar³

¹Department of Mathematics, University of Houston, Houston, Texas

²Department of Mathematics, University of Augsburg, Augsburg, Germany

³Basque Center for Applied Mathematics, Bilbao, Spain

Correspondence

Ronald H. W. Hoppe, Department of Mathematics, University of Houston, 651 P.G. Hoffman, Houston, TX 77204.

Email: rohop@math.uh.edu

Funding information

National Science Foundation, DMS-1520886.

We consider the numerical solution of a fourth-order total variation flow problem representing surface relaxation below the roughening temperature. Based on a regularization and scaling of the nonlinear fourth-order parabolic equation, we perform an implicit discretization in time and a C^0 Interior Penalty Discontinuous Galerkin (C^0 IPDG) discretization in space. The C^0 IPDG approximation can be derived from a mixed formulation involving numerical flux functions where an appropriate choice of the flux functions allows to eliminate the discrete dual variable. The fully discrete problem can be interpreted as a parameter dependent nonlinear system with the discrete time as a parameter. It is solved by a predictor corrector continuation strategy featuring an adaptive choice of the time step sizes. A documentation of numerical results is provided illustrating the performance of the C^0 IPDG method and the predictor corrector continuation strategy. The existence and uniqueness of a solution of the C^0 IPDG method will be shown in the second part of this paper.

KEYWORDS

C^0 interior penalty discontinuous Galerkin method, fourth-order total variation flow, surface relaxation

1 | INTRODUCTION

Surface relaxation by surface diffusion is about the relaxation of a high symmetry crystalline surface on which a particular profile has been imprinted such that the typical length scale of the imposed profile is much larger than the lattice constant (dimension of unit cells in the crystal lattice). Therefore,

surface relaxation is an important process in material sciences, in particular in the production of nanotechnology devices. The problem is to understand along which route the initial profile relaxes to a completely flat surface. One distinguishes between relaxation above and below the roughening temperature. Below the roughening temperature, the surface free energy has a cusp singularity. Several authors have suggested to model the dynamics by a total variation H^{-1} flow problem that can be formulated as a fourth-order total variation flow (TVF) problem (cf., e.g., [1–8]).

Given a bounded domain $\hat{\Omega} \subset \mathbb{R}^2$ with boundary $\hat{\Gamma} = \partial\hat{\Omega}$, the total variation- H^{-1} (TV- H^{-1}) minimization of the energy functional

$$E(w) = \beta \int_{\hat{\Omega}} |\nabla w| \, dx, \quad \beta > 0, \quad (1.1)$$

leads to the following fourth-order TVF problem

$$\frac{\partial w}{\partial t} + \beta \Delta \nabla \cdot \frac{\nabla w}{|\nabla w|} = 0 \quad \text{in } \hat{Q} := \hat{\Omega} \times (0, \hat{T}), \quad (1.2a)$$

$$\mathbf{n}_{\hat{\Gamma}} \cdot \beta \frac{\nabla w}{|\nabla w|} = \mathbf{n}_{\hat{\Gamma}} \cdot \nabla \nabla \left(\nabla \cdot \frac{\nabla w}{|\nabla w|} \right) = 0 \quad \text{on } \hat{\Sigma} := \hat{\Gamma} \times (0, \hat{T}), \quad (1.2b)$$

$$w(\cdot, 0) = w^0 \quad \text{in } \hat{\Omega}, \quad (1.2c)$$

where $\beta > 0$ is related to the mobility, $\hat{T} > 0$ is the final time, $\mathbf{n}_{\hat{\Gamma}}$ stands for the exterior unit normal at $\hat{\Gamma}$, and $w^0 \in L^2(\hat{\Omega})$ is some given initial data. The fourth-order Equation (1.2a) has to be interpreted as follows: On $H^{-1}(\hat{\Omega})$ we introduce an inner product according to

$$(w, z)_{-1, \hat{\Omega}} := (\nabla(-\Delta^{-1}w), \nabla(-\Delta^{-1}z))_{0, \hat{\Omega}},$$

where Δ^{-1} stands for the inverse of the Laplacian. For $E(w) = \beta \int_{\hat{\Omega}} |\nabla w| \, dx$, $w \in H^{-1}(\hat{\Omega})$, with $D(E) = \{w \in H^{-1}(\hat{\Omega}) \mid E(w) < \infty\}$, the subdifferential

$$\partial_{H^{-1}} E(w) = \{v \in H^{-1}(\hat{\Omega}) \mid (v, z - w)_{-1, \hat{\Omega}} \leq E(z) - E(w) \text{ for all } z \in H^{-1}(\hat{\Omega})\}$$

is given by (cf., e.g., [9])

$$\partial_{H^{-1}} E(w) = \{\Delta \nabla \cdot \xi \mid \xi(\hat{x}) \in \partial\Phi(\nabla w(\hat{x}))\},$$

where $\Phi(|\boldsymbol{\eta}|)$ and $\partial\Phi(|\boldsymbol{\eta}|)$ are given by

$$\Phi(\boldsymbol{\eta}) = \beta |\boldsymbol{\eta}|, \quad \partial\Phi(\boldsymbol{\eta}) = \begin{cases} \beta \boldsymbol{\eta} / |\boldsymbol{\eta}| & \text{if } \boldsymbol{\eta} \neq \mathbf{0} \\ \{\boldsymbol{\tau} \in \mathbb{R}^2 \mid |\boldsymbol{\tau}| \leq \beta\} & \text{if } \boldsymbol{\eta} = \mathbf{0} \end{cases}. \quad (1.3)$$

We thus obtain

$$-\frac{\partial w}{\partial t} \in \partial E_{H^{-1}}(w).$$

Initial-boundary value problems for fourth-order TVF problems have been considered mainly from an analytical point of view (cf., e.g., [9–13]).

Here, we consider the regularized TV- H^{-1} energy functional

$$E_{\text{reg}}(w) = \beta \int_{\hat{\Omega}} (\delta^2 + |\nabla w|^2)^{1/2} \, dx, \quad w \in H^{-1}(\hat{\Omega}),$$

where $\delta > 0$ is a regularization parameter. This leads to the regularized fourth-order TVF problem

$$\frac{\partial w}{\partial t} + \beta \Delta \nabla \cdot ((\delta^2 + |\nabla w|^2)^{-1/2} \nabla w) = 0 \quad \text{in } \hat{Q}, \quad (1.4a)$$

$$\mathbf{n}_{\widehat{\Gamma}} \cdot \beta(\delta^2 + |\nabla w|^2)^{-1/2} \nabla w = 0 \quad \text{on } \widehat{\Sigma}, \quad (1.4b)$$

$$\begin{aligned} \mathbf{n}_{\widehat{\Gamma}} \cdot \beta \nabla(\nabla \cdot (\delta^2 + |\nabla w|^2)^{-1/2} \nabla w) &= 0 \quad \text{on } \widehat{\Sigma}, \\ w(\cdot, 0) &= w^0 \quad \text{in } \widehat{\Omega}. \end{aligned} \quad (1.4c)$$

We further consider a scaling in both the time variable and the spatial variables according to

$$t = \delta \widehat{t}, \quad x_i = \delta \widehat{x}_i, \quad 1 \leq i \leq 2. \quad (1.5)$$

Setting $T := \delta \widehat{T}$, $\Omega := \delta \widehat{\Omega}$, $\Gamma := \partial\Omega$, $Q := \Omega \times (0, T)$, $\Sigma := \Gamma \times (0, T)$, and $u^0(x) = w^0(\delta^{-1}x)$, as well as

$$\omega(\nabla u) := 1 + |\nabla u|^2, \quad (1.6)$$

the scaled regularized fourth-order TVF problem reads as follows

$$\frac{\partial u}{\partial t} + \beta \delta^2 \Delta \nabla \cdot (\omega(\nabla u)^{-1/2} \nabla u) = 0 \quad \text{in } Q, \quad (1.7a)$$

$$\mathbf{n}_{\Gamma} \cdot \beta \delta^2 (\omega(\nabla u)^{-1/2} \nabla u) = \mathbf{n}_{\Gamma} \cdot \beta \delta^2 \nabla(\nabla \cdot (\omega(\nabla u)^{-1/2} \nabla u)) = 0 \quad \text{on } \Sigma, \quad (1.7b)$$

$$u(\cdot, 0) = u^0 \quad \text{in } \Omega. \quad (1.7c)$$

The numerical solution of the regularized fourth-order TVF problem with periodic boundary conditions has been considered in Kohn and Versieux [3] based on a mixed formulation of the implicitly in time discretized problem. At each time-step, this amounts to the solution of two second-order elliptic partial differential equations (PDEs) by standard Lagrangian finite elements with respect to a triangulation of the computational domain Ω [14]. On the other hand, Interior Penalty Discontinuous Galerkin (IPDG) methods for fourth-order elliptic boundary value problems, fourth-order and higher-order polyharmonic parabolic initial-boundary value problems have been studied in [15–28]. The advantage of the C^0 IPDG approach is that it directly applies to the fourth-order problem and thus only requires the numerical solution of one equation by using the same Lagrangian finite elements as in the mixed method.

Remark 1.1 We note that another example for a $TV\text{-}H^{-1}$ minimization problem is the minimization of the energy functional

$$E(w, \widehat{g}) = \int_{\widehat{\Omega}} |\nabla w| \, dx + \frac{\lambda}{2} \|w - \widehat{g}\|_{H^{-1}(\widehat{\Omega})}^2$$

which occurs in image recovery where w represents a true image, \widehat{g} describes a blurred and/or noisy image, and $\lambda > 0$ is a fidelity parameter (cf., [29–32]). The associated fourth-order TVF problem is given by the initial-boundary value problem

$$\frac{\partial w}{\partial t} + \lambda^{-1} \Delta \nabla \cdot \frac{\nabla w}{|\nabla w|} + w - \widehat{g} = 0 \quad \text{in } \widehat{Q}, \quad (1.8a)$$

$$\mathbf{n}_{\widehat{\Gamma}} \cdot \frac{\nabla w}{|\nabla w|} = \mathbf{n}_{\widehat{\Gamma}} \cdot \nabla \left(\nabla \cdot \frac{\nabla w}{|\nabla w|} \right) = 0 \quad \text{on } \widehat{\Sigma}, \quad (1.8b)$$

$$w(\cdot, 0) = w^0 \quad \text{in } \widehat{\Omega}. \quad (1.8c)$$

The paper is organized as follows: In Section 2, we will perform a discretization in time of the regularized and scaled fourth-order TVF problem and consider a reformulation in terms of the matrix of second-order partial derivatives of the unknown. Section 3 is devoted to the derivation of the C^0 IPDG

approximation based on an appropriate choice of numerical flux functions. In Section 4, we will show that the fully discrete system can be written as a parameter-dependent nonlinear system with the discrete time as a parameter. Since the choice of the time steps is crucial for the convergence of Newton's method, we suggest a predictor corrector continuation strategy with constant continuation as a predictor and Newton's method as a corrector featuring an adaptive choice of the time step sizes. This avoids a breakdown of Newton's method due to convergence failure because of too large time steps. Finally, in Section 5 we present numerical results illustrating the performance of the C⁰IPDG method and the predictor corrector continuation strategy.

Throughout the paper we will use the following notations and basic results. For vectors $\underline{\mathbf{x}} = (x_1, \dots, x_n)^T, \underline{\mathbf{y}} = (y_1, \dots, y_n)^T \in \mathbb{R}^n$ and for matrices $\underline{\underline{\mathbf{A}}} = (a_{ij})_{i,j=1}^n, \underline{\underline{\mathbf{B}}} = (b_{ij})_{i,j=1}^n \in \mathbb{R}^{n \times n}$ we denote by $\underline{\mathbf{x}} \cdot \underline{\mathbf{y}}$ and $\underline{\underline{\mathbf{A}}} : \underline{\underline{\mathbf{B}}}$ the Euclidean inner product $\underline{\mathbf{x}} \cdot \underline{\mathbf{y}} = \sum_{i=1}^n x_i y_i$ and the Frobenius inner product $\underline{\underline{\mathbf{A}}} : \underline{\underline{\mathbf{B}}} = \sum_{i,j=1}^n a_{ij} b_{ij}$. In particular, $|\underline{\mathbf{x}}| := (\underline{\mathbf{x}} \cdot \underline{\mathbf{x}})^{1/2}$ and $|\underline{\underline{\mathbf{A}}}| := (\underline{\underline{\mathbf{A}}} : \underline{\underline{\mathbf{A}}})^{1/2}$ refer to the Euclidean norm and the Frobenius norm, respectively.

We will further use standard notation from Lebesgue and Sobolev space theory (cf., e.g., [33]). In particular, for a bounded domain $D \subset \mathbb{R}^d, d \in \mathbb{N}$, we refer to $L^2(D)$ as the Hilbert space of square integrable functions on D with inner product $(\cdot, \cdot)_{0,D}$ and norm $\|\cdot\|_{0,D}$. Moreover, we denote by $H^m(D), m \in \mathbb{N}$, the Sobolev space with inner product $(\cdot, \cdot)_{m,D}$ and norm $\|\cdot\|_{m,D}$.

2 | IMPLICIT TIME-DISCRETIZATION

For the numerical solution of the regularized fourth-order TVF problem (Equation (1.7)) we perform a discretization in time with respect to a partition of the time interval $[0, T]$ into subintervals $[t_{m-1}, t_m]$ of length $\tau_m := t_m - t_{m-1}$. Denoting by u^m some approximation of u at time t_m , for $1 \leq m \leq M$ we have to solve the problems

$$u^m - u^{m-1} + \tau_m \beta \delta^2 \Delta \nabla \cdot (\omega(\nabla u^m)^{-1/2} \nabla u^m) = 0 \text{ in } \Omega, \quad (2.1a)$$

$$\mathbf{n}_\Gamma \cdot \beta \delta^2 (\omega(\nabla u^m)^{-1/2} \nabla u^m) = \mathbf{n}_\Gamma \cdot \beta \delta^2 \nabla (\nabla \cdot (\omega(\nabla u^m)^{-1/2} \nabla u^m)) = 0 \text{ on } \Gamma. \quad (2.1b)$$

Introducing the objective functional

$$J(v) := \frac{1}{2} \|v - u^{m-1}\|_{-1,\Omega}^2 + \tau_m \beta \delta^2 \int_{\Omega} (1 + |\nabla v|^2)^{1/2} dx, \quad (2.2)$$

it is easy to see that Equation (2.1) is related to the necessary and sufficient optimality condition for the minimization problem

$$J(u^m) = \inf_{v \in H^{-1}(\Omega)} J(v), \quad (2.3)$$

which has a unique solution, since the objective functional J is strictly convex, coercive, and lower semicontinuous. The fourth-order Equation (2.1a) can be reformulated in terms of the 2×2 matrix

$$D^2 u^m = \begin{pmatrix} \frac{\partial^2 u^m}{\partial x_1^2} & \frac{\partial^2 u^m}{\partial x_1 \partial x_2} \\ \frac{\partial^2 u^m}{\partial x_1 \partial x_2} & \frac{\partial^2 u^m}{\partial x_2^2} \end{pmatrix}.$$

of second partial derivatives of u^m . We note that the divergence of a matrix-valued function $\underline{\underline{\mathbf{q}}} = (q_{ij})_{i,j=1}^2$ with row vectors $\underline{\mathbf{q}}^{(i)} = (q_{i1}, q_{i2})^T, 1 \leq i \leq 2$, is defined by means of

$$\nabla \cdot \underline{\underline{\mathbf{q}}} := (\nabla \cdot \underline{\mathbf{q}}^{(1)}, \nabla \cdot \underline{\mathbf{q}}^{(2)})^T. \quad (2.4)$$

Theorem 2.1 *The fourth-order Equation (2.1a) is equivalent to*

$$u^m - u^{m-1} + \tau_m \beta \delta^2 \nabla \cdot \nabla \cdot (\omega(\nabla u^m)^{-3/2} \underline{\underline{\mathbf{M}}}(u^m) D^2 u^m) = 0, \quad (2.5)$$

where $\underline{\underline{\mathbf{M}}}(v)$ stands for the matrix-valued function

$$\underline{\underline{\mathbf{M}}}(v) := \begin{pmatrix} 1 + \left(\frac{\partial v}{\partial x_2} \right)^2 & -\frac{\partial v}{\partial x_1} \frac{\partial v}{\partial x_2} \\ -\frac{\partial v}{\partial x_1} \frac{\partial v}{\partial x_2} & 1 + \left(\frac{\partial v}{\partial x_2} \right)^2 \end{pmatrix}. \quad (2.6)$$

Proof. We reformulate the second term on the left-hand side of Equation (2.1a) according to

$$\begin{aligned} \Delta \nabla \cdot (\omega(\nabla u^m)^{-1/2} \nabla u^m) &= \nabla \cdot \nabla (\nabla \cdot (\omega(\nabla u^m)^{-1/2} \nabla u^m)) \\ &= \nabla \cdot \nabla \cdot \nabla (\omega(\nabla u^m)^{-1/2} \nabla u^m). \end{aligned} \quad (2.7)$$

Obviously, we have

$$\nabla (\omega(\nabla u^m)^{-1/2} \nabla u^m) = \begin{pmatrix} \frac{\partial}{\partial x_1} \\ \frac{\partial}{\partial x_2} \end{pmatrix} \left(\omega(\nabla u^m)^{-1/2} \begin{pmatrix} \frac{\partial u^m}{\partial x_1} \\ \frac{\partial u^m}{\partial x_2} \end{pmatrix} \right). \quad (2.8)$$

In particular,

$$\begin{aligned} \frac{\partial}{\partial x_1} \left(\omega(\nabla u^m)^{-1/2} \begin{pmatrix} \frac{\partial u^m}{\partial x_1} \\ \frac{\partial u^m}{\partial x_2} \end{pmatrix} \right) &= -\omega(\nabla u^m)^{-3/2} \left(\frac{\partial u^m}{\partial x_1} \frac{\partial^2 u^m}{\partial x_1^2} + \frac{\partial u^m}{\partial x_2} \frac{\partial^2 u^m}{\partial x_1 \partial x_2} \right) \begin{pmatrix} \frac{\partial u^m}{\partial x_1} \\ \frac{\partial u^m}{\partial x_2} \end{pmatrix} \\ &+ \omega(\nabla u^m)^{-1/2} \begin{pmatrix} \frac{\partial^2 u^m}{\partial x_1^2} \\ \frac{\partial^2 u^m}{\partial x_1 \partial x_2} \end{pmatrix} = \omega(\nabla u^m)^{-3/2} \begin{pmatrix} \left(1 + \left(\frac{\partial u^m}{\partial x_2} \right)^2 \right) \frac{\partial^2 u^m}{\partial x_1^2} - \frac{\partial u^m}{\partial x_1} \frac{\partial u^m}{\partial x_2} \frac{\partial^2 u^m}{\partial x_1 \partial x_2} \\ \left(1 + \left(\frac{\partial u^m}{\partial x_1} \right)^2 \right) \frac{\partial^2 u^m}{\partial x_1 \partial x_2} - \frac{\partial u^m}{\partial x_1} \frac{\partial u^m}{\partial x_2} \frac{\partial^2 u^m}{\partial x_1^2} \end{pmatrix}, \end{aligned} \quad (2.9)$$

and

$$\begin{aligned} \frac{\partial}{\partial x_2} \left(\omega(\nabla u^m)^{-1/2} \begin{pmatrix} \frac{\partial u^m}{\partial x_1} \\ \frac{\partial u^m}{\partial x_2} \end{pmatrix} \right) &= -\omega(\nabla u^m)^{-3/2} \left(\frac{\partial u^m}{\partial x_1} \frac{\partial^2 u^m}{\partial x_1 \partial x_2} + \frac{\partial u^m}{\partial x_2} \frac{\partial^2 u^m}{\partial x_2^2} \right) \begin{pmatrix} \frac{\partial u^m}{\partial x_1} \\ \frac{\partial u^m}{\partial x_2} \end{pmatrix} \\ &+ \omega(\nabla u^m)^{-1/2} \begin{pmatrix} \frac{\partial^2 u^m}{\partial x_1 \partial x_2} \\ \frac{\partial^2 u^m}{\partial x_2^2} \end{pmatrix} = \omega(\nabla u^m)^{-3/2} \begin{pmatrix} \left(1 + \left(\frac{\partial u^m}{\partial x_2} \right)^2 \right) \frac{\partial^2 u^m}{\partial x_1 \partial x_2} - \frac{\partial u^m}{\partial x_1} \frac{\partial u^m}{\partial x_2} \frac{\partial^2 u^m}{\partial x_1^2} \\ \left(1 + \left(\frac{\partial u^m}{\partial x_1} \right)^2 \right) \frac{\partial^2 u^m}{\partial x_2^2} - \frac{\partial u^m}{\partial x_1} \frac{\partial u^m}{\partial x_2} \frac{\partial^2 u^m}{\partial x_1 \partial x_2} \end{pmatrix}. \end{aligned} \quad (2.10)$$

Using Equations (2.9) and (2.10) in Equation (2.8), it follows that

$$\nabla (\omega(\nabla u^m)^{-1/2} \nabla u^m) = \omega(\nabla u^m)^{-3/2} \begin{pmatrix} 1 + \left(\frac{\partial u^m}{\partial x_2} \right)^2 & -\frac{\partial u^m}{\partial x_1} \frac{\partial u^m}{\partial x_2} \\ -\frac{\partial u^m}{\partial x_1} \frac{\partial u^m}{\partial x_2} & 1 + \left(\frac{\partial u^m}{\partial x_2} \right)^2 \end{pmatrix} D^2 u^m, \quad (2.11)$$

which can be written as

$$\nabla(\omega(\nabla u^m)^{-1/2} \nabla u^m) = \omega(\nabla u^m)^{-3/2} \underline{\underline{\mathbf{M}}}(u^m) D^2 u^m. \quad (2.12)$$

Remark 2.1 The matrix $\underline{\underline{\mathbf{M}}}(v)$ is symmetric positive definite with the eigenvalues

$$\lambda_{\min}(\underline{\underline{\mathbf{M}}}(v)) = 1, \quad \lambda_{\max}(\underline{\underline{\mathbf{M}}}(v)) = 1 + |\nabla v|^2. \quad (2.13)$$

For notational convenience we set

$$\underline{\underline{\mathbf{A}}}_1(v) := \omega(\nabla v)^{-3/2} \underline{\underline{\mathbf{M}}}(v). \quad (2.14)$$

The weak formulation of Equation (2.5) reads: Find

$$u^m \in V := \{v \in H^2(\Omega) \mid \mathbf{n}_\Gamma \cdot \beta \delta^2 \omega(\nabla v)^{-1/2} \nabla v = 0 \text{ on } \Gamma\}$$

such that for all $v \in V$ it holds

$$(u^m - u^{m-1}, v)_{0,\Omega} + \tau_m \beta \delta^2 \int_{\Omega} (\underline{\underline{\mathbf{A}}}_1(u^m) D^2 u^m) : D^2 v \, dx = 0. \quad (2.15)$$

Finally, we provide a mixed formulation of Equation (2.5), because the derivation of the C^0 IPDG method will be based on the discrete analogue of that mixed formulation. Introducing the matrix-valued function

$$\underline{\underline{\mathbf{p}}}^m := \omega(\nabla u^m)^{-1/4} D^2 u^m, \quad (2.16)$$

and the matrix

$$\underline{\underline{\mathbf{A}}}_2(v) := \omega(\nabla v)^{-5/4} \underline{\underline{\mathbf{M}}}(v). \quad (2.17)$$

the mixed formulation of Equations (2.1a) and (2.1b) reads as follows

$$\underline{\underline{\mathbf{p}}}^m - \omega(\nabla u^m)^{-1/4} D^2 u^m = 0 \text{ in } \Omega, \quad (2.18a)$$

$$u^m - u^{m-1} + \tau_m \beta \delta^2 \nabla \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u^m) \underline{\underline{\mathbf{p}}}^m = 0 \text{ in } \Omega, \quad (2.18b)$$

$$\mathbf{n}_\Gamma \cdot \beta \delta^2 \omega(\nabla u^m)^{-1/2} \nabla u^m = 0 \text{ on } \Gamma, \quad (2.18c)$$

$$\mathbf{n}_\Gamma \cdot \beta \delta^2 \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u^m) \underline{\underline{\mathbf{p}}}^m = 0 \text{ on } \Gamma. \quad (2.18d)$$

3 | C^0 INTERIOR PENALTY DISCONTINUOUS GALERKIN APPROXIMATION

Let \mathcal{T}_h be a geometrically conforming, uniform simplicial triangulation of Ω . We denote by $\mathcal{E}_h(\Omega)$ and $\mathcal{E}_h(\Gamma)$ the set of edges of \mathcal{T}_h in the interior of Ω and on the boundary Γ , respectively, and set $\mathcal{E}_h := \mathcal{E}_h(\Omega) \cup \mathcal{E}_h(\Gamma)$. For $K \in \mathcal{T}_h$ and $E \in \mathcal{E}_h$ we denote by h_K and h_E the diameter of K and the length of E , and we set $h := \max(h_K \mid K \in \mathcal{T}_h)$. Due to the assumptions on \mathcal{T}_h there exist constants $0 < c_R \leq C_R$, $0 < c_Q \leq C_Q$, and $0 < c_S \leq C_S$ such that for all $K \in \mathcal{T}_h$ it holds

$$c_R h_K \leq h_E \leq C_R h_K, \quad E \in \mathcal{E}_h(\partial K), \quad (3.1a)$$

$$c_Q h \leq h_K \leq C_Q h, \quad (3.1b)$$

$$c_S h_K^2 \leq \text{meas}(K) \leq C_S h_K^2. \quad (3.1c)$$

For two quantities A and B we write $A \lesssim B$, if there exists a constant $C > 0$ independent of h such that $A \leq CB$.

Denoting by $P_k(T)$, $k \in \mathbb{N}$, the linear space of polynomials of degree $\leq k$ on T , for $k \in \mathbb{N}$ we define

$$V_h := \{v_h \in C^0(\Omega) | v_h|_T \in P_k(T), \quad T \in \mathcal{T}_h\} \quad (3.2)$$

and note that $V_h \subset H^1(\Omega)$, but $V_h \not\subset H^2(\Omega)$. Further, we introduce

$$\underline{\underline{\mathbf{M}}}_h := \{\underline{\underline{\mathbf{q}}}_h \in L^2(\Omega)^{2 \times 2} | \underline{\underline{\mathbf{q}}}_h|_K \in P_k(K)^{2 \times 2}, \quad K \in \mathcal{T}_h\} \quad (3.3)$$

as the space of element-wise polynomial moment tensors.

For interior edges $E \in \mathcal{E}_h(\Omega)$ such that $E = K_+ \cap K_-$, $K_{\pm} \in \mathcal{T}_h$ and boundary edges on Γ we introduce the average and jump of ∇v_h according to

$$\{\nabla v_h\}_E := \begin{cases} \frac{1}{2}(\nabla v_h|_{E \cap K_+} + \nabla v_h|_{E \cap K_-}) & E \in \mathcal{E}_h(\Omega) \\ \nabla v_h|_E & E \in \mathcal{E}_h(\Gamma) \end{cases}, \quad (3.4a)$$

$$[\nabla v_h]_E := \begin{cases} \nabla v_h|_{E \cap K_+} - \nabla v_h|_{E \cap K_-} & E \in \mathcal{E}_h(\Omega) \\ \nabla v_h|_E & E \in \mathcal{E}_h(\Gamma) \end{cases}. \quad (3.4b)$$

The average $\{\Delta v_h\}_E$ and jump $[\Delta v_h]_E$ are defined analogously. We further denote by \mathbf{n}_E the unit normal vector on E pointing in the direction from K_+ to K_- .

In order to motivate the C^0 IPDG approximation, we will follow the approach taken in Arnold and co-workers [34] for second-order elliptic boundary value problems. For $\underline{\underline{\mathbf{p}}}^m \in \underline{\underline{\mathbf{M}}}_h$ and $u_h^m \in V_h$ we consider Equations (2.18a) and (2.18b) element-wise and Equations (2.18c) and (2.18d) edgewise, that is,

$$\underline{\underline{\mathbf{p}}}^m - \omega(\nabla u_h^m)^{-1/4} D^2 u_h^m = 0 \quad \text{in } K \in \mathcal{T}_h, \quad (3.5a)$$

$$u_h^m - u_h^{m-1} + \tau_m \beta \delta^2 \nabla \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}^m = 0 \quad \text{in } K \in \mathcal{T}_h, \quad (3.5b)$$

$$\mathbf{n}_E \cdot \beta \delta^2 \omega(\nabla u_h^m)^{-1/4} \nabla u_h^m = 0 \quad \text{on } E \in \mathcal{E}_h(\Gamma), \quad (3.5c)$$

$$\mathbf{n}_E \cdot \beta \delta^2 \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}^m = 0 \quad \text{on } E \in \mathcal{E}_h(\Gamma), \quad (3.5d)$$

with $u_h^0 = Q_h u^0$, where $Q_h : L^2(\Omega) \rightarrow V_h$ denotes the L^2 projection onto V_h . We multiply Equation (3.5a) by $\underline{\underline{\mathbf{q}}}_h \in \underline{\underline{\mathbf{M}}}_h$ and integrate over K :

$$\int_K \underline{\underline{\mathbf{p}}}^m : \underline{\underline{\mathbf{q}}}_h \, dx = \int_K (\omega(\nabla u_h^m)^{-1/4} D^2 u_h^m) : \underline{\underline{\mathbf{q}}}_h \, dx. \quad (3.6)$$

In view of Equation (2.11) Green's formula yields

$$\begin{aligned} \int_K (\omega(\nabla u_h^m)^{-1/4} D^2 u_h^m) : \underline{\underline{\mathbf{q}}}_h \, dx &= \int_K D^2 u_h^m : (\omega(\nabla u_h^m)^{-1/4} \underline{\underline{\mathbf{q}}}_h) \, dx \\ &= - \int_K \nabla u_h^m \cdot \nabla \cdot (\omega(\nabla u_h^m)^{-1/4} \underline{\underline{\mathbf{q}}}_h) \, dx + \int_{\partial K} \omega(\nabla u_h^m)^{-1/4} \nabla u_h^m \cdot \underline{\underline{\mathbf{q}}}_h \mathbf{n}_{\partial K} \, ds. \end{aligned} \quad (3.7)$$

On the other hand, we multiply Equation (3.5b) by $v_h \in V_h$ and integrate over K :

$$\int_K (u_h^m - u_h^{m-1}) v_h \, dx + \tau_m \beta \delta^2 \int_K \nabla \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}^m v_h \, dx = 0. \quad (3.8)$$

Applying Green's formula twice, we obtain

$$\begin{aligned}
& \int_K \nabla \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m v_h \, dx = - \int_K \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m \cdot \nabla v_h \, dx \\
& + \int_{\partial K} \mathbf{n}_{\partial K} \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m v_h \, ds = \int_K \underline{\underline{\mathbf{p}}}_h^m : \underline{\underline{\mathbf{A}}}_2(u_h^m) D^2 v_h \, dx \\
& - \int_{\partial K} \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m \mathbf{n}_{\partial K} \cdot \nabla v_h \, ds + \int_{\partial K} \mathbf{n}_{\partial K} \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m v_h \, ds.
\end{aligned} \tag{3.9}$$

Summing over all $K \in \mathcal{T}_h$ in Equations (3.7) and (3.9), we obtain the weak formulation of the mixed formulation Equations (3.5a)–(3.5d): Find $(u_h^m, \underline{\underline{\mathbf{p}}}_h^m) \in V_h \times \underline{\underline{\mathbf{M}}}_h$ such that for all $(v_h, \underline{\underline{\mathbf{q}}}_h) \in V_h \times \underline{\underline{\mathbf{M}}}_h$ it holds

$$\begin{aligned}
& \sum_{K \in \mathcal{T}_h} \int_K \underline{\underline{\mathbf{p}}}_h^m : \underline{\underline{\mathbf{q}}}_h \, dx + \sum_{K \in \mathcal{T}_h} \int_K \nabla u_h^m \cdot \nabla \cdot (\omega(\nabla u_h^m)^{-1/4} \underline{\underline{\mathbf{q}}}_h) \, dx \\
& - \sum_{K \in \mathcal{T}_h} \int_{\partial K} \omega(\nabla u_h^m)^{-1/4} \nabla u_h^m \cdot \underline{\underline{\mathbf{q}}}_h \mathbf{n}_{\partial K} \, ds = 0,
\end{aligned} \tag{3.10a}$$

$$\begin{aligned}
& \sum_{K \in \mathcal{T}_h} \int_K u_h^m v_h \, dx + \tau_m \beta \delta^2 \left(\sum_{K \in \mathcal{T}_h} \int_K \underline{\underline{\mathbf{p}}}_h^m : \underline{\underline{\mathbf{A}}}_2(u_h^m) D^2 v_h \, dx \right. \\
& \left. - \sum_{K \in \mathcal{T}_h} \int_{\partial K} \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m \mathbf{n}_{\partial K} \cdot \nabla v_h \, ds + \sum_{K \in \mathcal{T}_h} \int_{\partial K} \mathbf{n}_{\partial K} \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m v_h \, ds \right) \\
& = \sum_{K \in \mathcal{T}_h} \int_K u_h^{m-1} v_h \, dx.
\end{aligned} \tag{3.10b}$$

A general C^0 DG approximation of Equations (2.1b) and (2.1b) is based on the mixed formulation Equations (3.10a) and (3.10b) and characterized by numerical flux functions $\hat{\underline{\underline{\mathbf{u}}}}_{\partial K}^m$, $\hat{\underline{\underline{\mathbf{p}}}}_{\partial K}^m$, and $\hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m$ that are single-valued on $E \in \mathcal{E}_h(\Omega)$, that is, $\hat{\underline{\underline{\mathbf{u}}}}_{\partial K}^m|_{E_+} = \hat{\underline{\underline{\mathbf{u}}}}_{\partial K}^m|_{E_-}$, $\hat{\underline{\underline{\mathbf{p}}}}_{\partial K}^m|_{E_+} = \hat{\underline{\underline{\mathbf{p}}}}_{\partial K}^m|_{E_-}$, and $\hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m|_{E_+} = \hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m|_{E_-}$. We replace $\omega(\nabla u_h^m)^{-1/4} \nabla u_h^m \cdot \underline{\underline{\mathbf{q}}}_h \mathbf{n}_{\partial K}$ in Equation (3.10a) by $\hat{\underline{\underline{\mathbf{u}}}}_{\partial K}^m \cdot \underline{\underline{\mathbf{q}}}_h \mathbf{n}_{\partial K}$ and $\underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m \mathbf{n}_{\partial K} \cdot \nabla v_h = \underline{\underline{\mathbf{A}}}_2(u_h^m) D^2(u_h^m) \mathbf{n}_{\partial K} \cdot \omega(\nabla u_h^m)^{-1/4} \nabla v_h$ as well as $\mathbf{n}_{\partial K} \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m v_h$ in Equation (3.10b) by $\hat{\underline{\underline{\mathbf{p}}}}_{\partial K}^m \cdot \omega(\nabla u_h^m)^{-1/4} \nabla v_h$ and $\mathbf{n}_{\partial K} \cdot \hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m v_h$. In view of Equation (3.5d) we choose $\hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m|_E = \mathbf{n}_E \cdot \nabla \cdot \underline{\underline{\mathbf{A}}}_2(u_h^m) \underline{\underline{\mathbf{p}}}_h^m = 0$, $E \in \mathcal{E}_h(\Gamma)$. Then, no matter how we choose $\hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m|_E$ on $E \in \mathcal{E}_h(\Omega)$, due to $[\hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m]_E = 0$, $E \in \mathcal{E}_h(\Omega)$, and $v_h \in C^0(\Omega)$ we have

$$\sum_{K \in \mathcal{T}_h} \int_{\partial K} \mathbf{n}_{\partial K} \cdot \hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m v_h \, ds = \sum_{E \in \mathcal{E}_h(\Omega)} \mathbf{n}_E \cdot [\hat{\underline{\underline{\mathbf{s}}}}_{\partial K}^m]_E v_h \, ds = 0. \tag{3.11}$$

Consequently, observing Equation (3.11), in a general C^0 DG approximation, we are looking for a pair $(u_h^m, \underline{\underline{\mathbf{p}}}_h^m) \in V_h \times \underline{\underline{\mathbf{M}}}_h$ such that for all $(v_h, \underline{\underline{\mathbf{q}}}_h) \in V_h \times \underline{\underline{\mathbf{M}}}_h$ it holds

$$\begin{aligned}
& \sum_{K \in \mathcal{T}_h} \int_K \underline{\underline{\mathbf{p}}}_h^m : \underline{\underline{\mathbf{q}}}_h \, dx + \sum_{K \in \mathcal{T}_h} \int_K \nabla u_h^m \cdot \nabla \cdot (\omega(\nabla u_h^m)^{-1/4} \underline{\underline{\mathbf{q}}}_h) \, dx \\
& - \sum_{K \in \mathcal{T}_h} \int_{\partial K} \hat{\underline{\underline{\mathbf{u}}}}_{\partial K}^m \cdot \underline{\underline{\mathbf{q}}}_h \mathbf{n}_{\partial K} \, ds = 0,
\end{aligned} \tag{3.12a}$$

$$\sum_{K \in \mathcal{T}_h} \int_K u_h^m v_h \, dx + \tau_m \beta \delta^2 \left(\sum_{K \in \mathcal{T}_h} \int_K \underline{\underline{\mathbf{p}}}_h^m : \underline{\underline{\mathbf{A}}}_2(u_h^m) D^2 v_h \, dx \right.$$

$$- \sum_{K \in \mathcal{T}_h} \int_{\partial K} \hat{\mathbf{p}}_{\partial K}^m \cdot (\omega(\nabla u_h^m)^{-1/4} \nabla v_h) ds = \sum_{K \in \mathcal{T}_h} \int_K u_h^{m-1} v_h dx. \quad (3.12b)$$

In particular, for the C^0 IPDG approximation the numerical flux functions $\hat{\mathbf{u}}_{\partial K}^m$ and $\hat{\mathbf{p}}_{\partial K}^m$ are given by

$$\hat{\mathbf{u}}_{\partial K}^m|_E := \begin{cases} \{\omega(\nabla u_h^m)^{-1/4} \nabla u_h^m\}_E & E \in \mathcal{E}_h(\Omega) \\ \mathbf{0} & E \in \mathcal{E}_h(\Gamma) \end{cases}, \quad (3.13a)$$

$$\hat{\mathbf{p}}_{\partial K}^m|_E := \{\underline{\mathbf{A}}_2(u_h^m) D^2 u_h^m\}_E \mathbf{n}_E - \alpha h_E^{-1} [\omega(\nabla u_h^m)^{-1/4} \nabla u_h^m]_E, \quad E \in \mathcal{E}_h, \quad (3.13b)$$

where $\alpha > 0$ is a penalty parameter.

Theorem 3.1 *The particular choice (Equations (3.13a) and (3.13b)) of the numerical flux functions allows to eliminate the dual variable \mathbf{p}_h^m from the system (Equations (3.12a) and (3.12b)). We thus obtain the C^0 IPDG approximation: Find $u_h^m \in V_h$ such that for all $v_h \in V_h$ it holds*

$$(u_h^m, v_h)_{0,\Omega} + \tau_m \beta \delta^2 a_h^{IP}(u_h^m, v_h; u_h^m) = (u_h^{m-1}, v_h)_{0,\Omega}, \quad (3.14)$$

where for $z_h \in V_h$ the mesh-dependent C^0 IPDG form $a_h^{IP}(\cdot, \cdot; z_h) : V_h \times V_h$ is given by

$$\begin{aligned} a_h^{IP}(u_h, v_h; z_h) := & \sum_{K \in \mathcal{T}_h} (\underline{\mathbf{A}}_1(z_h) D^2 u_h, D^2 v_h)_{0,K} \\ & - \sum_{E \in \mathcal{E}_h} (\mathbf{n}_E \cdot \{\underline{\mathbf{A}}_2(z_h) D^2 u_h\}_E \mathbf{n}_E, \mathbf{n}_E \cdot [\omega(\nabla z_h)^{-1/4} \nabla v_h]_{0,E}) \\ & - \sum_{E \in \mathcal{E}_h} (\mathbf{n}_E \cdot \{\underline{\mathbf{A}}_2(z_h) D^2 v_h\}_E \mathbf{n}_E, \mathbf{n}_E \cdot [\omega(\nabla z_h)^{-1/4} \nabla u_h]_{0,E}) \\ & + \alpha \sum_{E \in \mathcal{E}_h} h_E^{-1} (\mathbf{n}_E \cdot [\omega(\nabla z_h)^{-1/4} \nabla u_h]_E, \mathbf{n}_E \cdot [\omega(\nabla z_h)^{-1/4} \nabla v_h]_{0,E}). \end{aligned} \quad (3.15)$$

Proof. If we choose $\mathbf{q}_h = \underline{\mathbf{A}}_2(u_h^m) D^2 v_h$ in Equations (3.12a) and use (3.13a), we obtain

$$\begin{aligned} & \sum_{K \in \mathcal{T}_h} \int_K \mathbf{p}_h^m : \underline{\mathbf{A}}_2(u_h^m) D^2 v_h dx \\ & + \sum_{K \in \mathcal{T}_h} \int_K \nabla u_h^m \cdot \nabla \cdot (\omega(\nabla u_h^m)^{-1/4} \underline{\mathbf{A}}_2(u_h^m) D^2 v_h) dx \\ & - \sum_{E \in \mathcal{E}_h(\Omega)} \int_E \{\omega(\nabla u_h^m)^{-1/4} \nabla u_h^m\}_E \cdot [\underline{\mathbf{A}}_2(u_h^m) D^2 v_h]_E \mathbf{n}_E ds = 0. \end{aligned} \quad (3.16)$$

For the second term on the left-hand side of Equation (3.16), applying Green's formula element-wise and observing Equations (2.14) and (2.17) yields

$$\begin{aligned} & \sum_{K \in \mathcal{T}_h} \int_K \nabla u_h^m \cdot \nabla \cdot (\omega(\nabla u_h^m)^{-1/4} \underline{\mathbf{A}}_2(u_h^m) D^2 v_h) dx \\ & = - \sum_{K \in \mathcal{T}_h} \int_K D^2 u_h^m : \underline{\mathbf{A}}_1(u_h^m) D^2 v_h dx \\ & + \sum_{K \in \mathcal{T}_h} \int_{\partial K} \omega(\nabla u_h^m)^{-1/4} \nabla u_h^m \cdot \underline{\mathbf{A}}_2(u_h^m) D^2 v_h \mathbf{n}_{\partial K} ds. \end{aligned} \quad (3.17)$$

Now, taking advantage of

$$\begin{aligned} & \sum_{K \in \mathcal{T}_h} \int_{\partial K} \underline{\mathbf{p}} \cdot \underline{\mathbf{q}} \, \mathbf{n}_{\partial K} \, ds \\ &= \sum_{E \in \mathcal{E}_h} \int_E [\underline{\mathbf{p}}]_E \cdot \{\underline{\mathbf{q}}\}_E \mathbf{n}_E \, ds + \sum_{E \in \mathcal{E}_h(\Omega)} \int_E \{\underline{\mathbf{p}}\}_E \cdot [\underline{\mathbf{q}}]_E \mathbf{n}_E \, ds \end{aligned} \quad (3.18)$$

with $\underline{\mathbf{p}} = \omega(\nabla u_h^m)^{-1/4} \nabla u_h^m$ and $\underline{\mathbf{q}} = \underline{\mathbf{A}}_2(u_h^m) D^2 v_h$, we get

$$\begin{aligned} & \sum_{K \in \mathcal{T}_h} \int_{\partial K} \omega(\nabla u_h^m)^{-1/4} \nabla u_h^m \cdot \underline{\mathbf{A}}_2(u_h^m) D^2 v_h \mathbf{n}_{\partial K} \, ds \\ &= \sum_{E \in \mathcal{E}_h} \int_E [\omega(\nabla u_h^m)^{-1/4} \nabla u_h^m]_E \cdot \{\underline{\mathbf{A}}_2(u_h^m) D^2 v_h\}_E \mathbf{n}_E \, ds \\ &+ \sum_{E \in \mathcal{E}_h(\Omega)} \int_E \{\omega(\nabla u_h^m)^{-1/4} \nabla u_h^m\}_E \cdot [\underline{\mathbf{A}}_2(u_h^m) D^2 v_h]_E \mathbf{n}_E \, ds. \end{aligned} \quad (3.19)$$

Using Equations (3.17) and (3.19) in Equation (3.16) it follows that

$$\begin{aligned} & \sum_{K \in \mathcal{T}_h} \int_K \underline{\mathbf{p}}_h^m : \underline{\mathbf{A}}_2(u_h^m) D^2 v_h \, dx \\ &= \sum_{K \in \mathcal{T}_h} \int_K D^2 u_h^m : \underline{\mathbf{A}}_1(u_h^m) D^2 v_h \, dx \\ &- \sum_{E \in \mathcal{E}_h} \int_E [\omega(\nabla u_h^m)^{-1/4} \nabla u_h^m]_E \cdot \{\underline{\mathbf{A}}_2(u_h^m) D^2 v_h\}_E \mathbf{n}_E \, ds \end{aligned} \quad (3.20)$$

On the other hand, inserting Equation (3.13b) into Equation (3.12b) and applying again Equation (3.18), we obtain

$$\begin{aligned} & \sum_{K \in \mathcal{T}_h} \int_K u_h^m v_h \, dx + \tau_m \beta \delta^2 \left(\sum_{K \in \mathcal{T}_h} \int_K \underline{\mathbf{p}}_h^m : \underline{\mathbf{A}}_2(u_h^m) D^2 v_h \, dx \right. \\ &- \sum_{K \in \mathcal{T}_h} \int_{\partial K} \hat{\underline{\mathbf{p}}}_{\partial K}^m \cdot \nabla v_h \, ds \Big) = \sum_{K \in \mathcal{T}_h} \int_K u_h^m v_h \, dx \\ &+ \tau_m \beta \delta^2 \left(\sum_{K \in \mathcal{T}_h} \int_K \underline{\mathbf{p}}_h^m : \underline{\mathbf{A}}_2(u_h^m) D^2 v_h \, dx \right. \\ &- \sum_{E \in \mathcal{E}_h} \int_E \{\underline{\mathbf{A}}_2(u_h^m) D^2 u_h^m\}_E \mathbf{n}_E \cdot [\omega(\nabla u_h^m)^{-1/4} \nabla v_h]_E \, ds \\ &+ \alpha \sum_{E \in \mathcal{E}_h} h_E^{-1} \int_E [\omega(\nabla u_h^m)^{-1/4} \nabla u_h^m]_E \cdot [\omega(\nabla u_h^m)^{-1/4} \nabla v_h]_E \, ds \Big) \\ &= \sum_{K \in \mathcal{T}_h} \int_K u_h^{m-1} v_h \, dx. \end{aligned} \quad (3.21)$$

The assertion follows from using Equations (3.20) in (3.21). \blacksquare

Remark 3.1 We note that the C^0 IPDG approximation (Equation (3.14)) can be derived from the boundary value problem (2.1a),(2.1b) (cf., e.g., [20] for a fourth-order boundary value problem related to the biharmonic problem). However, we have chosen the approach

via the mixed formulation, since the C^0 DG approximation (3.10a),(3.10b) is more general in so far as another choice of the numerical flux functions than (3.13a),(3.13b) may lead to a different C^0 DG approach as, for instance, a C^0 Local Discontinuous Galerkin (C^0 LDG) method.

4 | A PREDICTOR CORRECTOR CONTINUATION STRATEGY FOR THE NUMERICAL SOLUTION OF THE C^0 IPDG APPROXIMATION

The solution $u(x, t)$, $(x, t) \in Q$, of the fourth-order TVF problem (1.2) is characterized by

- the formation of facets around local extrema of the initial data with steep gradients at the interfaces,
- a finite extinction time $t_{ext} > 0$, that is, $u(x, t) = 0$, $x \in \Omega$, for $t \geq t_{ext}$.

The same behavior can be expected from the solution of the C^0 IPDG approximation (3.14). In particular, the appropriate choice of the time step is a crucial issue when solving the nonlinear system resulting from (3.14). Therefore, it is more advantageous to work with a variable time step $\tau_m = t_m - t_{m-1}$, $1 \leq m \leq M$, instead of a uniform time step $\Delta t = T/M$, $M \in \mathbb{N}$ and to choose τ_m such that convergence of a Newton-type method is guaranteed. This can be achieved by viewing (3.14) as a parameter dependent nonlinear system with the time as the parameter and to apply a predictor corrector continuation strategy featuring an adaptive choice of the time step sizes τ_m (cf., e.g., [35, 36]).

We assume $V_h = \text{span}\{\varphi_1, \dots, \varphi_{N_h}\}$, $N_h \in \mathbb{N}$, such that

$$u_h^m = \sum_{j=1}^{N_h} u_j^m \varphi_j.$$

Setting $\mathbf{u}^m := (u_1^m, \dots, u_{N_h}^m)^T$, the algebraic formulation of (3.14) leads to the nonlinear system

$$\mathbf{F}(\mathbf{u}^m, t_m) = \mathbf{0}, \quad (4.1)$$

where the components \mathbf{F}_i , $1 \leq i \leq N_h$, are given by

$$\begin{aligned} \mathbf{F}_i(\mathbf{u}^m, t_m) &= \sum_{j=1}^{N_h} u_j^m (\varphi_j, \varphi_i)_{0,\Omega} \\ &\quad + \tau_m \beta \delta^2 a_h^{DG} \left(\sum_{j=1}^{N_h} u_j^m \varphi_j, \varphi_i; \sum_{j=1}^{N_h} u_j^m \varphi_j \right) - \sum_{j=1}^{N_h} u_j^{m-1} (\varphi_j, \varphi_i)_{0,\Omega}. \end{aligned}$$

Given \mathbf{u}^{m-1} , the time step size $\tau_{m-1,0} = \tau_{m-1}$, and setting $k = 0$, where k is a counter for the predictor corrector steps, the predictor step for (4.1) consists of constant continuation leading to the initial guess

$$\mathbf{u}^{(m,k)} = \mathbf{u}^{m-1}, \quad t_m = t_{m-1} + \tau_{m-1,k}. \quad (4.2)$$

Setting $\nu = 0$ and $\mathbf{u}^{(m,k,\nu_1)} = \mathbf{u}^{(m,k)}$, for $\nu \leq \nu_{max}$, where $\nu_{max} > 0$ is a pre-specified maximal number, the Newton iteration

$$\mathbf{F}'(\mathbf{u}^{(m,k,\nu)}, t_m) \Delta \mathbf{u}^{(m,k,\nu)} = -\mathbf{F}(\mathbf{u}^{(m,k,\nu)}, t_m), \quad (4.3)$$

serves as a corrector whose convergence is monitored by the contraction factor

$$\Lambda^{(m,k,\nu)} = \frac{\|\overline{\Delta \mathbf{u}^{(m,k,\nu)}}\|}{\|\Delta \mathbf{u}^{(m,k,\nu)}\|}, \quad (4.4)$$

where $\overline{\Delta u^{(m,k,v)}}$ is the solution of the auxiliary Newton step

$$\mathbf{F}'(\mathbf{u}^{(m,k,v)}, t_m) \overline{\Delta u^{(m,k,v)}} = -\mathbf{F}(\mathbf{u}^{(m,k,v+1)}, t_m). \quad (4.5)$$

If the contraction factor satisfies

$$\Lambda^{(m,k,v)} < \frac{1}{2}, \quad (4.6)$$

we set $v = v + 1$. If $v > v_{\max}$, both the Newton iteration and the predictor corrector continuation strategy are terminated indicating non-convergence. Otherwise, we continue the Newton iteration (4.3). If (4.6) does not hold true, we set $k = k + 1$ and the time step is reduced according to

$$\tau_{m,k} = \max \left(\frac{\sqrt{2} - 1}{\sqrt{4\Lambda^{(m,v)} + 1} - 1} \tau_{m,k-1}, \tau_{\min} \right), \quad (4.7)$$

where $\tau_{\min} > 0$ is some pre-specified minimal time step. If $\tau_{m,k} \geq \tau_{\min}$, we go back to the prediction step (Equation (4.2)). Otherwise, the predictor corrector strategy is stopped indicating nonconvergence. The Newton iteration is terminated successfully, if for some $v^* > 0$ the relative error of two subsequent Newton iterates satisfies

$$\frac{\|\mathbf{u}^{(m,k,v^*)} - \mathbf{u}^{(m,k,v^*-1)}\|}{\|\mathbf{u}^{(m,k,v^*)}\|} < \varepsilon \quad (4.8)$$

for some prespecified accuracy $\varepsilon > 0$. In this case, we set

$$\mathbf{u}^m = \mathbf{u}^{(m,k,v_1^*)} \quad (4.9)$$

and predict a new time step according to

$$\tau_m = \min \left(\frac{(\sqrt{2} - 1) \|\Delta u^{(m,k,0)}\|}{2\Lambda^{(m,k,0)} \|\mathbf{u}^{(m,k,0)} - \mathbf{u}^m\|}, \text{amp} \right) \tau_{m,k}, \quad (4.10)$$

where $\text{amp} > 0$ is a prespecified amplification factor for the time step sizes. We set $m = m + 1$ and begin new predictor corrector iterations for the time interval $[t_m, t_{m+1}]$. The choice of the contraction factor (Equation (4.6)), the choice of the reduced time step (Equation (4.7)), and the choice of the enlarged time step (Equation (4.10)) are motivated by the affine covariant convergence theory of Newton's method (cf., e.g., [35, 36]).

5 | NUMERICAL RESULTS

We have implemented the C^0 IPDG method of Section 3 along with the predictor corrector continuation strategy of Section 5 for two examples. In both cases, we have chosen $\Omega = (0, 1)^2$, $\beta = 1.0 \cdot 10^{-7}$, polynomial degree $k = 2$ and penalization parameter $\alpha = 200.0$ in the C^0 IPDG method, and $v_{\max} = 50$, $\varepsilon = 1.0 \cdot 10^{-3}$, and $\tau_{\min} = 1.0 \cdot 10^{-8}$ for the predictor corrector continuation strategy.

Example 1 The first example is the same as in Kohn and Versieux [3], where the initial data u^0 has been chosen according to

$$u^0(x_1, x_2) = x_1(x_1 - 1)x_2(x_2 - 1) - \frac{1}{36}.$$

The C^0 IPDG approximation u_h^m has been computed for various regularization parameters δ and finite element mesh sizes h .




FIGURE 1 Example 1: Computed solution for $h = 1/10$ and $\delta = 2.5 \cdot 10^{-4}$ at initial time $t = 0$ s (top left), at time $t = 4.6 \cdot 10^{-5}$ s (top right), at time $t = 2.6 \cdot 10^{-3}$ s, and at time $t = 1.2 \cdot 10^{-2}$ s (bottom right) [Color figure can be viewed at wileyonlinelibrary.com]

For $\delta = 2.5 \cdot 10^{-4}$ and $h = 1/10$, Figure 1 displays the initial data u_h^0 at time $t = 0.0$ (top left), and the computed solutions at times $t = 4.6 \cdot 10^{-6}$ (top right), $t = 2.6 \cdot 10^{-3}$ (bottom left), and $t = 1.2 \cdot 10^{-3}$ (bottom right). We see that the solution develops facets around local extrema of the initial data with a narrow interface featuring steep gradients in between. The extinction time, that is, the time when the initial profile gets completely flat, is $t_{ext} = 1.1 \cdot 10^{-2}$.

For $\delta = 7.5 \cdot 10^{-3}$, and $h = 1/64$, Figure 2 shows the same behavior of the solution. However, due to the significantly smaller mesh size h the interface between the upper and lower facets is much better resolved. In this case, the extinction time turned out to be $t_{ext} = 2.2 \cdot 10^{-2}$.

The performance of the predictor corrector continuation strategy for $h = 1/10$ and $\delta = 2.5 \cdot 10^{-4}$ is shown in Figure 3 where the adaptive choice of the time steps τ_m is shown as a function of the iterations. We see that the time step sizes are gradually increasing in a step-like way where the individual steps correspond to the formation of the interface between the upper and the lower facets.




FIGURE 2 Example 1: Computed solution for $h = 1/64$ and $\delta = 7.5 \cdot 10^{-3}$ at initial time $t = 0$ s (top left), at time $t = 6.9 \cdot 10^{-7}$ s (top right), at time $t = 9.5 \cdot 10^{-5}$ s, and at time $t = 1.3 \cdot 10^{-3}$ s (bottom right) [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 3 Example 1: Performance of the predictor corrector continuation strategy for $h = 1/10$ and $\delta = 2.5 \cdot 10^{-4}$. Adaptive choice of time steps τ_m [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 4 Example 1: Performance of the predictor corrector continuation strategy for $h = 1/64$ and $\delta = 7.5 \cdot 10^{-3}$. Adaptive choice of time steps τ_m [Color figure can be viewed at wileyonlinelibrary.com]

Figure 4 shows the corresponding results for the predictor corrector continuation strategy in case $h = 1/64$ and $\delta = 7.5 \cdot 10^{-3}$. The convergence is expected to become a tougher issue for smaller mesh width h which is reflected by Figure 4.

As far as the convergence of the C^0 IPDG approximations u_h^m to the solution u^m , $m \geq 1$, of Equation (2.1) is concerned, we have computed an experimental convergence rate, since the exact solutions u^m are not explicitly known. The experimental convergence rate is given by

$$\text{err}(t_m) := \text{err}_2 \frac{\|u_h^m - u_{2h}^m\|}{\|u_{h/2}^m - u_h^m\|}, \quad (5.1)$$

where $\|\cdot\|$ stands for the norm

$$\|z_h^m\| := \left(\|z_h^m\|_{0,\Omega}^2 + \sum_{\ell=1}^m \tau_\ell \|z_h^\ell\|_{2,h/2,\Omega}^2 \right)^{1/2}. \quad (5.2)$$

Figure 5 displays the experimental convergence rate as a function of time for $h = 1/64$. Taking into account that the initial data is smooth, at the very beginning the rate drops significantly due to the formation of facets with sharp interfaces between the upper and the lower facet. Afterwards it stabilizes around 0.5.

Forthcoming work will be devoted to provide an a priori error estimate in the $\|\cdot\|$ -norm.

Example 2 The initial profile u^0 has been chosen according to

$$u^0(x_1, x_2) = \begin{cases} x_1 \left(\frac{1}{2} - x_1 \right) (1 - x_1) x_2 \left(\frac{1}{2} - x_2 \right) (1 - x_2) - \frac{1}{32} & 0 \leq x_1 < \frac{1}{2}, 0 \leq x_2 < \frac{1}{2} \\ x_1 \left(x_1 - \frac{1}{2} \right) (1 - x_1) x_2 \left(\frac{1}{2} - x_2 \right) (1 - x_2) - \frac{1}{32} & \frac{1}{2} \leq x_1 \leq 1, 0 \leq x_2 < \frac{1}{2} \\ x_1 \left(\frac{1}{2} - x_1 \right) (1 - x_1) x_2 \left(x_2 - \frac{1}{2} \right) (1 - x_2) - \frac{1}{32} & 0 \leq x_1 < \frac{1}{2}, \frac{1}{2} \leq x_2 \leq 1 \\ x_1 \left(x_1 - \frac{1}{2} \right) (1 - x_1) x_2 \left(x_2 - \frac{1}{2} \right) (1 - x_2) - \frac{1}{32} & \frac{1}{2} \leq x_1 \leq 1, \frac{1}{2} \leq x_2 \leq 1 \end{cases}.$$

Example 1: Experimental convergence rate




FIGURE 5 Example 1: Experimental convergence rate $\text{err}(t_m)$ for $h = 1/64$ [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 6 Example 2: Computed solution for $h = 1/128$ and $\delta = 4.5 \cdot 10^{-3}$ at initial time $t = 0$ s (top left), at time $t = 3.4 \cdot 10^{-5}$ s (top right), at time $t = 9.7 \cdot 10^{-3}$ s (bottom left), and at time $t = 7.1 \cdot 10^{-2}$ s (bottom right) [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 7 Example 2: Performance of the predictor corrector continuation strategy for $h = 1/128$ and $\delta = 4.5 \cdot 10^{-3}$. Adaptive choice of time steps τ_m [Color figure can be viewed at wileyonlinelibrary.com]

For $\delta = 4.5 \cdot 10^{-3}$ and $h = 1/128$, Figure 6 shows the route along which the initial profile becomes completely flat. We observe the development of four upper facets around the four maxima of the initial data and lower facets around the minima. Due to the small mesh size h , the formation of narrow interfaces with steep gradients between the upper and lower facets happens quickly. The extinction time is $t_{ext} = 1.2 \cdot 10^{-2}$.

Figure 7 displays the adaptive choice of the time steps. We see a similar behavior as in Example 1 for $h = 1/64$ and $\delta = 7.5 \cdot 10^{-3}$.

6 | CONCLUSION

We have derived a C^0 IPDG approximation of an implicitly in time discretized, regularized and scaled fourth-order TVF problem describing surface relaxation below the roughening temperature. We have further developed a predictor corrector continuation strategy for the numerical solution of the fully discretized problem featuring an adaptive choice of the time steps. Numerical results have been presented that illustrate the performance of the approach.

The existence and uniqueness of a solution of the C^0 IPDG method will be shown in part II of the paper.

ACKNOWLEDGMENTS

The work of the authors has been supported by the NSF grant DMS-1520886.

ORCID

Ronald H.W. Hoppe  <https://orcid.org/0000-0003-2287-6102>

REFERENCES

- [1] W. L. Chan et al., *Relaxation kinetics of nano-ripples on cu(001) surfaces*, Phys. Rev. B vol. 70 (2004) 245403.
- [2] J. Hager, H. Spohn, *Self-similar morphology and dynamics of periodic surface profiles below the roughening transition*, Surf. Sci. vol. 324 (1995) pp. 365–372.
- [3] R. V. Kohn, H. M. Versieux, *Numerical analysis of a steepest-descent PDE model for surface relaxation below the roughening temperature*, SIAM J. Numer. Anal. vol. 48 (2010) pp. 1781–1800.
- [4] D. Margetis, M. J. Aziz, H. A. Stone, *Continuum approach to self-similarity and scaling in morphological relaxation of a crystal with a facet*, Phys. Rev. B vol. 71 (2005) 165432.
- [5] M. V. Ramana Murty, *Morphological stability of nanostructures*, Phys. Rev. B vol. 62 (2000) pp. 17004–17011.
- [6] A. Rettori, J. Villain, *Flattening of grooves on a crystal surface: A method of investigation of surface roughness*, J. Phys. France vol. 49 (1988) pp. 257–267.
- [7] V. B. Shenoy, A. Ramasubramaniam, L. B. Freund, *A variational approach to nonlinear dynamics of nanoscale surface modulations*, Surf. Sci. vol. 529 (2003) pp. 365–383.
- [8] H. Spohn, *Surface dynamics below the roughening transition*, J. Phys. I. France vol. 3 (1993) pp. 69–81.
- [9] Y. Kashima, *A subdifferential formulation of fourth order singular diffusion equations*, Adv. Math. Sci. Appl. vol. 14 (2004) pp. 49–74.
- [10] M.-H. Giga, Y. Giga, *Very singular diffusion equations: Second and fourth order problems*, Jpn. J. Ind. Appl. Math. vol. 27 (2010) pp. 323–345.
- [11] Y. Giga, R. V. Kohn, *Scale-invariant extinction time estimates for some singular diffusion equations*, Discrete Contin. Dyn. Syst. Ser A vol. 30 (2011) pp. 509–535.
- [12] Y. Kashima, “*A variational approach to very singular gradient flow equations*,” in *Proceedings of Czech-Japanese seminar in applied mathematics 2004*, Czech Technical University in Prague, Prague, Czech Republic, 2004, pp. 79–84.
- [13] Y. Kashima, *Characterization of subdifferentials of a singular convex functional in Sobolev spaces of order minus one*, J. Funct. Anal. vol. 262 (2012) pp. 2833–2860.
- [14] P. G. Ciarlet, *The finite element method for elliptic problems*, SIAM, Philadelphia, 2002.
- [15] L. Beirao da Veiga, J. Niiranen, R. Stenberg, *A family of C^0 finite elements for Kirchhoff plates I: Error analysis*, Siam J. Numer. Anal. vol. 45 (2007) pp. 2047–2071.
- [16] L. Beirao da Veiga, J. Niiranen, R. Stenberg, *A family of C^0 finite elements for Kirchhoff plates II: Numerical results*, Comput. Meth. Appl. Mech. Eng. vol. 197 (2008) pp. 1850–1864.
- [17] D. Braess, R. H. W. Hoppe, C. Linsenmann, *A two-energies principle for the biharmonic equation and an a posteriori error estimator for an interior penalty discontinuous Galerkin approximation*, ESAIM: M2AN vol. 52 (2019) pp. 2479–2504.
- [18] S. C. Brenner et al., *A C^0 interior penalty method for a von Kármán plate*, Numer. Math. vol. 135 (2017) pp. 803–832.
- [19] S. C. Brenner et al., “*A C^0 interior penalty method for elliptic distributed optimal control problems in three dimensions with pointwise state constraints*,” in *Topics in numerical partial differential equations and scientific computing*, IMA volumes in mathematics and its applications, vol. 160, S. C. Brenner (Editor), Springer, New York, 2016, pp. 1–22.
- [20] S. C. Brenner, L.-Y. Sung, *C^0 interior penalty methods for fourth order elliptic boundary value problems on polygonal domains*, J. Sci. Comput. vol. 22(23) (2005) pp. 83–118.
- [21] G. Engel et al., *Continuous/discontinuous finite element approximations of fourth order elliptic problems in structural and continuum mechanics with applications to thin beams and plates, and strain gradient elasticity*, Comput. Methods Appl. Mech. Eng. vol. 191 (2002) pp. 3669–3750.
- [22] E. H. Georgoulis, P. Houston, *Discontinuous Galerkin methods for the biharmonic problem*, IMA J. Numer. Anal. vol. 29 (2009) pp. 573–594.
- [23] R. H. W. Hoppe, C. Linsenmann, “ *C^0 interior penalty discontinuous Galerkin approximation of a sixth order Cahn-Hilliard equation modeling microemulsification processes*,” in *Contributions to partial differential equations and applications*, Chapter 16, ECCOMAS series computational methods in applied sciences, B. N. Chetveruskin et al. (Editors), Springer, Berlin, 2019.
- [24] X. Huang, J. Huang, *A reduced local C^0 discontinuous Galerkin method for Kirchhoff plates*, Numer. Meth. Part. Diff. Equat. vol. 30 (2014) pp. 1902–1930.
- [25] I. Mozolevski, E. Süli, *A priori error analysis for the hp-version of the discontinuous Galerkin finite element method for the biharmonic equation*, Comput. Methods Appl. Math. vol. 3 (2003) pp. 1–12.
- [26] I. Mozolevski, E. Süli, P. R. Bösing, *Hp-version a priori error analysis of interior penalty discontinuous Galerkin finite element approximations to the biharmonic equation*, J. Sci. Comput. vol. 30 (2007) pp. 465–491.
- [27] E. Süli, I. Mozolevski, *Hp-version interior penalty DGFEMs for the biharmonic equation*, Comput. Methods Appl. Mech. Eng. vol. 196 (2007) pp. 1851–1863.
- [28] G. N. Wells, E. Kuhl, K. Garikipati, *A discontinuous Galerkin method for the Cahn-Hilliard equation*, J. Comput. Phys. vol. 218 (2006) pp. 860–877.

- [29] Y. Meyer, *Oscillating patterns in image processing and nonlinear evolution equations*. University Lecture Series No. 22, AMS, Providence, 2002.
- [30] S. Osher, A. Solé, L. A. Vese, *Image decomposition and restoration using total variation minimization and the H^{-1} Norm*, Multiscale Model. Simul. vol. 1 (2003) pp. 349–370.
- [31] L. A. Vese, S. Osher, *Modeling textures with total variation minimization and oscillating patterns in image processing*, J. Sci. Comput. vol. 19 (2003) pp. 553–572.
- [32] L. A. Vese, S. Osher, *Image denoising and decomposition with total variation minimization and oscillatory functions*, J. Math. Imaging Vision vol. 20 (2004) pp. 7–18.
- [33] L. Tartar, *Introduction to sobolev spaces and interpolation theory*, Springer, Berlin, 2007.
- [34] D. Arnold et al., *Unified analysis of discontinuous Galerkin methods for elliptic problems*, SIAM J. Numer. Anal. vol. 39 (2002) pp. 1749–1779.
- [35] P. Deufilhard, *Newton methods for nonlinear problems - Affine invariance and adaptive algorithms*, Springer, Berlin, 2004.
- [36] R. H. W. Hoppe, C. Linsenmann, *An adaptive Newton continuation strategy for the fully implicit finite element immersed boundary method*, J. Comput. Phys. vol. 231 (2012) pp. 4676–4693.