

From domain decomposition to homogenization theory

Daniel Peterseim, D. Varga, Barbara Verfürth

Angaben zur Veröffentlichung / Publication details:

Peterseim, Daniel, D. Varga, and Barbara Verfürth. 2020. "From domain decomposition to homogenization theory." *Lecture Notes in Computational Science and Engineering* 138: 29–40. https://doi.org/10.1007/978-3-030-56750-7_3.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



From Domain Decomposition to Homogenization Theory

Daniel Peterseim and Dora Varga and Barbara Verfürth

Abstract This paper rediscovers a classical homogenization result for a prototypical linear elliptic boundary value problem with periodically oscillating diffusion coefficient. Unlike classical analytical approaches such as asymptotic analysis, oscillating test functions, or two-scale convergence, the result is purely based on the theory of domain decomposition methods and standard finite elements techniques. The arguments naturally generalize to problems far beyond periodicity and scale separation and we provide a brief overview on such applications.

1 Introduction

Elliptic boundary value problems with oscillatory coefficients play a key role in the mathematical modelling and simulation of complex multiscale problems, for instance transport processes in porous media or the mechanical analysis of composite and multifunctional materials. The characteristic properties of such processes are determined by a complex interplay of effects on multiple non-separable length and time scales. The challenge is that the resolution of all details on all relevant scales may easily lead to a number of degrees of freedom and computational work in a direct numerical simulation which exceed today's computing resources by multiple orders of magnitude. The observation and prediction of physical phenomena from multiscale models, hence, requires insightful methods that effectively represent unresolved scales, i.e., multiscale methods.

Daniel Peterseim
Universität Augsburg e-mail: daniel.peterseim@math.uni-augsburg.de

Dora Varga
Universität Augsburg e-mail: dora.varga@math.uni-augsburg.de

Barbara Verfürth
Universität Augsburg e-mail: barbara.verfuerth@math.uni-augsburg.de

Homogenization is such a multiscale method. It seeks a simplified model that is able to capture the macroscopic responses of the process adequately by a few localized computations on the microscopic scale. Consider, e.g., prototypical second order linear elliptic model problems with highly oscillatory periodic diffusion coefficients that oscillate at frequency ε^{-1} for some small parameter $0 < \varepsilon \ll 1$. Then, the theory of homogenization shows that there exists a constant coefficient such that the corresponding diffusion process represents the macroscopic behaviour correctly. In practice, this yields a two- or multi-scale method that first computes the effective coefficient which is implicitly given through some PDE on the microscopic periodic cell, and then solves the macroscopic effective PDE. This is done for instance in the Multiscale Finite Element Method [8] or the Heterogeneous Multiscale Method [1]. In certain cases, the error of such procedures can be quantified in terms of the microscopic length scale ε . The approach and its theoretical foundation can be generalized to certain classes of non-periodic problems. However, the separation of scales, i.e., the separation of the characteristic frequencies of the diffusion coefficient and macroscopic frequencies of interest, seems to be essential for both theory and computation.

There is a more recent class of numerical homogenization methods that can deal with arbitrarily rough diffusion coefficients beyond scale separation [31, 23]. While, at first glance, these methods seemed to be only vaguely connected to classical homogenization theory, the recent paper [14] identifies them as a natural generalization of some new characterization of classical homogenization. Another deep connection, which was always believed to exist in the community of domain decomposition methods, is the one between homogenization and domain decomposition. This one was made precise only recently by Kornhuber and Yserentant [28, 26, 27]. By combining their iterative approach to homogenization and the results of [14], the present paper illuminates the role of domain decomposition in the theory of homogenization and provides homogenization limits without any advanced compactness arguments or two scale limits. In addition, compared with [14], we are able to drop a technical assumption on some artificial symmetries of the diffusion coefficient with respect to the periodic cell.

Our new construction of effective coefficients (see Sections 3–4) is not necessarily any easier than the classical one. For the simple diffusion model problem, this is merely an instance of mathematical curiosity and we do not mean to rewrite homogenization theory. However, the connection between homogenization theory and domain decomposition and, in particular, the method of proof turn out to be very interesting and, moreover, they unroll their striking potential for problems beyond scale separation and periodicity. Using this approach, new theoretical results could be derived and some of them are briefly discussed in Section 5.

2 Model problem and classical homogenization

For the sake of illustration we restrict ourselves to the simplest possible yet representative and relevant setting. Let $\Omega = [0, 1]^2$ be the unit square and $\varepsilon\Omega := [0, \varepsilon]^2$. Moreover, let $A_1 \in L^\infty(\Omega; \mathbb{R}^{2 \times 2})$ be a symmetric, uniformly elliptic, Ω -periodic (matrix-valued) coefficient and let $A_\varepsilon(x) := A_1(\frac{x}{\varepsilon})$, $x \in \Omega$. The extension to a cuboidal domain in $3D$ is straight forward. We denote by $V := H_\#^1(\Omega)/\mathbb{R}$ the equivalence class of Ω -periodic functions in $H^1(\Omega)$ factorised by constants, and similarly, by $V_\varepsilon := H_\#^1(\varepsilon\Omega)/\mathbb{R}$ for their ε -periodic counterparts. The model problem under consideration then reads: given $f \in L^2(\Omega)$, find a function $u_\varepsilon \in V$ such that

$$\int_{\Omega} A_\varepsilon(x) \nabla u_\varepsilon(x) \cdot \nabla v(x) \, dx = \int_{\Omega} f(x) v(x) \, dx, \quad (1)$$

for all $v \in V$. In order to ensure the well-posedness of the problem, we assume that $A_\varepsilon \in \mathcal{M}_{\alpha\beta}$, where $\mathcal{M}_{\alpha\beta}$ is defined as

$$\mathcal{M}_{\alpha\beta} := \{A \in L^\infty(\Omega) \mid \alpha|\xi|^2 \leq \xi \cdot A(x)\xi \leq \beta|\xi|^2 \text{ for all } \xi \in \mathbb{R}^2 \text{ and a.a. } x \in \Omega\}.$$

The idea behind classical homogenization is to look for a so-called effective (homogenized) coefficient $A_0 \in \mathcal{M}_{\alpha\beta}$ so that the solution $u_0 \in V$ of the problem

$$\int_{\Omega} A_0 \nabla u_0 \cdot \nabla v \, dx = \int_{\Omega} f v \, dx, \quad (2)$$

for all $v \in V$, represents the limit of the sequence $\{u_\varepsilon\}_{\varepsilon>0}$ of solutions of the problem (1). In general, explicit representations of effective coefficients are not known, except for the simple case of the one-dimensional or (locally) periodic setting. However, the so-called energy method of Murat and Tartar ([34]) or the two-scale convergence ([3]) provide us with the following form

$$(A_0)_{kj} = \int_{\Omega} \left(A_1(x)(e_j + \nabla w_j(x)) \right) \cdot \left(e_k + \nabla w_k(x) \right) \, dx, \quad (3)$$

where w_j are defined as the unique solutions in V of the so-called cell problems

$$\int_{\Omega} A_1(x) (\nabla w_j(x) - e_j) \cdot \nabla v(x) \, dx = 0,$$

for all $v \in V$, with the canonical basis $(e_j)_{j=1}^2$ of \mathbb{R}^2 . The substitution $x \mapsto \frac{x}{\varepsilon}$ yields

$$\begin{aligned} 0 &= \varepsilon^{-2} \int_{\varepsilon\Omega} A_1\left(\frac{x}{\varepsilon}\right) \underbrace{\left(\nabla w_j\left(\frac{x}{\varepsilon}\right) - e_j\right)}_{=: \hat{q}_j(x)} \cdot \underbrace{\nabla v\left(\frac{x}{\varepsilon}\right)}_{=: v_\varepsilon(x)} \, dx \\ &= \int_{\Omega} A_\varepsilon(x) (\nabla \hat{q}_j(x) - e_j) \cdot \nabla v_\varepsilon(x) \, dx. \end{aligned} \quad (4)$$

Since all functions v_ε in V_ε can be written as $v(\frac{x}{\varepsilon})$ for a certain function $v \in V$, equation (4) yields that the function $\hat{q}_j \in V_\varepsilon$ solves

$$\int_{\Omega} A_\varepsilon(x)(\nabla \hat{q}_j(x) - e_j) \cdot \nabla v_\varepsilon(x) \, dx = 0, \quad (5)$$

for all $v_\varepsilon \in V_\varepsilon$. Moreover, $\hat{q}_j \in V_\varepsilon \subset V$ solves the same problem in the space V , i.e.,

$$\int_{\Omega} A_\varepsilon(x)(\nabla \hat{q}_j(x) - e_j) \cdot \nabla v(x) \, dx = 0,$$

for all $v \in V$, since the solution of an elliptic model problem with periodic data (coefficient, source function) is also periodic, with the same period.

3 Novel characterization of the effective coefficient

In order to define the effective coefficient from the alternative perspective of finite elements, we first introduce the necessary notation on meshes, spaces, and interpolation operators.

We consider structured triangulations of $\Omega = [0, 1]^2$ as depicted in Figure 1, where the triangles T form the triangulation \mathcal{T}_H and the boldface squares Q are part of the square mesh Q_H . The theoretical arguments below require the triangulation to be aligned with the periodicity cells of the coefficient represented by the elements of Q_H . Moreover, \mathcal{T}_H should not introduce any nodes in the interiors of those cells. We shall emphasize that the general numerical homogenization method of Section 5 can deal with fairly general meshes. Denote the set of nodes by $\mathcal{N}_{\mathcal{T}_H} = \mathcal{N}_{Q_H}$. Since we are working with periodic boundary conditions, we will frequently understand Q_H and \mathcal{T}_H as periodic partitions (or partitions of the torus or partitions of the whole \mathbb{R}^2), i.e., we identify opposite faces of the unit square. The parameter H denotes the length of the quadrilaterals and is supposed to be not smaller than the microscopic length scale ε of the model problem.




Fig. 1 Admissible triangulations.

Let $\mathcal{P}_1(\mathcal{T}_H)$ denote the space of globally continuous piecewise affine functions on Ω with periodic boundary conditions. As in the continuous case with V , we also factor out the constants here, i.e., in fact we consider $(\mathcal{P}_1(\mathcal{T}_H))_{/\mathbb{R}}$, but still write $\mathcal{P}_1(\mathcal{T}_H)$ for simplicity. Since $\varepsilon \leq H$ is assumed, the finite element method with the space $\mathcal{P}_1(\mathcal{T}_H)$ does not yield faithful approximations of the solution u_ε to (1); see, e.g., [37, Sec. 1]. We introduce a bounded local linear projection operator $I_H: V \rightarrow \mathcal{P}_1(\mathcal{T}_H)$, which can be seen as a composition $I_H := E_H \circ \Pi_H$, where a function $v \in V$ is first approximated on every element $T \in \mathcal{T}_H$ by its L^2 -orthogonal projection Π_H onto the space of affine functions. Hence, a possibly globally discontinuous function $\Pi_H v$ is obtained. In the second step E_H , the values at the inner vertices of the triangulation are averages of the respective contributions from the single elements, i.e.,

$$E_H \circ \Pi_H(v)(z) := \frac{1}{\#\{T \in \mathcal{T}_H, z \in T\}} \sum_{\substack{T \in \mathcal{T}_H \\ z \in T}} \Pi_H(v)|_T(z)$$

for all vertices z , where the triangulation is understood in a periodic manner, see [35].

Let $W := \ker I_H$ be the kernel of the quasi-interpolation operator I_H . It can be seen as the set of rapidly oscillating functions, which cannot be captured by standard finite elements functions on the (coarse) mesh \mathcal{T}_H . Motivated by the reformulation (5) of the cell problems and the interpretation of W as rapidly oscillating functions, we define now the correctors $q_{Q,j}^\infty$ as the unique solutions in W of the following variational problems

$$\int_{\Omega} A_\varepsilon(x) \nabla q_{Q,j}^\infty(x) \cdot \nabla w(x) \, dx = \int_Q A_\varepsilon(x) e_j \cdot \nabla w(x) \, dx, \quad (6)$$

for all $w \in W$, and the correctors are defined for every $Q \in \mathcal{Q}_H$, $j = 1, 2$. We define the following w.r.t. \mathcal{Q}_H piecewise constant numerical coefficient A_H^∞ which will play the main role in Proposition 1:

$$\left[A_H^\infty|_Q \right]_{kj} = \frac{1}{|Q|} \int_Q A_\varepsilon(x) e_j \cdot e_k \, dx - \frac{1}{|Q|} \int_{\Omega} A_\varepsilon(x) \nabla q_{Q,j}^\infty(x) \cdot e_k \, dx, \quad (7)$$

for all $Q \in \mathcal{Q}_H$, $k, j = 1, 2$.

Proposition 1 *In the case that the mesh size H is an integer multiple of ε , the coefficient A_H^∞ coincides with the homogenized coefficient A_0 from classical homogenization defined in (3).*

Proof We will first show that the function $q_j := \sum_{Q \in \mathcal{Q}_H} q_{Q,j}^\infty$ coincides with the corrector $\hat{q}_j \in V_\varepsilon$, the unique solution of the problem (5). The crucial observation needed for the proof is the fact that the space of ε -periodic functions is contained in the kernel W of the quasi-interpolation operator I_H , in the case of the present setting with the triangulations \mathcal{T}_H and \mathcal{Q}_H . To see this we observe that, for an ε -periodic function $v_\varepsilon \in V_\varepsilon$, the values $I_H(v_\varepsilon)(z)$ coincide for all $z \in \mathcal{N}_{\mathcal{T}_H}$. That is,

$I_H(v_\varepsilon) \in \mathcal{P}_1(\mathcal{T}_H)$ is a global constant. As we factored out the constants, we can take the zero function as representative, i.e., $I_H(v_\varepsilon) = 0$.

Moreover, summing up the equations (6) over all $Q \in \mathcal{Q}_H$ and taking advantage of the symmetry of A_ε , we get that $q_j := \sum_{Q \in \mathcal{Q}_H} q_{Q,j}^\infty$ solves

$$\int_{\Omega} A_\varepsilon(x) \nabla q_j(x) \cdot \nabla w(x) \, dx = \int_{\Omega} A_\varepsilon(x) e_j \cdot \nabla w(x) \, dx \quad (8)$$

$$= \int_{\Omega} A_\varepsilon(x) \nabla w(x) \cdot e_j \, dx, \quad (9)$$

for all $w \in W$, and in particular for all $w \in V_\varepsilon$. The combination of (5) and (9) readily yields that $q_j \equiv \hat{q}_j$, $j = 1, 2$. Moreover, (9) with $w = q_{Q,k}^\infty$ implies

$$\begin{aligned} \int_{\Omega} A_\varepsilon(x) \nabla q_{Q,k}^\infty(x) \cdot \nabla e_j \, dx &= \int_{\Omega} A_\varepsilon(x) \nabla q_j(x) \cdot \nabla q_{Q,k}^\infty(x) \, dx \\ &= \int_{\Omega} A_\varepsilon(x) \nabla q_{Q,k}^\infty(x) \cdot \nabla q_j(x) \, dx \\ &= \int_Q A_\varepsilon(x) e_k \cdot \nabla q_j(x) \, dx. \end{aligned}$$

Hence, in the definition of A_H^∞ we can replace the second term, namely

$$\begin{aligned} \left(A_{H|Q}^\infty \right)_{kj} &= \frac{1}{|Q|} \int_Q A_\varepsilon(x) e_j \cdot e_k \, dx - \frac{1}{|Q|} \int_Q A_\varepsilon(x) e_j \cdot \nabla q_k(x) \, dx \\ &= \frac{1}{|Q|} \int_Q A_\varepsilon(x) e_j \cdot (e_k - \nabla \hat{q}_k(x)) \, dx \\ &= \left(A_0 \right)_{kj}, \end{aligned}$$

for $j, k = 1, 2$. □

4 Numerical effective coefficient by domain decomposition

The correctors $q_{Q,j}^\infty$ defined in the previous section require the solution of a global problem involving the oscillating coefficient A_ε . Employing domain decomposition, we introduce localized variants and then use arguments from the theory of iterative (domain decomposition) methods as presented in [26, 28] to show that the error decays exponentially in the number of iterations. With the localized correctors, we then introduce an effective localized coefficient A_H^ℓ which is piecewise constant on \mathcal{Q}_H .

Let ω_i be the union of all squares $Q \in \mathcal{Q}_H$ having the vertex z_i as a corner and let

$$W_i = \{v - I_H v \mid v \in H_0^1(\omega_i)\}.$$

We emphasize that ω_i is understood as a subset of \mathbb{R}^2 , i.e., it is continued over the periodic boundary. The functions in W_i vanish outside a small neighbourhood of the vertex z_i . The W_i are closed subspaces of the kernel W of I_H , see [26]. Let P_i be the a_ε -orthogonal projection from V to W_i , defined via the equation

$$a_\varepsilon(P_i v, w_i) = a_\varepsilon(v, w_i), \quad \forall w_i \in W_i.$$

Introducing the with respect to the bilinear form $a_\varepsilon(\cdot, \cdot)$ symmetric operator

$$P = P_1 + P_2 + \cdots + P_n,$$

the following properties are proved in [26]:

Lemma 1 *There are constants K_1 and K_2 , independent of H and ε , such that*

$$K_1^{-1} a_\varepsilon(v, v) \leq a_\varepsilon(Pv, v) \leq K_2 a_\varepsilon(v, v)$$

for all $v \in V$. Moreover, for an appropriate scaling factor ϑ only depending on K_1 and K_2 , there exists a positive constant $\gamma < 1$ such that

$$\| \text{id} - \vartheta P \|_{\mathcal{L}(V, V)} \leq \gamma. \quad (10)$$

Starting from $q_{Q,j}^0 = 0$, $j = 1, 2$, the localized correctors $q_{Q,j}^\ell$ are defined for all $Q \in \mathcal{Q}_H$ via

$$q_{Q,j}^{\ell+1} = q_{Q,j}^\ell + \vartheta P(x_j 1_Q - q_{Q,j}^\ell), \quad j = 1, 2, \quad (11)$$

where 1_Q denotes the characteristic function of Q and x_j denotes the j -th component of the (vector-valued) function $x \mapsto x$. The scaling factor ϑ is chosen as discussed in Lemma 1. The correction $P(x_j 1_Q - q_{Q,j}^\ell)$ is the sum of its components $C_i^\ell = P_i(x_j 1_Q - q_{Q,j}^\ell)$ in the subspaces W_i of W , where the C_i^ℓ solve the local equations

$$a_\varepsilon(C_i^\ell, w_i) = a_\varepsilon(x_j 1_Q, w_i) - a_\varepsilon(q_{Q,j}^\ell, w_i), \quad \forall w_i \in W_i. \quad (12)$$

The sloppy notation using 1_Q as argument in a_ε is to denote that the integration is over the element Q only, i.e., $a_\varepsilon(x_j 1_Q, w_i) = \int_Q A_\varepsilon e_j \cdot \nabla w_i \, dx$. Since the local projections P_i only slightly increase the support of a function, we deduce inductively that the support of $q_{Q,j}^\ell$ is contained in an ℓH -neighbourhood of Q . In particular, in each step of (11) only a few local problems of type (12) have to be solved.

We now replace $q_{Q,j}^\infty$ by its localized variant $q_{Q,j}^\ell$ in the definition of the numerical effective coefficient. This procedure is justified by an exponential error estimate in Proposition 2. We define the piecewise constant (on the mesh \mathcal{Q}_H) (localized) effective matrix A_H^ℓ via

$$(A_H^\ell|_Q)_{kj} = \frac{1}{|Q|} \int_Q A_\varepsilon(x) e_j \cdot e_k \, dx - \frac{1}{|Q|} \int_\Omega A_\varepsilon \nabla q_{Q,j}^\ell(x) \cdot e_k \, dx. \quad (13)$$

Since the numerical effective coefficient (7) is the “true” one in the sense that $A_H^\infty = A_0$, we simply need to estimate the error of the iterative approximation.

Proposition 2 *Let H be an integer multiple of ε and let the localization parameter ℓ be chosen of order $\ell \approx |\log H|$. Then,*

$$\|A_H^\infty - A_H^\ell\|_{L^\infty(\Omega)} \lesssim H. \quad (14)$$

Proof We first estimate the error between the correctors $q_{Q,j}^\infty$ and $q_{Q,j}^\ell$. Using the definition of $q_{Q,j}^\infty$ in (6), we deduce that $P(x_j 1_Q) = P(q_{Q,j}^\infty)$. Hence, we can characterize the error between the correctors $q_{Q,j}^\infty$ and their localized approximations $q_{Q,j}^\ell$ via

$$q_{Q,j}^\infty - q_{Q,j}^\ell = (\text{id} - \vartheta P)^\ell q_{Q,j}^\infty.$$

Using (10), this yields the exponential convergence of $q_{Q,j}^\ell$ towards $q_{Q,j}^\infty$, i.e.,

$$\|\nabla(q_{Q,j}^\infty - q_{Q,j}^\ell)\| \lesssim \gamma^\ell \|\nabla q_{Q,j}^\infty\| \lesssim \gamma^\ell |Q|^{1/2}. \quad (15)$$

By the definitions of A_H^∞ in (7) and A_H^ℓ in (13), we obtain

$$\begin{aligned} |(A_H^\infty|_Q)_{jk} - (A_H^\ell|_Q)_{jk}| &= |Q|^{-1} \left| \int_\Omega A_\varepsilon \nabla(q_{Q,j}^\ell - q_{Q,j}^\infty) \cdot e_k \, dx \right| \\ &\lesssim |Q|^{-1} \|e_k\|_{L^2(\Omega)} \|\nabla(q_{Q,j}^\ell - q_{Q,j}^\infty)\|_{L^2(\Omega)}. \end{aligned}$$

Estimate (15) and the choice $\ell \approx |\log H|$ readily imply the assertion. \square

The same estimate was previously derived in [14] with a slightly different localization strategy and with more restrictive conditions on the triangulation. There, the homogenization error in the L^2 -norm is quantified as follows. Let Ω be convex. Let $u_\varepsilon \in V$ solve (1) and let $u_0 \in V$ be the solution to (2). For sufficiently small ε , it holds that

$$\|u_\varepsilon - u_0\|_{L^2(\Omega)} \lesssim \varepsilon |\log \varepsilon|^2 \|f\|_{L^2(\Omega)}.$$

This estimate recovers the classical result that $u_\varepsilon \rightarrow u_0$ strongly in L^2 and furthermore states that the convergence is almost linear for right-hand sides $f \in L^2(\Omega)$. We shall emphasize that the proofs of [14] are solely based on standard techniques of finite elements. The authors believe that such a result is also possible in the slightly more general setup of this paper. However, it seems that there is no simple argument but the generalization requires to revise the analysis of [14] step by step which is far beyond the scope of this paper.

5 Beyond periodicity and scale separation

The numerical approach presented in Section 4 does not essentially rely on the assumption of periodicity or separation of scales (between the length scales of

the computational domain and the material structures). Of course, in such general situations, one cannot identify a constant effective coefficient. Instead the goal is to faithfully approximate the analytical solution by a (generalized) finite element method based on a (coarse) mesh, which does not need to resolve the fine material structures and thereby is computationally efficient.

For this generalization, note that the definition (11) can be formulated verbatim for any boundary value problem involving a potentially rough, but not necessarily periodic diffusion tensor $A \in L^\infty(\Omega)$. Moreover, the choice of the function $x_j 1_Q$ in the definition of $q_{Q,j}^\ell$ can be generalized to any function $v \in V$ in the following way. Define the operator $C_T^\ell : V \rightarrow W$ inductively via $C_T^0 = 0$ and

$$C_T^{\ell+1} = C_T^\ell + \vartheta P(\text{id}|_T - C_T^\ell)$$

for all $T \in \mathcal{T}_H$, see [26]. Instead of modifying the diffusion tensor as in the previous sections, we then modify the basis functions and define a generalized finite element method using the test and ansatz spaces $V_H^\ell := (\text{id} + C^\ell)\mathcal{P}_1(\mathcal{T}_H)$ with $C^\ell := \sum_{T \in \mathcal{T}_H} C_T^\ell$. This method is known as the Localized Orthogonal Decomposition (LOD) [23, 31, 19, 37] and originally arose from the concept of the Variational Multiscale Method [24, 25]. Note that mostly a slightly different definition of the correctors C_T^ℓ based on patches of diameter ℓH around the element T is used. The present approach via domain decomposition and iterative solvers was developed recently in [28, 26]. It has been shown in [31, 19] for instance, that the method approximates the analytical solution with an energy error of the order H even in the pre-asymptotic regime if the localization parameter ℓ is chosen of the order $\ell \approx |\log H|$ as in Proposition 2. Hence, the Localized Orthogonal Decomposition can efficiently treat general multiscale problems. Besides the above mentioned Galerkin-type ansatz with modified ansatz and test functions, Petrov-Galerkin formulations of the method [9] may have computational advantages [10] and even meshless methods are possible [21].

The Localized Orthogonal Decomposition is not restricted to elliptic diffusion problems and has underlined its potential in various applications and with respect to different (computational) challenges. Starting from the already mentioned application in the geosciences, we underline that the material coefficients are often characterized not only by rapid oscillations but also by a high contrast, i.e., the ratio β/α is large. Many error estimates, also for the standard LOD, are contrast-dependent, but a careful choice of the interpolation operator, see [17, 40], can overcome this effect. Apart from simple diffusion problems, porous media [7], elasticity problems [22] or coupling of those such as in poroelasticity [4] play important roles in these (and many other) applications. For instance in elasticity theory, not only heterogeneous materials are treated, but also the effect of locking can be reduced by the multiscale method in [22].

Another important area of research are acoustic and electromagnetic wave propagation problems, where the considered prototypical equations are the Helmholtz and Maxwell's equations. It is well known that standard finite element discretizations of the (indefinite) Helmholtz equation are only well-posed and converging under a

rather restrictive resolution condition between the mesh size and the wavenumber. In a series of paper [6, 13, 38], it was analysed that the LOD can relax this resolution condition if the localization parameter grows logarithmically with the wavenumber. For large wavenumbers, this is a great computational gain in comparison to standard numerical methods that even allows the simulation of physical phenomena in high contrast regimes [41]. Maxwell's equations, studied in [12, 42], on the other hand, pose a challenge as the involved curl-operator has a large kernel. Moreover, the natural finite element space are Nédélec's edge elements, for which stable interpolation operators are much less developed than for Lagrange finite elements. In the context of problems not based on standard Lagrange spaces, we also mention the mixed problem utilizing Raviart-Thomas spaces in [16]. Considering wave problems, the time-dependent wave equation with different time discretizations was studied in [2, 30]. Concerning time-dependency, an important question for the LOD construction is how to deal with time-dependent diffusion tensors. [18] presents an a posteriori error estimator in order to adaptively decide which correction to recompute in the next time step.

Apart from the treatment of multiscale coefficients in a variety of partial differential equations, the methodology can also be seen as a stabilization scheme similar as its origin in the variational multiscale methods. This has been exploited to deal with the pollution effect in Helmholtz problems mentioned above, for convection dominated diffusion problems [29] and, more importantly, to bypass CFL conditions in the context of explicit wave propagation on adaptive meshes [39].

Further unexpected applications are linear and nonlinear eigenvalue problems [32, 33], in particular the quantum-physical simulation based on the Gross-Pitaevskii equation. While the LOD can be employed to speed-up ground state computations for rather rough potentials [20], the underlying technique of localization by domain decomposition turned out to be of great value to provide (analytical) insight into the phenomenon of Anderson localization in this context. The recent paper [5] predicts and quantifies the emergence of localized eigenstates and might inspire progress regarding the understanding of localization effects which are observed for many other problems as well.

The present contribution aimed at unifying the view of the LOD and classical homogenization and domain decomposition. As already mentioned, close connections exist with [14] and its extension to stochastic homogenization [15]. Further applications involve a multilevel generalization of LOD named gambles [36] (due to a possible game-theoretic interpretation). This multilevel variant allows surprising results such as a sparse representation of the expected solution operator for random elliptic boundary value problems [11] which may inspire new computational strategies for uncertainty quantification in the future.

Acknowledgements This work was supported in part by the German Research Foundation (DFG) through the Priority Programme 1648 "Reliable simulation techniques in solid mechanics" under (PE2143/2-2).

References

1. Abdulle, A., E, W., Engquist, B., Vanden-Eijnden, E.: The heterogeneous multiscale method. *Acta Numer.* **21**, 1–87 (2012). DOI 10.1017/S0962492912000025
2. Abdulle, A., Henning, P.: Localized orthogonal decomposition method for the wave equation with a continuum of scales. *Math. Comp.* **86**(304), 549–587 (2017). DOI 10.1090/mcom/3114
3. Allaire, G.: Homogenization and two-scale convergence. *SIAM J. Math. Anal.* **23**(6), 1482–1518 (1992). DOI 10.1137/0523084
4. Altmann, R., Chung, E., Maier, R., Peterseim, D., Pun, S.: Computational multiscale methods for linear heterogeneous poroelasticity. *ArXiv preprint* **1801.00615** (2019). Accepted for publication in *J. Comput. Math.*
5. Altmann, R., Henning, P., Peterseim, D.: Quantitative anderson localization of schrödinger eigenstates under disorder potentials. *ArXiv preprint* **1803.09950** (2018)
6. Brown, D.L., Gallistl, D., Peterseim, D.: Multiscale Petrov-Galerkin method for high-frequency heterogeneous Helmholtz equations. In: *Meshfree methods for partial differential equations VIII, Lect. Notes Comput. Sci. Eng.*, vol. 115, pp. 85–115. Springer, Cham (2017)
7. Brown, D.L., Peterseim, D.: A multiscale method for porous microstructures. *Multiscale Model. Simul.* **14**(3), 1123–1152 (2016). DOI 10.1137/140995210
8. Efendiev, Y., Hou, T.Y.: Multiscale finite element methods, *Surveys and Tutorials in the Applied Mathematical Sciences*, vol. 4. Springer, New York (2009). Theory and applications
9. Elfverson, D., Ginting, V., Henning, P.: On multiscale methods in Petrov-Galerkin formulation. *Numer. Math.* **131**(4), 643–682 (2015). DOI 10.1007/s00211-015-0703-z
10. Engwer, C., Henning, P., Målqvist, A., Peterseim, D.: Efficient implementation of the localized orthogonal decomposition method. *Comput. Methods Appl. Mech. Engrg.* **350**, 123–153 (2019). DOI 10.1016/j.cma.2019.02.040
11. Feischl, M., Peterseim, D.: Sparse compression of expected solution operators. *ArXiv preprint* **1807.01741** (2018)
12. Gallistl, D., Henning, P., Verfürth, B.: Numerical homogenization of $\mathbf{H}(\text{curl})$ -problems. *SIAM J. Numer. Anal.* **56**(3), 1570–1596 (2018). DOI 10.1137/17M1133932
13. Gallistl, D., Peterseim, D.: Stable multiscale Petrov-Galerkin finite element method for high frequency acoustic scattering. *Comput. Methods Appl. Mech. Engrg.* **295**, 1–17 (2015). DOI 10.1016/j.cma.2015.06.017. URL <https://doi.org/10.1016/j.cma.2015.06.017>
14. Gallistl, D., Peterseim, D.: Computation of quasi-local effective diffusion tensors and connections to the mathematical theory of homogenization. *Multiscale Model. Simul.* **15**(4), 1530–1552 (2017). DOI 10.1137/16M1088533
15. Gallistl, D., Peterseim, D.: Numerical stochastic homogenization by quasi-local effective diffusion tensors. *ArXiv preprint* **1702.08858** (2019). Accepted for publication in *Communications in Mathematical Sciences*
16. Hellman, F., Henning, P., Målqvist, A.: Multiscale mixed finite elements. *Discrete Contin. Dyn. Syst. Ser. S* **9**(5), 1269–1298 (2016). DOI 10.3934/dcdss.2016051
17. Hellman, F., Målqvist, A.: Contrast independent localization of multiscale problems. *Multiscale Model. Simul.* **15**(4), 1325–1355 (2017). DOI 10.1137/16M1100460
18. Hellman, F., Målqvist, A.: Numerical homogenization of elliptic PDEs with similar coefficients. *Multiscale Model. Simul.* **17**(2), 650–674 (2019). DOI 10.1137/18M1189701
19. Henning, P., Målqvist, A.: Localized orthogonal decomposition techniques for boundary value problems. *SIAM J. Sci. Comput.* **36**(4), A1609–A1634 (2014). DOI 10.1137/130933198
20. Henning, P., Målqvist, A., Peterseim, D.: Two-level discretization techniques for ground state computations of Bose-Einstein condensates. *SIAM J. Numer. Anal.* **52**(4), 1525–1550 (2014). DOI 10.1137/130921520
21. Henning, P., Morgenstern, P., Peterseim, D.: Multiscale partition of unity. In: *Meshfree methods for partial differential equations VII, Lect. Notes Comput. Sci. Eng.*, vol. 100, pp. 185–204. Springer, Cham (2015)
22. Henning, P., Persson, A.: A multiscale method for linear elasticity reducing Poisson locking. *Comput. Methods Appl. Mech. Engrg.* **310**, 156–171 (2016). DOI 10.1016/j.cma.2016.06.034

23. Henning, P., Peterseim, D.: Oversampling for the multiscale finite element method. *Multiscale Model. Simul.* **11**(4), 1149–1175 (2013). DOI 10.1137/120900332
24. Hughes, T.J.R., Feijóo, G.R., Mazzei, L., Quincy, J.B.: The variational multiscale method—a paradigm for computational mechanics. *Comput. Methods Appl. Mech. Engrg.* **166**(1-2), 3–24 (1998). DOI 10.1016/S0045-7825(98)00079-6
25. Hughes, T.J.R., Sangalli, G.: Variational multiscale analysis: the fine-scale Green’s function, projection, optimization, localization, and stabilized methods. *SIAM J. Numer. Anal.* **45**(2), 539–557 (2007). DOI 10.1137/050645646
26. Kornhuber, R., Peterseim, D., Yserentant, H.: An analysis of a class of variational multiscale methods based on subspace decomposition. *Math. Comp.* **87**(314), 2765–2774 (2018). DOI 10.1090/mcom/3302
27. Kornhuber, R., Podlesny, J., Yserentant, H.: Direct and iterative methods for numerical homogenization. In: *Domain decomposition methods in science and engineering XXIII, Lect. Notes Comput. Sci. Eng.*, vol. 116, pp. 217–225. Springer, Cham (2017)
28. Kornhuber, R., Yserentant, H.: Numerical homogenization of elliptic multiscale problems by subspace decomposition. *Multiscale Model. Simul.* **14**(3), 1017–1036 (2016). DOI 10.1137/15M1028510
29. Li, G., Peterseim, D., Schedensack, M.: Error analysis of a variational multiscale stabilization for convection-dominated diffusion equations in two dimensions. *IMA J. Numer. Anal.* **38**(3), 1229–1253 (2018). DOI 10.1093/imanum/drx027
30. Maier, R., Peterseim, D.: Explicit computational wave propagation in micro-heterogeneous media. *BIT* **59**(2), 443–462 (2019). DOI 10.1007/s10543-018-0735-8
31. Målqvist, A., Peterseim, D.: Localization of elliptic multiscale problems. *Math. Comp.* **83**(290), 2583–2603 (2014). DOI 10.1090/S0025-5718-2014-02868-8
32. Målqvist, A., Peterseim, D.: Computation of eigenvalues by numerical upscaling. *Numer. Math.* **130**(2), 337–361 (2015). DOI 10.1007/s00211-014-0665-6
33. Målqvist, A., Peterseim, D.: Generalized finite element methods for quadratic eigenvalue problems. *ESAIM Math. Model. Numer. Anal.* **51**(1), 147–163 (2017). DOI 10.1051/m2an/2016019
34. Murat, F., Tartar, L.: H-convergence. *Séminaire d’Analyse Fonctionnelle et Numérique de l’Université d’Alger* (1978)
35. Ohlberger, M., Verfürth, B.: Localized Orthogonal Decomposition for two-scale Helmholtz-type problems. *AIMS Mathematics* **2**(3), 458–478 (2017)
36. Owhadi, H.: Multigrid with rough coefficients and multiresolution operator decomposition from hierarchical information games. *SIAM Rev.* **59**(1), 99–149 (2017). DOI 10.1137/15M1013894
37. Peterseim, D.: Variational multiscale stabilization and the exponential decay of fine-scale correctors. In: *Building bridges: connections and challenges in modern approaches to numerical partial differential equations, Lect. Notes Comput. Sci. Eng.*, vol. 114, pp. 341–367. Springer, [Cham] (2016)
38. Peterseim, D.: Eliminating the pollution effect in Helmholtz problems by local subscale correction. *Math. Comp.* **86**(305), 1005–1036 (2017). DOI 10.1090/mcom/3156
39. Peterseim, D., Schedensack, M.: Relaxing the CFL condition for the wave equation on adaptive meshes. *J. Sci. Comput.* **72**(3), 1196–1213 (2017). DOI 10.1007/s10915-017-0394-y
40. Peterseim, D., Scheichl, R.: Robust numerical upscaling of elliptic multiscale problems at high contrast. *Comput. Methods Appl. Math.* **16**(4), 579–603 (2016). DOI 10.1515/cmam-2016-0022
41. Peterseim, D., Verfürth, B.: Computational high frequency scattering from high contrast heterogeneous media. *ArXiv preprint* **1902.09935** (2019)
42. Verfürth, B.: Numerical homogenization for indefinite $H(\text{curl})$ -problems. In: *Proceedings of Equadiff 2017 conference*, pp. 137–146. Slovak University of Technology, Bratislava (2017)