

Methods that assist traction during endoscopic submucosal dissection of superficial gastrointestinal cancers: a systematic literature review

Georgios Tziatzios, Alanna Ebigbo, Stefan Karl Gölder, Andreas Probst, Helmut Messmann

Angaben zur Veröffentlichung / Publication details:

Tziatzios, Georgios, Alanna Ebigbo, Stefan Karl Gölder, Andreas Probst, and Helmut Messmann. 2020. "Methods that assist traction during endoscopic submucosal dissection of superficial gastrointestinal cancers: a systematic literature review." *Clinical Endoscopy* 53 (3): 286–301. <https://doi.org/10.5946/ce.2019.147>.

Open Access

Methods that Assist Traction during Endoscopic Submucosal Dissection of Superficial Gastrointestinal Cancers: A Systematic Literature Review

Georgios Tziatzios, Alanna Ebigo, Stefan Karl Gölder, Andreas Probst and Helmut Messmann

Department of Gastroenterology, University Hospital Augsburg, Augsburg, Germany

Endoscopic submucosal dissection (ESD) is a well-established method for the treatment of early-stage gastrointestinal neoplasms. Adequate submucosal exposure is one of the most significant factors related to an effective and safe dissection. The aim of this systematic review was to evaluate the efficacy and safety of various methods that assist traction during ESD of precancerous and early-stage neoplastic lesions of the gastrointestinal tract. We performed an electronic search of the MEDLINE and the Cochrane Controlled Trials Register databases for relevant studies published up to May 2019. Trials exclusively recruiting patients undergoing ESD for superficial gastrointestinal cancer were considered eligible for inclusion. Thirty-three articles including 3,134 patients met the inclusion criteria. The studies evaluated different approaches for widening the endoscopic view, including magnetic anchor-guided ESD (3 studies), use of a second endoscope (5 studies), clip-involving technique (21 studies), and miscellaneous methods (4 studies). Among them, only 6 were randomized controlled trials evaluating different approaches. Overall, the implementation of methods that assist traction during ESD significantly improved the operating time and R0 resection rate and decreased the rate of complications (bleeding and perforation). Interventions that assist traction seem efficacious in improving tissue traction, thus facilitating ESD performance. **Clin Endosc 2020;53:286-301**

Key Words: Dissection; Endoscopic; Method; Submucosal; Traction

INTRODUCTION

Endoscopic submucosal dissection (ESD) is a pioneer endoscopic method initially devised for the treatment of early-stage gastric neoplastic lesions.¹ To date, ESD has been established as an efficient method that achieves *en bloc* and R0 resection regardless of the lesion size, not only for early gastric cancerous lesions but also for lesions located in the colon or esophagus, overcoming the limitations of piecemeal endo-

scopic mucosal resection (EMR).^{2,3} Nonetheless, ESD remains a technically demanding and time-consuming procedure with a slow learning curve, particularly in Western countries.^{4,5} In addition, it is associated with higher rates of complications than EMR, including bleeding and perforation.⁶ The fundamental difficulty of the method lies in the accessibility of the submucosal layer during dissection. Meticulous identification of the dissection plane enables thorough recognition of the submucosal vessels and cutting line, thus reducing the risk of complications while increasing the possibility of achieving complete resection.⁷⁻¹¹ Traction is a method that could deliver satisfactory tissue tension within the submucosa and facilitate visualization of the dissection plane. Although traction has been the focus of many studies, most of the previous studies have included a single patient, or at most a very small number of patients, or involved non-human subjects, limiting the generalizability of their results to daily clinical practice. As these studies are beyond the scope of this systematic review,

Received: July 27, 2019 **Revised:** September 15, 2019

Accepted: September 24, 2019

Correspondence: Georgios Tziatzios
Department of Gastroenterology, University Hospital Augsburg, Stenglinstraße 2,
Augsburg 86156, Germany
Tel: +49-821-40001, **Fax:** +49-821-400-4585, **E-mail:** g_tziatzios@yahoo.gr
ORCID: <https://orcid.org/0000-0002-2945-6007>

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

we refer the readers to previous iterations.¹²⁻¹⁴ In this study, we addressed in a systematic manner all clinical studies involving only human participants, with the aim of providing further understanding of the traction methods that might substantially improve the safety and efficacy of ESD.

MATERIALS AND METHODS

Protocol

We conducted this review according to the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) recommendations (available in Supplementary 1).¹⁵ The protocol of the review can be accessed at the International Prospective Register of Systematic Reviews (PROSPERO registration number CRD42019135942).

Criteria for eligibility

The eligibility criteria were determined based on the PICO statement (P: patients undergoing ESD for gastrointestinal tract [GIT] precancerous/cancerous lesions; I: use of a method that enables traction to allow better exposure of the submucosal plane; C: comparison to conventional ESD; O: ESD outcomes including overall procedure time, curative resection [R0], resection speed, complication rate [bleeding or perforation], and recurrence rate as defined in each study). All types of trials published in the English language were considered eligible, whereas non-human studies, *ex vivo* or pilot studies, editorials, narrative reviews, case reports/series, video cases, and abstracts from conferences were excluded.

Identification and selection of studies

A computerized search of the MEDLINE and Cochrane Database of Systematic Reviews electronic databases for all relevant publications listed from database inception to May 2019 was performed. We combined the following search terms: “endoscopic submucosal dissection” and “traction” searched both as Medical Subject Headings and free-text terms. Moreover, the reference lists of the included original studies and pertinent reviews were manually searched for studies not initially identified. Our search strategy is detailed in Supplementary 2. Two members of the study team independently screened all initial abstracts. Subsequently, the full text of all eligible studies was independently assessed for eligibility. In case the full-text form of a study that appeared relevant could not be found, the corresponding author was contacted. If the author failed to provide missing information, the abstract was excluded. Any disagreements were resolved by consensus.

Extraction of data items

A structured form based on a Microsoft Excel sheet (Microsoft Co., Redmond, WA, USA) was used for data extraction. The following data were extracted from each study: first author name, study setting (publication year, origin), study design and primary outcome, type of intervention and comparator (if any), traction method, anatomical location of the ESD (esophagus, stomach, or colorectal area), level of endoscopist expertise (expert vs. non-expert), mean lesion size (as described in each study), presence of fibrosis, potential interference of the method during ESD (if provided by the study), endoscopist's subjective evaluation of the performance of each method, and outcomes (overall procedure time, curative resection [R0], resection speed, complication rate [bleeding or perforation], and recurrence rate as defined in each study).

Methodological quality of studies

The methodological quality of each study and the risk of bias of randomized and non-randomized studies were rated using the Cochrane Collaboration tool¹⁶ and Newcastle-Ottawa Scale,¹⁷ respectively. We used Review Manager 5.3 (The Nordic Cochrane Centre, The Cochrane Collaboration, Copenhagen, Denmark) software package to construct comparison plots.

RESULTS

Selection of studies

Our initial search generated 177 citations. After deduplication, 154 articles were retrieved and reviewed. Three more studies were identified after manual reference searching of the full-text articles. The inclusion criteria were met by 33 studies.¹⁸⁻⁵⁰ Fig. 1. illustrates the search flow.

Study characteristics

Table 1 summarizes the main characteristics of the included studies. The publication date of the studies ranged from 2006 to 2019, cumulatively enrolling 3,134 participants. All but 2 studies^{38,46} were conducted in a single-site setting. Twenty-three had a prospective design^{18-23,26,27,29,30,32,33,35,37-41,44-47,50} and 10 had a retrospective design.^{24,25,28,31,34,36,42,43,48,49} Among the prospective studies, 6 were randomized controlled trials (RCTs).^{27,30,32,39,45,46} The majority of the studies ($n=31$) were conducted in Eastern countries, whereas only 2 studies were conducted in Europe.^{38,48} With respect to the traction method, 3 studies evaluated magnetic anchor-guided ESD (MA-ESD),^{19,44,49} 5 evaluated the efficacy of double endoscopes,^{22,27,28,40,48} 21 investigated clip-involving methods for applying traction,^{24-26,29-39,41-43,45-47,50} and 4 assessed

miscellaneous methods.^{18,20,21,23} Concerning the anatomical location where ESD was performed, 6 studies evaluated ESD traction methods for esophageal lesions,^{21,26,32,38,41,43} 17 for stomach lesions,^{18-20,23,25,27-29,33,34,37,40,42,44,46-48} and 9 for colorectal lesions,^{22,24,30,35,36,39,45,49,50} whereas a single study³¹ enrolled patients with lesions in the stomach or colon. Data comparing the efficacy and safety of the traction methods between expert and non-expert endoscopists were available from only 5 studies,^{34,37,42,45,46} whereas ESD was performed by experts in all other studies. The mean lesion size, level of endoscopist expertise (expert vs. non-expert), endoscopist's subjective evaluation of the performance of the method, potential interference of the method with the ESD procedure, and the impact of each method on the main outcomes of the procedure are summarized in Table 2.

Study quality and risk of bias

The results of the risk of bias evaluation for the 6 RCTs^{27,30,32,39,45,46} are shown in Fig. 2. The overall quality can be characterized as questionable (detection bias and performance bias were the principal drawbacks). The results of the quality

assessment of observational studies according to the Newcastle-Ottawa scale are provided in Supplementary 3.

Magnetic anchor-guided endoscopic submucosal dissection

MA-ESD is a sophisticated method allowing dynamic tissue retraction with a rotatable external magnet (permanent magnet and electromagnet are the 2 available types). Traction is independent of the movement of the endoscope, thus serving as a “second hand” for the endoscopist.¹⁰ Compared with other methods that produce traction, the main advantage of this method is the lack of interference with endoscopic maneuvers during ESD while the constant movement of the external magnet changes the direction of the retraction, thus resulting in a dynamic traction phenomenon. Gotoda et al.¹⁹ investigated the feasibility of MA-ESD with an extracorporeal magnet in a study enrolling 25 subjects with early gastric cancer. All lesions were successfully removed *en bloc* without complications. No recurrence was observed in any of the patients after a median follow-up of 20 months. These results indicated, for the first time, that the technique is feasible and safe while offering excellent visualization for gastric ESD. To overcome




Fig. 1. Literature search flowchart and study selection.

Table 1. Characteristics of Studies Included in the Review

Study	Country	Study design	Period	Pts enrolled (total/ intervention/non- intervention, n)	Name of method	ESD site (LGI/ UGI, n)	Study's primary outcome
A. Magnetic - anchor guidance							
Gotoda et al. (2009) ¹⁹	Japan	Prospective, single-center	01/2005–05/2006	50/25/25	MA-ESD controlled by extracorporeal electro-magnet	0/50	Evaluate feasibility of MA-ESD for gastric lesions
Matsuzaki et al. (2018) ⁴⁴	Japan	Prospective, single-center	10/2016–06/2017	50/50/0	Neodymium magnets for MA-ESD	0/50	Evaluate feasibility of traction with good visualization
Ye et al. (2019) ⁴⁹	China	Retrospective, single-center, matched cohort	06/2017–01/2018	45/13/31	Gravity-based traction method - MBA-ESD	45/0	Compare safety and effectiveness of MBA-ESD and conventional ESD
B. Double endoscope							
Uraoka et al. (2010) ²²	Japan	Prospective, single-center	04/2006–10/2018	37/21/16	Thin endoscope-assisted ESD	37/0	Tumor size, <i>en bloc</i> resection rate, procedure time, complications rate
Ahn et al. (2013) ²⁷	Korea	RCT, single center	06/2010–08/2011	51/25/26	Transnasal endoscope assisted ESD	0/51	Procedure time, dropout rate, risk of perforation, uncontrolled bleeding, complications rate, evaluation by endoscopist
Higuchi et al. (2013) ²⁸	Japan	Retrospective, single center	10/2008–05/2012	57/30/27	Small-caliber upper gastrointestinal endoscope	0/30	Effectiveness and safety in patients with early gastric cancer accompanied by ulcer scars
Ogata et al. (2017) ⁴⁰	Japan	Prospective, single-center	1999–2015	122/122/0	DEILO	0/122	Effectiveness and safety of DEILO
Çolak et al. (2019) ⁴⁸	Turkey	Retrospective, single-center	01/2014–04/2018	22/22/0	Double endoscope	0/6	Evaluate feasibility of traction
C. Clip involving methods							
Okamoto et al. (2012) ^{24a)}	Japan	Retrospective, single-center	04/2010–05/2011	30/15/15	Cross-counter technique for colorectal tumors (thin tube, clip, nylon and balloon overtube)	30/0	Compare safety and efficacy of the new traction method, “cross-counter technique” for colorectal tumors
Okamoto et al. (2012) ^{25a)}	Japan	Retrospective, single-center	09/2009–08/2010	45/15/30	Cross-counter technique for early gastric cancer	0/15	Safety and efficacy of new method, for gastric tumors
Ota et al. (2012) ²⁶	Japan	Prospective, single-center	2005–2010	87/67/20	Clip traction	0/87	Evaluate usefulness in ESD for esophageal squamous cell carcinoma
Matsumoto et al. (2013) ²⁹	Japan	Prospective case control, single-center	09/2009–08/2010	74/37/37	Medical ring-assisted ESD	0/74	Evaluate usefulness for early gastric cancer

Table 1. Continued

Study	Country	Study design	Period	Pts enrolled (total/ intervention/non- intervention, n)	Name of method	ESD site (LGI/ UGI, n)	Study's primary outcome
Ritsuno et al. (2014) ³⁰	Japan	RCT, single-center	08/2010–12/2011	50/27/23	S-O clip-assisted ESD	50/0	Evaluate efficacy and safety for colorectal tumors ESD
Koike et al. (2015) ³²	Japan	RCT, single-center	05/2012–02/2013	40/20/20	Thread-traction method	0/40	Assess usefulness in esophageal ESD
Cai et al. (2015) ³¹	China	Retrospective, single-center	04/2014–08/2014	20/20/0	ESD-assisted dental floss	10/10	Evaluate the usefulness of ESD-assisted dental floss traction for the removal of gastrointestinal tumors
Suzuki et al. (2016) ³⁴	Japan	Retrospective, single-center	05/2012–12/2014	86/43/43	Dental floss and hemoclip	0/86	Evaluate the efficacy of a traction method for early gastric cancers that uses dental floss and a hemoclip
Yoshida et al. (2016) ³⁷	Japan	Prospective, single-center	08/2014–11/2014	190/95/95	Dental floss clip	0/195	Investigate efficacy for gastric ESD
Yamada et al. (2016) ³⁵	Japan	Prospective, single-center	11/2013–05/2014	140/17/123	Clip and snare assisted ESD	140/0	Evaluate the efficacy and safety of a clip and snare method with a prelooping technique for colorectal tumors
Yamasaki et al. (2016) ³⁶	Japan	Retrospective, single-center	10/2014–03/2015	23/23/0	Colonic ESD using clip and line	23/0	Evaluate the feasibility of traction-assisted colonic ESD using clip and line
Noda et al. (2016) ³³	Japan	Prospective, single-center	01/2014–03/2015	88/54/34	Thread-Traction with Polypectomy Snare Sheath	0/88	Compare the new method to conventional ESD for early gastric cancer
Xie et al. (2017) ⁴¹	Japan	Prospective, Case-matched comparative, single-center	03/2014–06/2015	100/50/50	Clip traction	0/100	Investigate efficacy and safety for early esophageal carcinoma
Mori et al. (2017) ³⁹	Japan	RCT, single-center	12/2015–07/2016	43/21/22	Ring-thread counter traction	43/0	Evaluate usefulness
Jacques et al. (2017) ³⁸	France	Prospective, multicenter	01/2015–12/2016	62/62/0	“Tunnel + clip” ESD	0/62	Investigate efficacy and safety of the tunnel + clip strategy for esophageal ESD
Kitagawa et al. (2018) ⁴³	Japan	Retrospective, single-center	03/2013–01/2017	103/103/0	Clip traction method	0/103	Evaluate safety and efficacy of ESD with clip traction for esophageal squamous cell carcinoma
Hashimoto et al. (2018) ⁴²	Japan	Retrospective, single-center	09/2016–11/2016	306/48/258	S-O clip-assisted ESD	0/306	Assess efficacy and safety for gastric neoplasms
Yoshida et al. (2018) ⁴⁶	Japan	RCT, multicenter	07/2015–09/2016	635/319/316	Dental floss clip assisted ESD	0/635	The study's primary endpoint was difference in total ESD procedure time between the two techniques for gastric neoplasms

Table 1. Continued

Study	Country	Study design	Period	Pts enrolled (total/ intervention/non- intervention, n)	Name of method	ESD site (LGI/ UGI, n)	Study's primary outcome
Yamasaki et al. (2018) ⁴⁵	Japan	RCT, single-center	08/2015–10/2016	85/42/43	Clip-and-thread technique	85/0	The study's primary endpoint was the procedure time between the two techniques, measured from beginning of sub mucosal injection until separating lesion from col-orectal wall
Zhang et al. (2019) ^{47(b)}	China	Prospective, single-center	08/2016–09/2018	54/54/0	Snare combined with endoclips assisted ESD	0/54	Evaluate feasibility of snare combined with endoclips for mucosal traction during gastroesophageal ESD
Zhang et al. (2019) ^{50(b)}	China	Prospective, single-center	01/2018–09/2018	50/50/0	Snare combined with endoclips assisted ESD	50/0	Evaluate feasibility and safety during colon and rectum ESD
D. Miscellaneous methods							
Imaeda et al. (2009) ²⁰	Japan	Prospective, single-center	10/2003–07/2008	252/252/0	External forceps	0/265	Evaluate feasibility of traction in various gastric locations
Yonezawa et al. (2006) ¹⁸	Japan	Prospective, single-center	03/2004–03/2005	60/20/40	Double-channel therapeutic endoscope ("R-scope")	0/20	Assess effectiveness and complications of the new scope compared to conventional ESD in various gastric locations
Hijikata et al. (2012) ²³	Japan	Prospective, single-center	N/A	68/25/43	Sheath-assisted counter traction ESD	0/68	Mean duration of procedure relative to tumor size, location, complications rate
Motohashi et al. (2009) ²¹	Japan	Prospective, single-center	N/A	9/9/0	Two-point fixed ESD	0/9	Evaluate feasibility of traction in early esophageal cancer >20 mm in diameter

DEILO, double endoscopic intraluminal operation; ESD, endoscopic submucosal dissection; LGI, lower gastrointestinal tract; MA-ESD, magnetic anchor-guided endoscopic submucosal dissection; MBA-ESD, magnetic bead-assisted endoscopic submucosal dissection; N/A, not applicable; RCT, randomized controlled trial; UGI, upper gastrointestinal tract.

^{a(b)}Studies conducted by the same authors group but refer to different lesion ESD site.

Table 2. Effect of Traction Methods on Endoscopic Submucosal Dissection Main Outcomes

Study	Method	Mean size	Anatomical location	Fibrosis degree	Expert vs. Non-expert	Endoscopist evaluation	Interference with ESD	Procedure time	Resection speed	Curative resection (R0)	Complications ^{b)}	Recurrence
A. Magnetic - anchor												
Gotoda et al. (2009) ^{19a)}	Magnetic anchor-guided ESD	55 mm (range, 33–125)	Stomach	Pts with fibrosis were excluded	2 experts	Supportive in 23/25 cases	–	80 min (range, 50–240)	N/A	100%	0%	0%
Matsuzaki et al. (2018) ^{44a)}	Neodymium magnets for magnetic anchor ESD	20 mm (range, 5–100)	Stomach	N/A	1 expert	N/A	–	49 min (range, 15–301)	N/A	100%	0%	N/A
Ye et al. (2019) ⁴⁹⁾	Gravity-based traction method - magnetic bead ESD	589 mm ² (range, 416–893)	Colorectal	Pts with fibrosis were excluded	1 expert	N/A	–	No impact	Lower	Similar	Lower	Similar
B. Double endoscope												
Uraoka et al. (2010) ²²⁾	Thin endoscope-assisted ESD	43.6 mm (SD, 16)	Colorectal	N/A	1 expert	Supportive	+	Lower	Lower	Higher	Lower	N/A
Ahn et al. (2013) ²⁷⁾	Transnasal endoscope assisted ESD	19.9 mm (SD, 7.2)	Stomach	N/A	2 non experts	Helpful for larger tumors	+	Higher	Similar	Similar	Similar	N/A
Higuchi et al. (2013) ²⁸⁾	Small-caliber upper gastrointestinal endoscope	45 mm (range, 28–70)	Stomach	N/A	Experts	Supportive	+	Lower	Higher	Higher	Similar	N/A
Ogata et al. (2017) ^{40a)}	Double endoscopic intraluminal operation	18 mm (range, 2–42)	Stomach	N/A	N/A	N/A	+	70.9 min (range, 20–207)	N/A	97.5%	6% (7/122)	1.6% (2/122)
Çolak et al. (2019) ^{48a)}	Double endoscope	N/A	Stomach	N/A	N/A	N/A	+	N/A	N/A	N/A	N/A	N/A
C. Clip involving methods												
Okamoto et al. (2012) ²⁴⁾	Cross-counter technique for colorectal	37.3 mm (SD, 9.3)	Colorectal	N/A	1 expert	N/A	–	Lower	Higher	Similar	Similar	Similar
Okamoto et al. (2012) ²⁵⁾	Cross-counter technique for early gastric cancer	15 mm (range, 10–35)	Stomach	N/A	2 non experts	N/A	–	Lower	Higher	Similar	Similar	Similar
Ota et al. (2012) ²⁶⁾	Clip traction	28.1 mm	Esophagus	N/A	2 experts	N/A	–	Lower	Higher	N/A	Lower	N/A
Matsumoto et al. (2013) ²⁹⁾	Medical ring-assisted ESD	39.8 mm (range, 21–90)	Stomach	N/A	5 non experts	N/A	–	Lower	Higher	Similar	Similar	N/A

Table 2. Continued

Study	Method	Mean size	Anatomical location	Fibrosis degree	Expert vs. Non-expert	Endoscopic evaluation	Interference with ESD	Procedure time	Resection speed	Curative resection (R0)	Complications ^{b)}	Recurrence
Ritsuno et al. (2014) ³⁰	S-O clip-assisted ESD	33.5 mm (SD, 12.5)	Colorectal	N/A	1 expert	N/A	–	Lower	Higher	Similar	Similar	N/A
Koike et al. (2015) ³²	Thread-traction method	35.5 mm (range, 21–100)	Esophagus	N/A	2 experts	Helpful for all cases	–	Lower	Higher	Similar	Similar	N/A
Cai et al. (2015) ^{31a)}	ESD-assisted dental floss traction	27.4 mm (range, 18–35)	Stomach and colorectal	N/A	1 expert	N/A	–	45 min (range, 30–100)	N/A	100%	0%	0%
Suzuki et al. (2016) ³⁴	Dental floss and a hemoclip (DFC-assisted ESD)	17.4 mm (SD, 11.8)	Stomach	N/A	Experts, trainees	N/A	–	Lower	Higher	Similar	Similar	N/A
Yoshida et al. (2016) ³⁷	Dental floss clip traction assisted ESD	17.0 mm (SD, 11.0)	Stomach	N/A	Experts, trainees	N/A	–	Lower	Higher	Similar	Similar	N/A
Yamada et al. (2016) ³⁵	Clip and snare assisted ESD	32.5 mm (SD, 10.9)	Colorectal	N/A	2 experts	N/A	–	Lower	Higher	Similar	Similar	N/A
Yamasaki et al. (2016) ^{36a)}	Colonic ESD using clip and line	27 mm (range, 20–44)	Colorectal	Pts after EMR were excluded	N/A	N/A	–	61 min (range, 18–172)	N/A	96%	4%	N/A
Noda et al. (2016) ³³	Thread-Traction with Polypectomy Snare Sheath	30 mm (range, 14–60)	Stomach	N/A	N/A	N/A	–	Lower	Higher	Similar	Similar	N/A
Xie et al. (2017) ⁴¹	Clip traction	9.60 cm ²	Esophagus	N/A	2 experts	N/A	–	Lower	Higher	Similar	Lower	N/A
Mori et al. (2017) ³⁹	Ring-thread counter traction	27.3 cm ² (range, 11.0–49.9)	Colorectal	N/A	3 experts	N/A	–	Lower	Higher	N/A	Similar	N/A
Jacques et al. (2017) ^{38a)}	“Tunnel + clip” ESD	220.6 cm ² (range, 47–68)	Esophagus	N/A	4 non experts	N/A	–	N/A	N/A	88.7%	0%	N/A
Kitagawa et al. (2018) ^{43a)}	Clip traction method	19 mm (range, 3–60)	Esophagus	N/A	3 experts	N/A	–	40 min (range, 13–230)	N/A	100%	0%	N/A
Hashimoto et al. (2018) ⁴²	S-O clip-assisted ESD	37.4 mm (SD, 12.1)	Stomach	N/A	Experts, trainees	N/A	–	Lower	Higher	Similar	Similar	N/A

Table 2. Continued

Study	Method	Mean size	Anatomical location	Fibrosis degree	Expert vs. Non-expert	Endoscopic evaluation	Interference with ESD	Procedure time	Resection speed	Curative resection (R0)	Complications ^{b)}	Recurrence
Yoshida et al. (2018) ⁴⁶	Dental floss clip assisted ESD	15.5 mm (SD, 8.9)	Stomach	N/A	Experts, trainees	N/A	–	Similar	N/A	Similar	Lower	N/A
Yamasaki et al. (2018) ⁴⁵	Clip-and-thread technique	30 mm (range, 20–60)	Colorectal	Pts after EMR were excluded	2 Experts, 2 trainees	N/A	–	Lower	N/A	Similar	Similar	N/A
Zhang et al. (2019) ⁴⁷	Snare combined with endoclips assisted ESD	42 mm (range, 20–80)	Stomach	N/A	1 expert	N/A	–	N/A	N/A	N/A	0%	N/A
Zhang et al. (2019) ⁵⁰	Snare combined with endoclips assisted ESD	45 mm (range, 20–90)	Colorectal	N/A	1 expert	N/A	–	32 min (range, 8–247)	N/A	100%	0%	N/A
D. Miscellaneous methods												
Imaeda et al. (2009) ^{20a)}	External grasping forceps	15.0 mm (range, 5–50)	Stomach	N/A	N/A	N/A	+	N/A	N/A	N/A	N/A	N/A
Yonezawa et al. (2006) ¹⁸	Double-channel therapeutic endoscope (“R-scope”)	18.5 mm (SD, 10.3)	Stomach	N/A	N/A	Supportive	–	Lower	Higher	Similar	Similar	Similar
Hijikata et al. (2012) ²³	Sheath-assisted counter traction ESD	20 mm	Stomach	N/A	N/A	Supportive	+	Lower	Lower	N/A	Similar	N/A
Motohashi et al. (2009) ^{21a)}	Two-point fixed ESD	27 mm (range, 11–56)	Esophagus	N/A	N/A	Supportive	–	N/A	N/A	100%	0%	0%

EMR, endoscopic mucosal resection; ESD, endoscopic submucosal dissection; N/A, not applicable; SD, standard deviation.

^{a)} Study enrolling only patients undergoing ESD with one technique; no comparative arm is available; results presented are only for the new method.

^{b)} Including bleeding (immediate/delayed), perforation, stricture formation.

the main drawback of the method (i.e., the large size of the external electromagnet), Matsuzaki et al.⁴⁴ conducted a feasibility study of MA-ESD with neodymium magnets for the treatment of gastric lesions. In the aforementioned trial, the magnetic traction consisted of an internal magnet attached to a clip. Although there were minor concerns about patient safety because of the low resistance to rust of the magnet after direct contact with human tissue, the authors reported obtaining adequate counter-traction in all cases, resulting in successful *en bloc* resection of lesions without adverse events or allergic reactions. Recently, Ye et al.⁴⁹ in their retrospective study compared the efficacy and safety of magnetic bead-assisted ESD (MBA-ESD) with those of standard ESD for large colorectal cancerous lesions. Despite the low enrollment in the study ($n=26$), the results supported the notion that MBA-ESD is equivalent to conventional ESD in terms of *en bloc* and R0 colonic resection as well as local recurrence. Furthermore, complications were totally absent (0% in the MBA-ESD group vs. 38.5% in the standard ESD group, $p=0.039$) while other

procedure-related parameters (i.e., dissection time and speed) also improved.

Use of double endoscopes

The implementation of a second endoscope to facilitate traction has been particularly considered for treating lesions that are difficult to resect with conventional ESD. As a rule, the initial circumferential incision is performed using the primary endoscope, followed by the insertion of a thinner endoscope that applies traction to the lesion through common grasping forceps passed through its working channel.¹⁴ The method is “operator friendly”, as it enhances the accessibility of the submucosa and increases the efficiency of submucosal dissection while reducing the risk of complications. Uraoka et al.²² reached the conclusion that the double-endoscope approach is technically easier and safer for large colorectal tumors. However, the method was applied only to rectal and rectosigmoid lesions, creating doubts about the efficacy of the method in more proximal colonic lesions. Moreover, the need for an additional person to operate the traction system and for a second light source represents further weaknesses. In an attempt to provide further clarifications, Higuchi et al.²⁸ conducted a retrospective study introducing an improved version of the method for early gastric cancer, in which a unique, switchable light source between the endoscopes was used. Although comparisons were made with historical control data, the cutting rate into specimens was improved (7% vs. 35%, $p=0.01$) with the double-endoscope method, with no serious adverse events noted. These results, however, were later refuted by the only RCT available on this matter.²⁷ Ahn et al.²⁷ found that the main outcomes of ESD were similar between endoscope-assisted ESD and standard ESD when used for gastric neoplastic lesions. Although the study was well designed, the enrollment of a low number of patients ($n=51$) and the performance of ESD procedures by inexperienced endoscopists are the main points of critique. Thereafter, results from 1 Western center and 1 Eastern were reported. The Western results, from a retrospective follow-up study originating from Turkey, showed no difference in the procedure duration and complication rate when using the double-endoscope method for upper gastrointestinal lesions.⁴⁸ Ogata et al.,⁴⁰ by enrolling relatively more subjects ($n=122$) with a long mean follow-up period (24 months), highlighted the safety and efficacy of a double-endoscopic intraluminal operation for precancerous gastric lesions. Taken together, these studies demonstrated somewhat conflicting results about the role of the double-endoscope method in ESD; however, they should be critically taken into account considering the low number of patients and the variable follow-up period.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)
Ahn et al. (2013) ²⁷	+	?	-	-	+	+
Koike et al. (2015) ³²	+	+	-	-	+	+
Mori et al. (2017) ³⁹	?	+	-	-	+	+
Ritsuno et al. (2014) ³⁰	?	+	-	-	+	+
Yamasaki et al. (2018) ⁴⁵	+	?	-	-	+	+
Yoshida et al. (2018) ⁴⁶	+	?	-	-	+	+

Fig. 2. Risk of bias among randomized controlled trials included in this review.

Clip-involving methods

Accumulating evidence has highlighted the impact of clip traction on the technical outcomes of ESD, as well as its therapeutic and prognostic implications. Early studies showed that traction produced by clips can significantly reduce the overall procedure time compared with standard ESD, with the results being consistent for both gastric and colonic precancerous lesions.^{24–26} This finding was also supported by other studies evaluating the efficacy of ESD with dental floss clip traction in several locations inside the GIT.^{31,34,36,37,41,43} On the other hand, data from RCTs attempting to investigate any potential benefit in terms of procedural outcomes when using clip-related methods seem somewhat conflicting. The earliest study³² showed that the thread-traction method resulted in significant shortening of the dissection time compared with conventional ESD in esophageal lesions (19.8 min vs. 31.8 min, $p=0.044$). Accordingly, Mori et al.³⁹ reported that ring-thread counter traction also optimized the total dissection time in colorectal ESD (130.0 [56.0–240.0] min vs. 80 [35.0–130.0] min, $p=0.001$). In another very recent Japanese RCT enrolling 84 patients with ≥ 20 mm superficial colorectal neoplasms, the clip-and-thread technique was related not only to a shorter procedure time (40 [11–86] min vs. 70 [30–180] min, $p<0.0001$) but also to a higher self-completion rate in non-experts (100% [39/39] vs. 90% [36/40], $p=0.04$) than conventional ESD.⁴⁵ However, the results published in the largest RCT thus far seem to refute the role of traction-assisted ESD for gastric neoplasms reported in the aforementioned studies. In that study,⁴⁶ 640 participants with gastric neoplasms were randomized at 14 centers across Japan to undergo standard ESD or ESD in which a dental floss clip was used to provide traction, with the ESD procedure time being the primary endpoint. No difference was observed in the mean ESD procedure time between the 2 methods (60.7 min vs. 58.1 min, $p=0.45$). However, perforation was less frequent in patients treated with dental floss clip traction-assisted ESD (2.2% vs. 0.3%, $p=0.04$). On the other hand, the new technique was particularly beneficial for lesions located in the greater curvature of the stomach, for which the mean procedure time was effectively reduced (104.1 min vs. 57.2 min, $p=0.01$). The clip-and-snare method has been reported to be another promising technique because it requires no special equipment to enable pushing and pulling movements without impairing flexibility. The technique was found to have a significant impact on the procedure time (45.6 min vs. 70.1 min [vs. control group], $p=0.047$), with a lower complication rate (5.9% vs. 8.1%, $p=1.00$).³⁵ Noda et al.³³ used a sheath of a polypectomy snare to create traction and achieved faster resection time, which was uniform among endoscopists of various skill levels. In keeping with the findings of those reports, the approach of inserting a selective snare along with

the endoscope has also been shown to be safe and feasible in ESD of colonic and gastric intraepithelial neoplasia.^{47,50} An alternative internal traction method using the S-O clip for treating colorectal lesions has been recommended. The device is advanced through the working channel of an endoscope and functions independently of endoscope movements. The safety and efficacy of the method were first evaluated in a prospective RCT, which showed that the mean ESD time for large colorectal tumors (>20 mm diameter) was significantly reduced when the S-O clip was used, compared with the standard technique (37.4 \pm 32.6 min vs. 67.1 \pm 44.1 min, $p=0.03$).³⁰ Similar results were also obtained in terms of the efficacy of the clip in the treatment of upper GIT epithelial neoplasms in a recent large retrospective Japanese study. The mean procedure time was significantly shorter in the S-O clip group (47.2 \pm 24.6 min vs. 69.2 \pm 67.1 min, $p=0.035$), with similar results in secondary outcomes to those of standard ESD (*en bloc* resection rate: 100% vs. 100%, $p=1.000$; perforation rate: 0% vs. 2.1%, $p=0.315$; and delayed bleeding rate: 2.1% vs. 4.3%, $p=0.558$).⁴² In another Japanese study, Matsumoto et al.²⁹ reported on the use of a “medical ring” during ESD in a prospective case-control study. The newly developed ring allowed adequate visualization, thus enhancing the performance of gastric ESD relative to the conventional procedure (3.18 \pm 2.29 dissection min/cm² vs. 6.3 \pm 3.6 dissection min/cm², $p<0.01$). Taking these observations into account, it is evident that more robust data are definitely needed in order to assess the performance of these novel methods overall and in specific populations.

Miscellaneous methods

We identified 4 studies^{18,20,21,23} evaluating various methods to improve traction during an ESD procedure. More than a decade ago, Yonezawa et al.,¹⁸ in a well-designed, 12-month follow-up prospective trial of 60 patients, showed that the use of a sophisticated double-channel endoscope (“R-scope”) resulted in significantly shorter operating time than standard ESD (57.9 \pm 29.7 min vs. 92.8 \pm 58.9 min, $p=0.016$), with similar efficacy and complication rates. Three years later, Motohashi et al.²¹ presented a novel method—the 2-point fixed ESD—for esophageal cancerous lesions that allowed traction to the site of dissection by using a hood fitted with a forceps channel. Their results implied that the technique is feasible for esophageal lesions. In an effort to optimize ESD procedures, authors from a large prospective Eastern trial used external biopsy forceps for early-stage gastric cancer.²⁰ Despite the promising results in terms of low mean procedure time, the method also had an important inherent shortcoming (i.e., inability to be performed when the lesion is located in other gastric sections such as the cardia and lesser curvature). Further insights into the clinical outcomes of traction-assisted ESD

were provided by a small, single-center study that investigated the utility of the novel sheath for ESD.²³ A total of 43 and 25 consecutive patients with early gastric carcinomas treated using the standard method and the new method, respectively, were compared. The use of the novel sheath was not only less time-consuming for lesions sized 20 mm and for all resected tumors regardless of location but was also technically simpler. Undoubtedly, the confirmation or rejection of these results requires more evidence because relevant prospective data are lacking.

Efficacy and safety of traction methods based on the anatomical site of endoscopic submucosal dissection

Although data comparing the efficacy and safety of various traction methods according to the anatomical location of ESD are lacking, clip-involving techniques and their modifications seem to be efficacious in providing sufficient traction in all GIT sites, offering particular benefits for lesions located in the esophagus and greater curvature of the gastric body. The use of a second endoscope has shown considerable efficacy in treating lesions located in the distal colon (sigmoid and rectum), whereas its usefulness for more proximal tumors is questionable. The method also seems advantageous for patients with gastric neoplasms. For magnet-assisted ESD, most of the data were derived from patients with upper GIT lesions, whereas its efficacy in colorectal tumor treatment remains to be established. Finally, as studies evaluating miscellaneous methods exclusively enrolled patients with tumors in the upper GIT (esophagus and stomach), it can be assumed that the efficacy of the methods is limited to those locations.

Efficacy of traction methods based on the expertise level of the endoscopist

Five studies^{34,37,42,45,46}—all evaluating clip-involving methods—provided data about the impact of the endoscopist's expertise level (expert vs. non expert) on ESD outcomes. Overall, the data showed that in terms of procedure time, the traction methods were mostly beneficial for expert endoscopists. Although the procedure time also improved when non-experts used traction methods, the difference did not reach significance in most studies. Notably, these findings were not verified by the largest RCT,⁴⁶ which reported no difference in procedural time regardless of the operator's experience.

Endoscopist's evaluation of the traction method and interference with the endoscopic submucosal dissection procedure

Eight studies^{18,19,21–23,27,28,32} reported on the endoscopist's subjective evaluation of the traction method performance. Over-

all, the results suggested that traction methods are supportive and helpful during ESD. The results on the interference of each method with the ESD procedure are presented in Table 2. The MA-ESD and clip-involving methods cause potentially no or minimal interference; however, this does not apply when double endoscopes are used, as significant interference between the 2 devices can be noted.

Overall comparison of methods

The use of magnets represents an innovative method of delivering traction during ESD. Compared with the other methods, the principal advantage of MA-ESD is that the traction applied to the ESD site is independent (i.e., easily adaptable without any interference with endoscope movement). On the other hand, the applicability of the method is limited by the significant decrease in force as the distance between the magnets increases (i.e., in cases of a thick abdominal wall). The large size of the magnets, scant data for the efficacy of the method for colorectal lesions, additional financial burden, and potential effects on the human body are also issues that should not be underestimated. The double-endoscope method can be a rewarding, handy option that allows traction to be effectively adjusted in any direction merely through standard maneuvering of a second endoscope—a procedure familiar to all endoscopists. However, the major flaw of this method is that the 2 endoscopes may interfere with each other. Although the use of a single transferable light source between the 2 endoscopes has been reported,²⁸ the method requires, in most cases, dual light sources, endoscopes, and physicians, making it an expensive approach. The method can be efficacious for upper GIT lesions, for which it has been mainly evaluated; however, in terms of colonic ESD, its competence is limited to distal colonic and rectal tumors. On the other hand, clip-involving methods and their modifications are, in most cases, relatively simple to use, can be applied to any site, do not require additional special equipment, and have a low cost; thus, they can be easily integrated into daily clinical practice. Notably, they can be particularly useful for some lesions such as those located in the greater curvature of the gastric body. However, they are not without caveats. Traction can be rather difficult to adjust as pulling is the only option. Moreover, the clip can be detached from tissue, necessitating repetition of the procedure, which is a troublesome task when the lesion lies in the proximal colon. The application of miscellaneous methods is limited owing to the paucity of data about their efficacy; thus, they cannot be recommended at a large scale. Taking these observations into account, it is evident that more robust data are definitely needed to assess the performance of these novel methods overall and in specific populations.

CONCLUSIONS

A large variety of endoscopic methods and novel devices has been introduced with the aim of improving traction during ESD. Despite the promising results, data about the overall and comparative efficacy of the methods remain scarce, thus preventing their integration into daily clinical practice. For the first time, we addressed in a systematic manner the knowledge deficit about the role of all available endoscopic methods and devices used to assist traction in ESD.

Our review indicated that multiple disparate endoscopic methods have the potential to improve tissue tension and facilitate visualization during ESD. Indeed, the majority of the studies uniformly showed that the implementation of any method results in significant improvement in the core quality features (overall operating time and complication rate) of each procedure relative to standard ESD. However, the studies reviewed here did not allow drawing firm conclusions on how to best achieve traction during ESD, as there was no evidence supporting the superiority of any of the strategies. Instead, endoscopists may consider adopting a method on a case-by-case basis, taking into account several factors such as lesion characteristics, proficiency level of the endoscopist, and availability of the method in a given health-care setting.

Undoubtedly, these technological advances can be considered adjuvant approaches to improve physician performance during ESD.¹⁴ On the other hand, it can be argued that their true benefit remains questionable at a time when a simple and costless alternative—traction by gravity—may be equally effective (Fig. 3). Gravity can induce traction, thus abolishing the need for any additional device by allowing controlling the direction of the traction force merely through changing the patient's position according to intraluminal fluid location.¹⁴ However, a head-to-head comparative evaluation between traction by gravity and any other traction technique is still lacking. Nonetheless, even traction by gravity is far from being the perfect concept, as the submucosal view may be not be sufficient during the initial stages of the procedure, when the

flap is not yet sufficiently exposed, or in non-expert hands.

An issue that remains to be elucidated is whether the efficacy and safety of the traction methods differ according to the anatomical location of the ESD procedure or the expertise level of the endoscopist. Although direct comparisons with respect to anatomical sites are totally absent, it seems that clip-involving methods have a beneficial effect on ESD outcomes throughout the GIT, particularly for the stomach. On the other hand, double-endoscope-assisted ESD is particularly efficacious in treating distal colonic (sigmoid, rectum) tumors. As far as the expertise level of the endoscopist is concerned, the majority of the studies showed that traction methods significantly improved ESD outcomes in expert hands. The latter seems to be also the case when the performance of non-experts comes into question. However, this finding was not consistent with the results of a large RCT⁴⁶ that refuted the impact of the operator's experience on procedure time, thus underlining the need for more data to verify this hypothesis.

Although the many technical interventions presented in this review seem to be beneficial in terms of improving ESD outcomes, concerns about the quality of the studies are raised. It should be underlined that most of the studies originated from expert centers, which limits the external validity (i.e., generalizability) of the results.^{2,12} Additional studies are warranted to examine whether the results also apply to different hospital settings. Moreover, most of the studies had a single-center and retrospective design, enrolled populations with divergent ESD indications, or only aimed to prove the feasibility of a new method rather than to offer comparative results. Even results from published RCTs should be interpreted with caution because of serious study limitations. Surprisingly, the largest RCT meeting our eligibility criteria⁴⁶ suggested that dental floss clip-assisted ESD traction does not reduce the procedure time.

Our review could also have significant implications on future research directions. Future studies should systematically assess endoscopist-related (e.g., variable level of skill), patient-related (e.g., location and type of the lesion), and



Fig. 3. Traction methods: (A) double-endoscope method, (B) clip-with-line method, and (C) traction by gravity.

setting-related factors in an effort to reach solid conclusions about the real-world effectiveness of the techniques. Likewise, efforts should be made to perform comparative evaluations of different approaches, as a further step toward optimization of the ESD procedure. Finally, future studies will benefit from larger sample sizes, which will provide not only additional evidence for efficacy but also adequate statistical power to detect changes in consequential clinical and procedural outcomes.

Effective and safe endoscopic removal or GIT precancerous lesions always remains a focal point. New medical devices that provide and facilitate tissue retraction in special cases have already emerged. Among them, a new device consisting of a double-balloon platform and a sheath (DiLumen Endolumenal Interventional Platform; Lumendi Ltd., High Wycombe, UK) has recently been shown to facilitate exposure of difficult-to-access lesions owing to poor endoscope maneuverability and loop presence.⁸ Another endoscopic system (ORISE TRS; Boston Scientific Co., Marlborough, MA, USA) has been reported to facilitate challenging colorectal ESD because of significant fibrosis from previous tattooing, by providing constant traction and adequate view of the field to be dissected.⁹ Finally, an exterior supplementary working channel (Ovesco, Tübingen, Germany) device mounted on the tip of a standard endoscope was recently introduced, allowing an additional endoscopic tool to be used for traction.¹¹ The adjustable distance between working channels makes the device suitable not only for ESD but also for EMR without the need for a dual-channel endoscope.

A number of strengths of this study could be cited. We used a rigorous methodology and performed a recursive bibliographic search including a detailed search of all pertinent bibliographies. The independent assessment of eligibility and strict evaluation of study quality are considered additional advantages. Our effort adds to the existing literature by addressing the topic of traction during ESD based on data exclusively from human studies, adding to the understanding of how existing studies, despite their inherent caveats, may still guide clinical practice with the eventual aim of highlighting potential fields for future research.

Our work also has a number of limitations that merit further discussion. First, we did not perform a meta-analysis; however, the presence of statistical and clinical heterogeneities among the studies (different designs, variable indications and endoscopist skill levels, distinct techniques evaluated) would have made a quantitative meta-analysis inappropriate. In this regard, the possibility of publication bias cannot be ruled out. Furthermore, the lack of additional database search (i.e., EMBASE), English-language restriction, and Eastern origin of the majority of the studies also represent weaknesses of this review. Finally, the studies tend to have limited validity given

the setting where they were conducted.

In summary, evidence from this systematic review suggests that several methods, including MA-ESD, use of double endoscopes, clip-involving techniques, and others, seem to be effective in improving the performance of ESD in patients with variable gastrointestinal lesions. Each method presents distinct advantages but also has considerable drawbacks, as outlined in Tables 1 and 2. Although none of the methods have sufficient evidence to be recommended at a large scale, interventional endoscopists should be aware of them in the pursuit of strategies for improving ESD performance.

Conflicts of Interest

Georgios Tziatzios is a scholar of the Hellenic Society of Gastroenterology. The other authors have no financial conflicts of interest.

Author Contributions

Conceptualization: Helmut Messmann
Data curation: Georgios Tziatzios, Alanna Ebigo, Stefan Karl Gölder, Andreas Probst, HM
Formal analysis: GT, AE, SKG, AP, HM
Investigation: GT, AE, SKG, AP, HM
Methodology: GT, AE, SKG, AP, HM
Project administration: GT, AE, SKG, AP, HM
Resources: GT, AE, SKG, AP, HM
Software: GT, AE, SKG, AP, HM
Supervision: GT, AE, SKG, AP, HM
Validation: GT, AE, SKG, AP, HM
Visualization: GT, AE, SKG, AP, HM
Writing-review&editing: GT, AE, SKG, AP, HM

ORCID

Alanna Ebigo: <https://orcid.org/0000-0001-7765-035X>
Stefan Karl Gölder: <https://orcid.org/0000-0003-4507-5770>
Andreas Probst: <https://orcid.org/0000-0002-1486-9298>
Helmut Messmann: <https://orcid.org/0000-0002-4026-6837>

REFERENCES

1. Fuccio L, Ponchon T. Colorectal endoscopic submucosal dissection (ESD). *Best Pract Res Clin Gastroenterol* 2017;31:473-480.
2. Pimentel-Nunes J, Dinis-Ribeiro M, Ponchon T, et al. Endoscopic submucosal dissection: European Society of Gastrointestinal Endoscopy (ESGE) guideline. *Endoscopy* 2015;47:829-854.
3. Saito Y, Fukuzawa M, Matsuda T, et al. Clinical outcome of endoscopic submucosal dissection versus endoscopic mucosal resection of large colorectal tumors as determined by curative resection. *Surg Endosc* 2010;24:343-352.
4. Probst A, Golger D, Anthuber M, Märkl B, Messmann H. Endoscopic submucosal dissection in large sessile lesions of the rectosigmoid: learning curve in a European center. *Endoscopy* 2012;44:660-667.
5. Probst A, Ebigo A, Märkl B, et al. Endoscopic submucosal dissection for early rectal neoplasia: experience from a European center. *Endoscopy* 2017;49:222-232.
6. Nishizawa T, Yahagi N. Endoscopic mucosal resection and endoscopic submucosal dissection: technique and new directions. *Curr Opin Gastroenterol* 2017;33:315-319.

7. Probst A, Golger D, Arnholdt H, Messmann H. Endoscopic submucosal dissection of early cancers, flat adenomas, and submucosal tumors in the gastrointestinal tract. *Clin Gastroenterol Hepatol* 2009;7:149-155.
8. Jacques J, Albouys J, Guyot A, et al. Endoscopic submucosal dissection of a laterally spreading tumor in the right colon with a gastroscope after shortening the colon using a new double-balloon platform. *Endoscopy* 2019;51:E364-E365.
9. Jawaid S, Yang D, Draganov PV. Tissue retractor system-assisted endoscopic submucosal dissection of a large rectal tumor with significant fibrosis from direct tattooing. *VideoGIE* 2019;4:84-86.
10. Mortagy M, Mehta N, Parsi MA, et al. Magnetic anchor guidance for endoscopic submucosal dissection and other endoscopic procedures. *World J Gastroenterol* 2017;23:2883-2890.
11. Walter B, Schmidbaur S, Krieger Y, Meining A. Improved endoscopic resection of large flat lesions and early cancers using an external additional working channel (AWC): a case series. *Endosc Int Open* 2019;7:E298-E301.
12. Oyama T. Counter traction makes endoscopic submucosal dissection easier. *Clin Endosc* 2012;45:375-378.
13. Imaeda H, Hosoe N, Kashiwagi K, et al. Advanced endoscopic submucosal dissection with traction. *World J Gastrointest Endosc* 2014;6:286-295.
14. Tsuji K, Yoshida N, Nakanishi H, Takemura K, Yamada S, Doyama H. Recent traction methods for endoscopic submucosal dissection. *World J Gastroenterol* 2016;22:5917-5926.
15. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
16. Higgins JPT, Green S. *Cochrane handbook for systematic reviews of interventions* version 5.1.0 [Internet]. London: Cochrane Training; c2011 [updated 2011 Mar; cited 2019 Feb 14]. Available from: <https://training.cochrane.org/handbook/archive/v5.1/>.
17. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010;25:603-605.
18. Yonezawa J, Kaise M, Sumiyama K, Goda K, Arakawa H, Tajiri H. A novel double-channel therapeutic endoscope ("R-scope") facilitates endoscopic submucosal dissection of superficial gastric neoplasms. *Endoscopy* 2006;38:1011-1015.
19. Gotoda T, Oda I, Takakawa K, Ueda H, Kobayashi T, Kakizoe T. Prospective clinical trial of magnetic-anchor-guided endoscopic submucosal dissection for large early gastric cancer (with videos). *Gastrointest Endosc* 2009;69:10-15.
20. Imaeda H, Hosoe N, Ida Y, et al. Novel technique of endoscopic submucosal dissection using an external grasping forceps for superficial gastric neoplasia. *Dig Endosc* 2009;21:122-127.
21. Motohashi O, Nishimura K, Nakayama N, Takagi S, Yanagida N. Endoscopic submucosal dissection (two-point fixed ESD) for early esophageal cancer. *Dig Endosc* 2009;21:176-179.
22. Uraoka T, Ishikawa S, Kato J, et al. Advantages of using thin endoscope-assisted endoscopic submucosal dissection technique for large colorectal tumors. *Dig Endosc* 2010;22:186-191.
23. Hijikata Y, Ogasawara N, Sasaki M, et al. Endoscopic submucosal dissection with sheath-assisted counter traction using a novel sheath for early gastric cancers. *Hepatogastroenterology* 2012;59:353-356.
24. Okamoto K, Muguruma N, Kitamura S, Kimura T, Takayama T. Endoscopic submucosal dissection for large colorectal tumors using a cross-counter technique and a novel large-diameter balloon overtube. *Dig Endosc* 2012;24 Suppl 1:96-99.
25. Okamoto K, Okamura S, Muguruma N, et al. Endoscopic submucosal dissection for early gastric cancer using a cross-counter technique. *Surg Endosc* 2012;26:3676-3681.
26. Ota M, Nakamura T, Hayashi K, et al. Usefulness of clip traction in the early phase of esophageal endoscopic submucosal dissection. *Dig Endosc* 2012;24:315-318.
27. Ahn JY, Choi KD, Lee JH, et al. Is transnasal endoscope-assisted endoscopic submucosal dissection for gastric neoplasm useful in training beginners? A prospective randomized trial. *Surg Endosc* 2013;27:1158-1165.
28. Higuchi K, Tanabe S, Azuma M, et al. Double-endoscope endoscopic submucosal dissection for the treatment of early gastric cancer accompanied by an ulcer scar (with video). *Gastrointest Endosc* 2013;78:266-273.
29. Matsumoto K, Nagahara A, Ueyama H, et al. Development and clinical usability of a new traction device "medical ring" for endoscopic submucosal dissection of early gastric cancer. *Surg Endosc* 2013;27:3444-3451.
30. Ritsuno H, Sakamoto N, Osada T, et al. Prospective clinical trial of traction device-assisted endoscopic submucosal dissection of large superficial colorectal tumors using the S-O clip. *Surg Endosc* 2014;28:3143-3149.
31. Cai SL, Shi Q, Chen T, Zhong YS. Dental floss traction assists in treating gastrointestinal mucosal tumors by endoscopy. *J Laparoendosc Adv Surg Tech A* 2015;25:571-576.
32. Koike Y, Hirasawa D, Fujita N, et al. Usefulness of the thread-traction method in esophageal endoscopic submucosal dissection: randomized controlled trial. *Dig Endosc* 2015;27:303-309.
33. Noda H, Ogasawara N, Koshino A, et al. Thread-traction with a sheath of polypectomy snare facilitates endoscopic submucosal dissection of early gastric cancers. *Gastroenterol Res Pract* 2016;2016:9415497.
34. Suzuki S, Gotoda T, Kobayashi Y, et al. Usefulness of a traction method using dental floss and a hemoclip for gastric endoscopic submucosal dissection: a propensity score matching analysis (with videos). *Gastrointest Endosc* 2016;83:337-346.
35. Yamada S, Doyama H, Ota R, et al. Impact of the clip and snare method using the prelooping technique for colorectal endoscopic submucosal dissection. *Endoscopy* 2016;48:281-285.
36. Yamasaki Y, Takeuchi Y, Uedo N, et al. Traction-assisted colonic endoscopic submucosal dissection using clip and line: a feasibility study. *Endosc Int Open* 2016;4:E51-E55.
37. Yoshida M, Takizawa K, Ono H, et al. Efficacy of endoscopic submucosal dissection with dental floss clip traction for gastric epithelial neoplasia: a pilot study (with video). *Surg Endosc* 2016;30:3100-3106.
38. Jacques J, Legros R, Rivory J, et al. The "tunnel + clip" strategy standardised and facilitates oesophageal ESD procedures: a prospective, consecutive bi-centric study. *Surg Endosc* 2017;31:4838-4847.
39. Mori H, Kobara H, Nishiyama N, Fujihara S, Matsunaga T, Masaki T. Novel effective and repeatedly available ring-thread counter traction for safer colorectal endoscopic submucosal dissection. *Surg Endosc* 2017;31:3040-3047.
40. Ogata K, Yanai M, Kuriyama K, et al. Double endoscopic intraluminal operation (DEILO) for early gastric cancer: outcome of novel procedure for endoscopic submucosal dissection. *Anticancer Res* 2017;37:343-347.
41. Xie X, Bai JY, Fan CQ, et al. Application of clip traction in endoscopic submucosal dissection to the treatment of early esophageal carcinoma and precancerous lesions. *Surg Endosc* 2017;31:462-468.
42. Hashimoto R, Hirasawa D, Iwaki T, et al. Usefulness of the S-O clip for gastric endoscopic submucosal dissection (with video). *Surg Endosc* 2018;32:908-914.
43. Kitagawa Y, Suzuki T, Hara T, Yamaguchi T. Safety and efficacy of endoscopic submucosal dissection using IT knife nano with clip traction method for early esophageal squamous cell carcinoma. *Surg Endosc* 2018;32:450-455.
44. Matsuzaki I, Hattori M, Hirose K, et al. Magnetic anchor-guided endoscopic submucosal dissection for gastric lesions (with video). *Gastrointest Endosc* 2018;87:1576-1580.
45. Yamasaki Y, Takeuchi Y, Uedo N, et al. Efficacy of traction-assisted colorectal endoscopic submucosal dissection using a clip-and-thread technique: a prospective randomized study. *Dig Endosc* 2018;30:467-476.
46. Yoshida M, Takizawa K, Suzuki S, et al. Conventional versus traction-assisted endoscopic submucosal dissection for gastric neoplasms:

- a multicenter, randomized controlled trial (with video). *Gastrointest Endosc* 2018;87:1231-1240.
47. Zhang Q, Yao X, Cai JQ, Zhang BH, Wang Z. Snare combined with endoclips in endoscopic submucosal dissection with mucosal traction for gastroesophageal neoplasia. *J Gastroenterol Hepatol* 2019;34:1049-1057.
48. Çolak Ş, Gürbulak B, Çakar E, Bektaş H. Resection of mucosal and submucosal gastrointestinal lesions and a double endoscope experience. *JSLs* 2019;23.
49. Ye L, Yuan X, Pang M, et al. Magnetic bead-assisted endoscopic submucosal dissection: a gravity-based traction method for treating large superficial colorectal tumors. *Surg Endosc* 2019;33:2034-2041.
50. Zhang Q, Xing TY, Wang Z. A snare combined with endoclips to assist in endoscopic submucosal dissection for intraepithelial neoplasia in the entire colon and rectum. *Scand J Gastroenterol* 2019;54:114-121.

Supplementary 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	4
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	5
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	5-6 Supplementary 2
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	6
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	6-7
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	9
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	7
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	8

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	9
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	9
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	10
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	10, Table 1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	11
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	11-12
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	12-17
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	12, Figure 2, Supplementary 3
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	14
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	18-21
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	20
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	20-21
FUNDING	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	

Supplementary 2. Search Strategy

PubMed (Date Run: 18/05/19)			
Step	Search strategy	Found	Time
#1	Search “endoscopic mucosal resection”[MeSH Terms] OR (“endoscopic”[All Fields] AND “mucosal”[All Fields] AND “resection”[All Fields]) OR “endoscopic mucosal resection”[All Fields] OR (“endoscopic”[All Fields] AND “submucosal”[All Fields] AND “dissection”[All Fields]) OR “endoscopic submucosal dissection”[All Fields]	7,036	11:30:06
#2	Search “traction”[MeSH Terms] OR “traction”[All Fields]	22,511	11:33:29
#3	#1 OR #2 AND English[lang]	146	11:34:50

Cochrane Central Register of Clinical Trials (Date Run: 18/05/19)		
ID	Search strategy	Found
#1	(“endoscopic submucosal dissection”) (Word variations have been searched):ti,ab,kw	687
#2	(“traction”) (Word variations have been searched)	1,838
#3	#1 AND #2	31

Supplementary 3. Newcastle-Ottawa Scale

Study	A. Selection			B. Comparability		C. Outcome		Score	
	Representativeness of exposed cohort	Selection of non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of the study	Comparability of cohorts on the basis of the design or analysis	Assessment of outcome	Was follow-up long enough for outcomes to occur		Adequacy of follow up of cohorts
A. Magnetic - anchor									
Gotoda et al. (2009) ¹⁹	*	*	*	*	—	*	*	*	7
Matsuzaki et al. (2018) ⁴⁴	*	—	*	*	—	*	—	—	4
Ye et al. (2019) ⁴⁹	*	*	*	*	**	*	*	*	9
B. Double endoscope									
Uraoka et al. (2010) ²²	*	*	*	*	**	*	*	*	9
Higuchi et al. (2013) ²⁸	*	*	*	*	**	*	*	*	9
Ogata et al. (2017) ⁴⁰	*	—	*	*	—	*	*	*	6
Çolak et al. (2019) ⁴⁸	*	—	*	*	—	*	—	—	4
C. Clip involving methods									
Okamoto et al. (2012) ²⁴	*	*	*	*	**	*	—	—	7
Okamoto et al. (2012) ²⁵	*	*	*	*	**	*	—	—	7
Ota et al. (2012) ²⁶	*	*	*	*	**	*	—	—	7
Matsumoto et al. (2013) ²⁹	*	*	*	*	**	*	—	—	7
Cai et al. (2015) ³¹	*	—	*	*	—	*	*	*	6
Suzuki et al. (2016) ³⁴	*	*	*	*	**	*	—	—	7
Yoshida et al. (2016) ³⁷	*	*	*	*	**	*	—	—	7
Yamada et al. (2016) ³⁵	*	*	*	*	**	*	—	—	7
Yamasaki et al. (2016) ³⁶	*	—	*	*	—	*	—	—	4
Noda et al. (2016) ³³	*	*	*	*	**	*	—	—	7
Xie et al. (2017) ⁴¹	*	*	*	*	**	*	—	—	7
Jacques et al. (2017) ³⁸	*	—	*	*	—	*	—	—	4
Kitagawa et al. (2018) ⁴³	*	—	*	*	—	*	*	*	6
Hashimoto et al. (2018) ⁴²	*	*	*	*	**	*	—	—	7
Zhang et al. (2019) ⁴⁷	*	—	*	*	—	*	*	*	6
Zhang et al. (2019) ⁵⁰	*	—	*	*	—	*	*	*	6

Study	A. Selection			B. Comparability		C. Outcome		Score
	Representativeness of exposed cohort	Selection of non-exposed cohort	Ascertainment of exposure	Demonstration that outcome of interest was not present at start of the study	Comparability of cohorts on the basis of the design or analysis	Assessment of outcome	Adequacy of follow up of cohorts	
D. Miscellaneous methods								
Imaeda et al. (2009) ²⁰	*	—	*	*	—	*	—	4
Yonezawa et al. (2006) ¹⁸	*	*	*	*	**	*	*	9
Hijikata et al. (2012) ²³	*	*	*	*	—	*	—	5
Motohashi et al. (2009) ²¹	*	—	*	*	—	*	—	4

According to the scale's "star system", each study receives a score based upon three different perspectives: study selection groups; comparability; and ascertainment of outcome of interest.

A. Selection

1. Representativeness of intervention cohort—a] Selection from general population or general hospital*; b] only selected group of patients; c] no description of inclusion/exclusion criteria
2. Selection of non-intervention cohort—a] drawn from same community as intervention cohort*; b] drawn from different source; c] no description of the derivation of the non-intervention cohort
3. Ascertainment of outcome—a] examination protocol or department's archive*; b] no description
4. Demonstration that outcome was not present at start of study—a] yes*; b] no

B. Comparability

1. Comparability of cohorts on basis of design or analysis—a] study controls for participants' age and gender*; b] study controls for examinations' indication*

C. Outcome

1. Assessment of outcome—a] independent physician's assessment*; b] record linkage*; c] self-report; d] no description
2. Was follow up long—a] yes*; b] no

Adequacy of follow up of cohort—a] complete follow up*; b] minimal loss to follow up ($\leq 20\%$) all ages included, all diseases, or description of those lost suggesting no difference from those followed*; c] follow up rate $< 80\%$ and no description of losses to follow up or description suggesting differences from those followed; d] no statement.