

## Serum uromodulin is associated with but does not predict type 2 diabetes in elderly KORA F4/FF4 study participants

Cornelia Then, Holger Then, Christa Meisinger, Margit Heier, Annette Peters, Wolfgang Koenig, Wolfgang Rathmann, Jürgen Scherberich, Jochen Seissler

### Angaben zur Veröffentlichung / Publication details:

Then, Cornelia, Holger Then, Christa Meisinger, Margit Heier, Annette Peters, Wolfgang Koenig, Wolfgang Rathmann, Jürgen Scherberich, and Jochen Seissler. 2019. "Serum uromodulin is associated with but does not predict type 2 diabetes in elderly KORA F4/FF4 study participants." *The Journal of Clinical Endocrinology & Metabolism* 104 (9): 3795–3802. <https://doi.org/10.1210/jc.2018-02557>.

### Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

**Deutsches Urheberrecht**

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



# Serum Uromodulin Is Associated With But Does Not Predict Type 2 Diabetes in Elderly KORA F4/FF4 Study Participants

Cornelia Then,<sup>1,2</sup> Holger Then,<sup>3</sup> Christa Meisinger,<sup>4,5</sup> Margit Heier,<sup>6</sup> Annette Peters,<sup>6,7</sup> Wolfgang Koenig,<sup>8,9,10</sup> Wolfgang Rathmann,<sup>11</sup> Jürgen Scherberich,<sup>12</sup> and Jochen Seissler<sup>1,2</sup>

<sup>1</sup>Medizinische Klinik und Poliklinik IV, Klinikum der Universität München, LMU München, 80336 Munich, Germany; <sup>2</sup>Clinical Cooperation Group Diabetes, Ludwig-Maximilians-Universität München and Helmholtz Zentrum München, 80336 Munich, Germany; <sup>3</sup>Freie Waldorfschule Augsburg, 86169 Augsburg, Germany; <sup>4</sup>Independent Research Group Clinical Epidemiology, Helmholtz Zentrum München – German Research Center for Environmental Health (GmbH), 85764 Neuherberg, Germany; <sup>5</sup>Chair of Epidemiology at UNIKAT, 86156 Augsburg, Ludwig-Maximilians-Universität München, Germany; <sup>6</sup>Institute of Epidemiology, Helmholtz Zentrum München – German Research Center for Environmental Health (GmbH), 85764 Neuherberg, Germany; <sup>7</sup>Research Unit of Molecular Epidemiology, German Research Center for Environmental Health, 85764 Neuherberg, Germany; <sup>8</sup>Institute of Epidemiology and Biostatistics, University of Ulm, 89081 Ulm, Germany; <sup>9</sup>Deutsches Herzzentrum München, Technische Universität München, 80636 Munich, Germany; <sup>10</sup>DZHK (German Centre for Cardiovascular Research), partner site Munich Heart Alliance, 80636 Munich, Germany; <sup>11</sup>German Diabetes Center, Leibniz Institute at Heinrich Heine University Düsseldorf, Institute of Biometrics and Epidemiology, 40225 Düsseldorf, Germany; and <sup>12</sup>Klinikum München-Harlaching, Teaching Hospital of the Ludwig-Maximilians-Universität, 81545 Munich, Germany

**ORCID numbers:** [0000-0001-5180-1973](https://orcid.org/0000-0001-5180-1973) (C. Then).

**Aims:** Serum uromodulin has recently emerged as promising biomarker for kidney function and was suggested to be associated with type 2 diabetes (T2D) in patients with coronary heart disease. Here, we analyzed the association of serum uromodulin with T2D in the population-based KORA F4/FF4 study.

**Methods:** In 1119 participants of the KORA F4 study aged 62 to 81 years, serum uromodulin was measured, and the association of serum uromodulin with T2D was assessed using logistic and linear regression models stratified for sex. After a mean follow-up time of 6.5 years, 635 participants were re-evaluated. Glucose tolerance status was determined by oral glucose tolerance test at baseline and at the follow-up examination except in cases of known T2D.

**Results:** Serum uromodulin was inversely associated with T2D in the crude analysis and after adjustment for age and body mass index in men ( $P < 0.001$ ) and in women ( $P < 0.05$ ). After further adjustment for estimated glomerular filtration rate, serum uromodulin was significantly inversely associated with T2D in men ( $P < 0.001$ ) but not in women. Serum uromodulin was not associated with prediabetes after multivariate adjustment and did not predict T2D in men or in women after the follow-up time of  $6.5 \pm 0.3$  years.

**Conclusions:** In participants of the KORA F4 study, serum uromodulin is independently associated with T2D in men but is not a predictor of future development of T2D. (*J Clin Endocrinol Metab* 104: 3795–3802, 2019)

---

Abbreviations: BMI, body mass index; eGFR, glomerular filtration rate; HOMA-IR, homeostatic model assessment of insulin resistance; hsCRP, high-sensitivity C-reactive protein; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; sUmod, serum uromodulin; T2D, type 2 diabetes mellitus.

**T**ype 2 diabetes mellitus (T2D) entails a roughly doubled cardiovascular risk (1), but the link between metabolic and vascular disturbances is incompletely understood, and very little is known about potentially protective factors, one of which might be the glycosylphosphatidylinositol-anchored protein uromodulin. Uromodulin is synthesized in tubular cells of the ascending limb of Henle's loop and released into the urine by proteolytic cleavage at a rate of 75 to 200 mg/24 h (2–4). In the urinary tract, the highly abundant uromodulin exerts antilithogenic, anti-infective, and immunomodulatory functions (5–12). The relevance of uromodulin for kidney integrity becomes obvious in the case of mutations of the uromodulin-coding gene leading to severe kidney damage, such as cystic kidney disease, recurrent urinary tract infections, familial juvenile hyperemic nephropathy, and congenital nephrolithiasis (4, 7, 9, 13–15). In addition to these severe uromodulin defects, even small changes in uromodulin concentration and/or function, caused for instance by uromodulin loci variants, may provoke and/or accelerate kidney disease (16–22). *Vice versa*, kidney disease with loss of uromodulin-producing cells due to other reasons entails decreased uromodulin secretion and might thus aggravate the decline of kidney function (23, 24).

Other circumstances may influence uromodulin secretion. A decreased urinary uromodulin was described in patients with diabetes (25, 26). In diabetes, urinary uromodulin may additionally feature an altered glycation pattern, possibly influencing its functionality (27). Recently, a decreased serum uromodulin (sUmod) was described in patients with diabetes who had coronary heart disease (28). These observations suggest a link between diabetes and uromodulin. However, the causal nature of the relation of circulatory uromodulin and diabetes remains to be clarified.

The presence of uromodulin in serum has long been recognized (29) but has gained little interest until recently. Uromodulin concentration in serum is about 1000-fold lower than in the urine (30). Nevertheless, sUmod does not seem to be a “byproduct” only. Uromodulin is actively secreted by endocytotic vesicles from the basolateral side of tubular cells into the interstitial space and circulation (30–32). The physiological functions and significance of uromodulin in the circulation remain largely unknown. Because T2D is often associated with kidney disease, we here aimed to assess the association of sUmod with T2D and its predictive value for T2D in the population-based KORA F4/FF4 cohort.

## Methods

### Study participants

The KORA (Cooperative Health Research in the Region of Augsburg) S4/F4/FF4 cohort study originally involved 4261

participants from the general community in the health survey S4 (1999–2001). Follow-up examinations took place from 2006 to 2008 (KORA F4 with 3080 participants) and from 2013 to 2014 (KORA FF4 with 2279 participants). The loss of participants from S4 to FF4 was due to participants' death ( $n = 455$ ), relocation ( $n = 296$ ), refusal ( $n = 570$ ), illness or lack of time ( $n = 504$ ), and no contact possible ( $n = 157$ ). Recruitment and eligibility criteria for the KORA studies have been described previously (33). The study design, standardized sampling methods, and data collection (medical history, medication, anthropometric measurements, blood pressure) have been described in detail elsewhere (34, 35). All study participants gave written informed consent. The study was approved by the Ethics Committees of the Bavarian Medical Association and the Bayerische Landesärztekammer in adherence to the declaration of Helsinki and Istanbul. In 1119 randomly selected participants aged 62 to 81 years in the KORA F4 cohort, sUmod was measured. Criteria for a preknown diagnosis of diabetes mellitus were a validated medical diagnosis or current use of glucose-lowering agents. After an overnight fasting period, all nondiabetic participants underwent a standard 75-g oral glucose tolerance test. Newly diagnosed diabetes, impaired glucose tolerance (IGT), impaired fasting glucose (IFG), and normal glucose tolerance were defined according to the 1999 World Health Organization diagnostic criteria based on both fasting and postchallenge glucose values (T2D:  $\geq 7.0$  mmol/L fasting and/or  $\geq 11.1$  mmol/L 2-hour glucose; IFG:  $\geq 6.1$  mmol/L and  $< 7.0$  mmol/L; IGT:  $\geq 7.8$  to  $< 11.1$  mmol/L 2-hour glucose). Prediabetes was defined as IFG and/or IGT (2-hour glucose  $\geq 7.8$  to  $< 11.1$  mmol/L). Arterial hypertension was defined as systolic blood pressure  $\geq 140$  mm Hg, diastolic blood pressure  $\geq 90$  mm Hg, or known hypertension with the use of antihypertensive drugs. Of the 1119 participants with sUmod measurements in KORA F4, 130 died, and 354 could not be contacted or declined participation in the FF4 survey. Thus, the study sample in the FF4 examination comprised 635 participants. The follow-up time amounted to  $6.5 \pm 0.3$  years.

### Laboratory measurements

Blood was collected after an overnight fast of at least 8 hours without stasis, and the samples were kept at 4°C until centrifugation. Serum samples were stored at  $-80^{\circ}\text{C}$  until assayed. Except for 2-hour glucose, all blood parameters were determined in fasting blood samples. Measurements of serum creatinine and cystatin C were performed as described elsewhere (36). Glomerular filtration rate (eGFR) was calculated using the Chronic Kidney Disease Epidemiology Collaboration Equation (2009) based on serum creatinine, cystatin C, or both parameters. sUmod was determined using the commercially available uromodulin ELISA kit (Euroimmun AG, Lübeck, Germany) with a lower detection limit of 2 ng/mL, an intra-assay precision of 1.8% to 3.2%, and an interassay precision of 6.6%. The measurement procedure was performed as described previously (32).

### Statistical analyses

Characteristics of the study participants were compared between men and women using *t* tests in case of normally distributed variables. For skewedly distributed variables, Mann-Whitney *U* tests were performed;  $\chi^2$  tests were used to compare binomial proportions. The associations of sUmod

with the outcomes of interest were assessed in logistic regression models in the case of categorical dependent variables and in linear regression models in the case of continuous dependent variables. sUmod was considered an independent variable. In multinomial logistic and linear regression analyses, the association of sUmod with the respective dependent variables was adjusted for confounders as indicated for each observation. The models were fitted for the whole study group and for women and men. Participants with missing covariables were excluded from the respective analyses. The number of participants included in the fully adjusted model is given for each analysis. The level of statistical significance was set at 5%. All calculations were performed using the statistical environment R, version 3.4.4.

## Results

### Study population characteristics

Baseline characteristics of the study population stratified for sex are shown in Table 1. Men had an adverse metabolic and cardiovascular risk profile compared with women, including significantly higher waist circumference, higher systolic and diastolic blood pressure, higher triglycerides, lower high-density lipoprotein cholesterol, and a higher proportion of subjects with T2D. However, women in the current study group had a higher body mass index (BMI), low-density lipoprotein cholesterol, and high-sensitivity C-reactive protein (hsCRP). Women displayed significantly higher sUmod levels than men.

### Association of sUmod with sex, age, height, weight, and BMI

The association of sUmod with sex was highly significant ( $P < 0.001$ ) and largely independent of possible confounders (Table 2). sUmod declined with advancing age. This inverse association was attenuated after correction for eGFR but remained significant ( $P < 0.001$ ) (Table 2). Adjustments for other possible confounders did not further substantially weaken the association of age and decreased sUmod. sUmod was not associated with height after correction for age in men or women (data not shown), whereas weight was inversely related to sUmod in both sexes in the crude analysis and after adjustment for age (men:  $-0.17 \pm 0.04$ ; women:  $-0.26 \pm 0.05$ ;  $P < 0.001$  for both observations). sUmod correlated inversely with BMI in the total study cohort, in men and in women without and with adjustment for age (total cohort:  $-0.23 \pm 0.03$ ; men:  $-0.18 \pm 0.04$ ; women:  $-0.26 \pm 0.05$ ;  $P < 0.001$  for each observation).

### Association of sUmod with prediabetes, diabetes mellitus, and parameters of glucose regulation

Study participants with prediabetes or T2D displayed lower sUmod levels compared with subjects with normal glucose tolerance (Fig. 1). sUmod was inversely associated with prediabetes in the total study cohort and in men but not in women (Table 3). After adjustment for age,

**Table 1. Characteristics of Study Participants**

	All Subjects	Men	Women	$P^a$
n	1119	569	550	
Age, y	70.3 $\pm$ 5.5	70.4 $\pm$ 5.6	70.3 $\pm$ 5.4	n.s. <sup>b</sup>
BMI, kg/m <sup>2</sup>	28.7 $\pm$ 4.5	28.4 $\pm$ 3.9	29.0 $\pm$ 5.0	0.018 <sup>b</sup>
Waist circumference, cm	98.2 $\pm$ 12.2	102.8 $\pm$ 10.4	93.4 $\pm$ 12.0	<0.001 <sup>b</sup>
Systolic blood pressure, mm Hg	128.5 $\pm$ 19.6	132.2 $\pm$ 19.2	124.8 $\pm$ 19.4	<0.001 <sup>b</sup>
Diastolic blood pressure, mm Hg	74.0 $\pm$ 10.0	75.4 $\pm$ 10.3	72.5 $\pm$ 9.7	<0.001 <sup>b</sup>
Arterial hypertension, %	699 (62)	380 (67)	319 (58)	0.002 <sup>c</sup>
HDL cholesterol, mmol/L	1.45 $\pm$ 0.37	1.33 $\pm$ 0.33	1.57 $\pm$ 0.36	<0.001 <sup>b</sup>
LDL cholesterol, mmol/L	3.64 $\pm$ 0.93	3.52 $\pm$ 0.9	3.76 $\pm$ 0.95	<0.001 <sup>b</sup>
Triglycerides, mmol/L	1.27 (0.94; 1.77)	1.35 (0.94; 1.90)	1.23 (0.93; 1.66)	0.009 <sup>d</sup>
Prediabetes, %	290 (26)	151 (27)	139 (25)	n.s. <sup>c</sup>
T2D, %	218 (19)	131 (23)	87 (16)	0.006 <sup>c</sup>
HbA1c, mmol/mol	38 (36; 41)	38 (36; 42)	38 (36; 41)	n.s. <sup>d</sup>
HbA1c, %	5.6 (5.4; 5.9)	5.6 (5.4; 6.0)	5.6 (5.4; 5.9)	n.s. <sup>d</sup>
hsCRP, nmol/L	14.7 (7.5; 30.4)	13.8 (7.0; 29.0)	15.9 (8.6; 32.3)	0.046 <sup>d</sup>
eGFR, mL/min/1.73 m <sup>2</sup>	77.9 (67.2; 87.8)	79.1 (68.2; 87.4)	76.8 (66.0; 88.1)	n.s. <sup>d</sup>
Physically inactive, <sup>e</sup> %	556 (50)	293 (51)	263 (48)	n.s. <sup>c</sup>
sUmod, ng/mL	152.5 (110.4; 207.6)	138.6 (103.3; 187.6)	169.0 (120.4; 224.7)	<0.001 <sup>d</sup>

Values are means  $\pm$  SD, median (first quartile; third quartile), or proportion (%).

Abbreviations: HDL, high-density lipoprotein; LDL, low-density lipoprotein; n.s., not significant.

<sup>a</sup>The  $P$  value is related to the null hypothesis of no sex differences.

<sup>b</sup>Student  $t$  test.

<sup>c</sup>Chi square test.

<sup>d</sup>Mann-Whitney  $U$  test.

<sup>e</sup>Physically active: participating in sports  $\geq 1$  h/wk in winter and summer.

**Table 2. Association of sUmod With Age and Sex: Results of Multinomial Linear/Logistic Regression Models**

Age	Sex (Female)
Without adjustment	
-3.032 ± 0.392 <sup>a</sup>	26.242 ± 4.336 <sup>a</sup>
Adjustment for eGFR	
-1.666 ± 0.412 <sup>a</sup>	26.900 ± 4.114 <sup>a</sup>
Multivariate adjustment <sup>b</sup>	
-1.313 ± 0.418 <sup>d</sup>	Multivariate adjustment <sup>c</sup> 23.723 ± 4.207 <sup>a</sup>

In the fully adjusted model, n = 1065.

<sup>a</sup>P < 0.001.

<sup>b</sup>Adjusted for sex, BMI, diabetes, hypertension, low-density lipoprotein, eGFR, hsCRP, myocardial infarction or stroke, smoking behavior, and physical activity.

<sup>c</sup>Adjusted for age, BMI, diabetes, hypertension, low-density lipoprotein, eGFR, hsCRP, myocardial infarction or stroke, smoking behavior, and physical activity.

<sup>d</sup>P < 0.002.

sex, and BMI, the associations of sUmod with prediabetes were no longer significant. sUmod correlated inversely with T2D in the total cohort in men and in women without adjustment and after adjustment for sex, age, and BMI (Table 3). However, in women, further adjustment for eGFR attenuated the association to nonsignificance.

After adjustment for age, sex, BMI, and eGFR, sUmod was inversely associated with HbA1c as a continuous


**Table 3. Relation of sUmod and Prevalent Prediabetes and T2D: Results of Multinomial Logistic Regression Models**

	Prediabetes	T2D
Without adjustment		
Total study cohort	0.8 (0.69–0.92) <sup>a</sup>	0.55 (0.47–0.65) <sup>b</sup>
Men	0.79 (0.65–0.98) <sup>a</sup>	0.55 (0.44–0.69) <sup>b</sup>
Women	0.82 (0.66–1.01)	0.57 (0.44–0.73) <sup>b</sup>
Adjustment for age, sex, and BMI		
Total study cohort	0.91 (0.78–1.07)	0.67 (0.56–0.81) <sup>b</sup>
Men	0.88 (0.71–1.1)	0.63 (0.5–0.81) <sup>b</sup>
Women	0.95 (0.76–1.2)	0.73 (0.55–0.97) <sup>a</sup>
Adjustment for age, sex, BMI, and eGFR		
Total study cohort	0.89 (0.75–1.05)	0.68 (0.56–0.82) <sup>b</sup>
Men	0.83 (0.65–1.05)	0.62 (0.48–0.81) <sup>b</sup>
Women	0.95 (0.75–1.21)	0.75 (0.56–1.01)
Adjustment for age, sex, BMI, eGFR, and hsCRP		
Total study cohort	0.91 (0.77–1.08)	0.68 (0.56–0.83) <sup>b</sup>
Men	0.84 (0.66–1.07)	0.63 (0.48–0.82) <sup>b</sup>
Women	0.98 (0.77–1.24)	0.76 (0.56–1.01)

Adjusted ORs (95% CI) for glucose regulation (prediabetes and T2D with normal glucose tolerance as reference) as dependent variable and sUmod as independent variable (per standard deviation). In the fully adjusted model (total cohort; men; women), n = 872; 426; 446 for the analysis of prediabetes (prediabetes yes: 288; 150; 138); in the fully adjusted model (total cohort; men; women), n = 797; 405; 392 for the analysis of T2D (T2D yes: 213; 129; 84).

<sup>a</sup>P < 0.05.

<sup>b</sup>P < 0.001.



**Figure 1.** sUmod levels according to glucose tolerance status. Given are medians, interquartile ranges, and 10th to 90th percentiles. \*\*P < 0.01; \*\*\*P < 0.001.

variable in the total study cohort (coefficient -0.09 ± 0.03; P = 0.003) and in men (-0.14 ± 0.05; P = 0.005) but not in women (-0.053 ± 0.06; P = 0.13). Compatible with the lack of an association of sUmod with prediabetes, the relation of sUmod with moderately elevated HbA1c levels in the prediabetic range (39 to 46 mmol/mol; 5.7% to 6.4%) was not significant in women and was no longer significant in the total study cohort and in men after adjustment for age and BMI. HbA1c values in the diabetic range (≥48 mmol/mol; 6.5%) were inversely associated with sUmod in the total cohort and in men after adjustment for age, BMI, and eGFR. In women, the inverse relation of sUmod and HbA1c values ≥48 mmol/mol (6.5%) was no longer significant after adjustment for age and BMI (37).

sUmod was associated with a lower fasting glucose and a lower insulin resistance as determined by homeostatic model assessment of insulin resistance (HOMA-IR). In women, the association with HOMA-IR was attenuated to nonsignificance after adjustment for age and BMI. After adjustment for age, BMI, and eGFR, fasting insulin was weakly but significantly inversely associated with sUmod in men but not in women, whereas a relation of proinsulin with sUmod could not be established after correction for age, sex, BMI, and eGFR (37).

sUmod was inversely related to hsCRP in men after adjustment for age, BMI, and eGFR ( $-0.10 \pm 0.04$ ;  $P < 0.05$ ). In women, hsCRP and sUmod correlated without correction for confounders but not after multivariate adjustment (data not shown). The risk reduction for T2D in study participants with higher sUmod values was largely independent of hsCRP because correction for hsCRP did not change the level of significance (Table 3).

#### Lacking association of sUmod with T2D in follow-up

sUmod was not a predictor for prediabetes or T2D in men or in women after multivariate adjustment in the follow-up examination after  $6.5 \pm 0.3$  years (OR for T2D in men: 1.35; 95% CI, 0.87 to 2.11; women: OR, 0.77; 95% CI, 0.51 to 1.17) (Table 4). Fittingly, HbA1c (men:  $-0.001 \pm 0.046$ ; women:  $-0.071 \pm 0.038$ ), fasting glucose (men:  $0.056 \pm 0.049$ ; women:  $-0.031 \pm 0.038$ ), fasting insulin (men:  $-0.119 \pm 0.064$ ; women:  $-0.073 \pm 0.044$ ), and HOMA-IR (men:  $-0.070 \pm 0.059$ ; women:  $-0.052 \pm 0.033$ ) were not significantly associated with baseline sUmod in men or in women in the follow-up survey after adjustment for age, BMI, eGFR, and the respective baseline parameters.

## Discussion

Results from this study show the association of sUmod with T2D in a population-based cohort. Our results describing a strong link of T2D with sUmod are in line with previous studies that predominantly involved participants with kidney and/or coronary disease (28, 38). In contrast to previous data (28), we here also show an association of sUmod with HOMA-IR and fasting insulin. The stratification for sex in our study revealed a highly significant association of sUmod with T2D in men but not in women. To our knowledge, a different

association of uromodulin and T2D depending on sex has not been described. Another important finding of our study is that sUmod is not an independent marker of prediabetes as defined by increased fasting glucose, impaired glucose tolerance, and/or moderately increased HbA1c levels. In our cohort, participants with prediabetes were asymptomatic, and prediabetes was mainly detected by a screening test. Thus, the majority of individuals with prediabetes may not have developed diabetic nephropathy, which may explain the unchanged uromodulin serum levels.

The current data clearly demonstrate that sUmod is not a predictor of future prediabetes or T2D, which is in line with the lack of an association with prediabetes but contradicts the results of Leisher *et al.* (28), which describe lower sUmod values in individuals developing diabetes within a follow-up time of 3.5 years. The discrepant results may arise from differences in the study populations [in the study of Leisher *et al.* (28), 55% of the participants suffered from advanced coronary heart disease] and from the multivariate adjustments of the longitudinal analyses of our study. Regarding our current data, it can be largely excluded that sUmod plays a causative role in diabetes pathogenesis. The inverse association of sUmod with diabetes may be due to a suppression of uromodulin secretion. Experimental data investigating this possibility are currently lacking. Another possibility for the eGFR-independent relation of sUmod and diabetes is an early diabetes-related kidney injury that is underestimated by GFR calculations. In early kidney injury, a sUmod decrease may precede the eGFR decline determined by the endogenous filtration markers creatinine or cystatin C because a progressive nephron loss may initially be compensated for by an increased hydraulic pressure and subsequent glomerular hyperfiltration (31, 32). Therefore, in early kidney

**Table 4. Relation of sUmod and Incident Prediabetes and T2D: Results of Multinomial Logistic Regression Models**

	Prediabetes	T2D	Prediabetes or T2D
	Without adjustment		
Total study cohort	0.90 (0.69–1.18)	0.83 (0.64–1.08)	0.81 (0.66–1.00)
Men	0.97 (0.65–1.45)	1.07 (0.73–1.56)	0.85 (0.60–1.17)
Women	0.91 (0.62–1.34)	0.66 (0.45–0.96) <sup>a</sup>	0.88 (0.67–1.16)
	Adjustment for age, sex, BMI, eGFR, and hsCRP		
Total study cohort	1.03 (0.77–1.38)	1.0 (0.75–1.35)	1.01 (0.80–1.27)
Men	1.03 (0.68–1.57)	1.35 (0.87–2.11)	1.00 (0.70–1.44)
Women	1.06 (0.7–1.6)	0.77 (0.51–1.17)	1.02 (0.75–1.39)
n <sup>b</sup>	112; 61; 51	72; 41; 31	184; 102; 82

Adjusted ORs (95% CI) for glucose regulation (prediabetes and T2D with normal glucose tolerance as reference) as dependent variable and sUmod as independent variable (per standard deviation).

<sup>a</sup> $P < 0.05$ .

<sup>b</sup>Number of participants included in the fully adjusted model who met criteria (total cohort; men; women).

disease, correction for eGFR may not exclude that the sUmod decline is in fact primarily related to the reduction in nephron number. This presumption may be especially true for diabetic nephropathy, in which hyperfiltration is typical.

In our cohort aged 62 to 81 years, sUmod was lower in older participants. The decline of uromodulin with advancing age is well known (32, 39, 40) and is mostly attributed to declining kidney function and a reduction in nephron numbers at higher age. In our study, eGFR reduction only partially explained the age-related sUmod decrease. The attribution of sUmod values to nephron mass as the only determinant remains open to further study considering the lack of an association of sUmod with height, the inverse relation to weight, and the considerably higher sUmod levels in women compared with men in our study population. Although sex-related data regarding nephron numbers in healthy women and men are lacking, higher sUmod in women is probably not due to a (relatively) higher nephron mass. Rather, hormonal and receptor-related factors may be involved. We assume that higher sUmod values in women may mirror a sex-related nephroprotective role of the molecule because women suffering from chronic kidney disease reach end-stage renal disease 3 to 5 years later than men (41). However, our data suggest that sUmod is also subjected to regulation processes beside kidney mass and function. sUmod joins various other, well-known metabolic parameters featuring more favorable profiles in women than in men. Interestingly, the inverse association of sUmod with prediabetes, T2D, HbA1c, fasting insulin, and insulin resistance was stronger in men than in women. The reasons for sex-related differences in the link between sUmod and T2D are unknown. Because women have generally higher sUmod values, the possible suppressive effect of diabetes may be less obvious. Hormonal parameters might influence uromodulin quantity as well as uromodulin function. For example, uromodulin isolated from the urine of pregnant women has different immunomodulatory properties compared with uromodulin from men and nonpregnant women, probably due to an altered uromodulin glycosylation state (42). Whether such differences are also detectable in sUmod remains to be clarified.

Overall, the physiological functions of uromodulin in the circulation and thus possible causal interferences of a reduced sUmod in diabetes with outcome are largely unknown. Similar to the known uromodulin capacities in urine, immunomodulatory properties in the blood stream have been proposed (43–46). Uromodulin receptors in the body remain to be determined for the most part but possibly include Toll-like receptor 4, by which uromodulin may activate myeloid dendritic cells (47). In the

current study, sUmod correlated inversely with hsCRP in men but not significantly in women after multivariate adjustment. Because two studies involving patients admitted for coronary angiography detected an inverse association of sUmod and cardiovascular events and mortality (38, 48), it is tempting to speculate that sUmod is one link between diabetes and cardiovascular complications. Another possibility is that very early kidney affection is not adequately assessable to date and plays a more prominent role for cardiovascular complications than previously assumed.

### Study strengths and limitations

The major strengths of our study are the population-based design with a large, well-characterized, community-based cohort and the follow-up time of 6.5 years. All participants without prediagnosed diabetes underwent an oral glucose tolerance test at baseline and in the follow-up examination. However, only participants aged 62 to 81 years were included; therefore, the relation of sUmod with metabolic disturbances in younger subjects remains to be confirmed.

### Conclusions

We assessed the association of sUmod with T2D. The current data indicate that sUmod is a valuable biomarker for T2D in elderly individuals but is not a predictor for future T2D, suggesting that the presence of T2D causes a decrease of uromodulin serum levels. Preclinical studies are needed to clarify the mechanistic basis of the reduced sUmod levels in T2D. Further clinical studies will determine whether the reduced sUmod in T2D predicts diabetic nephropathy and cardiovascular complications in individuals with diabetes.

### Acknowledgments

We thank Victor Herbst, Matthias Block, and Wolfgang Schlumberger for the gift of the uromodulin assay and quality control. The KORA study was initiated and financed by the Helmholtz Zentrum München – German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research (BMBF) and by the State of Bavaria. The KORA-Study Group consists of A. Peters (speaker), H. Schulz, L. Schwettmann, R. Leidl, M. Heier, K. Strauch, and their co-workers, who are responsible for the design and conduct of the KORA studies. We gratefully acknowledge the contribution of all field staff members conducting the KORA F4 study and thank all study participants.

**Financial Support:** The study was supported by the Deutsche Forschungsgemeinschaft (RA-45913/3-1), the Virtual Diabetes Institute (Helmholtz Zentrum München), the Clinical Cooperation Group Diabetes (Ludwig-Maximilians-Universität München and Helmholtz Zentrum München), and the Federal Ministry of Health and the Ministry of Innovation, Science,

Research and Technology of the state North Rhine Westphalia. The study sponsor was not involved in the design of the study; the collection, analysis, and interpretation of data; writing the report; or the decision to submit the report for publication.

**Author Contributions:** C.M., M.H., A.P., W.K., W.R., J. Scherberich, and J. Seissler conceived and designed the study. C.T., C.M., M.H., A.P., W.K., W.R., J. Scherberich, and J. Seissler collected the data. C.T., H.T., J. Scherberich, and J. Seissler analyzed the data, interpreted the results, and wrote the manuscript. All authors revised the manuscript critically for important intellectual content and approved the final version. C.T. and J. Seissler are the guarantors of this work.

**Correspondence and Reprint Requests:** Cornelia Then, MD, Medizinische Klinik und Poliklinik IV – Klinikum der Ludwig-Maximilians-Universität, Ziemssenstraße 1, 80336 München, Germany. E-mail: [cornelia.then@med.uni-muenchen.de](mailto:cornelia.then@med.uni-muenchen.de).

**Disclosure Summary:** The authors have nothing to disclose.

## References

- Huang D, Refaat M, Mohammedi K, Jayyousi A, Al Suwaidi J, Abi Khalil C. Macrovascular complications in patients with diabetes and prediabetes. *BioMed Res Int*. 2017;2017:7839101.
- Cavallone D, Malagolini N, Serafini-Cessi F. Mechanism of release of urinary Tamm-Horsfall glycoprotein from the kidney GPI-anchored counterpart. *Biochem Biophys Res Commun*. 2001; 280(1):110–114.
- Schaeffer C, Santambrogio S, Perucca S, Casari G, Rampoldi L. Analysis of uromodulin polymerization provides new insights into the mechanisms regulating ZP domain-mediated protein assembly. *Mol Biol Cell*. 2009;20(2):589–599.
- Devuyst O, Olinger E, Rampoldi L. Uromodulin: from physiology to rare and complex kidney disorders. *Nat Rev Nephrol*. 2017; 13(9):525–544.
- Pak J, Pu Y, Zhang ZT, Hasty DL, Wu XR. Tamm-Horsfall protein binds to type 1 fimbriated *Escherichia coli* and prevents *E. coli* from binding to uroplakin Ia and Ib receptors. *J Biol Chem*. 2001; 276(13):9924–9930.
- Raffi HS, Bates JM, Jr, Laszik Z, Kumar S. Tamm-horsfall protein protects against urinary tract infection by proteus mirabilis. *J Urol*. 2009;181(5):2332–2338.
- Bates JM, Raffi HM, Prasad K, Mascarenhas R, Laszik Z, Maeda N, Hultgren SJ, Kumar S. Tamm-Horsfall protein knockout mice are more prone to urinary tract infection: rapid communication. *Kidney Int*. 2004;65(3):791–797.
- Serafini-Cessi F, Monti A, Cavallone D. N-Glycans carried by Tamm-Horsfall glycoprotein have a crucial role in the defense against urinary tract diseases. *Glycoconj J*. 2005;22(7-9):383–394.
- Mo L, Huang HY, Zhu XH, Shapiro E, Hasty DL, Wu XR. Tamm-Horsfall protein is a critical renal defense factor protecting against calcium oxalate crystal formation. *Kidney Int*. 2004;66(3): 1159–1166.
- Kreft B, Jabs WJ, Laskay T, Klinger M, Solbach W, Kumar S, van Zandbergen G. Polarized expression of Tamm-Horsfall protein by renal tubular epithelial cells activates human granulocytes. *Infect Immun*. 2002;70(5):2650–2656.
- Säemann MD, Weichhart T, Hörl WH, Zlabinger GJ. Tamm-Horsfall protein: a multilayered defence molecule against urinary tract infection. *Eur J Clin Invest*. 2005;35(4):227–235.
- Darisipudi MN, Thomasova D, Mulay SR, Brech D, Noessner E, Liapis H, Anders H-J. Uromodulin triggers IL-1 $\beta$ -dependent innate immunity via the NLRP3 inflammasome. *J Am Soc Nephrol*. 2012; 23(11):1783–1789.
- Hart TC, Gorry MC, Hart PS, Woodard AS, Shihabi Z, Sandhu J, Shirts B, Xu L, Zhu H, Barmada MM, Bleyer AJ. Mutations of the UMOD gene are responsible for medullary cystic kidney disease 2 and familial juvenile hyperuricaemic nephropathy. *J Med Genet*. 2002;39(12):882–892.
- Dahan K, Devuyst O, Smaers M, Vertommen D, Loute G, Poux JM, Viron B, Jacquot C, Gagnadoux MF, Chauveau D, Büchler M, Cochat P, Cosyns JP, Mougnot B, Rider MH, Antignac C, Verellen-Dumoulin C, Pirson Y. A cluster of mutations in the UMOD gene causes familial juvenile hyperuricemic nephropathy with abnormal expression of uromodulin. *J Am Soc Nephrol*. 2003; 14(11):2883–2893.
- Williams SE, Reed AAC, Galvanovskis J, Antignac C, Goodship T, Karet FE, Kotanko P, Lhotta K, Morinière V, Williams P, Wong W, Rorsman P, Thakker RV. Uromodulin mutations causing familial juvenile hyperuricaemic nephropathy lead to protein maturation defects and retention in the endoplasmic reticulum. *Hum Mol Genet*. 2009;18(16):2963–2974.
- Garimella PS, Biggs ML, Katz R, Ix JH, Bennett MR, Devarajan P, Kestenbaum BR, Siscovick DS, Jensen MK, Shlipak MG, Chaves PHM, Sarnak MJ. Urinary uromodulin, kidney function, and cardiovascular disease in elderly adults. *Kidney Int*. 2015;88(5): 1126–1134.
- Olden M, Corre T, Hayward C, Toniolo D, Ulivi S, Gasparini P, Pistis G, Hwang S-J, Bergmann S, Campbell H, Cocca M, Gandin I, Giroto G, Glaudemans B, Hastie ND, Loffing J, Polasek O, Rampoldi L, Rudan I, Sala C, Traglia M, Vollenweider P, Vuckovic D, Youhanna S, Weber J, Wright AF, Kutalik Z, Bochud M, Fox CS, Devuyst O. Common variants in UMOD associate with urinary uromodulin levels: a meta-analysis. *J Am Soc Nephrol*. 2014;25(8): 1869–1882.
- Bleyer AJ, Hart PS, Kmoch S. Hereditary interstitial kidney disease. *Semin Nephrol*. 2010;30(4):366–373.
- Ekici AB, Hackenbeck T, Morinière V, Pannes A, Buettner M, Uebe S, Janka R, Wiesener A, Hermann I, Grupp S, Hornberger M, Huber TB, Isbel N, Mangos G, McGinn S, Soreth-Rieke D, Beck BB, Uder M, Amann K, Antignac C, Reis A, Eckardt KU, Wiesener MS. Renal fibrosis is the common feature of autosomal dominant tubulointerstitial kidney diseases caused by mutations in mucin 1 or uromodulin. *Kidney Int*. 2014;86(3):589–599.
- Reznichenko A, Böger CA, Snieder H, van den Born J, de Borst MH, Damman J, van Dijk MCRF, van Goor H, Hepkema BG, Hillebrands JL, Leuvenink HGD, Niesing J, Bakker SJL, Seelen M, Navis G; REGaTTA (Renal GeneTics TrAnplantation) Groningen group. UMOD as a susceptibility gene for end-stage renal disease. *BMC Med Genet*. 2012;13(1):78.
- Köttgen A, Hwang S-J, Larson MG, Van Eyk JE, Fu Q, Benjamin EJ, Dehghan A, Glazer NL, Kao WHL, Harris TB, Gudnason V, Shlipak MG, Yang Q, Coresh J, Levy D, Fox CS. Uromodulin levels associate with a common UMOD variant and risk for incident CKD. *J Am Soc Nephrol*. 2010;21(2):337–344.
- Lhotta K. Uromodulin and chronic kidney disease. *Kidney Blood Press Res*. 2010;33(5):393–398.
- Chakraborty J, Below AA, Solaiman D. Tamm-Horsfall protein in patients with kidney damage and diabetes. *Urol Res*. 2004;32(2): 79–83.
- Tsai C-Y, Wu T-H, Yu C-L, Lu J-Y, Tsai Y-Y. Increased excretions of beta2-microglobulin, IL-6, and IL-8 and decreased excretion of Tamm-Horsfall glycoprotein in urine of patients with active lupus nephritis. *Nephron*. 2000;85(3):207–214.
- Bernard AM, Ouled AA, Lauwerys RR, Lambert A, Vandeleene B. Pronounced decrease of Tamm-Horsfall proteinuria in diabetics. *Clin Chem*. 1987;33(7):1264.
- Holmquist P, Torffvit O, Jørgensen PE, Tørring N, Nexø E, Sjöblad S. Early urinary changes in Tamm-Horsfall protein and epidermal growth factor in diabetic children. *Pediatr Nephrol*. 2001;16(6):488–492.
- Chang C-C, Chen C-Y, Huang C-H, Wu C-L, Wu H-M, Chiu P-F, Kor C-T, Chen T-H, Chang G-D, Kuo C-C, Wen H-C, Huang C-Y,

- Chang C-H. Urinary glycosylated uromodulin in diabetic kidney disease. *Clin Sci (Lond)*. 2017;131(15):1815–1829.
28. Leiherer A, Muendlein A, Saely CH, Kinz E, Brandtner EM, Fraunberger P, Drexel H. Serum uromodulin is associated with impaired glucose metabolism. *Medicine (Baltimore)*. 2017;96(5):e5798.
  29. Thornley C, Dawnay A, Cattell WR. Human Tamm-Horsfall glycoprotein: urinary and plasma levels in normal subjects and patients with renal disease determined by a fully validated radioimmunoassay. *Clin Sci (Lond)*. 1985;68(5):529–535.
  30. El-Achkar TM, McCracken R, Liu Y, Heitmeier MR, Bourgeois S, Rysse J, Wu X-R. Tamm-Horsfall protein translocates to the basolateral domain of thick ascending limbs, interstitium, and circulation during recovery from acute kidney injury. *Am J Physiol Renal Physiol*. 2013;304(8):F1066–F1075.
  31. Scherberich JE, Gruber R, Nockher WA, Christensen EI, Schmitt H, Herbst V, Block M, Kaden J, Schlumberger W. Serum uromodulin: a marker of kidney function and renal parenchymal integrity. *Nephrol Dial Transplant*. 2018;33(2):284–295.
  32. Steubl D, Block M, Herbst V, Nockher WA, Schlumberger W, Satanovskij R, Angermann S, Hasenau A-L, Stecher L, Heemann U, Rengers L, Scherberich J. Plasma uromodulin correlates with kidney function and identifies early stages in chronic kidney disease patients. *Medicine (Baltimore)*. 2016;95(10):e3011.
  33. Holle R, Happich M, Löwel H, Wichmann HE; MONICA/KORA Study Group. KORA: a research platform for population based health research. *Gesundheitswesen*. 2005;67(Suppl 1):S19–S25.
  34. Rathmann W, Strassburger K, Heier M, Holle R, Thorand B, Giani G, Meisinger C. Incidence of Type 2 diabetes in the elderly German population and the effect of clinical and lifestyle risk factors: KORA S4/F4 cohort study. *Diabet Med*. 2009;26(12):1212–1219.
  35. Meisinger C, Rückert IM, Rathmann W, Döring A, Thorand B, Huth C, Kowall B, Koenig W. Retinol-binding protein 4 is associated with prediabetes in adults from the general population: the Cooperative Health Research in the Region of Augsburg (KORA) F4 Study. *Diabetes Care*. 2011;34(7):1648–1650.
  36. Seissler J, Feghelm N, Then C, Meisinger C, Herder C, Koenig W, Peters A, Roden M, Lechner A, Kowall B, Rathmann W. Vaso-regulatory peptides pro-endothelin-1 and pro-adrenomedullin are associated with metabolic syndrome in the population-based KORA F4 study. *Eur J Endocrinol*. 2012;167(6):847–853.
  37. Then C, Then H, Meisinger C, Heier M, Peters A, Koenig W, Rathmann W, Scherberich J, Seissler J. Data from: Serum uromodulin is associated with but does not predict type 2 diabetes in elderly KORA F4/FF4 study participants. Open Data LMU. Deposited 15 February 2019. <https://doi.org/10.5282/ubm/data.142>.
  38. Delgado GE, Kleber ME, Scharnagl H, Krämer BK, März W, Scherberich JE. Serum uromodulin and mortality risk in patients undergoing coronary angiography. *J Am Soc Nephrol*. 2017;28(7):2201–2210.
  39. Risch L, Lhotta K, Meier D, Medina-Escobar P, Nydegger UE, Risch M. The serum uromodulin level is associated with kidney function. *Clin Chem Lab Med*. 2014;52(12):1755–1761.
  40. Fedak D, Kuźniewski M, Fugiel A, Wiczorek-Surdacka E, Przepiórkowska-Hoyer B, Jasik P, Miarka P, Dumnicka P, Kapusta M, Solnica B, Sułowicz W. Serum uromodulin concentrations correlate with glomerular filtration rate in patients with chronic kidney disease. *Pol Arch Med Wewn*. 2016;126(12):995–1004.
  41. Helve J, Kramer A, Abad-Diez JM, Couchoud C, de Arriba G, de Meester J, Evans M, Glaudet F, Grönhagen-Riska C, Heaf JG, Lezaic V, Nordio M, Palsson R, Pechter Ü, Resic H, Santamaria R, Santiuste de Pablos C, Massy ZA, Zurriaga Ó, Jager KJ, Finne P. Factors associating with differences in the incidence of renal replacement therapy among elderly: data from the ERA-EDTA Registry. *Nephrol Dial Transplant*. 2018;33(8):1428–1435.
  42. Easton RL, Patankar MS, Clark GF, Morris HR, Dell A. Pregnancy-associated changes in the glycosylation of tamm-horsfall glycoprotein. Expression of sialyl Lewis(x) sequences on core 2 type O-glycans derived from uromodulin. *J Biol Chem*. 2000;275(29):21928–21938.
  43. Rhodes DCJ, Hinsman EJ, Rhodes JA. Tamm-Horsfall glycoprotein binds IgG with high affinity. *Kidney Int*. 1993;44(5):1014–1021.
  44. Rhodes DCJ. Binding of Tamm-Horsfall protein to complement 1q measured by ELISA and resonant mirror biosensor techniques under various ionic-strength conditions. *Immunol Cell Biol*. 2000;78(5):474–482.
  45. Hession C, Decker JM, Sherblom AP, Kumar S, Yue CC, Mattaliano RJ, Tizard R, Kawashima E, Schmeissner U, Heletky S, Chow EP, Burne CA, Shaw A, Muchmore AV. Uromodulin (Tamm-Horsfall glycoprotein): a renal ligand for lymphokines. *Science*. 1987;237(4821):1479–1484.
  46. Wu TH, Li KJ, Yu CL, Tsai CY. Tamm-Horsfall protein is a potent immunomodulatory molecule and a disease biomarker in the urinary system. *Molecules*. 2018;23(1):E200.
  47. Säemann MD, Weichhart T, Zeyda M, Staffler G, Schunn M, Stuhlmeier KM, Sobanov Y, Stulnig TM, Akira S, von Gabain A, von Ahsen U, Hörl WH, Zlabinger GJ. Tamm-Horsfall glycoprotein links innate immune cell activation with adaptive immunity via a Toll-like receptor-4-dependent mechanism. *J Clin Invest*. 2005;115(2):468–475.
  48. Leiherer A, Muendlein A, Saely CH, Ebner J, Brandtner EM, Fraunberger P, Drexel H. Serum uromodulin is a predictive biomarker for cardiovascular events and overall mortality in coronary patients. *Int J Cardiol*. 2017;231:6–12.