

Comment on “Measurability of nonequilibrium thermodynamics in terms of the Hamiltonian of mean force”

Peter Talkner, Peter Hänggi

Angaben zur Veröffentlichung / Publication details:

Talkner, Peter, and Peter Hänggi. 2020. “Comment on ‘Measurability of nonequilibrium thermodynamics in terms of the Hamiltonian of mean force.’” *Physical Review E* 102 (6): 066101. <https://doi.org/10.1103/physreve.102.066101>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



Comment on “Measurability of nonequilibrium thermodynamics in terms of the Hamiltonian of mean force”

Peter Talkner¹ and Peter Hänggi^{1,2,3}

¹Department of Physics, University of Augsburg, D-86135 Augsburg, Germany

²Nanosystems Initiative Munich, D-80799 München, Germany

³Department of Physics, National University of Singapore, Singapore 117546



(Received 29 June 2020; accepted 23 November 2020; published 21 December 2020)

In a recent paper [Phys. Rev. E **101**, 050101(R) (2020)] an attempt is presented to formulate the nonequilibrium thermodynamics of an open system in terms of the Hamiltonian of mean force. The purpose of the present comment is to clarify severe restrictions of this approach and to stress that recently noted ambiguities [Phys. Rev. E **94**, 022143 (2016)] of fluctuating thermodynamic potentials cannot be removed in the suggested way.

DOI: 10.1103/PhysRevE.102.066101

The Hamiltonian of mean force [1], such as its classical precursor, the potential of mean force [2,3], encompasses the complete information on the equilibrium thermodynamical behavior of an open system. It is defined in terms of the bath-renormalized Boltzmann factor at the inverse temperature β , given by the expression,

$$e^{-\beta H^*(\beta, \mu)} = Z_B^{-1}(\beta, \mu) \text{Tr}_B e^{-\beta H_{\text{tot}}(\mu)}, \quad (1)$$

where $H^*(\beta, \mu)$ and $H_{\text{tot}}(\mu) = H_S(\mu) + H_{SB}(\mu) + H_B(\mu)$ denote the Hamiltonian of mean force, the Hamiltonian of the system $H_S(\mu)$, and the environment $H_B(\mu)$, including the mutual interaction $H_{SB}(\mu)$, respectively. With μ , all externally controllable parameters, including those parameters λ , exclusively acting on the considered open system, but also including all other globally acting parameters, such as electric and magnetic fields, are specified. Furthermore, Tr_B indicates the partial trace over the environmental Hilbert space for quantum systems and, for a classical system, the integral over the environmental phase space with respect to a properly permutation-symmetry-adapted dimensionless volume element. With the normalization by the partition function of the environment, $Z_B(\beta, \mu) = \text{Tr}_B e^{-\beta H_B(\mu)}$, the Hamiltonian of mean force agrees with the bare system Hamiltonian for isolated systems; for further details see Refs. [4,5]. With the partition function for the Hamiltonian of mean force $Z_S(\beta, \mu) = \text{Tr}_S e^{-\beta H^*(\beta, \mu)} = Z_{\text{tot}}(\beta, \mu)/Z_B(\beta, \mu)$ and the resulting Helmholtz free energy $F_S(\beta, \mu) = -\beta^{-1} \ln Z_S(\beta, \mu)$, the equilibrium thermodynamics of the open system becomes accessible. Likewise, for a quantum system, the density matrix and, for a classical system, the respective phase space probability density function (pdf) are specified in terms of the Hamiltonian of mean force according to

$$\rho(\beta, \mu) = Z_S^{-1}(\beta, \mu) e^{-\beta H^*(\beta, \mu)}. \quad (2)$$

On the other hand, from the knowledge of $\rho(\beta, \mu)$, which can, in principle, be inferred from a system-intrinsic point of view, it is not possible to separate $\rho(\beta, \mu)$ into the Hamiltonian of mean force and the system's partition function in

conflict with a central dogma of stochastic thermodynamics [6–8] which postulates that all necessary information can be obtained from observations of the system without recourse to data from the environment. As a resort, Strasberg and Esposito argue in Ref. [8] that only those differences of free energies or, equivalently, those ratios of the system's partition functions are relevant for the description of thermodynamic processes that are taken at different values of the direct system control parameters λ . With this assumption, the large class of thermodynamic processes accompanied by changes in global parameters as well as of temperature and pressure is excluded and most relevant thermodynamic quantities, such as specific heat, isobaric thermal expansion, magnetization, and polarization, magnetic, and electric susceptibilities, and compression factors are not accessible within such an approach. Hence, even though the description of special processes may not require extra information about the environment, in general, a system intrinsic description, i.e., one that is exclusively based on the observation of the system, be it by means of quantum tomography or of a monitoring of the stochastic trajectories of a classical open system, is not possible [4,5]. In contrast to Strasberg and Esposito [8] being concerned with the nonequilibrium dynamics of a special class of problems, our main aim in Refs. [4,5] is the characterization of the thermal equilibrium of open systems in its dependence on *all* relevant parameters. In this context, the modification of the Hamiltonian of mean force suggested by Strasberg and Esposito [8] cannot be qualified as an addition of an “irrelevant constant” that would have “no thermodynamic consequences”; the quotations are taken from Ref. [8]. For general thermodynamic processes it would rather lead to erroneous conclusions. Even in the restricted situation of isothermal processes at *constant* global parameters it is not sufficient to know the reduced density matrix or the pdf of the open system to infer changes in thermodynamic quantities. Additionally, process-specific relations, such as the Jarzynski equality or second-law-like relations must be imposed. This further narrows the predictive power of the method suggested by Ref. [8]. Moreover, the

inference of free energy differences on the basis of the Jarzynski equality often requires an unrealistically large amount of data [9,10]. The same kind of problem must be expected also for the other methods, in particular, when the tomography of time-dependent states is required to estimate, e.g., an effective system Hamiltonian on the basis of Eq. (23) in Ref. [8].

As explained in Refs. [4,5], the concept of fluctuating thermodynamic potentials suffers already in thermal equilibrium from ambiguities that can be subsumed as the set of all functions with a vanishing equilibrium average value with respect to the respective Gibbs state of the total system. For transient and other nonequilibrium processes, the respective set of functions is characterized by a vanishing average with respect to the actual time-dependent state of the system. It, hence, changes with time in dependence of the initial pdf of the system as it follows from Eqs. (53) and (54) in combination with Eqs. (50) and (51) of Ref. [4], but it does *not* collapse to an empty set as Strasberg and Esposito wrongly conclude in Footnote [60] of Ref. [8]. We further note that even the specification of a particular fluctuating free energy as performed in Eq. (5) of Ref. [11] does still leave the fluctuating internal energy and entropy largely unspecified as can be seen from the Eqs. (107) and (108) in Ref. [5]. In the particular case of an equilibrium system the supposedly fluctuating free energy defined in (5) of Ref. [11] yields the nonfluctuating equilibrium free energy [4].

Furthermore, we would like to clarify two misleading literal citations in Ref. [8] which are taken out of their original context. Our statement in Ref. [5] that “...presents in practice

an impossible task” does not refer to the Hamiltonian of mean force as insinuated by Strasberg and Esposito [8] but to the reconstruction of the total system’s Hamiltonian solely based on open system’s trajectories. Our observation that the first law of thermodynamics for quantum open systems interacting with their environments at a finite strength is doubtful is based on the fact that then the respective observables that determine work and heat do not commute. Hence, their simultaneous measurement is excluded by the laws of quantum mechanics. The condition that measurements “need to be error free”, cited from Ref. [5], refers to one of the mathematical properties that generalized energy measurements must satisfy in order to yield the Crooks and the Jarzynski fluctuation relations [12,13] but has not been made in the context of the first law.

Finally, we would like to stress that, by its very definition, the Hamiltonian of mean force describes thermodynamic *equilibrium*. Expressions, such as the nonequilibrium free energy in Eq. (5) of Ref. [8] or the corresponding fluctuating nonequilibrium free energy in Eq. (5) of Ref. [11], based on the Hamiltonian of mean force and the system’s pdf at time t and at the position of a random trajectory, are mere postulates without a deeper rationale. The latter object, resulting in the described way from the pdf, is by construction a function of the starting point of the considered random trajectory and a functional of the random force having acted up to time t . This very construct does not follow as a transformation of a pdf according to the proper rules of probability theory. It, therefore, is not normalized with respect to the starting point and, hence, has no obvious probabilistic meaning, see the Supplemental Material [14].

-
- [1] M. Campisi, P. Talkner, and P. Hänggi, Fluctuation Theorem for Arbitrary Open Quantum Systems, *Phys. Rev. Lett.* **102**, 210401 (2009).
 - [2] L. Onsager, Theories of concentrated electrolytes, *Chem. Rev.* **13**, 73 (1933).
 - [3] G. Kirkwood, Statistical mechanics of fluid mixtures, *J. Chem. Phys.* **3**, 300 (1935).
 - [4] P. Talkner and P. Hänggi, Open system trajectories specify fluctuating work but not heat, *Phys. Rev. E* **94**, 022143 (2016).
 - [5] P. Talkner and P. Hänggi, Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical, *Rev. Mod. Phys.* **92**, 041002 (2020).
 - [6] U. Seifert, Stochastic thermodynamics, fluctuation theorems and molecular machines, *Rep. Prog. Phys.* **75**, 126001 (2012).
 - [7] U. Seifert, From stochastic thermodynamics to thermodynamic inference, *Annu. Rev. Condens. Matter Phys.* **10**, 171 (2019).
 - [8] P. Strasberg and M. Esposito, Measurability of nonequilibrium thermodynamics in terms of the Hamiltonian of mean force, *Phys. Rev. E* **101**, 050101(R) (2020).
 - [9] S. Kim, Y. W. Kim, P. Talkner, and J. Yi, Comparison of free-energy estimators and their dependence on dissipated work, *Phys. Rev. E* **86**, 041130 (2012).
 - [10] J. W. Deng, A. M. Tan, P. Hänggi, and J. B. Gong, Merits and qualms of work fluctuations theorems in classical fluctuation theorems, *Phys. Rev. E* **95**, 012106 (2017).
 - [11] See Supplemental Material of Ref. [8].
 - [12] B. P. Venkatesh, G. Watanabe, and P. Talkner, Transient quantum fluctuation theorems and generalized measurements, *New J. Phys.* **16**, 015032 (2014).
 - [13] K. Ito, P. Talkner, B. P. Venkatesh, and G. Watanabe, Generalized energy measurements and quantum work compatible with fluctuation theorems, *Phys. Rev. A* **99**, 032117 (2019).
 - [14] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevE.102.066101> for the discussion of a particular exemplary case, which includes Refs. [15–17].
 - [15] U. Seifert, Entropy Production Along a Stochastic Trajectory and an Integral Fluctuation Theorem, *Phys. Rev. Lett.* **95**, 040602 (2005).
 - [16] M. C. Wang and G. E. Uhlenbeck, On the theory of the Brownian motion, *Rev. Mod. Phys.* **17**, 323 (1945).
 - [17] D. Ludwig, Persistence of dynamical systems under random perturbations, *SIAM Rev.* **17**, 605 (1975).