

## Interacting fermions: correlation functions obtained with the Gutzwiller wave function

Florian Gebhard, Dieter Vollhardt

### Angaben zur Veröffentlichung / Publication details:

Gebhard, Florian, and Dieter Vollhardt. 1987. "Interacting fermions: correlation functions obtained with the Gutzwiller wave function." *Japanese Journal of Applied Physics* 26 (S3-1): 481–82. <https://doi.org/10.7567/jjaps.26s3.481>.

# Interacting Fermions: Correlation Functions Obtained with the Gutzwiller Wave Function

Florian GEBHARD\*

Physik Department, T 30, Technische Universität München, D-8046 Garching, F.R. Germany

Dieter VOLLHARDT\*

Max-Planck-Institut für Physik und Astrophysik, D-8000 München 40, F.R. Germany

A recently developed analytic approach is used to calculate correlation functions for interacting spin-1/2 fermions on a lattice in terms of the Gutzwiller wave function. In one dimension the evaluations are performed analytically and without any approximation. Whenever possible comparison with exact results (Hubbard-model, spinless fermions) is made.

The well-known difficulties involved in the theoretical treatment of interacting Fermi systems, especially those with a strong, short-range repulsive interaction, have made variational methods particularly attractive [1]. In order to study a lattice model of itinerant electrons with an on-site interaction [2,3] ("Hubbard-model") Gutzwiller [2] introduced a variational wave function  $|\psi_G\rangle$ , which controls local (on-site) density fluctuations in the ground state wave function of the non-interacting Fermi gas,  $|\psi_0\rangle$ ,

$$|\psi_G\rangle = \prod_i [1 - (1-g)D_i] |\psi_0\rangle \quad (1)$$

Here  $D_i = n_{i\uparrow}n_{i\downarrow}$  is the number operator for double occupancy of a lattice site ("D-site") and  $0 \leq g \leq 1$  is a variational parameter. The projection operator in (1) merely reduces the number of D-sites (where the interactions occur) of the spin configurations contributing to  $|\psi_0\rangle$ . Although  $|\psi_G\rangle$  is simple in structure, exact evaluations of expectation values  $\langle X \rangle = \langle \psi_G | X | \psi_G \rangle / \langle \psi_G | \psi_G \rangle$  of an operator  $X$  have not been possible for a long time. Therefore expansion techniques [4], mean field-type approximations [5] and numerical methods were employed [6-9]. Most recently, however, a new, analytically tractable approach to the problem was developed by Metzner and Vollhardt [10], which is based on a somewhat unconventional combination of Wick's theorem and well-known diagrammatic techniques. Expectation values are expressed as a series in powers of  $1-g^2$ , whose coefficients can be calculated to any order - at least in one dimension. In higher dimensions numerical methods have to be used. In this way the ground state energy of the  $d=1$  Hubbard-model in terms of  $|\psi_G\rangle$  has been obtained without approximation [10].

We now go on to calculate correlation functions (CFs) [11]. In lattice models with itinerant up/down spins a site may be occupied by single spins, D-sites and empty sites ("holes"). Introducing number operators at site  $i$  for the spin ( $S_i^z = n_{i\uparrow} - n_{i\downarrow}$ ), density ( $N_i = n_{i\uparrow} + n_{i\downarrow}$ ), D-sites ( $D_i = n_{i\uparrow}n_{i\downarrow}$ ) and holes ( $H_i = (1-n_{i\uparrow})(1-n_{i\downarrow})$ ) we want to calculate the CFs

$$C_j^{XY} = \frac{1}{L} \sum_i \langle X_i Y_{i+j} \rangle - \langle X \rangle \langle Y \rangle \quad (2)$$

where  $L$  is the number of lattice sites,  $X = L^{-1} \sum_i X_i$  etc. and  $X_i, Y_i = S_i^z, N_i, D_i, H_i$ . This defines 10 CFs. For  $n_\uparrow = n_\downarrow = n/2$ , with  $n = N/L \leq 1$  as the particle density, there remain only seven, four of which are independent. We choose  $C^{SS}, C^{NN}, C^{HH}$  and  $C^{DH}$  since they have a direct physical significance. We note that for  $X = Y$  the  $\vec{q} = 0$  limit of the Fourier transform of (2),  $C^{XX}(q=0) = \langle X^2 \rangle - \langle X \rangle^2$ , is a measure for the fluctuations in the number  $\langle X \rangle$  around its average. In the present case, where a fixed number of particles  $\langle N \rangle = n$  and a total spin  $\langle S^z \rangle = 0$  is considered,  $C^{SS}(\vec{q}=0) = C^{NN}(\vec{q}=0) = 0$ , while  $C^{DD}(\vec{q}=0) \neq 0$  because there is no conservation law for D-sites or holes. -

In the following we limit our discussion to  $d = 1$  dimension and  $n = 1$  unless stated otherwise.

1) The spin-CF  $C_j^{SS}$  is of particular interest, because the results obtained with (1) may be compared with the exact results for  $j = 1, 2$  in the  $d = 1$  Hubbard-model in the atomic limit [12]. One obtains  $C_j^{SS}(q) = -(1-g^2)^{-1} \ln F(Q)$  where  $F(x) = 1 - (1-g^2)x$  and  $Q = |q|/\pi$ . In the atomic ( $g = 0$ ),  $C_j^{SS}(q=2k_F)$  is seen to diverge logarithmically, implying an anti-ferromagnetic transition. In real space

$$C_j^{SS} = - \frac{1}{\pi j} \int_0^1 dy \frac{\sin(\pi j y)}{F(y)}, \quad j > 0 \quad (3)$$

so that for  $g = 0$  one finds

$$C_j^{SS} \Big|_{g=0} = (-1)^j \frac{\text{Si}(\pi j)}{\pi j} \quad (4)$$

In Fig. 1  $C_j^{SS}$  is plotted for different values of the correlation parameter  $g$ . The  $(-1)^j/j$  dependence has already been suggested earlier [7]. The numerical calculations for  $j = 1$  [6-8] and  $j = 2$  [7] are seen to be very accurate. The results obtained from (4) are also in very good agreement with exact [12] and numerical [13] results for the antiferromagnetic Heisenberg chain. Hence we see that for  $n = 1$  the Gutzwiller wave function describes spin correlations in the interacting system very well.

\*Address as of Oct. 1, 1987: Institut für Theoretische Physik C, Technische Hochschule Aachen, 5100 Aachen, FRG.




Fig 1. The magnitude of the spin-correlation function  $C_j^{SS}$  vs. separation  $j$  for different correlation parameters  $g$ .




Fig. 2. The probability for finding an empty site at separation  $j$  from a doubly occupied site, normalized to the non-interacting case, for different correlation parameters  $g$ .

2) The density-CF is found as  $C^{NN}(q) = [g^2/(1-g^2)] \ln F(-Q/g^2)$ . The additional  $g^2$ -factor, as compared to  $C^{SS}$ , reflects the suppression of density fluctuations for  $g \rightarrow 0$ .

3) For  $g = 0$  and  $n < 1$  holes act as spinless fermions whose number is conserved [8]. In this limit the hole-CF  $C^{HH}$  is given by  $C_{g=0}^{HH} = C_{g=0}^{NN}$ , while its exact value is given by  $C_{g=1}^{HH}$  with  $n, k_F$  replaced by the hole concentration  $n_h = 1-n$  and  $k_F^h = \pi n_h$ , respectively. It turns out [11] that  $C_{g=0}^{HH}$  obtained with  $|\psi_G\rangle$  describes the overall features of the exact result (correlation hole of width  $1/2n_h$ , oscillations etc.) very well as was already concluded from Monte-Carlo calculations [8]. -

4) For  $n = 1$  the CF between D-sites and holes is given by  $C_j^{DH} = C_j^{HH} - \frac{1}{2} C_j^{NN}$ , such that for  $g \rightarrow 0$   $C_j^{DH}(q) \approx -\frac{1}{2} C_j^{NN}(q)$ . Since correlations

in (1) merely tend to smoothen out the distribution of particles on the lattice, the latter result implies that there is only an average correlation between D-sites and holes, resembling the non-interacting case. In Fig. 2 we show  $\tilde{C}_j^{DH} = (C_j^{DH} + d^2)/d$ , the probability for finding a hole at distance  $j$  from a D-site, normalised to the non-interacting case, which clearly shows this feature. On the other hand, for strong interactions this probability should be higher than in the uncorrelated case, because this would make a decay of a high-energy D-site easier. As discussed earlier [6] the lack in spatial correlation appears to be the main origin for the rather high ground state energy of the  $d = 1$  Hubbard-model obtained with  $|\psi_G\rangle$  at strong interactions  $U$ , which is

caused by logarithmic corrections to the usual  $-t^2/U$ -dependence [10].

The above results may be directly used to diagonalize Hubbard-type models with more complicated than on-site interactions. -

We thank W. Metzner for useful discussions.

## REFERENCES

- 1) E. Feenberg, *Theory of Quantum Fluids* (Academic Press, New York, 1968).
- 2) M.C. Gutzwiller, Phys. Rev. Lett. **10** 159 (1963), Phys. Rev. **134A**, 923 (1964); **137A** 1726 (1965).
- 3) J. Hubbard, Proc. R. Soc. London, Ser. A **276**, 238 (1963).
- 4) See, for example, D. Baeriswyl and K. Maki, Phys. Rev. **B31** 6633 (1985).
- 5) For a review, see D. Vollhardt, Rev. Mod. Phys. **56** 99 (1984).
- 6) T.A. Kaplan, P. Horsch and P. Fulde, Phys. Rev. Lett. **49** 889 (1982).
- 7) P. Horsch and T.A. Kaplan, J. Phys. **C16** L1203 (1983).
- 8) C. Gros, R. Joynt and T.M. Rice, Phys. Rev. B (in press).
- 9) H. Yokoyama and H. Shiba, Technical Report of ISSP, Ser. A., No. 1729 (1986).
- 10) W. Metzner and D. Vollhardt, MPI-preprint PAE/PTH 14/87.
- 11) F. Gebhard and D. Vollhardt, MPI-preprint PAE/PTH 29/87.
- 12) H. Bethe, Z. Phys. **71** 205 (1931); M. Takahashi, J. Phys. **C10** 1289 (1977).
- 13) T.A. Kaplan, P. Horsch and J. Borysowicz, Phys. Rev. **B35** 1877 (1987).