



An unusual case of intracerebral Non-Langerhans cell histiocytosis with review of the literature

K. Mueller, M. Trippel, Ansgar Berlis, W. Janzarik, B. Volk

Angaben zur Veröffentlichung / Publication details:

Mueller, K., M. Trippel, Ansgar Berlis, W. Janzarik, and B. Volk. 2007. "An unusual case of intracerebral Non-Langerhans cell histiocytosis with review of the literature." *Diagnostic Pathology* 2 (Suppl 1): S24. https://doi.org/10.1186/1746-1596-2-s1-s24.

Diagnostic Pathology



Oral presentation Open Access

An unusual case of intracerebral Non-Langerhans cell histiocytosis with review of the literature

K Mueller*1, M Trippel2, A Berlis3, W Janzarik4 and B Volk1

Address: ¹Abt. Neuropathologie, Universität Freiburg, Germany, ²Abt. Stereotaktische Neurochirurgie, Universität Freiburg, Germany, ³Abt. Neurochirurgie, Sektion Neuroradiologie, Universität Freiburg, Germany and ⁴Abt. Neurologie, Universität Freiburg, Germany

from 35te Tagung der Pathologen am Oberrhein/35th Meeting of Pathologists of the Upper Rhine Region (PATOR) The Institute of Pathology, University Hospital Freiburg, Germany. I July 2006

Published: 14 March 2007

Diagnostic Pathology 2007, 2(Suppl 1):S24 doi:10.1186/1746-1596-2-S1-S24

© 2007 Mueller et al; licensee BioMed Central Ltd.

Aims

In the face of an intracerebral case of Non-Langerhans cell histiocytosis (NLCH), the diagnostic procedure and the clinical handling of this rare histiocytic tumorous lesion are demonstrated and differential diagnoses as Rosai-Dorfman or Erdheim Chester diseases are discussed.

Methods

A stereotactic serial biopsy of frontal brain was taken from a patient with unclear contrast enhancing masses around the sellar region, the hypothalamus, and a wide lining intraventricular. Smear preparations, conventional histology, immunohistochemistry and electron microscopy were done to rule out the etiology of the tumorous lesion.

Results

The morphological analysis of our case revealed a histiocytic proliferation, which, in view of the immunohistochemical and ultrastructural findings, was classified as Non-Langerhans cell histiocytosis. No infectious agents could be detected and no lymphoma-like features could be observed.

Conclusion

This case shows the unusual manifestation of a primary intracerebral Non-Langerhans cell histiocytosis, which begins as a solid mass around the sellar region and shows a remarkable extension as a flat lining of inner and outer liquor spaces. Stereotactic brain biopsy technique today is a valuable high tech tool to clarify intracerebral tumorous lesions with a minimum of risk for the patient and with the possibility to perform all necessary tissue investigations.

^{*} Corresponding author