

Relationship between periodontitis and psoriasis: a two sample Mendelian randomization study





Hansjörg Baurecht, Dennis Freuer, Christine Welker, Lam C. Tsoi, James T. Elder, Benjamin Ehmke, Michael F. Leitzmann, Birte Holtfreter, Sebastian Edgar Baumeister

Angaben zur Veröffentlichung / Publication details:

Baurecht, Hansjörg, Dennis Freuer, Christine Welker, Lam C. Tsoi, James T. Elder, Benjamin Ehmke, Michael F. Leitzmann, Birte Holtfreter, and Sebastian Edgar Baumeister. 2022. "Relationship between periodontitis and psoriasis: a two sample Mendelian randomization study." *Journal of Clinical Periodontology* 49 (6): 573–79.
<https://doi.org/10.1111/jcpe.13620>.

ORIGINAL ARTICLE

Relationship between periodontitis and psoriasis: A two-sample Mendelian randomization study

Hansjörg Baurecht¹  | Dennis Freuer² | Christine Welker¹ | Lam C. Tsoi^{3,4,5} |
James T. Elder^{3,6} | Benjamin Ehmke⁷  | Michael F. Leitzmann¹ |
Birte Holtfreter⁸  | Sebastian-Edgar Baumeister⁹ 

¹Department of Epidemiology and Preventive Medicine, University of Regensburg, Regensburg, Germany

²Chair of Epidemiology, University of Augsburg, University Hospital Augsburg, Augsburg, Germany

³Department of Dermatology, University of Michigan Medical School, Ann Arbor, Michigan, USA

⁴Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, Michigan, USA

⁵Department of Biostatistics, Center for Statistical Genetics, University of Michigan, Ann Arbor, Michigan, USA

⁶Ann Arbor Veterans Affairs Hospital, Ann Arbor, Michigan, USA

⁷Clinic for Periodontology and Conservative Dentistry, University of Münster, Münster, Germany

⁸Department of Restorative Dentistry, Periodontology, Endodontology, and Preventive and Pediatric Dentistry, University Medicine Greifswald, Greifswald, Germany

⁹Institute of Health Services Research in Dentistry, University of Münster, Münster, Germany

Correspondence

Hansjörg Baurecht, Department of Epidemiology and Preventive Medicine, University of Regensburg, Franz-Josef-Strauss Allee 11, 93057 Regensburg, Germany.
Email: hansjoerg.baurecht@ukr.de

Funding information

Arthritis National Research Foundation; National Psoriasis Foundation; the Dermatology Foundation; Ann Arbor VA Hospital; National Institutes of Health, Grant/Award Numbers: K01AR072129, R01AR065183, R01AR063611,

Abstract

Aim: Observational research suggests that periodontitis affects psoriasis. However, observational studies are prone to reverse causation and confounding, which hampers drawing causal conclusions and the effect direction. We applied the Mendelian randomization (MR) method to comprehensively assess the potential bi-directional association between periodontitis and psoriasis.

Materials and Methods: We used genetic instruments from the largest available genome-wide association study of European descent for periodontitis (17,353 cases, 28,210 controls) to investigate the relationship with psoriasis (13,229 cases, 21,543 controls), and vice versa. Causal Analysis Using Summary Effect (CAUSE) estimates and inverse variance-weighted (IVW) MR analyses were used for the primary analysis. Robust MR approaches were used for sensitivity analyses.

Results: Both univariable methods, CAUSE and IVW MR analyses, did not reveal any impact of periodontitis on psoriasis (CAUSE odds ratio [OR] = 1.00, $p = 1.00$; IVW OR = 1.02, $p = .6247$), or vice versa (CAUSE OR = 1.01, $p = .5135$; IVW OR = 1.00, $p = .7070$). The null association was corroborated by pleiotropy-robust methods with ORs close to 1 and p -values $>.59$. Overall, MR analyses did not suggest any effect of periodontitis on psoriasis. Similarly, there was no evidence to support an effect of psoriasis on periodontitis.

Conclusions: Within the limitations of this MR study, the outcomes supported neither periodontitis affecting psoriasis nor psoriasis affecting periodontitis.

KEYWORDS

epidemiology, Mendelian randomization, periodontitis, psoriasis

Clinical Relevance

Scientific rationale for study: Previous observational research had suggested a relationship between periodontitis and psoriasis. Observational estimates on the association of periodontitis and psoriasis might be subject to confounding or reverse causation.

R01AR054966, R01AR050511,
R01AR042742

Principal findings: The application of complementary Mendelian randomization approaches to investigate causality did not indicate an association between periodontitis and psoriasis, or vice versa.

Practical implications: The study results do not suggest that treatment for periodontal disease or psoriasis would improve comorbid disease activity.

1 | INTRODUCTION

Periodontitis is a microbe-associated inflammatory disease of the tooth-supporting tissue that affects approximately 50% of the adult population, with 10% suffering from the severe form of the disease (Bernabe et al., 2020). Periodontitis is associated with the presence of a dysbiotic microbial community in a susceptible host (Hajishengallis et al., 2020). Although bacteria are required in the pathogenesis, it is the host inflammatory response to the microbial challenge that drives immune-cell-mediated self-degradation of the periodontal tissue, resulting in tooth loss (Van Dyke et al., 2020). Psoriasis is a chronic, papulosquamous inflammatory skin disease often accompanied with concomitant comorbidities and is prevalent in approximately 2%–4% of the general population (Christophers, 2001). A distinct configuration of the skin microbiota has been shown in psoriasis patients characterized by the co-occurrence of bacterial communities (Fyhrquist et al., 2019). Though the exact role of bacteria, streptococci in particular, in the pathogenesis of psoriasis remains unresolved, an abnormal immune response to dysbiotic skin microbiota has been proposed as the mechanistic link (Langan et al., 2018, 2019; Liang et al., 2021).

Whereas an imbalance in the periodontal microbiota and the inflammatory response of the host explains local tissue destruction, it is unclear whether this imbalanced relationship can link periodontal disease with extra-oral comorbidities such as psoriasis. Several observational studies have proposed an association between periodontitis and psoriasis, which has been substantiated by meta-analyses (Ungprasert et al., 2017; Zhang et al., 2020) estimating pooled odds ratios (ORs) ranging from 1.55 (95% confidence interval [CI], 1.35–1.77) to 2.87 (95% CI, 1.75–4.69). In both periodontitis and psoriasis, the predominant immune response is neutrophilic (Christophers, 2017), and further common etiopathogenesis pathways have been hypothesized (Falcao & Bullón, 2019). Pro-inflammatory cytokines play key roles in both periodontitis and plaque psoriasis. Available evidence from immunological and genetic studies has highlighted the immunological circuits that converge on adaptive immune pathways involving IL-17, IL-23, TH17, TNF- α , and TNF- γ in psoriasis and periodontitis (Hawkes et al., 2017; Marchesan et al., 2020; Griffiths et al., 2021). IL-17, TNF- α , and IL-36 contribute to the amplification of self-sustaining inflammatory circuits and influx of neutrophils in psoriasis.

The emerging findings that inflammatory processes in different peripheral tissues may be interlinked may improve our mechanistic understanding of comorbidities. A recent meta-analysis including eight studies reported a significantly worse periodontal health among psoriasis patients (Qiao et al., 2019).

However, both diseases share several risk factors such as smoking, alcohol consumption, stress, and immune depression (Sharma et al., 2015), which might act as confounders leading to spurious correlation and are difficult to control in observational studies. In particular, smoking and alcohol consumption have been recently reported to be causally associated with periodontitis (Baumeister et al., 2021).

Mendelian randomization (MR) is an alternative approach to account for observational bias (Burgess et al., 2018, 2019). MR relies on the natural random assortment of genetic variants during meiosis, yielding a random distribution of genetic variants in a population. Individuals are naturally assigned at birth to inherit a genetic variant that affects a risk factor or disease susceptibility, or not inherit such a variant. The natural randomization that occurs in the generation of each individual's genetic make-up is analogous to the design of a randomized trial. MR uses instrumental variable (IV) analysis where the genetic variants are the instruments, analogous to the random assignment to treatment and control groups in a randomized trial. Because the genetic variants are typically unassociated with confounders, differences in the outcome between those who carry the variant and those who do not can be attributed to differences in risk factor or disease susceptibility. Thus, MR provides a reliable understanding of the influence of modifiable exposures on the trait of interest compared to traditional observational studies which are susceptible to confounding or reverse causation (Richmond & Davey Smith, 2021). We carried out a bidirectional MR study on the association of periodontitis and psoriasis, as well as a series of sensitivity analyses, to account for pleiotropic single nucleotide polymorphisms (SNPs) associated with potential confounding factors.

2 | MATERIALS AND METHODS

In MR analyses, genetic variants strongly associated with an exposure will be exploited as IVs to investigate the potential causal effect of an exposure with an outcome of interest (Burgess et al., 2018). Because of the random assignment of genetic variants at conception, MR estimates are not biased by confounding, reverse causation, and measurement error. Most commonly, inference is based on SNPs as IVs identified by genome-wide association studies (GWAS) (Burgess et al., 2018). For the validity of each IV, three main assumptions have to be fulfilled: (1) there is robust association between the instrument and the exposure ("relevance"); (2) instruments affects the outcome through the exposure ("exclusion restriction"); and (3) the genetic variant is not associated with confounders of the exposure–outcome

association (“exchangeability”) (Labrecque & Swanson, 2018). To satisfy the first MR assumption, we selected SNPs associated with the exposure at a significance threshold of $p < 5 \times 10^{-6}$ for periodontitis and $p < 5 \times 10^{-8}$ for psoriasis. From these set of SNPs, we selected only independent instruments with the lowest p -value by calculating pair-wise linkage disequilibrium and omitting SNPs with $r^2 \geq 0.001$ (Burgess et al., 2019). Further verification of the first assumption was made by computing the F -statistic and the proportion of the explained phenotypic variance by all SNPs (Burgess & Thompson, 2011).

We performed a two-sample MR analysis based on summary statistics from the largest available GWAS on periodontitis and psoriasis including 17,353 periodontitis cases and 28,210 controls (Shungin et al., 2019) based on the GLIDE consortium (Shungin et al., 2015) as well as 13,229 psoriasis cases and 21,543 controls (Tsoi et al., 2017). Both GWAS were conducted in people of European Caucasian descent. Periodontitis cases were classified by either the Centers for Disease and Control and Prevention/American Academy of Periodontology case definition (CDC/AAP) (Page & Eke, 2007) or the Community Periodontal Index (CPI) (World Health Organization, 2013), which is based on probing depths and the number of deep periodontal pockets (Shungin et al., 2015). Psoriasis was diagnosed by experienced physicians. All included GWAS obtained ethical review board approval and informed consent as described in the respective original manuscript (Tsoi et al., 2017; Shungin et al., 2019).

For analyses, genetic variants from different studies were harmonized with regard to their effects, and palindromic SNPs (i.e., SNPs whose alleles consist of a base and its complementary base) were excluded. In order to maintain consistency in SNPs used as IVs across different analysis, we used only variants available for all examined traits and did not replace missing variants by proxies.

2.1 | Statistical analyses

Statistical power was calculated according to Brion et al. (2013), which exploits asymptotic theory and derives the power via the non-centrality parameter from the respective asymptotic χ^2 -distribution. As primary analysis we applied the Causal Analysis Using Summary Effect Estimates (CAUSE) approach (Morrison et al., 2020), which has been demonstrated to outperform other established methods to detect causal relationships in the presence of pleiotropy (Wang et al., 2021). In addition, we carried out a series of sensitivity analyses including pleiotropy-robust methods.

First, we harmonized summary statistics results to ensure effect size alignment and to prohibit strand mismatch. Then we carried out CAUSE, which incorporates all genetic variants after clumping and thereby increases statistical power, with periodontitis as exposure and psoriasis as outcome, and vice versa. CAUSE used genome-wide summary statistics to disentangle causality (i.e., SNPs affect psoriasis through their effect on periodontitis) from correlated pleiotropy (i.e., violation of the MR exchangeability assumption whereby SNPs are associated with periodontitis and psoriasis through a shared heritable factor), while taking into account uncorrelated horizontal

pleiotropy (an exclusion restriction violation where the SNPs associate with periodontitis through separate mechanisms). It uses Bayesian modelling to assess whether the sharing model (the model that fixes the causal effect at zero) fits the data at least as well as the causal model (the model that allows a causal effect different from zero) (Morrison et al., 2020).

To complete the picture, we carried out the standard inverse variance-weighted (IVW) MR with the multiplicative random effects model (Burgess et al., 2019) for which valid estimates of the exposure on the outcome are based on all three fulfilled assumptions (Labrecque & Swanson, 2018). Weak instrument-exposure association might reduce the plausibility of MR relevance assumption. Therefore, we examined F -statistics of the instruments and calculated the explained phenotypic variance. Violations of assumptions (2) and (3) can occur through horizontal pleiotropy, that is, instruments exerting an effect on the outcome independent of the exposure. To investigate pleiotropy, we searched for previously reported associations of instruments with the shared risk factors, in particular smoking and alcohol consumption, in PhenoScanner (www.phenoscaner.medschl.cam.ac.uk/) and the GWAS Catalog (www.ebi.ac.uk/gwas/). In addition, we performed a suite of pleiotropy-robust methods (weighted median, robust adjusted profile score [RAPS], radial regression, MR Pleiotropy RESidual Sum and Outlier [MR-PRESSO]) to address the issue of pleiotropy (Slob & Burgess, 2020). Finally, we examined the heterogeneity of the ratio estimators using Cochran's Q , I^2 statistic, and the MR-Egger intercept, and performed leave-one SNP-out analysis.

All analyses were performed using the packages CAUSE (1.2.0), MendelianRandomization (0.5.1), TwoSampleMR (0.5.6), and MR-PRESSO (1.0) in R, version 4.0.5. Reporting follows the STROBE-MR statement (Skrivankova et al., 2021). The study protocol was not pre-registered.

3 | RESULTS

Descriptive statistics on phenotypes from studies included in the GWAS on exposure and outcome, including the number of individuals suffering from periodontitis classified by CDC/AAP and CPI, are presented in Table S1.

CAUSE analysis showed no effect of periodontitis on psoriasis (OR = 1.0, 95% credible interval [CredIn] = 0.16–6.17, $p = 1.00$). Similarly, CAUSE did not indicate an effect of psoriasis on periodontitis (OR = 1.01, 95% CredIn = 0.97–1.04, $p = .5135$). Similar effect estimates were produced when the IVW method was used (Table 1, Figure 1).

Pleiotropy-robust methods corroborated the findings of no effect of periodontitis on psoriasis ($.5918 \leq p \leq .6868$), or vice versa ($.7014 \leq p \leq .9837$, Table 1). The 76 and 46 SNPs selected as IVs for periodontitis and psoriasis explained 3.1% and 14.6% of the variance, respectively (Table S2). The F -statistics ranged between 16.5–25.1 and 29.9–918.3 for periodontitis and psoriasis (Table S2). Both the explained phenotypic variance and the high F -statistics of the variants

Outcome	Exposure	Method	OR	95% CI	p-Value
Psoriasis	Periodontitis	IVW	1.017	0.95–1.088	.6247
		Weighted median	1.020	0.926–1.123	.6868
		Robust adjusted profile score	1.019	0.947–1.098	.6098
		IVW radial	1.017	0.956–1.082	.5918
		MR-PRESSO	1.017	0.956–1.082	.5935
Periodontitis	Psoriasis	IVW	0.995	0.967–1.023	.7070
		Weighted median	1.000	0.959–1.044	.9837
		Robust adjusted profile score	0.995	0.965–1.026	.7471
		IVW radial	0.995	0.967–1.023	.7014
		MR-PRESSO	0.995	0.967–1.023	.7033

Abbreviations: CI, confidence interval; IVW, inverse variance-weighted; MR-PRESSO, MR Pleiotropy RESidual Sum and Outlier; OR, odds ratio.

TABLE 1 Mendelian randomization estimates for the relationship between genetically instrumented periodontitis and psoriasis, and vice versa




FIGURE 1 Mendelian randomization estimates for the relationship between genetically instrumented periodontitis and psoriasis, and vice versa. CI, confidence interval; IVW, inverse variance-weighted; MR-PRESSO, MR pleiotropy RESidual Sum and Outlier; OR, odds ratio; RAPS, robust adjusted profile score

TABLE 2 Heterogeneity of Wald ratios and MR-Egger test for directional pleiotropy

Exposure	Heterogeneity				
	Outcome	Q	df	I ²	p-Value
Periodontitis	Psoriasis	59.1	71	0.20	.8424
Psoriasis	Periodontitis	42.3	44	0.04	.5443
Exposure	MR-Egger test for directional pleiotropy				p-Value
	Outcome	Intercept	SE		
Periodontitis	Psoriasis	−0.0082	0.0079		.3060
Psoriasis	Periodontitis	−0.0009	0.0054		.8680

Abbreviations: df, degree of freedom; MR, Mendelian randomization; Q, heterogeneity statistic Q.

verify the MR assumption (1). No substantial heterogeneity between Wald ratios in the IVW estimate was observed (Table 2), and leave-one-out analysis did not reveal any leverage points with high influence (Table S3). Using the PhenoScanner and GWAS Catalog, we also did

not find any genetic instrument of periodontitis and psoriasis to be associated with any potential confounders (Table S4). Pleiotropy-robust MR methods showed similar results to IVW (Table 1, Figure 1). Low or no heterogeneity, no indication of potential confounding SNPs, and similar results by pleiotropy-robust analyses implicitly reassure that MR assumptions (2) and (3) hold.

On a 5% significance level, our analyses had a power of >87% to detect a causal effect of OR = 1.2 of periodontitis on psoriasis and a power of >96% for a causal effect of OR = 1.1 of psoriasis on periodontitis (Table S5).

4 | DISCUSSION

The results of the MR study do not support evidence for an effect of periodontitis on psoriasis, or vice versa. The MR analysis, CAUSE in particular, was powered to detect small effects, and produced consistent estimates using different MR techniques. Our MR estimates contradict the available observational studies, which suggested a

bidirectional association of periodontitis and psoriasis (Ungprasert et al., 2017; Qiao et al., 2019; Zhang et al., 2020). Although several pathogenic links between both diseases have been suggested because of the shared common immunological response (Hawkes et al., 2017; Dalmády et al., 2020; Marchesan et al., 2020; Griffiths et al., 2021), our MR study does not support a causal link in either direction, as the estimated relative risks were close to zero. One explanation could be that previously observed associations between periodontitis and psoriasis are coincidental or thwarted by an unknown confounder. Moreover, a causal link between periodontitis and psoriasis, or the converse, cannot be established in observational studies because most patients with periodontitis exhibit systemic health problems. Periodontitis is often accompanied by a range of systemic diseases (Holmstrup et al., 2017; Hajishengallis & Chavakis, 2021; Teles et al., 2021) and most psoriatic patients suffer from numerous comorbidities (Pearce et al., 2005) including rheumatological, cardiovascular, and psychiatric complications (Greb et al., 2016). Therefore, it is possible that these comorbidities, and in particular those sharing inflammatory pathways, contribute to the association between periodontitis and psoriasis.

Nevertheless, the applied complementary MR approaches yielded homogeneous results with point estimates slightly above and below 1 for both hypotheses—psoriasis influencing periodontitis, and vice versa—with highly overlapping CIs. Consequently, it is unlikely that periodontitis and psoriasis are causally related. If this is the case, the strength of the effects is unlikely to be of clinical significance. The study had limitations. MR based on genetic summary statistics limits the range of analyses. However, based on the observed and consistent negative results from several complementary approaches with effect estimates close to 1, it is unlikely that the finding is distorted by any form of bias. MR analyses are less susceptible to bias involving violation of MR assumptions and provide robust evidence when effects are very small or absent (VanderWeele et al., 2014). The CAUSE analysis exploits the full range of genetic variants with no evidence for a causal relationship between periodontitis and psoriasis. The wide credible intervals for these estimates may suggest low power to rule out very small effects. However, the null association was corroborated by univariable and pleiotropy-robust MR approaches. The genetic instruments explained 3.4% and 14.6% of the phenotypic variance of periodontitis and psoriasis with minimum *F*-statistics of 16.48 and 29.94, consistent with the absence of weak instrument bias. One should keep in mind that MR studies are located at the interface between observational and interventional studies and therefore in the evidence-based pyramid below randomized clinical trials (RCTs) and systematic reviews of RCTs (Davies et al., 2018). Another limitation is that the partial CPI commonly underestimates severe periodontitis and over-estimates less severe periodontitis (Baelum et al., 1993), which mis-identified the numerator of the MR Wald ratio estimator.

5 | CONCLUSION

Our study does not indicate that periodontitis causally affects psoriasis, or vice versa. However, caution is warranted before generalization of the findings and further studies are needed to finally clarify the

subject. We stress the importance of combining evidence from multiple lines of observational and experimental research that align to strengthen causal inference (Munafò et al., 2021). In vivo studies of animal models would therefore further strengthen the evidence on putative bidirectional causation between periodontitis and psoriasis.

ACKNOWLEDGEMENTS

This work received no specific financial support. James T. Elder was supported by awards R01AR042742, R01AR050511, R01AR054966, R01AR063611, and R01AR065183 from the National Institutes of Health, and by the Ann Arbor VA Hospital. Lam C. Tsoi was supported by an award (K01AR072129) from the National Institutes of Health, and by awards from the Dermatology Foundation, the National Psoriasis Foundation, and the Arthritis National Research Foundation.

CONFLICT OF INTEREST

All authors disclose no conflicts of interest.

ETHICS STATEMENT

Ethical approval was granted for each of the cohorts and informed consent was obtained from all participants prior to participation.

AUTHOR CONTRIBUTIONS

Study design: Hansjörg Baurecht and Sebastian-Edgar Baumeister. *Data analysis and interpretation:* Hansjörg Baurecht, Dennis Freuer, and Sebastian-Edgar Baumeister. *Manuscript writing and critical revision:* Hansjörg Baurecht, Dennis Freuer, Christine Welker, Lam C. Tsoi, James T. Elder, Benjamin Ehmke, Michael F. Leitzmann, Birte Holtfreter, and Sebastian-Edgar Baumeister. *Data acquisition and curation:* Hansjörg Baurecht, Christine Welker, and Lam C. Tsoi.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Hansjörg Baurecht  <https://orcid.org/0000-0002-9265-5594>

Benjamin Ehmke  <https://orcid.org/0000-0002-2418-6765>

Birte Holtfreter  <https://orcid.org/0000-0002-6541-3127>

Sebastian-Edgar Baumeister  <https://orcid.org/0000-0002-9391-6602>

REFERENCES

- Baelum, V., Fejerskov, O., Manji, F., & Wanzala, P. (1993). Influence of CPITN partial recordings on estimates of prevalence and severity of various periodontal conditions in adults. *Community Dentistry and Oral Epidemiology*, 21(6), 354–359. <https://doi.org/10.1111/j.1600-0528.1993.tb01098.x>
- Baumeister, S.-E., Freuer, D., Nolde, M., Kocher, T., Baurecht, H., Khazaei, Y., Ehmke, B., & Holtfreter, B. (2021). Testing the association between tobacco smoking, alcohol consumption, and risk of periodontitis: A Mendelian randomization study. *Journal of Clinical Periodontology*, 48(11), 1414–1420.
- Bernabe, E., Marcenes, W., Hernandez, C. R., Bailey, J., Abreu, L. G., Alipour, V., Amini, S., Arabloo, J., Arefi, Z., Arora, A., Ayanore, M. A.,

- Bärnighausen, T. W., Bijani, A., Cho, D. Y., Chu, D. T., Crowe, C. S., Demoz, G. T., Demsie, D. G., Forooshani, Z. S. D., ... Kassebaum, N. J. (2020). Global, regional, and national levels and trends in burden of oral conditions from 1990 to 2017: A systematic analysis for the global burden of disease 2017 study. *Journal of Dental Research*, 99(4), 362–373. <https://doi.org/10.1177/0022034520908533>
- Brion, M.-J. A., Shakhbazov, K., & Visscher, P. M. (2013). Calculating statistical power in Mendelian randomization studies. *International Journal of Epidemiology*, 42(5), 1497–1501. <https://doi.org/10.1093/ije/dyt179>
- Burgess, S., Davey Smith, G., Davies, N. M., Dudbridge, F., Gill, D., Glymour, M. M., Hartwig, F. P., Holmes, M. V., Minelli, C., Relton, C. L., & Theodoratou, E. (2019). Guidelines for performing Mendelian randomization investigations. *Wellcome Open Research*, 4, 186. <https://doi.org/10.12688/wellcomeopenres.15555.2>
- Burgess, S., Foley, C. N., & Zuber, V. (2018). Inferring causal relationships between risk factors and outcomes from genome-wide association study data. *Annual Review of Genomics and Human Genetics*, 19, 303–327. <https://doi.org/10.1146/annurev-genom-083117-021731>
- Burgess, S., & Thompson, S. G. (2011). Avoiding bias from weak instruments in Mendelian randomization studies. *International Journal of Epidemiology*, 40(3), 755–764. <https://doi.org/10.1093/ije/dyr036>
- Christophers, E. (2001). Psoriasis—epidemiology and clinical spectrum. *Clinical and Experimental Dermatology*, 26(4), 314–320. <https://doi.org/10.1046/j.1365-2230.2001.00832.x>
- Christophers, E. (2017). Periodontitis and risk of psoriasis: Another comorbidity. *Journal of the European Academy of Dermatology and Venereology*, 31(5), 757–758. <https://doi.org/10.1111/jdv.14249>
- Dalmády, S., Kemény, L., Antal, M., & Gyulai, R. (2020). Periodontitis: A newly identified comorbidity in psoriasis and psoriatic arthritis. *Expert Review of Clinical Immunology*, 16(1), 101–108. <https://doi.org/10.1080/1744666X.2019.1700113>
- Davies, N. M., Holmes, M. V., & Davey Smith, G. (2018). Reading Mendelian randomisation studies: A guide, glossary, and checklist for clinicians. *BMJ (Clinical Research Ed.)*, 362, k601. <https://doi.org/10.1136/bmj.k601>
- Falcao, A., & Bullón, P. (2019). A review of the influence of periodontal treatment in systemic diseases. *Periodontology 2000*, 79(1), 117–128. <https://doi.org/10.1111/prd.12249>
- Fyrhquist, N., Muirhead, G., Prast-Nielsen, S., Jeanmougin, M., Olah, P., Skoog, T., Jules-Clement, G., Feld, M., Barrientos-Somarribas, M., Sinkko, H., van den Bogaard, E. H., Zeeuwen, P. L. J. M., Rikken, G., Schalkwijk, J., Niehues, H., Däubener, W., Eller, S. K., Alexander, H., Pennino, D., ... Alenius, H. (2019). Microbe-host interplay in atopic dermatitis and psoriasis. *Nature Communications*, 10(1), 4703. <https://doi.org/10.1038/s41467-019-12253-y>
- Greb, J. E., Goldminz, A. M., Elder, J. T., Lebwohl, M. G., Gladman, D. D., Wu, J. J., Mehta, N. N., Finlay, A. Y., & Gottlieb, A. B. (2016). Psoriasis. *Nature Reviews. Disease Primers*, 2, 16082. <https://doi.org/10.1038/nrdp.2016.82>
- Griffiths, C. E. M., Armstrong, A. W., Gudjonsson, J. E., & Barker, J. N. W. N. (2021). Psoriasis. *Lancet (London, England)*, 397(10281), 1301–1315. [https://doi.org/10.1016/S0140-6736\(20\)32549-6](https://doi.org/10.1016/S0140-6736(20)32549-6)
- Hajishengallis, G., & Chavakis, T. (2021). Local and systemic mechanisms linking periodontal disease and inflammatory comorbidities. *Nature Reviews. Immunology*, 21(7), 426–440. <https://doi.org/10.1038/s41577-020-00488-6>
- Hajishengallis, G., Chavakis, T., & Lambris, J. D. (2020). Current understanding of periodontal disease pathogenesis and targets for host-modulation therapy. *Periodontology 2000*, 84(1), 14–34. <https://doi.org/10.1111/prd.12331>
- Hawkes, J. E., Chan, T. C., & Krueger, J. G. (2017). Psoriasis pathogenesis and the development of novel targeted immune therapies. *The Journal of Allergy and Clinical Immunology*, 140(3), 645–653. <https://doi.org/10.1016/j.jaci.2017.07.004>
- Holmstrup, P., Damgaard, C., Olsen, I., Klinge, B., Flyvbjerg, A., Nielsen, C. H., & Hansen, P. R. (2017). Comorbidity of periodontal disease: two sides of the same coin? An introduction for the clinician. *Journal of Oral Microbiology*, 9(1), 1332710. <https://doi.org/10.1080/20002297.2017.1332710>
- Labrecque, J., & Swanson, S. A. (2018). Understanding the assumptions underlying instrumental variable analyses: A brief review of falsification strategies and related tools. *Current Epidemiology Reports*, 5(3), 214–220. <https://doi.org/10.1007/s40471-018-0152-1>
- Langan, E. A., Griffiths, C. E. M., Solbach, W., Knobloch, J. K., Zillikens, D., & Thaçi, D. (2018). The role of the microbiome in psoriasis: Moving from disease description to treatment selection? *The British Journal of Dermatology*, 178(5), 1020–1027. <https://doi.org/10.1111/bjd.16081>
- Langan, E. A., Kunstner, A., Miodovnik, M., Zillikens, D., Thaçi, D., Baines, J. F., Ibrahim, S. M., Solbach, W., & Knobloch, J. K. (2019). Combined culture and metagenomic analyses reveal significant shifts in the composition of the cutaneous microbiome in psoriasis. *The British Journal of Dermatology*, 181(6), 1254–1264. <https://doi.org/10.1111/bjd.17989>
- Liang, X., Ou, C., Zhuang, J., Li, J., Zhang, F., Zhong, Y., & Chen, Y. (2021). Interplay between skin microbiota dysbiosis and the host immune system in psoriasis: Potential pathogenesis. *Frontiers in Immunology*, 12, 764384. <https://doi.org/10.3389/fimmu.2021.764384>
- Marchesan, J. T., Girnary, M. S., Moss, K., Monaghan, E. T., Egnatz, G. J., Jiao, Y., Zhang, S., Beck, J., & Swanson, K. V. (2020). Role of inflammasomes in the pathogenesis of periodontal disease and therapeutics. *Periodontology 2000*, 82(1), 93–114. <https://doi.org/10.1111/prd.12269>
- Morrison, J., Knoblauch, N., Marcus, J. H., Stephens, M., & He, X. (2020). Mendelian randomization accounting for correlated and uncorrelated pleiotropic effects using genome-wide summary statistics. *Nature Genetics*, 52(7), 740–747. <https://doi.org/10.1038/s41588-020-0631-4>
- Munafò, M. R., Higgins, J. P. T., & Smith, G. D. (2021). Triangulating evidence through the inclusion of genetically informed designs. *Cold Spring Harbor Perspectives in Medicine*, 11(8), a040659. <https://doi.org/10.1101/cshperspect.a040659>
- Page, R. C., & Eke, P. I. (2007). Case definitions for use in population-based surveillance of periodontitis. *Journal of Periodontology*, 78(7 Suppl), 1387–1399. <https://doi.org/10.1902/jop.2007.060264>
- Pearce, D. J., Morrison, A. E., Higgins, K. B., Crane, M. M., Balkrishnan, R., Fleischer, A. B., & Feldman, S. R. (2005). The comorbid state of psoriasis patients in a university dermatology practice. *The Journal of Dermatological Treatment*, 16(5–6), 319–323. <https://doi.org/10.1080/09546630500335977>
- Qiao, P., Shi, Q., Zhang, R., Eingling, L., Wang, P., Wang, J., & Liu, H. (2019). Psoriasis patients suffer from worse periodontal status—a meta-analysis. *Frontiers in Medicine*, 6, 212. <https://doi.org/10.3389/fmed.2019.00212>
- Richmond, R. C., & Davey Smith, G. (2021). Mendelian randomization: Concepts and scope. *Cold Spring Harbor Perspectives in Medicine*, 12(1), a040501. <https://doi.org/10.1101/cshperspect.a040501>
- Sharma, A., Raman, A., & Pradeep, A. R. (2015). Association of chronic periodontitis and psoriasis: Periodontal status with severity of psoriasis. *Oral Diseases*, 21(3), 314–319. <https://doi.org/10.1111/odi.12271>
- Shungin, D., Cornelis, M. C., Divaris, K., Holtfreter, B., Shaffer, J. R., Yu, Y.-H., Barros, S. P., Beck, J. D., Biffar, R., Boerwinkle, E. A., Crout, R. J., Ganna, A., Hallmans, G., Hindy, G., Hu, F. B., Kraft, P., McNeil, D. W., Melander, O., Moss, K. L., ... Franks, P. W. (2015). Using genetics to test the causal relationship of total adiposity and periodontitis: Mendelian randomization analyses in the Gene-Lifestyle Interactions and Dental Endpoints (GLIDE) Consortium. *International Journal of Epidemiology*, 44(2), 638–650. <https://doi.org/10.1093/ije/dyv075>
- Shungin, D., Haworth, S., Divaris, K., Agler, C. S., Kamatani, Y., Keun Lee, M., Grinde, K., Hindy, G., Alaraudanjoki, V., Pesonen, P.,

- Teumer, A., Holtfreter, B., Sakaue, S., Hirata, J., Yu, Y.-H., Ridker, P. M., Giulianini, F., Chasman, D. I., Magnusson, P. K. E., ... Johansson, I. (2019). Genome-wide analysis of dental caries and periodontitis combining clinical and self-reported data. *Nature Communications*, 10(1), 2773. <https://doi.org/10.1038/s41467-019-10630-1>
- Skrivankova, V. W., Richmond, R. C., Woolf, B. A. R., Davies, N. M., Swanson, S. A., VanderWeele, T. J., Higgins, J. P. T., Timpson, N. J., Dimou, N., Langenberg, C., Golub, R. M., Loder, E. W., Gallo, V., Tybjaerg-Hansen, A., Smith, G. D., Egger, M., & Richards, J. B. (2021). Strengthening the reporting of observational studies in epidemiology using mendelian randomisation (STROBE-MR): Explanation and elaboration. *BMJ*, 375, n2233. <https://doi.org/10.1136/bmj.n2233>
- Slob, E. A. W., & Burgess, S. (2020). A comparison of robust Mendelian randomization methods using summary data. *Genetic Epidemiology*, 44(4), 313–329. <https://doi.org/10.1002/gepi.22295>
- Teles, F., Wang, Y., Hajishengallis, G., Hasturk, H., & Marchesan, J. T. (2021). Impact of systemic factors in shaping the periodontal microbiome. *Periodontology 2000*, 85(1), 126–160. <https://doi.org/10.1111/prd.12356>
- Tsoi, L. C., Stuart, P. E., Tian, C., Gudjonsson, J. E., Das, S., Zawistowski, M., Ellinghaus, E., Barker, J. N., Chandran, V., Dand, N., Duffin, K. C., Enerbäck, C., Esko, T., Franke, A., Gladman, D. D., Hoffmann, P., Kingo, K., Köks, S., Krueger, G. G., ... Elder, J. T. (2017). Large scale meta-analysis characterizes genetic architecture for common psoriasis associated variants. *Nature Communications*, 8, 15382. <https://doi.org/10.1038/ncomms15382>
- Ungprasert, P., Wijarnpreecha, K., & Wetter, D. A. (2017). Periodontitis and risk of psoriasis: A systematic review and meta-analysis. *Journal of the European Academy of Dermatology and Venereology*, 31(5), 857–862. <https://doi.org/10.1111/jdv.14051>
- Van Dyke, T. E., Bartold, P. M., & Reynolds, E. C. (2020). The nexus between periodontal inflammation and dysbiosis. *Frontiers in Immunology*, 11, 511. <https://doi.org/10.3389/fimmu.2020.00511>
- VanderWeele, T. J., Tchetgen Tchetgen, E. J., Cornelis, M., & Kraft, P. (2014). Methodological challenges in mendelian randomization. *Epidemiology*, 25, 427–435. <https://doi.org/10.1097/EDE.0000000000000081>
- Wang, L., Gao, B., Fan, Y., Xue, F., & Zhou, X. (2021). Mendelian randomization under the omnigenic architecture. *Briefings in Bioinformatics*, 22(6), bbab322. <https://doi.org/10.1093/bib/bbab322>
- World Health Organization. (2013). *Oral health surveys: Basic methods* (5th ed.). Nonserial Publications.
- Zhang, X., Gu, H., Xie, S., & Su, Y. (2020). Periodontitis in patients with psoriasis: A systematic review and meta-analysis. *Oral Diseases*, 28(1), 33–43. <https://doi.org/10.1111/odi.13617>

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Baurecht, H., Freuer, D., Welker, C., Tsoi, L. C., Elder, J. T., Ehmke, B., Leitzmann, M. F., Holtfreter, B., & Baumeister, S.-E. (2022). Relationship between periodontitis and psoriasis: A two-sample Mendelian randomization study. *Journal of Clinical Periodontology*, 49(6), 573–579. <https://doi.org/10.1111/jcpe.13620>