

Anomalous Josephson effect of s-wave pairing states in chiral double layers

Klaus G. Ziegler, Andreas Sinner, Yurii E. Lozovik

Angaben zur Veröffentlichung / Publication details:

Ziegler, Klaus G., Andreas Sinner, and Yurii E. Lozovik. 2022. "Anomalous Josephson effect of s-wave pairing states in chiral double layers." *Physical Review Letters* 128 (15): 157001.
<https://doi.org/10.1103/physrevlett.128.157001>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under these conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publiz/>



Anomalous Josephson Effect of s -Wave Pairing States in Chiral Double Layers

Klaus Ziegler¹, Andreas Sinner¹, and Yurii E. Lozovik^{2,3}

¹*Institut für Physik, Universität Augsburg, D-86135 Augsburg, Germany*

²*Institute of Spectroscopy, Russian Academy of Sciences, 142190 Troitsk, Moscow, Russia*

³*Moscow Institute of Electronics and Mathematics, National Research University Higher School of Economics, 101000 Moscow, Russia*



(Received 20 April 2021; accepted 22 March 2022; published 12 April 2022)

We consider s -wave pairing in a double layer of two chiral metals due to interlayer Coulomb interaction and study the Josephson effect near a domain wall, where the sign of the order parameter jumps. The domain wall creates two evanescent modes at the exceptional zero-energy point, whose superposition is associated with currents flowing in different directions in the two layers. Assuming a toroidal geometry, the effective Josephson current winds around the domain walls, whose direction is determined by the phase difference of the complex coefficients of the superimposed zero-energy modes. Thus, the zero-energy mode is directly linked to a macroscopic current. This result can be understood as an interplay of the conventional Josephson current perpendicular and the edge current parallel to a domain wall in a double layer of two chiral metals. As a realization we suggest the surface of a ring-shaped topological insulator. The duality between electron-electron and electron-hole double layers indicates that this effect should also be observable in excitonic double layers.

DOI: [10.1103/PhysRevLett.128.157001](https://doi.org/10.1103/PhysRevLett.128.157001)

A Josephson junction is an important tool for probing paired electron states, since it is sensitive to the phase difference of the pairing order parameter on both sides of the junction. The latter generates a current that flows perpendicular to the Josephson junction. This effect has been studied in many systems with paired states, including conventional s -wave superconductors [1,2], as well as unconventional and topological superconductors [3–5].

In this Letter, we will consider the internal (intralayer) Josephson effect in a two-dimensional electronic double layer with interlayer s -wave pairing, assuming that both layers have a Dirac spectrum consisting of two bands and a spectral node (see Fig. 1). Such conditions can be realized, for instance, on the surface of a 3D topological insulator [6,7].

The nodal spectrum influences strongly the formation of local currents due to edge modes along the Josephson junction. They compete with the conventional Josephson current that tends to cross the junction perpendicularly. As a result, we expect an anomalous Josephson effect, where the effective Josephson current has a component that flows parallel to the junction. The details of these competing effects will be studied in this Letter. For this purpose, the Josephson junction is simplified as a domain wall (i.e., it is an infinitesimally narrow Josephson junction). Zero-energy edge modes are created along the domain wall, causing local currents. Finally, we impose periodic boundary conditions to obtain a toroidal geometry with two domain walls.

Our approach is motivated by the fact that double layers of chiral materials have rich properties due to the

combination of electronic interlayer pairing and quasiparticle edge modes. The interplay of a superconducting Josephson current and edge currents represents an intimate connection of the Josephson effect and topology, which does not exist in conventional superconductors. Although the edge modes depend on the sample geometry, the anomalous Josephson effect is robust and determined only by topology, which is characterized by the number of edges and domain walls. As a special example, we will consider in this Letter the case of a torus with two domain walls, which has no edges. Such a geometry can be realized with two metallic layers, separated by a dielectric sheet [8] and connected by a metallic boundary, similar to Fig. 2. The internal Josephson current in the double layer is induced by an external current through inductive coupling. Since the edge currents are directly associated with the zero-energy quasiparticle states, this gives also a direct access to these




FIG. 1. Electronic double layer with domain wall, which is given by a sign jump of the pairing order parameter. The currents (blue arrows) flow in the same (opposite) direction in the two layers parallel (perpendicular) to the domain wall.




FIG. 2. After gluing the double layer of Fig. 1 to form a torus, the current $\mathbf{I} = (j_x + j_x^s, j_y + j_y^s)$ winds along the two domain walls around the torus.

states through the external current. In particular, it enables us to change the direction of the current relative to the domain wall. Thus, the sensitivity of the anomalous Josephson effect to an external current provides a method by which one can probe and control the internal superconducting properties. This opens a wide field for new experiments on superconducting layered materials.

The theory of the electronic double layer is dual to that of an electron-hole double layer due to a duality transformation discussed previously by us [9]. This relation connects the electronic double layer physics with the excitonic physics. In particular, the duality suggests that the interlayer Josephson effect should also exist for the electron-electron double layer when interlayer hopping is present, as it was studied before for electron-hole double layers [10,11]. In that case, the Josephson currents are homogeneous in each layer. In the present Letter, this effect will not appear due to the absence of interlayer hopping. On the other hand, we expect an anomalous intralayer Josephson effect in electron-hole layers when we implement a Josephson junction inside the layers, as visualized in Fig. 1.

Model.—We consider an electronic double layer with interlayer Coulomb repulsion but without interlayer tunneling. The layers themselves are chiral metals, described by Dirac Hamiltonians with opposite chirality. Such a system can be defined by the tight-binding Hamiltonian

$$\mathcal{H}_{ee} = \sum_{\mathbf{r}, \mathbf{r}'} \sum_{s=\uparrow, \downarrow} \sum_{\mu, \mu'} H_{\mathbf{r}\mathbf{r}', s, \mu\mu'} c_{\mathbf{r}, s, \mu}^\dagger c_{\mathbf{r}', s, \mu'} + \sum_{\mathbf{r}, \mathbf{r}'} \sum_{\mu, \mu'} V_{\mathbf{r}\mathbf{r}'} c_{\mathbf{r}, \uparrow, \mu}^\dagger c_{\mathbf{r}, \uparrow, \mu} c_{\mathbf{r}', \downarrow, \mu'}^\dagger c_{\mathbf{r}', \downarrow, \mu'}, \quad (1)$$

where $\mu = 1, 2$ is the band index of the two bands in each layer. $H_{\mathbf{r}\mathbf{r}', \uparrow, \mu\mu'}$ ($H_{\mathbf{r}\mathbf{r}', \downarrow, \mu\mu'}$) is the hopping matrix element in the top (bottom) layer, and $c_{\mathbf{r}, \uparrow, \mu}^\dagger$ creates an electron at site \mathbf{r} in the top layer in the band with index μ . An example is the honeycomb lattice, which is bipartite and consists of two triangular sublattices. In this case, $\mu = 1, 2$ refers to the two triangular lattices and the coordinate \mathbf{r} refers only to one of the two triangular lattices. There exists a duality transformation $c_{\mathbf{r}, \downarrow, \mu} \rightarrow d_{\mathbf{r}, \downarrow, \mu}^\dagger$ (i.e., we replace the electrons in the bottom layer by holes). This implies a transformation $\mathcal{H}_{ee} \rightarrow \mathcal{H}_{eh}$, where \mathcal{H}_{eh} is the Hamiltonian

of an electron-hole gas [9]. The latter interacts via an attractive Coulomb interaction, a system that has been studied intensively in terms of excitons [10,12–21].

Mean-field approximation.—First, we briefly discuss the BCS approach for the electron-hole double layer. For this purpose, we introduce the BCS order parameter

$$\Delta_{\mathbf{r}\mathbf{r}', \mu\mu'} = V_{\mathbf{r}\mathbf{r}'} \langle c_{\mathbf{r}, \uparrow, \mu} d_{\mathbf{r}', \downarrow, \mu'} \rangle, \quad (2)$$

which describes Cooper pairing of electrons and holes by forming excitons [10]. Then the interaction term of \mathcal{H}_{eh} reads in BCS approximation

$$- \sum_{\mathbf{r}, \mathbf{r}'} \sum_{\mu, \mu'} V_{\mathbf{r}\mathbf{r}'} c_{\mathbf{r}, \uparrow, \mu}^\dagger c_{\mathbf{r}, \uparrow, \mu} d_{\mathbf{r}', \downarrow, \mu'}^\dagger d_{\mathbf{r}', \downarrow, \mu'} \approx - \sum_{\mathbf{r}, \mathbf{r}'} \sum_{\mu, \mu'} (d_{\mathbf{r}', \downarrow, \mu'}^\dagger c_{\mathbf{r}, \uparrow, \mu}^\dagger \Delta_{\mathbf{r}\mathbf{r}', \mu\mu'} + \text{H.c.}). \quad (3)$$

The attractive interaction between the electrons in the top layer and the holes in the bottom layer causes electron-hole interlayer pairing despite the fact that the electronic Coulomb interaction $V_{\mathbf{r}\mathbf{r}'}$ is repulsive. This leads to the formation of a BCS state because the electron-hole pairs (excitons) can condense. The interlayer pairing has some similarity with the resonating valence bond idea [22,23], where in the present case the bond consists of the two layers.

With a uniform order parameter Δ , we get for the quasiparticles the Bogoliubov–de Gennes (BdG) Hamiltonian matrix

$$\langle \mathcal{H}_{eh} \rangle \approx H_{\text{BdG}} = \begin{pmatrix} H_\uparrow & \Delta \\ \Delta^\dagger & -H_\downarrow^* \end{pmatrix}, \quad (4)$$

where the 2×2 matrix structure refers to the top and bottom layer, while $H_{\uparrow, \downarrow}$ and Δ are 2×2 matrices with respect to the band index μ . This mean-field (MF) result can be used to transform back $d_{\mathbf{r}, \downarrow, \mu}^\dagger \rightarrow c_{\mathbf{r}, \downarrow, \mu}$, such that we have again electrons in both layers. It gives us the effective quasiparticle Hamiltonian matrix

$$\langle \mathcal{H}_{ee} \rangle \approx H_{\text{MF}} = \begin{pmatrix} H_\uparrow & \Delta \\ \Delta^\dagger & H_\downarrow \end{pmatrix}, \quad (5)$$

which reads for two layers with opposite chiralities

$$H_{\text{MF}} = \begin{pmatrix} h_1 \sigma_1 + h_2 \sigma_2 & \Delta \sigma_2 \\ \Delta \sigma_2 & h_1 \sigma_1 - h_2 \sigma_2 \end{pmatrix}. \quad (6)$$

Here we have assumed that the antisymmetric hopping elements h_1 and h_2 are the same in both layers. σ_j is the Pauli matrix with respect to the sublattice structure, and Δ is a real scalar pairing order parameter. For this Hamiltonian, we will discuss the effect of a domain wall.

In Fourier representation, $h_{1,2}$ and Δ can depend on the 2D wave vector $\mathbf{k} = (k_x, k_y)$. This Hamiltonian belongs to the symmetry class DIII according to Ref. [24], and it has two degenerate bands with first Chern numbers ± 1 . Its gapped quasiparticle dispersion reads $E_{\mathbf{k}} = \pm \sqrt{h_1^2 + h_2^2 + \Delta^2}$. The latter agrees with the dispersion of the Hamiltonian with identical layers [9]

$$H'_{\text{MF}} = \begin{pmatrix} h_1\sigma_1 + h_2\sigma_2 & \Delta\sigma_3 \\ \Delta\sigma_3 & h_1\sigma_1 + h_2\sigma_2 \end{pmatrix}. \quad (7)$$

Thus, the difference in terms of chirality can only be seen in the eigenfunctions, but not in their spectra. It should be noted that in this case the pairing takes place between the same subbands, whereas in the case with opposite chirality the pairing occurs between different subbands.

The dispersion $E_{\mathbf{k}} = \pm \sqrt{h_1^2 + h_2^2 + \Delta^2}$ can vanish when h_1 and/or h_2 are imaginary. This is the case for evanescent modes. The phenomenon is known from conventional Josephson junctions, where evanescent modes exist inside the gap.

Zero-energy modes at a domain wall.—An inhomogeneous order parameter Δ breaks translational invariance and divides the layers in different regions. For a Josephson junction [1], we typically choose a region around $x = 0$, where the order parameter vanishes. For our purpose, we can consider the simple case [23,25] that $\text{sgn}(\Delta)$ jumps at the domain wall along the y direction at $x = 0$, as visualized in Fig. 1. Such a discontinuous phase change suppresses the Andreev states except for the zero-energy modes. It should be kept in mind here that the domain wall is created by a potential, and the corresponding change of the order parameter should be obtained via the BCS-like equation. In practice, this requires some tedious calculations and we simply focus here on the domain wall in terms of $\Delta(x)$, following the recipe proposed in Ref. [25]. Next, we analyze the effect of the domain wall. The system is translational invariant in the y direction, such that we can use Fourier components with respect to k_y . The domain wall breaks the translational invariance in the x direction. Considering only the low-energy MF Hamiltonian, we can write $h_1 \sim i\partial_x$, $h_2 \sim k_y$, and

$$H_{\text{MF}} = \begin{pmatrix} i\partial_x\sigma_1 + h_2\sigma_2 & \Delta(x)\sigma_2 \\ \Delta(x)\sigma_2 & i\partial_x\sigma_1 - h_2\sigma_2 \end{pmatrix}. \quad (8)$$

For the zero-energy mode, we can make the ansatz $\Psi_{k_2}(x) = \psi_{k_2} e^{-bx}$, where b depends on the sign of x . Then we get the eigenmode equation

$$\begin{pmatrix} -ib\sigma_1 + h_2\sigma_2 & \Delta(x)\sigma_2 \\ \Delta(x)\sigma_2 & -ib\sigma_1 - h_2\sigma_2 \end{pmatrix} \Psi_{k_2}(x) = 0. \quad (9)$$

Solving this equation for $x < 0$ and for $x > 0$ and using the matching condition at $x = 0$, the evanescent solutions require $b = \text{sgn}(x)\sqrt{h_2^2 + \Delta^2}$ and $h_2 = 0$, such that $b = \text{sgn}(x)|\Delta|$ and the zero-energy modes read

$$\Psi_1 = \frac{1}{\mathcal{N}} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} e^{-|\Delta||x|}, \quad \Psi_2 = \frac{1}{\mathcal{N}} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix} e^{-|\Delta||x|}, \quad (10)$$

with the normalization $\mathcal{N} = \sqrt{2/|\Delta|}$. Ψ_1 has the same wave function on the top and the bottom layer, whereas Ψ_2 has an opposite sign on the two layers.

For a more general phase change $\exp(-i\theta/2) \rightarrow \exp(i\theta/2)$ at $x = 0$, the matching condition reads

$$b_- = b_+ e^{-i\theta} = b_+ (\cos\theta - i\sin\theta), \quad (11)$$

where b_+ (b_-) refers to the right (left) side with respect to the domain wall. To get exponentially decaying functions $\exp(-b_{\pm}x)$ on both sides of the domain wall, the sign of $\text{Re}(b_-)$ [$\text{Re}(b_+)$] must be negative (positive). This requires $\cos\theta < 0$ due to Eq. (11); i.e., for $\pi/2 < \theta < 3\pi/2$. The decay length of the bound state is $-1/|\Delta| \cos\theta$ in units of the lattice spacing. For $\cos\theta \geq 0$ there is no bound state at the domain wall.

Symmetries and degenerate zero-energy modes.—The characteristic polynomial of the MF Hamiltonian has four degenerate zero solutions for $h_2 = 0$ and $b(x) = \text{sgn}(x)|\Delta|$. We consider the block-diagonal matrices $S_j = \text{diag}(\sigma_j, \sigma_j)$ and $T_j = \text{diag}(\sigma_j, -\sigma_j)$ for $j = 1, 2, 3$. First, for $h_2 = 0$ the MF Hamiltonian is invariant under the transformation $H_{\text{MF}} \rightarrow T_1 H_{\text{MF}} T_1$, which creates a new zero mode Ψ_2 from Ψ_1 as $\Psi_2 = T_1 \Psi_1$. Moreover, T_2 as a sublattice transformation is also a particle-hole transformation, since $T_2 H_{\text{MF}} T_2 = -H_{\text{MF}}$ and, therefore, $T_2 \Psi_E = \Psi_{-E}$. Thus, T_1 and T_2 create from Ψ_1 three more zero modes. In particular, the fact that the transformation matrices obey the following rules

$$S_j^2 = T_j^2 = \mathbf{1}, \quad T_1 T_2 = S_1 S_2 = iS_3 \quad (12)$$

and $S_3 \Psi_1 = \Psi_1$, $T_2 \Psi_1 = i\Psi_2$, and $T_2 T_1 \Psi_1 = S_2 S_1 \Psi_1 = i\Psi_1$ reflects that the zero-energy is an exceptional point with a two-dimensional eigenspace [26]. In the context of line defects in the BdG Hamiltonian, the appearance of exceptional points has been discussed recently [27]. It should be noted that the zero eigenmodes of H_{MF} in Eq. (10) are real. But any superposition of the two zero modes $\Phi = a_1 \Psi_1 + a_2 \Psi_2$ with complex coefficients $a_j = |a_j| e^{i\varphi_j}$ and normalization $|a_1|^2 + |a_2|^2 = 1$ is also a zero mode. Thus, the zero eigenmodes are complex, in general, but can also be chosen as real. Since both zero eigenmodes decay exponentially with $|x|$, we must employ an additional

condition that creates a unique solution for the physical system. The y direction does not select valid solutions because we need $h_2 = 0$ for the matching condition; i.e., the solution must be constant in that direction, which can only be satisfied by periodic boundary conditions. A possible solution is to impose a condition on the physical properties, for instance, by fixing the current density. This will be discussed subsequently.

Currents.—The current densities induced by the zero modes are expressed separately for the top and for the bottom layer, where we have the wave functions

$$\Phi_{\uparrow} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \frac{e^{-|\Delta||x|}}{\mathcal{N}}, \quad \Phi_{\downarrow} = \begin{pmatrix} a_1 \\ -a_2 \end{pmatrix} \frac{e^{-|\Delta||x|}}{\mathcal{N}}. \quad (13)$$

Using the quasiparticle continuity equation [28], we get for the top layer

$$\partial_t \Phi_{\uparrow} \cdot \Phi_{\uparrow} + \partial_x j_{x\uparrow} = i\Delta(x) \Phi_{\uparrow} \cdot \sigma_2 \Phi_{\downarrow} \quad (14)$$

and for the bottom layer

$$\partial_t \Phi_{\downarrow} \cdot \Phi_{\downarrow} + \partial_x j_{x\downarrow} = i\Delta(x) \Phi_{\downarrow} \cdot \sigma_2 \Phi_{\uparrow}, \quad (15)$$

where the scalar product contains an implicit complex conjugation: $\Phi_{\uparrow} \cdot \Phi_{\uparrow} \equiv |\Phi_{\uparrow 1}|^2 + |\Phi_{\uparrow 2}|^2$. The current densities are related to the current operator $(e/i\hbar)[H_{\text{MF}}, r_{\mu}]$, projected onto the top or bottom layer, respectively,

$$j_{x\uparrow} = -j_{x\downarrow} = |\Delta| |a_1 a_2| \cos(\varphi_2 - \varphi_1) e^{-2|\Delta||x|}. \quad (16)$$

The terms on the right-hand side of Eqs. (14) and (15) represent the source and drain provided by the pairing condensate [28], which can be identified with the supercurrents $j_{x\sigma}^s$ through the relation

$$\partial_x j_{x\uparrow}^s = -i\Delta(x) \Phi_{\downarrow} \cdot \sigma_2 \Phi_{\uparrow} \quad (17)$$

and accordingly for $j_{x\downarrow}^s$. An x integration of this equation from the domain wall to some position x then provides

$$j_{x\uparrow,\downarrow}^s(x) \sim \pm |\Delta| |a_1 a_2| \cos(\varphi_2 - \varphi_1) (1 - e^{-2|\Delta||x|}). \quad (18)$$

Since the wave function is constant with respect to the y component, the corresponding current densities

$$j_{y\uparrow} = j_{y\downarrow} = |\Delta| |a_1 a_2| \sin(\varphi_2 - \varphi_1) e^{-2|\Delta||x|} \quad (19)$$

do not appear in the continuity equations. The properties $j_{y\uparrow} = j_{y\downarrow}$ ($j_{x\uparrow} = -j_{x\downarrow}$) reflect the fact that the currents in the two layers are (anti)correlated (cf. Fig. 1). This effect should be experimentally observable, since the interlayer current-current correlation is associated with the drag effect [9,29].

A nonvanishing current requires that both zero modes contribute (i.e., $a_1, a_2 \neq 0$). This implies that according to Eq. (13) the eigenvectors Φ_{\uparrow} and Φ_{\downarrow} are linearly independent. The currents of the two layers in the y direction are not balanced in contrast to the currents in the x direction; i.e., they do not cancel each other in the two layers. Thus, there is a net current along the y direction in the double layer (cf. Fig. 1). Since the layers are charge separated, there is no charge current between them. Therefore, the currents must be conserved in each layer.

The picture in Fig. 1 is incomplete though because the edges of the layers have not been included. But since the wave functions decay exponentially away from the domain wall and the order parameter is the same in both layers, we have effectively periodic boundary conditions in x and y directions, resulting in the toroidal geometry of Fig. 2.

After preparing the wave function $\Phi = (\Phi_{\uparrow}, \Phi_{\downarrow})$ we create a current density $(j_{x,\uparrow}, j_{y,\uparrow})$ in the top and $(-j_{x,\uparrow}, j_{y,\uparrow})$ in the bottom layer. By changing the wave function Φ through the coefficients a_j these current densities also change according to the above relations. This change can be achieved experimentally with an external current source that couples inductively to the double layer. By choosing the direction of the current (i.e., the angle $\varphi_2 - \varphi_1$), we excite the corresponding wave function Φ .

Discussion.—The existence of edge modes, a consequence of the chiral metallic layers, affects the Josephson currents near the domain wall. A measure of the interplay between the conventional Josephson current and the edge current is the direction of the quasiparticle current with respect to the domain walls in Fig. 2. Such edge modes appear on all edges including the sample boundaries. In Fig. 1, the latter have not been depicted because an infinite 2D sample was assumed. On the other hand, it is well known that a consistent description of edge modes requires a compact manifold [23]. Figure 2 presents a compact version of the double layer as a single layer on a torus (i.e., for periodic boundary conditions), which has two domain walls. Such a geometry can be realized as the surface of a ring-shaped topological insulator.

The existence of two degenerate zero modes requires an additional physical constraint to lift the degeneracy and to obtain a unique solution. In our case, this was achieved by considering the current in the sample. Depending on that current, we get a specific linear combination of the two zero modes. From a physical perspective this means that we induce a current density in the system by coupling it to an external current, which excites the corresponding quasiparticle state. The exponential decay of quasiparticle modes and their corresponding currents away from the domain walls on the scale $1/|\Delta|$ implies the existence of a supercurrent due to charge conservation [28]. In other words, the quasiparticle currents near the domain wall are part of a stationary current inside the entire torus, winding around the domain walls (cf. Fig. 2). This current, which is

proportional to $|\Delta|$, could be measured through inductance, e.g., by using a coil. Very accurate measurements of the current can be performed with a superconducting quantum interference device [30]. A more direct probe of the zero modes would be possible with an electronic Mach-Zehnder interferometer [31,32].

Our calculation was performed for the special example of a jump of the order parameter phase $\pm\Delta$. Similar calculations can be done for other shapes of $\Delta = |\Delta|e^{i\alpha(x)}$, provided that the order parameter phase represents a kink with a global phase change from one boundary to the other in the x direction. Although the anomalous Josephson effect will not be changed qualitatively, the decay of the wave functions and of the current densities is increased to $-1/|\Delta| \cos \theta$ for a phase change θ . Moreover, a broader Josephson junction might have different modes (cf. discussion in Ref. [33]).

In conclusion, we found an anomalous Josephson current caused by the superposition of two zero-energy modes in the vicinity of the domain wall. The direction of this current is determined by the coefficients of the superposition. Conversely, an external current in a specific direction can induce a certain superposition of the zero modes on the surface of a ring-shaped topological insulator.

This research was supported by a grant of the Julian Schwinger Foundation for Physics Research. Y. E. L. was supported through the Grants No. RFBR 20-02-00410 and No. 20-52-00035.

-
- [1] B. Josephson, *Phys. Lett.* **1**, 251 (1962).
 - [2] P. W. Anderson and J. M. Rowell, *Phys. Rev. Lett.* **10**, 230 (1963).
 - [3] Y. Tanaka, T. Hirai, K. Kusakabe, and S. Kashiwaya, *Phys. Rev. B* **60**, 6308 (1999).
 - [4] Y. Asano, Y. Tanaka, M. Sigrist, and S. Kashiwaya, *Phys. Rev. B* **67**, 184505 (2003).
 - [5] H.-J. Kwon, K. Sengupta, and V. M. Yakovenko, *Eur. Phys. J. B* **37**, 349 (2004).
 - [6] X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).
 - [7] A. A. Burkov, *J. Phys. Condens. Matter* **27**, 113201 (2015).
 - [8] A. K. Geim and I. V. Grigorieva, *Nature (London)* **499**, 419 (2013).
 - [9] A. Sinner, Yurii E. Lozovik, and K. Ziegler, *Phys. Rev. Research* **2**, 033085 (2020).
 - [10] Y. E. Lozovik and V. I. Yudson, *Sov. J. Exp. Theor. Phys.* **44**, 389 (1976).
 - [11] Y. E. Lozovik and A. V. Poushnov, *Phys. Lett. A* **228**, 399 (1997).
 - [12] B. Halperin and R. Englman, *Solid State Commun.* **66**, 711 (1988).
 - [13] J. P. Eisenstein and A. H. MacDonald, *Nature (London)* **432**, 691 (2004).
 - [14] A. V. Balatsky, Y. N. Joglekar, and P. B. Littlewood, *Phys. Rev. Lett.* **93**, 266801 (2004).
 - [15] P. Pieri, D. Neilson, and G. C. Strinati, *Phys. Rev. B* **75**, 113301 (2007).
 - [16] J.-J. Su and A. H. MacDonald, *Nat. Phys.* **4**, 799 (2008).
 - [17] D. K. Efimkin, Y. E. Lozovik, and A. A. Sokolik, *Phys. Rev. B* **86**, 115436 (2012).
 - [18] C. Zhang and G. Jin, *Appl. Phys. Lett.* **103**, 202111 (2013).
 - [19] P. Pieri, D. Neilson, and A. R. Hamilton, *Phys. Rev. Lett.* **110**, 146803 (2013).
 - [20] J. I. A. Li, T. Taniguchi, K. Watanabe, J. Hone, and C. R. Dean, *Nat. Phys.* **13**, 751 (2017).
 - [21] Z. Wang, D. A. Rhodes, K. Watanabe, T. Taniguchi, J. C. Hone, J. Shan, and K. F. Mak, *Nature (London)* **574**, 76 (2019).
 - [22] P. W. Anderson, *Science* **235**, 1196 (1987).
 - [23] E. Fradkin, *Field Theories of Condensed Matter Physics*, (Cambridge University Press, Cambridge, England, 2013).
 - [24] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, *Phys. Rev. B* **78**, 195125 (2008).
 - [25] N. Read and D. Green, *Phys. Rev. B* **61**, 10267 (2000).
 - [26] T. Kato, *Perturbation Theory for Linear Operators*, Grundlehren der mathematischen Wissenschaften A Series of Comprehensive Studies in Mathematics, 2nd ed. (Springer, Berlin, 1976).
 - [27] I. Mandal, *Europhys. Lett.* **110**, 67005 (2015).
 - [28] G. E. Blonder, M. Tinkham, and T. M. Klapwijk, *Phys. Rev. B* **25**, 4515 (1982).
 - [29] G. Vignale and A. H. MacDonald, *Phys. Rev. Lett.* **76**, 2786 (1996).
 - [30] S. Das Sarma, M. Freedman, and C. Nayak, *Phys. Rev. Lett.* **94**, 166802 (2005).
 - [31] Y. Ji, Y. Chung, D. Sprinzak, M. Heiblum, D. Mahalu, and H. Shtrikman, *Nature (London)* **422**, 415 (2003).
 - [32] A. R. Akhmerov, J. Nilsson, and C. W. J. Beenakker, *Phys. Rev. Lett.* **102**, 216404 (2009).
 - [33] J. A. Sauls, *Phil. Trans. R. Soc. A* **376**, 20180140 (2018).