

The methyl rotational potentials of various isotopic methyl iodides and their coupling to phonons

M. Prager, J. Stanislawski, Wolfgang Häusler

Angaben zur Veröffentlichung / Publication details:

Prager, M., J. Stanislawski, and Wolfgang Häusler. 1987. "The methyl rotational potentials of various isotopic methyl iodides and their coupling to phonons." In *Quantum aspects of molecular motions in solids: proceedings of an ILL-IFF Workshop, Grenoble, France, September 24–26, 1986*, edited by Anton Heidemann, Andreas Magerl, Dieter Richter, Michael Prager, and Tasso Springer, 107–12. Berlin: Springer.
https://doi.org/10.1007/978-3-642-71914-1_20.



The Methyl Rotational Potentials of Various Isotopic Methyl Iodides and Their Coupling to Phonons

M. Prager¹, J. Stanislawski¹, and W. Häusler²

¹Institut für Festkörperforschung der Kernforschungsanlage Jülich, Postfach 1913, D-5170 Jülich, Fed. Rep. of Germany

²Institut für Theoretische Physik der Universität Erlangen-Nürnberg, Glückstr. 6, D-8520 Erlangen, Fed. Rep. of Germany

The rotational potentials of methyl groups in protonated and deuterated methyl iodide were determined from the observed tunnel splittings and the librational energies. The potential becomes stronger in the deuterated matrix by about 10 %. For pure CD_3I the isotope effect on the librational modes shows a considerable further increase. No rotor-rotor coupling is found. The coupling to low-frequency phonons can explain the unusual increase of the tunnel splitting with temperature observed for $T < 15 \text{ K}$.

1. Introduction

The methyl halides are the simplest molecules containing CH_3 groups and thus among the most simple molecules of all. They represent a symmetric prolate top, with the small momentum of inertia being equal to that of the CH_3 group. Methyl iodide has only one solid phase: Pnma with $Z=4$ molecules at c-sites /2/. The molecules are arranged in the a-c-plane as chains along the c-direction with alternating orientation of the molecules.

Various reasons make the methyl iodide interesting from the point of view of methyl group rotation: i) because of the molecular symmetry there is no intramolecular hindering potential. The methyl rotation is identical with the overall rotation of the whole molecule around its small axis of inertia. The rotational potential fully originates from the interaction with neighbouring molecules. ii) The temperature behaviour of the tunnelling modes, i.e. their coupling to phonons /3,4/, might be especially simple because the libration of the methyl group is an external mode. In force models available for molecular crystals /5,6/ it can be found out by symmetry analysis with what phonons at all the methyl rotation can interact. Thus not the full density of states /4/ but individual phonon branches might be of importance. This could stimulate new theoretical efforts. iii) The lattice dynamics of molecular crystals consisting of polar molecules is an actually very active field /6/. Due to the simplicity of the molecules and the structure of the methyl halides CH_3I might be a candidate for a model system in this field.

1) This work is a part of a more comprehensive study of the methyl halides /1/

While methyl iodide is already intensively investigated with the Raman and FIR technique /7,8/ there is only one incoherent neutron scattering experiment known to us /9/ which was mainly concentrating on multiphonon scattering and not on the methyl group rotation. It is the scope of the present experiment, covering the energy range from 0.4 μeV to 60 meV and temperatures from 4K to 100K for various isotopic species, to determine the rotational potential of the methyl groups, the effect of isotopic substitution on this potential and the coupling of the quantum rotor to phonons.

2. Experiments and Results

The high-resolution experiments were performed at the backscattering spectrometer IN10 at the ILL, Grenoble, using the best obtainable energy resolution $\delta E_e = 0.4 \mu\text{eV}$. The energy range $2 < E [\text{meV}] < 60$ was investigated with the thermal time-of-flight spectrometer SV22 at the FRJ2 in Jülich using various combinations of incoming neutron energy and energy resolution.

The high-resolution spectrum of CH_3I shows a tunnelling peak of resolution width at an energy transfer $\hbar\omega_t = 2.44 \mu\text{eV}$ (fig 1). With temperature this peak behaves quite unusually (fig. 2): Its frequency increases up to a temperature $T \sim 20 \text{ K}$. Thereafter it decreases as usually observed. The broadening, described by a Lorentzian, follows in the whole temperature range very well an Arrhenius behaviour with an activation energy $E_a^\Gamma = 14 \text{ meV}$. Tunnelling spectra were also recorded for 2 % and 50 % CH_3I in CD_3I . A linear decrease with concentration is found yielding for infinitely low concentration $\hbar\omega_t = 2.02 \mu\text{eV}$.

The prominent feature of the spectra in the meV region is a strong doublet line. Fig. 3 shows spectra of CH_3I , CH_3I in CD_3I and CD_3I . With increasing temperature




Fig. 1 Tunnelling spectrum of CH_3I .
Energy resolution $\delta E = 0.4 \mu\text{eV}$.
Momentum transfer $Q = 1.88 \text{ \AA}^{-1}$.




Fig. 2 Temperature dependence of the tunnel splitting in CH_3I . The solid line is a fit with the model described in the text.




Fig. 3 Spectra of a) CH_3I , b) CD_3I and c) CH_3I in CD_3I (the spectrum of CD_3I is subtracted to show only the transition energies of the protonated molecules) in the meV region. The low-energy peak of each doublet represents the libration of the methyl groups. See the text.

all transition energies shift to lower values and the line intensities diminish. The lines at 13.27 meV in CH_3I , 13.8 meV in CH_3I diluted in CD_3I and 10.6 meV in CD_3I are faster damped with temperature than each second strong peak.

3. Discussion

As a consequence of the structure [1] all methyl groups in solid methyl iodide are equivalent. The relative intensities of the tunnelling and elastic lines (fig. 1) are in agreement with the theoretically predicted values. No effect of coupling, suggestable from the head-to-head arrangement of the molecules and as found in lithium acetate [10], is seen in the tunnelling spectrum. We first also believed that the doublet line in the meV region is a consequence of coupling. However, because of the different temperature dependences of the two lines and because of the different isotope effects (see below), they must be of different origin.

Since no effects of rotor-rotor coupling could be found we describe the methyl rotational energies by a single particle rotational potential:

$$V(\varphi) = -\frac{V_3}{2} \cos 3\varphi + \frac{V_6}{2} \cos 6\varphi. \quad (1)$$

From the eigenvalues $\hbar\omega_l = 2.44 \mu\text{eV}$ and $E_{01} = 13.27 \text{ meV}$ we obtain

$$V_5 = V_3 + V_6 = 24.9 \text{ meV}, \quad \delta = \frac{V_3}{V_5} = 0.92. \quad (2)$$

A second solution with $V_3, V_6 < 0$ can be excluded from the measured isotope effect (see below). Thus, in agreement with the general experience /10/, the deviation from a purely threefold cosine potential is small. The potential also describes rather well the second excited librational state found at 24.2 meV (calculated energy 24.6 meV). The second strong peak at 14.54 meV is assigned to a libration R_x, R_y of the molecule perpendicular to its symmetry axis.

Beside the temperature dependence, the isotope effect is useful to identify the character of the modes related with the observed transition energies. Generally the absolute strength of the potentials changes not or very little by deuteration, and the eigenvalues of the deuterated compound are obtained by simply scaling the masses. For librational modes the effect is roughly of the order $\sqrt{2}$.

In CH_3I we expect exactly 1.35 by scaling the potential (2). However, in CD_3I we find only a shift by a factor 1.25 to 10.6 meV. The mode at 10.6 meV was identified as libration via its temperature dependence. This means that the rotational potential must have considerably increased by deuteration. Keeping fixed the shape of the potential (2), characterized by δ , we find $V_S = 29.5$ meV or an increase by 25 %, probably caused by a relaxation of the CD_3I molecule into a new equilibrium orientation in the a-c-plane. Because of this strong isotope effect, earlier Raman and FIR lines were wrongly assigned /6/. The R_x, R_y libration has little decreased to 13.1 meV as expected.

An independent determination of this large isotope effect was achieved by using CH_3I as a probe in the matrix of CD_3I . Now two independent transition energies, $E_{01} = 2.02$ meV and $E_{01} = 13.8$ meV are available to determine the rotational potential. The shape is found to be unchanged, while the strength has increased by 10 % to $V_S = 27.5$ meV compared to pure CH_3I . It is reasonable that the potential is weaker than for pure CD_3I since the properties of both, the surrounding and the 'Auf'-molecule, enter in the interaction (e.g. octopole-octopole interaction in methane).

The tunnel splitting in methyl iodide shows a very unusual increase with temperature before it decreases in the well-known way /11/. A similar behaviour was recently found in some other materials also, e.g. /12/ and /13/. The large isotope effect has already shown that the molecular orientation in methyl iodide is sensitive to shallow changes of the interaction. Thus we expect that it also might be influenced by the temperature and change the rotational potential. Such a temperature behaviour can be described in a model which couples the quantum rotor to low-energy phonons via an interaction Hamiltonian

$$H_{\text{int}}^C = g^C \sqrt{2m\omega/\hbar} x \cos 3\varphi \quad (3)$$

not considered so far /3,4/. Here g^C is the coupling constant, x the oscillator coordinate and φ the rotational angle. The Hamiltonian of the whole system is

$$H = H_{\text{rot}} + H_{\text{osc}} + H_{\text{int}}^C \quad (4)$$

with

$$H_{\text{rot}} = \frac{L^2}{2I} + V_3 \cos 3\varphi,$$

$$H_{\text{osc}} = \frac{p^2}{2m} + \frac{m}{2} \tilde{\omega}^2 x^2.$$

Using the formalism developed in ref. 14 the tunnelling frequency corresponding to the excitation of n and $n + 1$ oscillator levels is found to change as

$$\hbar \omega_t(n+1) / \hbar \omega_t(n) = 1 + A^C, \quad \text{with } A^C > 0$$

for moderate coupling g^C . If the decrease of the tunnelling frequency at higher temperatures is taken into account phenomenologically, we get the following expression for the temperature dependence of the tunnelling frequency

$$\hbar \omega_t(T) = \hbar \omega_t(0) \left\{ \frac{1 - \exp(-E^C/k_B T)}{1 - (1 + A^C) \exp(-E^C/k_B T)} - A^S \exp(-E^S/k_B T) \right\}. \quad (5)$$

The solid curve in fig. 2 is a fit using this formula with parameters $A^C = 0.22$, $A^S = 49.06$, $E^C = 2.15$ meV and $E^S = 12.94$ meV. While E^S is close to E_{01} as usually found, E^C represents the phonon energy. Indeed the density of states shows increased intensity in this energy range (fig. 3). The coupling parameter derived from these values is $g^C = 2.9$ meV, the strength of the rotational potential $V_3 = 18.2$ meV.

4. Conclusions

Rotational tunnelling and librations of methyl iodide could be well explained as a single particle rotation. No methyl rotor-rotor coupling, suggested from the structure, can be observed: the two transitions around 14 meV belong to two different molecular rotations (around and perpendicular to the symmetry axis) and are not a consequence of coupling. With deuteration the rotational potential of the methyl group increases by 25 %, probably due to a change of the equilibrium orientation. This large isotope effect beside the simple mass effect is unambiguously confirmed by using CH_3I as a probe in CD_3I . The very unusual increase of the tunnel splitting with temperature for $T < 15$ K could be explained by a coupling to low-frequency phonons via an interaction Hamiltonian $H_{\text{int}}^C \sim x \cos 3\varphi$ not considered so far.

It would be very useful for a more thorough interpretation of our data to investigate the lattice dynamics of methyl iodide.

References

1. M. Prager, J. Stanislawski and W. Häusler
submitted to J. Chem. Phys.
2. T. Kawaguchi, M. Hijikigawa, Y. Hayafiyi, M. Ikeda, R. Fukishima
and Y. Tomie, Bull. Chem. Soc. Jpn. 46, 53 (1973)

3. A. Hüller, Z. Phys. B36, 215 (1980)
4. A.C. Hewson, J. Phys. C15, 3841 and 3855 (1982)
5. P. Brüesch, Phonons: Theory and Experiments I
Springer Series in Solid State Sciences Vol. 34 (Springer, Berlin,
Heidelberg, New York 1982)
6. K.H. Link, H. Grimm, B. Dorner, H. Zimmermann, H. Stiller and
P. Bleckmann, J. Phys. Chem. Solids 46, 135 (1985)
7. H. Takeuchi, J.L. Bribes, I. Hirada and T. Shimanouchi,
J. Raman Spectrosc. 4, 235 (1976)
8. O.S. Binbrek, A. Anderson, B. Andrews and B.H. Torrie,
J. Raman Spectrosc. 15, 406 (1984)
9. J.A. Janik, A. Bajorek, J.M. Janik, I. Natkaniec, K. Parlinski and
M. Sudnik, Acta Physica Polonica XXXIII, 419 (1968)
10. S. Clough, A. Heidemann, A.H. Horsewill and M.N.J. Paley
Z. Phys. B55, 1 (1984)
11. W. Press, Springer Tracts in Modern Physics, Vol. 92,
(Springer, Berlin, Heidelberg, New York 1981)
12. M. Prager and W. Müller-Warmuth, Z. Naturforschung 39a, 1187 (1984)
13. L.P. Ingman, M. Punkkinen, A. H. Vuorimäki and E.E. Ylinen,
J. Phys. C18, 5033 (1985)
14. W. Häusler and A. Hüller, Z. Phys. B59, 177 (1985)