

Evaluation of Ventricular Repolarization Variability in Patients With Nonischemic Dilated Cardiomyopathy From Vectorcardiography

Martin Schmidt¹, Filip Karisik², Sebastian Zaunseder³, Axel Linke⁴, Hagen Malberg¹, Mathias Baumert²

¹ Institute of Biomedical Engineering, TU Dresden, Dresden, Germany

² School of Electrical and Electronic Engineering, University of Adelaide, Adelaide, Australia

³ Faculty of Information Technology, University of Applied Science and Arts Dortmund, Dortmund, Germany

⁴ Department for Internal Medicine and Cardiology, Herzzentrum Dresden, TU Dresden, Dresden, Germany

Abstract

To investigate the predictive value of Ventricular repolarization variability (VRV) in patients with nonischemic dilated cardiomyopathy, we analyzed the Defibrillator in Non-Ischemic Cardiomyopathy Treatment Evaluation trial (DEFINITE). The Telemetric and Holter ECG Warehouse (THEW) data set E-HOL-03-0401-017 comprises 393 recordings from 236 patients. All patients had a left ventricular ejection fraction < 36 % and were randomized to receiving standard medical therapy with or without an ICD. 24h-Holter 3-lead (Frank lead system) ECGs were performed at enrollment and after up to 5 years' follow-up. The all-cause mortality during the follow-up period was 4.8 %. We analyzed three-dimensional variability of the T-loop and QT interval variability on a single lead basis by employing three-dimensional signals adaptation and two-dimensional signal warping, respectively, to quantify VRV. To assess the predictive value of VRV parameters, Kaplan-Meier survival curves of baseline Holter ECGs were calculated. Our results showed significant association to survival ($P < 0.01$ by the log-rank test) for T wave amplitude corrected QT interval variability index ($cQTV_i$) on single lead basis. Low $cQTV_i$ group showed no mortality for the entire observation period. We found no associations between $cQTV_i$ groups and patient-specific parameters.

1. Introduction

Ventricular repolarization variability (VRV) has predictive value for mortality in patients with ischemic heart disease [1], where the mode of death is likely malignant ventricular arrhythmia [2]. However, the potential for risk stratification in patients with nonischemic cardiomyopathy

remains unclear [2].

QT interval variations of the surface electrocardiogram (ECG) reflect beat-to-beat fluctuations of the ventricular repolarization. Several studies have shown that temporal VRV as indicated by an increased QT interval variability (QTV) is associated with cardiac mortality [2, 3]. However, measuring subtle beat-to-beat changes in QT interval remains challenging. Although novel QTV techniques have improved sensitivity and robustness [4, 5], conventional QTV measures still lack in insightful QTV description [2]. Alongside the improved QTV measurement, new techniques are able to capture VRV from vectorcardiography (VCG) [6] and thus offer new feature sets for the characterization of the ventricular repolarization.

In this study we investigate the predictive value of VRV for all-cause mortality in patients with nonischemic dilated cardiomyopathy. Therefore, we analyzed the Defibrillator in Non-Ischemic Cardiomyopathy Treatment Evaluation trial (DEFINITE) to demonstrate (1) the applicability of new methods to characterize VRV using the VCG and (2) to investigate the predictive value of VRV parameters from VCG for risk stratification of all-cause mortality in the DEFINITE study.

2. Material and Methods

2.1. Data Material

The Telemetric and Holter ECG Warehouse (THEW) data set E-HOL-03-0401-017 was used for analysis [7]. The data set comprises 393 recordings from 236 patients of the DEFINITE study (67 women, age 60 ± 14 years; 168 men, age: 58 ± 12 years; 1 record not specified). All patients had a left ventricular ejection fraction < 36% and were randomized to receiving standard medical therapy

with or without an ICD (see Tab. 1). 24h-Holter 3-lead (Frank lead system) ECGs were performed at enrollment and after up to 5 years' follow-up with a sampling frequency of 500 Hz and an amplitude resolution of $1 \mu\text{V}$ [8]. The all-cause mortality during the follow-up period was 4.8%.

Table 1. Characteristics of DEFINITE patients used (THEW data) at baseline. Percentages are in normal brackets and ranges are in square brackets.

Characteristic	Therapy	
	Standard	ICD
Patients	103 (44 %)	132 (56 %)
Sex	male	70 (30 %)
	female	33 (14 %)
Age, yr	mean 58.0 [22-80]	58.3 [26-82]
NYHA	I	25 (11 %)
	II	56 (24 %)
	III	22 (9 %)
LVEF, %	mean 21.8 [10-36]	20.6 [9-35]

2.2. VRV Quantification

We analyzed three-dimensional variability of the T loop and QT interval variability on a single lead basis by employing three-dimensional signal adaptation (3DSA) [6] and iterative two-dimensional signal warping (i2DSW) [5, 9], respectively, to quantify VRV.

Using orthogonal ECG leads, a vector ECG was generated. Its loops are characterized by 3DSA. This adaptation takes place in different phases, the global and the local template adaptation. Templates of interest are first globally translated, scaled and rotated using statistical methods in the form of Procrustes analysis. In the global phase, the template is roughly adapted via Procrustes' analysis in order to be subsequently adapted to the local morphology changes via a kernel ridge regression. To account for complex noise behavior encountered in quasi-periodic signals, 3DSA ensues to complete the framework with an adaptive noise model. VRV was evaluated by 3DSA using the two parameters mean-loop-length (MLL) and distance variability (DV) from the T loop. MML is calculated as the sum of the distances between adjacent points on a loop and DV contains a vector of distances between related points on the template loop and the matched template loop. Mean (DVmean) and standard deviation (DVsd) of DV were used. [6]

Single lead analysis was performed in pseudo-orthogonal lead bipolar Z because of the highest T wave amplitude and thus maximized signal-to-noise ratio [10]. By applying i2DSW we were able to account for subtle inhomogeneous shape variations of the ECG waveform

in time and amplitude [5, 9]. i2DSW automatically generates a template beat with common features of interest (e.g. the PQ, QRS or QT intervals or amplitude related information) based on ensemble averaging of appropriate beats. To adapt the template, a 2D mesh of warping points is superimposed. By shifting warping points in time- and amplitude-direction and minimizing the Euclidean distance between template and the beat under consideration, the optimal template adaptation is calculated. From the adapted template, changes of annotated points can be tracked from beat to beat in time and amplitude. Next to the adapted template we used QT intervals and T wave amplitudes for further analysis. Automatic beat rejection [5] was applied to exclude noisy heart beats. We used standard QTV parameters including standard deviation of QT intervals (SDQT) and the QT interval variability index (QTVi) [11]. To consider the inverse relationship between QTV and T wave amplitude, we calculated the T wave amplitude-corrected QTV measures of cSDQT (cSDQT) and cQTVi (cQTVi) [10].

All parameters were calculated from 5-min epochs throughout the 24-h Holter ECGs. To characterize daily VRV we calculated the median, max quantile (highest 95 %) and min quantile (lowest 5 %).

2.3. Survival Analysis

To assess the predictive value of VRV parameters, Kaplan-Meier survival curves of baseline Holter ECGs were calculated for trichotomized median, min and max of SDQT, QTVi, cSDQT, cQTVi, MLLT, DVmean, and DVsd. Significant parameters were further investigated for their correlations with patient-specific characteristics to show their independent value. Furthermore, Spearman's correlation analysis was performed for a clearer understanding of the relationships between the analyzed parameters.

3. Results and Discussion

Kaplan-Meier survival analysis shows significant association between survival and cQTVi ($P < 0.01$ by the log-rank test; Fig. 1). At year one the mortality was zero in the low and the medium cQTVi group and 5.0 % in the group with high cQTVi. After 2 years, it was 0.0 % in the low and the medium cQTVi group and 10.7 % in the group with high cQTVi. At the end of survival estimation (after 3 years) cQTVi was 0.0 % in the low cQTVi group, 6.3 % in the medium cQTVi group and 10.7 % in the group with high cQTVi. We found qualitatively similar results when using uncorrected QTVi, but the difference in survival across tertiles was not statistically significant ($P = 0.17$ by the log-rank test). Also, all other analyzed parameters have not been able to achieve significant pre-




Figure 1. Hazard functions of Kaplan–Meier survival analysis; Estimated hazard of DEFINITE patients for trichotomized cQTVi (left) and trichotomized DVmeanMax (right) for 3 years. Last observations of a patient are marked in the curves with a cross (censored data).

diction results. However, max of DVmean, extracted from the VCG, also shows qualitatively similar behavior for predicting mortality (see Fig. 1).

Table 2 shows the patient-specific characteristics of the trichotomized cQTVi. No relevant or even statistically significant differences were found between cQTVi groups with respect to sex, age, therapy, NYHA class, and LVEF.

Table 2. Patient characteristics according to trichotomized cQTVi for sex, age, therapy, NYHA class, and LVEF at baseline. Ranges are shown in square brackets.

		cQTVi		
		low	median	high
Sex	male	41	44	46
	female	18	15	15
Age, yr	mean	56.8	60.8	57.4
		[26-82]	[27-80]	[22-81]
Therapy	Standard	23	27	29
	ICD	36	32	31
NYHA	I	18	14	16
	II	32	33	32
	III	9	12	12
LVEF, %	mean	20.8	22.0	20.0
		[10-34]	[10-35]	[9-34]

Results of the correlation analysis between the studied parameters are shown in Figure 2. We found medium effects ($|\rho| > 0.3$) between of DVmean and DVsd (median, min and max) and SDQT and QTVi (median and max). The largest effect ($|\rho| > 0.5$) is between max of SDQT and max of DVmean. Correlations between the VCG parameters and the single channel parameters are lower after T wave amplitude-correction.

Low cQTVi group showed zero mortality for the entire




Figure 2. Correlation analysis between single lead ECG (SDQT, cSDQT, QTVi, cQTVi) and VCG (MLLT, DVmean, DVsd) VRV parameters (highlighted in red). Absolute values of Spearman's ρ are shown.

observation period. No associations were found between cQTVi groups and patient-specific parameters; therefore, this noninvasive parameter provides additional prognostic value for risk estimation. In the low cQTVi group, 36 patients received an ICD implantation (see Tab. 2). Since the risk of mortality is zero based on the survival analysis shown here, implantation implies additional risks for the patient.

4. Conclusion

In this study, we used two techniques to quantify VRV in VCG and on a single derivative basis. Our results show that Patients with non-ischemic cardiomyopathy and

a high cQTVi have a significant higher all-cause mortality than patients with low cQTVi. Hence, cQTVi could contribute to a treatment decision, as in this group the implantation of an ICD could be avoided, thus reducing costs and risk for the patient. In upcoming treatment decision processes, VRV parameters may provide additional information besides established ones such as LVEF or NYHA class to enable improvement of patient survival. However, it should be noted the small number of deaths in the study investigated, and therefore further studies are needed.

Further investigations will focus on the comparison of risk stratification based on VCG and two-dimensional single lead ECG [12] VRV parameters.

Acknowledgments

Dataset E-HOL-03-0401-017 used for this research was provided by the Telemetric and Holter ECG Warehouse of the University of Rochester (THEW), NY. This work was supported by the Australian Government Research Training Program (RTP) Scholarship and by grants of the Graduate Academy of the TU Dresden (Lab2Lab, Postdoc Starter Kit). We thank the ZIH of the TU Dresden for providing of computational power.

References

- [1] El-Hamad FJ, Bonabi SY, Müller A, Steger A, Schmidt G, Baumert M. Augmented Oscillations in QT Interval Duration Predict Mortality Post Myocardial Infarction Independent of Heart Rate. *Frontiers in Physiology* 2020;11:1442. ISSN 1664-042X.
- [2] Baumert M, Porta A, Vos MA, Malik M, Couderc JP, Laguna P, Piccirillo G, Smith GL, Tereshchenko LG, Volders PGA. QT Interval Variability in Body Surface ECG: Measurement, Physiological Basis, and Clinical Value: Position Statement and Consensus Guidance Endorsed by the European Heart Rhythm Association Jointly with the ESC Working Group on Cardiac Cellular Electrophysiology. *Europace* June 2016;18(6):925–944.
- [3] Schmidt M, Baumert M, Penzel T, Malberg H, Zaunseder S. Nocturnal Ventricular Repolarization Lability Predicts Cardiovascular Mortality in the Sleep Heart Health Study. *Am J Physiol Heart Circ Physiol* December 2018;316(3):H495–H505.
- [4] Baumert M, Starc V, Porta A. Conventional QT Variability Measurement vs. Template Matching Techniques: Comparison of Performance Using Simulated and Real ECG. *PLoS ONE* July 2012;7(7):e41920.
- [5] Schmidt M, Baumert M, Porta A, Malberg H, Zaunseder S. Two-Dimensional Warping for One-Dimensional Signals-Conceptual Framework and Application to ECG Processing. *IEEE Trans Signal Process* November 2014; 62(21):5577–5588.
- [6] Karisik F, Baumert M. Inhomogeneous Template Adaptation of Temporal Quasi-Periodic Three-Dimensional Signals. *IEEE Trans Signal Process* December 2019; 67(23):6067–6077. ISSN 1053-587X, 1941-0476.
- [7] Couderc JP, Zareba W, McNitt S, Maison-Blanche P, Moss AJ. Repolarization Variability in the Risk Stratification of MADIT II Patients. *Europace* September 2007;9(9):717–723.
- [8] Rashba EJ, Estes NM, Wang P, Schaechter A, Howard A, Zareba W, Couderc JP, Perkiomaki J, Levine J, Kadish A. Preserved Heart Rate Variability Identifies Low-risk Patients With Nonischemic Dilated Cardiomyopathy: Results From the Definite Trial. *Heart Rhythm* March 2006; 3(3):281–286. ISSN 1547-5271.
- [9] Schmidt M, Baumert M, Malberg H, Zaunseder S. Iterative Two-Dimensional Signal Warping—Towards a Generalized Approach for Adaption of One-Dimensional Signals. *Biomed Signal Process Control* May 2018;43:311–319.
- [10] Schmidt M, Baumert M, Malberg H, Zaunseder S. T Wave Amplitude Correction of QT Interval Variability for Improved Repolarization Lability Measurement. *Front Physiol* 2016;7.
- [11] Berger RD, Kasper EK, Baughman KL, Marban E, Calkins H, Tomaselli GF. Beat-to-beat QT Interval Variability - Novel Evidence for Repolarization Lability in Ischemic and Nonischemic Dilated Cardiomyopathy. *Circulation* September 1997;96(5):1557–1565.
- [12] Schmidt M, Dunker R, Malberg H, Zaunseder S. Quantification of Ventricular Repolarization Fluctuations in Patients With Myocardial Infarction. In *2020 Computing in Cardiology*. September 2020; 1–4.

Address for correspondence:

Martin Schmidt
 Institute of Biomedical Engineering, TU Dresden
 Fetscherstraße 29, 01307 Dresden, Germany
 martin.schmidt@tu-dresden.de