

Identifying the intracellular response in surface acoustic wave-stimulated wound healing [Abstract]

Kathrin Baumgartner, Paul Taeufer, Manuel Schleicher, Hanna Engelke, Manuel Brugger, Christoph Westerhausen

Angaben zur Veröffentlichung / Publication details:

Baumgartner, Kathrin, Paul Taeufer, Manuel Schleicher, Hanna Engelke, Manuel Brugger, and Christoph Westerhausen. 2023. "Identifying the intracellular response in surface acoustic wave-stimulated wound healing [Abstract]." *European Biophysics Journal* 52 (Supplement 1): S70. <https://doi.org/10.1007/s00249-023-01668-7>.



P-15**Identifying the intracellular response in surface acoustic wave-stimulated wound healing**

Kathrin Baumgartner^{1,2,3,4}, Paul Täufer², Manuel Schleicher^{1,2}, Hanna Engelke⁵, Manuel Brugger², Christoph Westerhausen^{1,2,3,6}

¹ Institute for Theoretical Medicine, Physiology, University of Augsburg, 86159 Augsburg, Germany, ² Institute of Physics, Experimental Physics I, University of Augsburg, 86159 Augsburg, Germany, ³ Center for NanoScience (CeNS), Ludwig-Maximilians-Universität Munich, 80799 Munich, Germany, ⁴ Hanns-Seidel-Stiftung e.V., 80636 Munich, Germany, ⁵ Institute for Pharmaceutical Sciences, Pharmaceutical Cell Biology, University of Graz, 8010 Graz, Austria, ⁶ Augsburg Center for Innovative Technologies (ACIT), 86159 Augsburg, Germany

Surface Acoustic Waves (SAW) are a powerful tool for biophysical applications like manipulation, sensing and stimulation of cellular properties or functions. We here stimulate epithelial wound healing assays on LiNbO₃ substrates with an interdigital transducer (IDT), generating travelling Rayleigh waves with a frequency of $f = 160$ MHz. By variation of the SAW intensities and pulse parameters, we reveal an optimal SAW stimulation intensity range between $P = 4$ and 8 mW, accelerating wound healing up to 235 %. The stimulation efficiency increases with increasing pulse width. Following the hypothesis that the cytoskeleton is responsible for faster cell migration in SAW-stimulated wound healing, we show that the orientation of actin fibres is significantly enhanced parallel to the SAW propagation direction. Ultimately, we aim to identify the contribution of the protein YAP and SPPL3 membrane proteases to the stimulation process. Thereby, we hope to unravel the underlying biophysical and biochemical mechanisms in SAW-stimulated wound healing and gain a better understanding of the impact of SAW on cellular processes.