

**85-Plat****STED Nanoscopy in Living Cells using Live Cell Compatible Markers****Katrin I. Willig<sup>1</sup>**, Birka Hein<sup>1</sup>, U. Valentin Nägerl<sup>2</sup>, Stefan W. Hell<sup>1</sup>.<sup>1</sup>Max-Planck-Institute for Biophysical Chemistry, Goettingen, Germany,<sup>2</sup>Max Planck Institute of Neurobiology, Mißingen-Martinsried, Germany.

We demonstrate far-field optical imaging with subdiffraction resolution in the interior of living mammalian cells and tissue by applying stimulated emission depletion (STED) microscopy.

Utilizing a yellow fluorescent protein (YFP) to image individual structural elements of the endoplasmatic reticulum (ER) and the tubular network (Fig.1) revealed a focal plane (x,y) resolution < 50 nm inside the living cell, corresponding to a 4-fold improvement over that of a confocal microscope and a 16-fold reduction in the focal spot cross-sectional area. Time lapse STED imaging of dendritic spines of YFP-positive hippocampal neurons in organotypic slices outperforms confocal microscopy in revealing important structural details. As an alternative to the fluorescent protein we employed a genetically encoded protein tag which can be stained *in vivo* with modified organic dyes. Using a rhodamine dye of high photostability enabled us to image structures in the living cell with a resolution of ~ 40 nm.

Thus nanoscale imaging of structures in the interior of living cells greatly expands the scope of light microscopy in cell biology.




Fig. 1. Citrine labeled microtubules.