



Universität Augsburg

Institut für
Mathematik

Dirk Blömker, Wei Wang

**Qualitative Properties of Local Random Invariant Manifolds for
SPDEs with Quadratic Nonlinearity**

Preprint Nr. 33/2008 — 01. Dezember 2008

Institut für Mathematik, Universitätsstraße, D-86135 Augsburg

<http://www.math.uni-augsburg.de/>

Impressum:

Herausgeber:

Institut für Mathematik

Universität Augsburg

86135 Augsburg

<http://www.math.uni-augsburg.de/forschung/preprint/>

ViSdP:

Dirk Blömker

Institut für Mathematik

Universität Augsburg

86135 Augsburg

Preprint: Sämtliche Rechte verbleiben den Autoren © 2008

Qualitative Properties of Local Random Invariant Manifolds for SPDEs with Quadratic Nonlinearity

Dirk Blömker
Institut für Mathematik
Universität Augsburg
86135 Augsburg
dirk.bloemker@math.uni-augsburg.de

Wei Wang
School of Mathematics
University of Adelaide
5005 Adelaide Australia
w.wang@adelaide.eu.au
&
Department of Mathematics
Nanjing University
210093 Nanjing China
wangweinju@yahoo.com.cn

November 27, 2008

Abstract

The qualitative properties of local random invariant manifolds for stochastic partial differential equations with quadratic nonlinearities and multiplicative noise is studied by a cut off technique. By a detail estimates on the Perron fixed point equation describing the local random invariant manifold, the structure near a bifurcation is given.

1 Introduction

Stochastic partial differential equations SPDEs arise as macroscopic mathematical models of complex systems under random influences. There have been rapid progresses in this area [17, 20, 14, 24, 4, 18]. More recently, SPDEs have been investigated in the context of random dynamical systems (RDS) [1]; see [8, 10, 11, 12, 13, 21, 15, 16, e.g.], among others.

Invariant manifolds are special invariant sets locally represented by graphs in state spaces (function spaces) where the solution process of the system is defined. A random invariant manifold provides a geometric structure to reduce stochastic dynamics. In fact the dynamics of the system is completely determined by that on the globally attracting invariant manifold. Nevertheless, due to the inherit non-autonomous nature of the SPDE, these manifolds move in time. So one can reduce the system to a lower dimensional system on the invariant manifold together with the dynamics of that manifold.

There are some results on the existence of random invariant manifold for a class of stochastic partial differential equations with Lipschitz nonlinearity, [15, 16, 19] etc. In [8] an estimate of dimension of the local invariant manifold is given, in order to study a pitchfork type bifurcation for stochastic reaction-diffusion equations with cubic nonlinearity.

In the recent work [23], by a detail study of the properties of flow, an invariant manifold reduction of a class stochastic partial differential equations with Lipschitz nonlinearity is obtained. The reduced equation is a stochastic differential equation defined on a finite dimensional invariant manifold.

There is a difficulty in reducing infinite dimensional stochastic systems with non-Lipschitz nonlinearity to a lower dimensional system on an invariant manifold. But it can be done near an equilibrium solution by introducing a cut-off function near the equilibrium solution such that the nonlinearity becomes globally Lipschitz in an approximate space. Then the approach in [23] could be followed and the result will be a local one. However, the invariant manifold is still moving in the infinite dimensional space. It is in general difficult to see clearly the dynamics of the reduces system even though it is essential finite dimensional.

At the same time an amplitude equation is an important tool to describe qualitatively the dynamics of stochastic systems near a change stability, which has been studied heavily for additive noise ([2, 3, 5, e.g.]) and for multiplicative noise (see [4, 6]). In contrast to random invariant manifolds, which move in time, the dynamics studied via amplitude equations approximates the dynamics on a given fixed vector space, and the essential dynamics is given by a stochastic ordinary differential equation on dominant modes. The drawback in this case is that the results hold only with high probability.

Results that hold almost surely are not possible to obtain.

In this paper we study the properties of the random invariant manifold locally. In fact we consider properties of local invariant manifolds for equations near a change of stability with quadratic nonlinearities and multiplicative noise. One of the simplest examples is a Burgers-type equation

$$\partial_t u = \partial_x^2 u + u + \nu u + \frac{1}{2} \partial_x u^2 + \sigma u \circ \dot{W} \quad (1)$$

subject to Dirichlet boundary conditions on $[0, \pi]$ with one-dimensional multiplicative noise of Stratonovic-type.

There are many examples of SPDEs near its first change of stability, where our results apply. For instance, we can consider the Kuramoto-Sivashinsky equation, a model from surface growth (cf. [7]), Navier-Stokes equations with an additional linear term, or the Rayleigh-Benard system near the convective instability.

For our main results we reduce the cut-off equation of (1) to a stochastic differential equation on an attracting random invariant manifold in a small cut-off ball. Our results extend [9], where the local dimension of a random invariant manifold near a fixed point is studied. Using the linearization in our case the local dimension is one, but here we show that with high probability the manifold is locally the graph of a quadratic function over a one-dimensional space (cf. Theorem 3).

Let us compare our results with amplitude equations. Typically, one assumes there the scaling $\sigma = \varepsilon$, $\nu = \nu_0 \varepsilon^2$ for some small ε with $|\nu_0| \leq 1$. In that scaling our Burgers-type equation reads

$$\partial_t u = \partial_x^2 u + u + \nu_0 \varepsilon^2 u + \frac{1}{2} \partial_x u^2 + \varepsilon u \circ \dot{W}. \quad (2)$$

In [4] it was shown that for solutions of size $\mathcal{O}(\varepsilon)$ with probability higher than $1 - C_p \varepsilon^p$ for all $p > 1$

$$u(t, x) \approx \varepsilon a(\varepsilon^2 t) \sin$$

on intervals of length $\mathcal{O}(\varepsilon^{-2})$, where

$$\partial_T a = \nu_0 a - \frac{1}{12} a^3 + a \circ \dot{W}$$

Our main results of this paper show for ν and σ small, but without scaling assumption, that there is a time dependent random Lipschitz map $\phi(t, \omega, \cdot) : \mathbb{R} \rightarrow \mathbb{R}$ describing locally the random invariant manifold such that near 0 the flow along the manifold is

$$\partial_t a = \nu a + a \phi(t, a) + \sigma a \circ \dot{W}. \quad (3)$$

Moreover, for a given t the probability that

$$\phi(t, a) \approx -\frac{1}{12}a^2$$

is larger than $1 - C \exp\{-1/\varepsilon\}$.

The rest of the paper is organized as follows. In Section 2, assumptions and main results are presented. Section 3 states the results on existence of random invariant manifolds using a cut-off technique. While Section 4 provides results on the local structure of the manifold.

2 Setting and Results

Consider the following abstract equation

$$\partial_t u = -Lu + \nu u + B(u, u) + \sigma u \circ \dot{W}, \quad u(0) = u_0 \in H \quad (4)$$

where W is a standard real valued Brownian motion, and the noise is in the Stratonovic sense. The real constants ν and σ describe the distance from the bifurcation and the noise strength, respectively. Assume H is a real separable Hilbert space with norm $\|\cdot\|$ and scalar product $\langle \cdot, \cdot \rangle$.

We make the following assumptions.

Assumption A₁ *Let L be a non-negative unbounded operator with compact resolvent in the Hilbert space H , and suppose that $\{e_k\}_{k=1}^\infty$ and $\{\lambda_k\}_{k=1}^\infty$ is an orthonormal basis of eigenfunctions and the corresponding ordered eigenvalues ($\lambda_{k+1} \geq \lambda_k$).*

Assumption A₂ *The kernel of L , $H_c := \text{span}\{e_1, \dots, e_N\}$, is finite dimensional, i.e. $\lambda_1 = \dots = \lambda_N = 0$ and $\lambda_* = \lambda_{N+1} > 0$.*

Denote by P_c the orthogonal projection from H to H_c . Furthermore, $P_s = I - P_c$.

Remark 1. *Note that the assumption that the eigenfunctions form an orthonormal set is for simplicity. We could relax this, assuming that P_c is a continuous projection commuting with L , which is no longer self-adjoint in that case.*

Let us now introduce the interpolation spaces H^α , $\alpha > 0$, as the domain of $L^{\alpha/2}$ endowed with scalar product $\langle u, v \rangle_\alpha = \langle u, (1+L)^\alpha v \rangle$ and corresponding norm $\|\cdot\|_\alpha$. Furthermore, we identify $H^{-\alpha}$ as the dual of H^α with respect to the inner product in H .

We make the following assumption on the nonlinearity.

Assumption A₃

For some $\alpha \in (0, 1)$ let $B : H \times H \rightarrow H^{-\alpha}$ be a bounded bilinear and symmetric operator, i.e. $B(u, \bar{u}) = B(\bar{u}, u)$ and there is a constant $C_B > 0$ such that $\|B(u, \bar{u})\|_{-\alpha} \leq C_B \|u\| \|\bar{u}\|$. Suppose that for $B(u) \triangleq B(u, u)$ we have

$$\langle B(u), u \rangle = 0 \quad \text{for all } u \in H^\alpha.$$

From bilinearity and boundedness, we immediately obtain a local Lipschitz condition for B . To be more precise, for all $R > 0$

$$\|B(u) - B(\bar{u})\|_{-\alpha} = \|B(u - \bar{u}, u + \bar{u})\|_{-\alpha} \leq 2RC_B \|u - \bar{u}\| \quad (5)$$

for all $u, \bar{u} \in H$ with $\|u\| \leq R$ and $\|\bar{u}\| \leq R$.

Random Dynamical Systems

Before giving our main result, we recall some basic theory of random dynamical systems. We will work on the canonical probability space $(\Omega_0, \mathcal{F}_0, \mathbb{P})$ where the sample space Ω_0 consists of the sample paths of $W(t)$. To be more precise W is the identity on Ω_0 , with

$$\Omega_0 = \{w \in C([0, \infty), \mathbb{R}) : w(0) = 0\},$$

and \mathbb{P} the Wiener measure. For more details see [1].

Let $\theta_t : (\Omega_0, \mathcal{F}_0, \mathbb{P}) \rightarrow (\Omega_0, \mathcal{F}_0, \mathbb{P})$ be a metric dynamical system (driven system), that is,

- $\theta_0 = id$,
- $\theta_t \theta_s = \theta_{t+s}$, for all $s, t \in \mathbb{R}$,
- the map $(t, \omega) \mapsto \theta_t \omega$ is measurable and $\theta_t \mathbb{P} = \mathbb{P}$ for all $t \in \mathbb{R}$.

On Ω_0 the map θ_t is the shift

$$\theta_t \omega(\cdot) = \omega(\cdot + t) - \omega(t), \quad t \in \mathbb{R}, \omega \in \Omega_0. \quad (6)$$

Definition 1. Let (\mathcal{X}, d) be a metric space with Borel σ -algebra \mathcal{B} , then a random dynamical system on (\mathcal{X}, d) over θ_t on $(\Omega_0, \mathcal{F}_0, \mathbb{P})$ is a measurable map

$$\begin{aligned} \varphi : \mathbb{R}^+ \times \Omega_0 \times \mathcal{X} &\rightarrow \mathcal{X} \\ (t, \omega, x) &\mapsto \varphi(t, \omega)x \end{aligned}$$

having the following cocycle property

$$\varphi(0, \omega)x = x, \quad \varphi(t, \theta_\tau \omega) \circ \varphi(\tau, \omega)x = \varphi(t + \tau, \omega)x$$

for $t, \tau \in \mathbb{R}^+$, $x \in \mathcal{X}$ and $\omega \in \Omega$.

A RDS φ is continuous or differentiable if $\varphi(t, \omega) : \mathcal{X} \rightarrow \mathcal{X}$ is continuous or differentiable (see [1] for more details on RDS).

It is well known that in order to show that (4) generates an RDS, one can rely on a random transformation to a random PDE. For this we introduce the following real-valued stationary process $z(t) = z(\theta_t \omega)$ on Ω_0 , where

$$z(\omega) = \int_{-\infty}^0 e^s \omega(s) ds. \quad (7)$$

Now the mapping $t \mapsto z(\theta_t \omega)$ is continuous and solves

$$dz + zdt = \sigma d\omega \quad \text{or} \quad z(\theta_t \omega) = z(\omega) - \int_0^t z(\theta_s \omega) ds + \sigma \omega(t). \quad (8)$$

Moreover,

$$\lim_{t \rightarrow \pm\infty} \frac{|z(\theta_t \omega)|}{|t|} = 0 \quad \text{and} \quad \lim_{t \rightarrow \pm\infty} \frac{1}{t} \int_0^t z(\theta_\tau \omega) d\tau = 0 \quad \text{for a.e. } \omega \in \Omega_0.$$

The above properties hold in a θ_t invariant set $\Omega \subset \Omega_0$ of full probability, see [15].

By the transformation

$$v(t) = e^{-z(t)} u(t) \quad (9)$$

Equation (4) becomes

$$\partial_t v = -Lv + zv + \nu v + e^z B(v, v), \quad (10)$$

which is an evolutionary equation with random stationary coefficient. Then for almost all $\omega \in \Omega$, by the same discussion for the wellposedness to deterministic evolutionary equation [22], for any $t_0 < T \in \mathbb{R}$ and $v_0 \in H$ there is a unique solution $v(t, \omega; t_0, v_0) \in C(t_0, T; H)$ of equation (10) with $v(t_0) = v_0$ and the map $v_0 \mapsto v(t, \omega; t_0, v_0)$ is continuous for all $t \geq t_0$. Then by the stationary transformation (9) we have the following result.

Theorem 1. *Under Assumptions $A_1 - A_3$, (4) generates a continuous RDS $\varphi(t, \omega)$ on H .*

For a continuous random dynamical system φ on \mathcal{X} given by Definition 1, we need the following notions to describe its dynamical behavior.

Definition 2. *A collection $M = M(\omega)_{\omega \in \Omega}$ of non-empty closed sets $M(\omega) \subset \mathcal{X}$, $\omega \in \Omega$ is a random set, if*

$$\omega \mapsto \inf_{y \in M(\omega)} d(x, y)$$

is a real valued random variable for any $x \in \mathcal{X}$.

Definition 3. A random set $\mathcal{B}(\omega)$ is called a tempered absorbing set for a random dynamical system φ if for any bounded set $K \subset \mathcal{X}$ there exists $t_K(\omega)$ such that $\forall t \geq t_K(\omega)$

$$\varphi(t, \theta_{-t}\omega, K) \subset \mathcal{B}(\omega).$$

and for all $\varepsilon > 0$

$$\lim_{t \rightarrow \infty} e^{-\varepsilon t} d(\mathcal{B}(\theta_{-t}\omega)) = 0, \text{ for a.e. } \omega \in \Omega,$$

where $d(\mathcal{B}) = \sup_{x \in \mathcal{B}} d(x, 0)$, with $0 \in \mathcal{X}$, is the diameter of \mathcal{B} , if $0 \in \mathcal{B}$.

For more details about random set we refer to [12].

Definition 4. A random set $M(\omega)$ is called a positive invariant set for a random dynamical system $\varphi(t, \omega, x)$ if

$$\varphi(t, \omega, M(\omega)) \subset M(\theta_t\omega), \text{ for } t \geq 0.$$

If $M(\omega) = \{x_1 + \psi(x_1, \omega) | x_1 \in \mathcal{X}_1\}$ is the graph of a random Lipschitz mapping

$$\psi(\cdot, \omega) : \mathcal{X}_1 \rightarrow \mathcal{X}_2$$

with $\mathcal{X} = \mathcal{X}_1 \oplus \mathcal{X}_2$, then $M(\omega)$ is called a Lipschitz invariant manifold of φ .

For further details about the random invariant manifold theory, see [15].

Definition 5. We say that (4) has a local random invariant manifold (LRIM) with radius R , if there is a random set $\mathcal{M}^R(\omega)$, which is defined by the graph of a continuous function, and for any $u_0 \in \mathcal{M}^R(\omega)$,

$$\varphi(t, \theta_{-t}\omega)u_0 \in \mathcal{M}^R(\omega)$$

for all $t \in (0, \tau_0(\omega))$ with

$$\tau_0(\omega) = \inf\{t \geq 0 : \varphi(t, \theta_{-t}\omega)u_0 \notin B_R(0)\}. \quad (11)$$

Main Results

We prove the following theorem in Section 3. See Theorems 4 and 5.

Theorem 2. (Existence) Under Assumptions $A_1 - A_3$, the random dynamical system $\varphi(t, \omega)$ defined by (4) has a LRIM $\mathcal{M}^R(\omega)$ for sufficiently small $R > 0$. This manifold is given as the graph of a random Lipschitz map $h(\omega, \cdot) : H_c \rightarrow H_s$:

$$\mathcal{M}^R(\omega) = \{(\xi, e^{z(\omega)}h(\omega, e^{-z(\omega)}\xi)) : \xi \in H_c\}.$$

Moreover, if $\lambda_* > 4\nu$, then the LRIM $\mathcal{M}^R(\omega)$ is locally exponentially attracting almost surely in the small ball $B_R(0)$. That is for any $\|u_0\| < R$

$$\text{dist}(\varphi(t, \theta_{-t}\omega)u_0, \mathcal{M}^R(\omega)) \leq 2RD(t, \theta_{-t}\omega)e^{-\lambda_*t}$$

for all $t < \tau_0(\omega)$ (see (11)) with $D(t, \omega)$ is a tempered increasing process, see (28).

We are interested in the property of solutions on the LRIM $\mathcal{M}^R(\omega)$ and the dynamics of $\mathcal{M}^R(\omega)$ itself near the first bifurcation $\nu = 0$.

It is proved that

Theorem 3. (Local Shape) *Let Assumptions $A_1 - A_3$ be true. Suppose $\sigma > 0$, $|\nu| < \sigma$ and $R \leq 1$, and let h be the fixed point given by Theorem 4.*

Then for $\sigma \rightarrow 0$ with probability larger than $1 - C \exp\{-1/\sqrt{\sigma}\}$ we have

$$\|e^{z(\omega)}h(e^{-z(\omega)}\xi) - L_s^{-1}B_s(\xi, \xi)\| \leq C(\|\xi\| + R^2 + \sqrt{\sigma}) \cdot \|\xi\|^2, \quad (12)$$

for all $\|\xi\| \leq \frac{1}{2}R$.

Let us finally comment on the flow on the manifold. Using Theorem 2, it is easy to describe the flow along the invariant manifold $\mathcal{M}(\omega)$ on a small ball of radius $R/2$ around 0 by the following equation

$$\partial_t u_c = \nu u_c + P_c B^{(R)}(u_c + e^z h(e^{-z} u_c)) + \sigma u_c \circ \dot{W}(t).$$

By Theorem 3 we can replace $e^z h(e^{-z} u_c)$ by

$$e^z L_s^{-1} B_s(e^{-z} u_c, e^{-z} u_c) + r(t), \quad (*)$$

where the probability of $r(t)$ being large is bounded by $C \exp\{-1/\sqrt{\sigma}\}$.

This finally yields an equation, where

$$\partial_t u_c = \nu u_c + P_c B^{(R)}(u_c + e^z h(e^{-z} u_c)) + \sigma u_c \circ \dot{W}(t).$$

This rederives an amplitude equation for the equation. Nevertheless, for a detailed analysis of the flow on \mathcal{M} , we would need bound on r , which are uniform in time. This will be postponed to future work.

To conclude the presentation of the main results, we remark that it should be straightforward to generalize the presented results to higher, but finite dimensional noise.

3 Existence of Local Invariant Manifold

In this section we construct a LRIM for system (4) for small $R > 0$ and prove its exponentially attracting property. For this we rely on a cut-off technique given by the following definition. The LRIM is as usual given by the RIM for the cut-off system.

Definition 6. (cut-off) *Let $\chi : H \rightarrow \mathbb{R}$ be a bounded smooth function such that $\chi(u) = 1$ if $\|u\| \leq 1$ and $\chi(u) = 0$ if $\|u\| \geq 2$. For any $R > 0$, we define $\chi_R(u) = \chi(u/R)$ for all $u \in H$.*

Given a radius $R > 0$ we define

$$B^{(R)}(u) = \chi_R(u)B(u, u).$$

Now by Assumption (A₃) for given $1 > \alpha > 0$, the operator $B^{(R)}$ is globally Lipschitz-continuous from space H to $H^{-\alpha}$ with Lipschitz constant

$$\text{Lip}_{H, H^{-\alpha}}(B^{(R)}) = L_R := 2RC_B. \quad (13)$$

Consider now the following cut-off system

$$\partial_t u = -Lu + \nu u + B^{(R)}(u) + \sigma u \circ \dot{W}, \quad u(0) = u_0. \quad (14)$$

As before by the transformation $v = ue^{-z}$, we have

$$v_t = -Lv + zv + \nu v + e^{-z}B^{(R)}(e^z v), \quad v(0) = u_0 e^{-z(0)}. \quad (15)$$

In order to obtain a random invariant manifold for random dynamical system $\varphi^R(t, \omega)$ defined by the above stochastic equation (14) we study the transformed equation (15). By the projections P_c and P_s equation (15) is split into

$$\begin{aligned} \partial_t v_c &= \nu v_c + z v_c + P_c e^{-z} B^{(R)}(e^z v), & v_c(0) &= P_c u_0 e^{-z(0)}, \\ \partial_t v_s &= -L_s v_s + \nu v_s + z v_s + P_s e^{-z} B^{(R)}(e^z v), & v_s(0) &= P_s u_0 e^{-z(0)}. \end{aligned}$$

We use the Lyapunov–Perron method on the following random space with random norm depending on $\omega \in \Omega$. The process z was defined in (8).

Definition 7. *For $-\nu < \eta < \lambda_* - \nu$ define the Banach space*

$$C_\eta^- = \left\{ v \in C((-\infty, 0], H) : \sup_{t \leq 0} \{ e^{\eta t - \int_0^t z(s) ds} \|v(t)\| \} < \infty \right\} \quad (16)$$

with norm

$$\|v\|_{C_\eta^-} = \sup_{t \leq 0} \{ e^{\eta t - \int_0^t z(s) ds} \|v(t)\| \} < \infty.$$

Definition 8. Define the nonlinear operator \mathcal{T} on C_η^- for given $\xi \in H_c$ and $\omega \in \Omega_0$ as

$$\begin{aligned} \mathcal{T}(v, \xi)(t) &= e^{\nu t + \int_0^t z(s) ds} \xi + \int_0^t e^{\nu(t-\tau) + \int_\tau^t z(r) dr} e^{-z(\tau)} P_c B^{(R)} (v(\tau) e^{z(\tau)}) d\tau \\ &\quad + \int_{-\infty}^t e^{(-L_s + \nu)(t-\tau) + \int_\tau^t z(r) dr} e^{-z(\tau)} P_s B^{(R)} ((v(\tau) e^{z(\tau)}) d\tau. \end{aligned} \quad (17)$$

By the Lipschitz property of $B^{(R)}$ (cf. (13)) it can be verified directly that for any $\xi \in H_c$ and $\omega \in \Omega_0$, $\mathcal{T}(\cdot, \xi) : C_\eta^- \rightarrow C_\eta^-$. Further, a short calculation shows that it is a Lipschitz continuous map. To be more precise, for any $v, \bar{v} \in C_\eta^-$

$$\begin{aligned} &[\mathcal{T}(v)(t) - \mathcal{T}(\bar{v})(t)] e^{-\int_0^t z(r) dr} \\ &= + \int_0^t e^{\nu(t-\tau) + \int_0^\tau z(r) dr - z(\tau)} [P_c B^{(R)} (v(\tau) e^{z(\tau)}) - P_c B^{(R)} (\bar{v}(\tau) e^{z(\tau)})] d\tau \\ &\quad + \int_{-\infty}^t e^{(-L_s + \nu)(t-\tau) - \int_0^\tau z(r) dr - z(\tau)} [P_s B^{(R)} ((v(\tau) e^{z(\tau)}) - P_s B^{(R)} ((\bar{v}(\tau) e^{z(\tau)})] d\tau. \end{aligned}$$

Now we use that for all $t > 0$

$$\|e^{Lt} u\| \leq M_{\alpha, \lambda} e^{-t\lambda} t^{-\alpha} \|u\|_{-\alpha} \quad (18)$$

for some $\lambda < \lambda_*$ sufficiently close, together with (13) and the estimate $\|P_c v\| \leq C_\alpha \|P_c\|_{-\alpha}$ for some constant C_α .

We obtain for $0 < \eta + \nu < \lambda$

$$\begin{aligned} \|\mathcal{T}(v) - \mathcal{T}(\bar{v})\|_{C_\eta^-} &\leq \int_0^t e^{(\nu+\eta)(t-\tau)} C_\alpha L_R \|v - \bar{v}\|_{C_\eta^-} d\tau \\ &\quad + \int_{-\infty}^t e^{(-\lambda+\nu+\eta)(t-\tau)} M_{\alpha, \lambda} L_R \|v - \bar{v}\|_{C_\eta^-} d\tau \\ &\leq L_R \left[\frac{C_\alpha}{\eta + \nu} + M_{\alpha, \lambda} \frac{\Gamma(1-\alpha)}{(\lambda - \eta - \nu)^{1-\alpha}} \right] \|v - \bar{v}\|_{C_\eta^-}. \end{aligned}$$

Note that our bound on $\text{Lip}(\mathcal{T})$ is actually independent of ξ and ω .

Now for all $\xi \in H_c$ and $\omega \in \Omega_0$ the operator $\mathcal{T} : C_\eta^- \rightarrow C_\eta^-$ is a contraction provided R is sufficiently small such that

$$L_R \left[\frac{C_\alpha}{\eta + \nu} + M_{\alpha, \lambda} \frac{\Gamma(1-\alpha)}{(\lambda - \eta - \nu)^{1-\alpha}} \right] < 1. \quad (19)$$

The celebrated theorem of Banach yields the following theorem.

Theorem 4. *Suppose $0 < \eta + \nu < \lambda_*$ and (19). Then the operator \mathcal{T} has a unique fixed point $v^* = v^*(\omega, \xi) \in C_\eta^-$.*

Define $h(\omega, \xi) = P_s v^*(0, \omega; \xi)$. Thus, by Definition 8

$$h(\omega, \xi) = \int_{-\infty}^0 e^{(L_s - \nu)\tau + \int_\tau^0 z(r)dr} e^{-z(\tau)} P_s B^{(R)}(v^*(\tau, \xi) e^{z(\tau)}) d\tau. \quad (20)$$

Then by the same discussion as in [16],

$$\mathcal{M}_{cut}^R(\omega) = \{(\xi, e^{z(\omega)} h(\omega, e^{-z(\omega)} \xi)) : \xi \in H_c\} \quad (21)$$

is a random invariant manifold for the random dynamical system $\varphi^R(t, \omega)$, which is the graph of $e^{z(\omega)} h(\omega, e^{-z(\omega)} \xi)$. Moreover, on $B(0, R)$

$$\mathcal{M}^R(\omega) = \mathcal{M}_{cut}^R(\omega) \quad (22)$$

defines a LRIM of the random dynamical system $\varphi(t, \omega)$.

Now we prove the attracting property of the random invariant manifold $\mathcal{M}_{cut}^R(\omega)$ for equation (14). We follow the approach of [23]. It is enough to prove the cone invariance property:

Lemma 1. *Fix $\delta > 0$ and define the cone*

$$\mathcal{K}_\delta = \{u \in H : \|u_s\| < \delta \|u_c\|\}.$$

Suppose that R is sufficiently small such that

$$\lambda_* \geq 2(1 + \frac{1}{\delta})^2 L_R^2 + 4(1 + \delta) L_R \quad (23)$$

and

$$\lambda_* > 4\nu + 2L_R^2(1 + \frac{1}{\delta})^2. \quad (24)$$

Let v, \bar{v} be two solutions of (15) with initial value $v_0 = u_0 e^{z(\omega)}$ and $\bar{v}_0 = \bar{u}_0 e^{z(\omega)}$.

If $v(t_0) - \bar{v}(t_0) \in \mathcal{K}_\delta$, then $v_c(t) - \bar{v}_c(t) \in \mathcal{K}_\delta$ for all $t \geq t_0$.

Moreover, if $v - \bar{v}$ is outside \mathcal{K}_δ at some time t_0 then

$$\begin{aligned} & \|v_s(t, \theta_{-t}\omega) - \bar{v}_s(t, \theta_{-t}\omega)\|^2 \\ & \leq \|u_0 - \bar{u}_0\|^2 \exp \left\{ -\frac{1}{2} \lambda_* t + z(\omega) + 2 \int_0^t z(\theta_{\tau-t}\omega) d\tau \right\} \end{aligned} \quad (25)$$

for all $t \in [0, t_0]$.

In the proof we will see that the 4ν is not optimal in (24), but for simplicity of proof, we keep the 4ν .

Proof. Define

$$p = v_c - \bar{v}_c \quad \text{and} \quad q = v_s - \bar{v}_s,$$

then

$$\begin{aligned} \partial_t p &= \nu p + zp + e^{-z} P_c B^{(R)}(ve^z) - e^{-z} P_c B^{(R)}(\bar{v}e^z) \\ \partial_t q &= -L_s q + \nu q + zq + e^{-z} P_s B^{(R)}(ve^z) - e^{-z} P_s B^{(R)}(\bar{v}e^z). \end{aligned}$$

By the property of operator L and the Lipschitz property of $B^{(R)}$ we obtain for some positive constant c_1 depending on λ_* and α

$$\frac{1}{2} \frac{d}{dt} \|p\|^2 \geq \nu \|p\|^2 + z \|p\|^2 - L_R \|p\|^2 - L_R \|p\| \|q\| \quad (26)$$

and

$$\frac{1}{2} \frac{d}{dt} \|q\|^2 \leq -\|q\|_1^2 + \nu \|q\|^2 + z \|q\|^2 + L_R \|q\| \|q\|_\alpha + L_R \|p\| \|q\|_\alpha. \quad (27)$$

Thus

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} (\|q\|^2 - \delta^2 \|p\|^2) \\ & \leq -\|q\|_1^2 + \nu \|q\|^2 + z \|q\|^2 + L_R \|q\| \|q\|_\alpha + L_R \|p\| \|q\|_\alpha \\ & \quad - \nu \delta^2 \|p\|^2 - z \delta^2 \|p\|^2 + \delta^2 L_R \|p\|^2 + \delta^2 L_R \|p\| \|q\|. \end{aligned}$$

Now suppose that $v - \bar{v} \in \mathcal{K}_\delta$ (i.e. $\delta \|p\| = \|q\|$) for some t . Then

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} (\|q\|^2 - \delta^2 \|p\|^2) \\ & \leq -\|q\|_1^2 + L_R \|q\| \|q\|_\alpha + L_R \|p\| \|q\|_\alpha + \delta^2 L_R \|p\|^2 + \delta^2 L_R \|p\| \|q\| \\ & \leq -\|q\|_1^2 + L_R (\|p\| + \|q\|) \|q\|_1 + \delta^2 L_R \|p\|^2 + \delta^2 L_R \|p\| \|q\| \\ & \leq -\|q\|_1^2 + L_R (1 + \frac{1}{\delta}) \|q\| \|q\|_1 + (1 + \delta) L_R \|q\|^2 \\ & \leq -\frac{1}{2} \|q\|_1^2 + \left(\frac{1}{2} L_R^2 (1 + \frac{1}{\delta})^2 + (1 + \delta) L_R \right) \|q\|^2 \end{aligned}$$

where we used Young inequality in the last step.

By (23) we obtain using Poincare inequality

$$\frac{d}{dt} (\|q\|^2 - \delta^2 \|p\|^2) \leq -\frac{1}{2} \lambda_* \|q\|^2,$$

which yields the desired cone invariance.

For the second claim consider now that if $p+q$ is outside the cone at time t_0 (i.e., $\|q(t_0)\| > \delta\|p(t_0)\|$). Then by the first result we have $\|q(t)\| > \delta\|p(t)\|$ for $t \in [0, t_0]$. Then by (27) we derive

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|q\|^2 &\leq -\|q\|_1^2 + (\nu + z)\|q\|^2 + L_R\|q\|\|q\|_1 + L_R\|p\|\|q\|_1 \\ &\leq -\|q\|_1^2 + (\nu + z)\|q\|^2 + L_R(1 + \frac{1}{\delta})\|q\|\|q\|_1 \\ &\leq -\frac{1}{2}\|q\|_1^2 + \left(\nu + z + \frac{1}{2}L_R^2(1 + \frac{1}{\delta})^2\right)\|q\|^2 \end{aligned}$$

By (24)

$$\frac{d}{dt} \|q(t, \omega)\|^2 \leq (-\frac{1}{2}\lambda_* + 2z(\theta_t\omega))\|q(t, \omega)\|^2 .$$

As the result holds for all ω , a comparison principle and replacing ω by $\theta_{-t}\omega$ yields

$$\|q(t, \theta_{-t}\omega)\|^2 \leq \|q(0, \omega)\|^2 \exp \left\{ -\frac{1}{2}\lambda_*t + 2 \int_0^t z(\theta_{\tau-t}\omega)ds \right\} .$$

Finally, $\|q(0, \omega)\| = \|v_s(0, \omega) - \bar{v}_s(0, \omega)\| \leq \|u_0 - \bar{u}_0\|e^{z(\omega)}$ yields the result. \square

To finish the proof of Theorem 2, we also need the following lemma.

Lemma 2. *For any given $T > 0$, the following initial value problem*

$$\begin{aligned} \dot{v}_c &= \nu v_c + z v_c + P_c e^{-z} B^{(R)}((v_c + v_s)e^z), \quad v_c(T) = \xi \in H_c \\ \dot{v}_s &= (-L_s + \nu + z)v_s + P_s e^{-z} B^{(R)}((v_c + v_s)e^z), \quad v_s(0) = h^s(v_c(0)) \end{aligned}$$

has a unique solution $(v_c(t, \theta_{-t}\omega), v_s(t, \theta_{-t}\omega)) \in C(0, T; H_c \times H_s)$ which lies on the manifold $\mathcal{M}_{cut}^R(\omega)$ a.a..

Proof. This proof is same as Lemma 3.3 and Lemma 3.8 of [15] except that one should be careful of the nonlinearity is not Lipschitz on space H in proving the contraction property of the solving operator. This step is same as the proof of contraction property of operator \mathcal{T} defined in (17). \square

Now following the contradicting discussion in [23] we verify the following attracting property of the random invariant manifold.

Theorem 5. *Assume (19), (23) and (24). For any solution $u(t, \theta_{-t}\omega)$ of cut-off system (14), there is one orbit $U(t, \theta_{-t}\omega)$ on $\mathcal{M}_{cut}^R(\omega)$ with $P_c U(t)$ solves the following equation*

$$\partial_t u_c = \nu u_c + P_c B^{(R)}(u_c + e^z h(e^{-z} u_c)) + \sigma u_c \circ \dot{W}(t)$$

such that

$$\|u(t, \theta_{-t}\omega) - U(t, \theta_{-t}\omega)\| \leq D(t, \theta_{-t}\omega) \|u(0, \omega) - U(0, \omega)\| e^{-\lambda_* t}$$

where $D(t, \theta_{-t}\omega)$ is a tempered increasing process defined by

$$D(t, \theta_{-t}\omega) = e^{z(\omega) + \int_{-t}^0 z(\theta_s \omega) ds}. \quad (28)$$

Then Theorem 2 is a direct result of Theorem 5. Moreover we have the following result.

Corollary 1. *There is a LRIM $\mathcal{M}(\omega)$ for system (4) in a small ball $B(0, R)$ with R and ν satisfy (19), (23), and (24). Moreover for $\|u_0\| < R$*

$$\text{dist}(\varphi(t, \theta_{-t}\omega)u_0, \mathcal{M}(\omega)) \leq 2RD(t, \theta_{-t}\omega)e^{-\lambda_* t}$$

for all $t \in [0, \tau_0]$ where τ_0 is defined by (11).

4 Shape of the Invariant Manifold

In this section we show that the LRIM \mathcal{M}_{cut}^R , see (21), which denotes the random invariant manifold of equation (14) is locally quadratic. That is the Lipschitz-function h (cf. Theorem 4) describing the manifold is quadratic near the fixed point 0. Suppose σ and ν are small and choose the cut-off R also small. Consider first the LRIM of the transformed equation (15). It is given by

$$\tilde{\mathcal{M}}_{cut}^R(\omega) = \text{Graph}(h) = \{(\xi, h(\omega, \xi)) : \xi \in H_c\},$$

where

$$h(\omega, \xi) = P_s v^*(0, \omega, \xi) = v_s^*(0, \omega, \xi)$$

with v^* the unique fixed point of the contraction \mathcal{T} in the space C_η^- (i.e., $v^* = \mathcal{T}(v^*)$, cf. Theorem 4). Thus for any $\xi \in H_c$ and $t \leq 0$

$$\begin{aligned} v^*(t) &= e^{\nu t + \int_0^t z(s) ds} \xi + \int_0^t e^{\nu(t-\tau) + \int_\tau^t z(r) dr} e^{-z(\tau)} B_c^{(R)}(e^{z(\tau)} v^*(\tau)) d\tau \\ &\quad + \int_{-\infty}^t e^{(-L_s + \nu)(t-\tau) + \int_\tau^t z(r) dr} e^{-z(\tau)} B_s^{(R)}(e^{z(\tau)} v^*(\tau)) d\tau. \end{aligned} \quad (29)$$

The local invariant manifold for the original equation (4) is given by (cf. (22))

$$\mathcal{M}^R(\omega) = \{(\xi, e^{z(\omega)} h(\omega, e^{-z(\omega)} \xi)) : \xi \in H_c\}.$$

In order to describe the shape of the LRIM, we first establish two bounds on v^* in C_η^- .

Lemma 3. *Under Assumptions $A_1 - A_3$ suppose $\lambda_* > \nu + \eta > 0$. Then there is a constant $C > 0$ such that for all $\xi \in H_c$*

$$\|v^*\|_{C_\eta^-} \leq C\|\xi\| \quad \text{and} \quad \|v_s^*\|_{C_\eta^-} \leq CR\|\xi\| .$$

Proof. For the first bound note that $\mathcal{T}(0) = e^{\nu t + \int_0^t z(s) ds} \xi$. Thus

$$\|\mathcal{T}(0)\|_{C_\eta^-} = \sup_{t \leq 0} e^{(\nu+\eta)t} \|\xi\| = \|\xi\| .$$

Now as \mathcal{T} is a contraction

$$\|v^*\|_{C_\eta^-} \leq \|\mathcal{T}(v^*) - \mathcal{T}(0)\|_{C_\eta^-} + \|\mathcal{T}(0)\|_{C_\eta^-} \leq \text{Lip}(\mathcal{T})\|v^*\|_{C_\eta^-} + \|\xi\| .$$

We obtain the first claim with $C = 1/(1 - \text{Lip}(\mathcal{T}))$.

For the second claim we bound (29) by using the semigroup estimate from (18) together with the fact that

$$\|B_s^{(R)}(ve^z)\|_{-\alpha} \leq CR\|v\|e^z , \quad (30)$$

for $v \in H$, which follows immediately from Definition 6 and Assumption A_3 . Thus

$$\begin{aligned} \|v_s^*(t)\| &= \|P_s v^*(t)\| & (31) \\ &= \left\| \int_{-\infty}^t e^{(-L_s + \nu)(t-\tau) + \int_\tau^t z(r) dr} e^{-z(\tau)} B_s^{(R)}(e^{z(\tau)} v^*(\tau)) d\tau \right\| \\ &\leq C \int_{-\infty}^t (t-\tau)^{-\alpha} e^{-(\lambda-\nu)(t-\tau) + \int_\tau^t z(r) dr} R \|v^*(\tau)\| d\tau \\ &\leq CR e^{\int_0^t z(r) dr} \int_{-\infty}^t (t-\tau)^{-\alpha} e^{-(\lambda-\nu)(t-\tau)} e^{-\eta\tau} d\tau \|v^*\|_{C_\eta^-} . \end{aligned}$$

Then using the first claim yields

$$\|v_s^*\|_{C_\eta^-} \leq CR\|\xi\| \int_{-\infty}^0 |\tau|^{-\alpha} e^{-(\lambda-\nu-\eta)|\tau|} d\tau .$$

Choosing $\lambda > \nu + \eta$ finishes the proof. \square

Now we prove the first reduction step, which removes the explicit dependence of h on v_s^* . Define for $t \leq 0$ the cut-off $\chi_R^*(t) = \chi_R(v^*(t)e^{z(t)})$ and the approximation

$$g_1^*(t) = \int_{-\infty}^t e^{(-L_s + \nu)(t-\tau) + \int_\tau^t z(r) dr} \chi_R^*(\tau) B_s(v_c^*(\tau), v_c^*(\tau)) e^{z(\tau)} d\tau . \quad (32)$$

Lemma 4. Under Assumptions $A_1 - A_3$ suppose $\lambda_* > \eta + \nu > 0$, then

$$\|v_s^* - g_1^*\|_{C_{\bar{\eta}}} \leq CR^2 \|\xi\| \quad \text{for all } \xi \in H_c.$$

Proof. We use

$$\begin{aligned} e^{-z} B_s^{(R)}(e^z v^*) &= \chi_R^*(t) B_s(v^*, v^*) e^z \\ &= \chi_R^*(t) [B_s(v_c^*, v_c^* e^z) + 2B_s(v_s^*, v_c^* e^z) + B_s(v_s^*, v_s^* e^z)]. \end{aligned}$$

Now from

$$\|\chi_R^* v_c^* e^z\| \leq R \quad \text{and} \quad \|\chi_R^* v_s^* e^z\| \leq R$$

we obtain

$$\|e^{-z} B_s^{(R)}(e^z v^*) - \chi_R^*(t) B_s(v_c^*, v_c^* e^z)\|_{-\alpha} \leq CR \|v_s^*\|.$$

Analogously to (31) we derive with $\lambda \in (\eta + \nu, \lambda_*)$ that

$$\begin{aligned} \|v_s^*(t) - g_1^*(t)\| &\leq C \int_{-\infty}^t (t - \tau)^{-\alpha} e^{-(\lambda - \nu)(t - \tau) + \int_{\tau}^t z(r) dr} R \|v_s^*(\tau)\| d\tau \\ &\leq C \int_{-\infty}^0 |\tau|^{-\alpha} e^{-(\lambda - \nu - \eta)|\tau|} d\tau \cdot e^{\int_0^t z(r) dr} e^{-\eta t} R \|v_s^*\|_{C_{\bar{\eta}}}. \end{aligned}$$

Thus using Lemma 3, the claim follows. \square

Recall that in order to bound h , we only need to consider $v_s^*(0)$. From (32)

$$g_1^*(0) = \int_{-\infty}^0 e^{L_s \tau} e^{-\nu \tau} e^{\int_{\tau}^0 z(r) dr} \chi_R^*(\tau) B_s(v_c^*(\tau), v_c^*(\tau)) e^{z(\tau)} d\tau,$$

where

$$v_c^*(\tau) e^{-\nu \tau} e^{\int_{\tau}^0 z(r) dr} - \xi = \int_0^{\tau} e^{-\nu r + \int_r^0 z(t) dt} e^{z(r)} \chi_R^*(r) B_c(v^*(r), v^*(r)) dr.$$

Using $B_c(v_c^*, v_c^*) = 0$ and thus

$$e^z \chi_R^* B_c(v^*, v^*) = 2B_c(v_s^*, e^z \chi_R^* v_c^*) + B_c(v_s^*, e^z \chi_R^* v_s^*)$$

we obtain for $\tau \leq 0$

$$\begin{aligned} &\left\| v_c^*(\tau) e^{-\nu \tau} e^{\int_{\tau}^0 z(r) dr} - \xi \right\| \\ &\leq \int_{\tau}^0 e^{-\nu r + \int_r^0 z(t) dt} R \|v_s^*(r)\| dr \\ &\leq R \int_{\tau}^0 e^{-(\nu + \eta)r} dr \|v_s^*\|_{C_{\bar{\eta}}} \leq CR^2 \|\xi\| e^{-(\nu + \eta)\tau}. \end{aligned} \tag{33}$$

Hence,

$$\left\| v_c^*(\tau) - \xi e^{\nu\tau} e^{\int_0^\tau z(r)dr} \right\| \leq CR^2 \|\xi\| e^{-\eta\tau} e^{\int_0^\tau z(r)dr}.$$

Thus for

$$g_2^* := \int_{-\infty}^0 e^{L_s\tau} \chi_R^*(\tau) B_s(\xi, v_c^*(\tau)) e^{z(\tau)} d\tau \quad (34)$$

we obtain using (33) and estimates analogous to (31)

$$\begin{aligned} \|g_1^*(0) - g_2^*\| &\leq C \int_{-\infty}^0 |\tau|^{-\alpha} e^{-\lambda|\tau|} R \cdot \|e^{-\nu\tau} e^{\int_\tau^0 z(r)dr} v_c^*(\tau) - \xi\| d\tau \\ &\leq CR^3 \|\xi\| \int_{-\infty}^0 |\tau|^{-\alpha} e^{-\lambda|\tau|} e^{-(\nu+\eta)\tau} d\tau. \end{aligned}$$

Thus together with Lemma 4 we have

Theorem 6. *Under Assumptions $A_1 - A_3$ suppose $R \leq 1$, and $\lambda_* > \nu + \eta > 0$. Then*

$$\|h(\xi) - g_2^*\| \leq CR^2 \|\xi\| \quad \text{for all } \xi \in H_c.$$

This theorem is how far we can get without ω -dependent deterministic bounds. In the following we will rely on the following lemma (cf. (8)).

Lemma 5. *There is a random variable $K(\omega)$ such that $K(\omega) - 1$ has a standard exponential distribution and for all $t \leq 0$*

$$\int_0^t z(r)dr + z(t) = z(0) + \sigma\omega(t) \leq \sigma(K(\omega) + |t|).$$

Proof. Define

$$\tilde{K}(\omega) = \sup_{s \leq 0} \{\omega(s) + s\}. \quad (35)$$

For example from [25] we know that $2\tilde{K}$ has a standard exponential distribution. Obviously,

$$\sigma\omega(t) \leq \sigma(\tilde{K}(\omega) + |t|) \quad \text{for all } t \leq 0.$$

Moreover, by definition

$$\begin{aligned} z(0) &= \sigma \int_{-\infty}^0 e^s \omega(s) ds \\ &\leq \sigma \int_{-\infty}^0 e^s ds \tilde{K}(\omega) - \sigma \int_{-\infty}^0 s e^s ds = \sigma(\tilde{K}(\omega) + 1). \end{aligned} \quad (36)$$

□

Remark 2. For $s < 0$ we have

$$|\omega(s)| \leq \max\{\omega(s), -\omega(s)\} \leq K^\pm(\omega) + |s|,$$

where

$$K^\pm(\omega) = K(\omega) + K(-\omega)$$

and $K(-\omega)$ has the same law as $K(\omega)$. Furthermore, for $|z(0)|$ a similar estimate is true.

Define

$$g_3^* := \int_{-\infty}^0 e^{L_s \tau} \chi_R^*(\tau) e^{z(\tau)} e^{\nu \tau} e^{\int_0^\tau z(r) dr} d\tau B_s(\xi, \xi). \quad (37)$$

Using (33) yields

$$\begin{aligned} \|g_2^* - g_3^*\| &\leq C \int_{-\infty}^0 |\tau|^{-\alpha} e^{-\lambda|\tau|} e^{z(\tau)} \|\xi\| \cdot \|v_c^*(\tau) - \xi e^{\nu \tau} e^{\int_0^\tau z(r) dr}\| d\tau \\ &\leq CR^2 \|\xi\|^2 \int_{-\infty}^0 |\tau|^{-\alpha} e^{-\lambda|\tau|} e^{\int_0^\tau z(r) dr} e^{z(\tau)} e^{-\eta \tau} d\tau. \end{aligned}$$

Then by Lemma 5 we obtain for $\sigma < \frac{1}{2}(\lambda - \eta)$

$$\begin{aligned} \|g_2^* - g_3^*\| &\leq CR^2 \|\xi\|^2 \int_{-\infty}^0 |\tau|^{-\alpha} e^{(\eta - \lambda + 2\sigma)|\tau|} e^{2\sigma K(\omega)} d\tau \\ &\leq CR^2 \|\xi\|^2 e^{2\sigma K(\omega)}. \end{aligned} \quad (38)$$

Now we eliminate the cut-off. Suppose $\|\xi\| e^{z(0)} \leq R$, then we can use the smoothness of χ_R (mean value theorem) to obtain

$$\begin{aligned} |\chi_R^*(\tau) - 1| &= |\chi_R(v^*(\tau) e^{z(\tau)}) - \chi_R(\xi e^{z(0)})| \\ &\leq \frac{C}{R} \|v_s^*(\tau) e^{z(\tau)}\| + \frac{C}{R} \|v_c^*(\tau) e^{z(\tau)} - \xi e^{z(0)}\| \\ &\leq C(1 + R) \|\xi\| e^{-\eta \tau + z(\tau) + \int_0^\tau z(r) dr} + \frac{C}{R} \|\xi\| \cdot |e^{z(0)} - e^{\nu \tau + \int_0^\tau z(r) dr}| \end{aligned}$$

where we used Lemma 3 and (33).

Now from Lemma 5, and the fact that $z(t) + \int_0^t z(r) dr = z(0) + \sigma \omega(t)$ we derive

$$|\chi_R^*(\tau) - 1| \leq C(1 + R) \|\xi\| e^{-\eta \tau + \sigma |\tau| + \sigma K(\omega)} + C |1 - e^{\nu \tau + \sigma \omega(\tau)}| e^{z(0)}.$$

Thus

$$|\chi_R^*(\tau) - 1| \leq C(\|\xi\|(1 + R) + \sigma K_2(\omega)) e^{-\eta \tau} e^{\sigma |\tau| + \sigma K(\omega)}, \quad (39)$$

where we define

$$K_2(\omega) := \sup_{\tau \leq 0} \left| \frac{1 - e^{\nu\tau + \sigma\omega(\tau)}}{\sigma e^{\eta|\tau|}} \right|. \quad (40)$$

For this random constant K_2 we obtain the following relation to $K(\omega)$. Note that this result is by far not optimal, but it is rather simple to derive.

Lemma 6. *Suppose $|\nu| + |\sigma| < \frac{\eta}{2}$ and $|\nu| \leq |\sigma|$, then there is a constant depending only on η such that*

$$K_2(\omega) \leq C e^{|\sigma|K^\pm(\omega)} (1 + K^\pm(\omega)).$$

Proof. We use the rather crude estimate $|1 - e^x| \leq |x|e^{|x|}$ for all $x \in \mathbb{R}$. Recall Remark 2 to obtain

$$\begin{aligned} K_2(\omega) &\leq \sup_{\tau \leq 0} \frac{|\nu||\tau| + |\sigma||\omega|}{|\sigma|} e^{|\nu||\tau| + |\sigma\omega(\tau)| - \eta|\tau|} \\ &\leq \sup_{\tau \leq 0} \frac{(|\nu| + |\sigma|)|\tau| + |\sigma|}{|\sigma|K^\pm(\omega)} e^{|\sigma|K^\pm(\omega)} e^{|\tau|(|\nu| + |\sigma| - \eta)} \\ &\leq 2e^{|\sigma|K^\pm(\omega)} \sup_{\tau \leq 0} (|\tau| + K^\pm(\omega)) e^{-\frac{1}{2}|\tau|\eta}. \end{aligned}$$

□

Now we proceed to bound g_3^*

$$\begin{aligned} &\left\| g_3^* - \int_{-\infty}^0 e^{L_s\tau} e^{z(\tau)} e^{\nu\tau} e^{\int_0^\tau z(r)dr} d\tau B_s(\xi, \xi) \right\| \\ &\leq C \int_{-\infty}^0 |\tau|^{-\alpha} e^{(-\lambda + 2\sigma)|\tau|} e^{\sigma K(\omega)} |1 - \chi_R^*(\tau)| d\tau \|\xi\|^2 \\ &\stackrel{(39)}{\leq} C \int_{-\infty}^0 |\tau|^{-\alpha} e^{(-\lambda + \eta + 2\sigma)|\tau|} e^{2\sigma K(\omega)} d\tau (\|\xi\| (1 + R) + \sigma K_2(\omega)) \|\xi\|^2 \\ &\leq C e^{2\sigma K(\omega)} (\|\xi\| + \sigma K_2(\omega)) \|\xi\|^2 \end{aligned} \quad (41)$$

for $R \leq 1$ and $\eta - \lambda + 2\sigma < 0$.

For the final step note that

$$\int_{-\infty}^0 e^{L_s\tau} e^{z(\tau)} e^{\nu\tau} e^{\int_0^\tau z(r)dr} d\tau = e^{z(0)} \int_{-\infty}^0 e^{L_s\tau} e^{\nu\tau} e^{\sigma\omega(\tau)} d\tau.$$

We finally bound

$$\begin{aligned}
& \left\| \int_{-\infty}^0 e^{L_s \tau} e^{\nu \tau} e^{\sigma \omega(\tau)} d\tau B_s(\xi, \xi) - L_s^{-1} B_s(\xi, \xi) \right\| \\
&= \left\| \int_{-\infty}^0 e^{L_s \tau} (1 - e^{\nu \tau} e^{\sigma \omega(\tau)}) d\tau B_s(\xi, \xi) \right\| \\
&\leq C \int_{-\infty}^0 |\tau|^{-\alpha} e^{(-\lambda + \eta)|\tau|} \sigma K_2(\omega) d\tau \|\xi\|^2 \\
&\leq C \sigma K_2(\omega) \|\xi\|^2
\end{aligned} \tag{42}$$

for $\lambda > \eta$.

Combining (38), (41), and (42), we proved the following theorem:

Theorem 7. *Let Assumptions $A_1 - A_3$ be true. Suppose $\lambda_* > \eta + 2\sigma$, $\lambda_* > \nu + \eta$, $\sigma > 0$, and $\|\xi\| e^{z(0)} \leq R \leq 1$, and let h be the fixed point given by Theorem 4. Then*

$$\|h(\xi) - e^{z(0)} L_s^{-1} B_s(\xi, \xi)\| \leq C(e^{2\sigma K(\omega)} + 1) \cdot (\|\xi\| + R^2 + \sigma K_2(\omega)) \cdot \|\xi\|^2,$$

where K and K_2 are defined in (35) and (40).

This is the final main Theorem of this section. The main Theorem 3 is now a simple corollary.

Proof of Theorem 3. Define

$$\Omega_K = \left\{ \omega \in \Omega_0 : K^\pm(\omega) > 1/\sqrt{\sigma} \right\}.$$

By Lemma 5 this set has probability less than $C \exp\{-1/\sqrt{\sigma}\}$. Moreover, on the complement Ω_K^c we have

$$K \leq \frac{1}{\sqrt{\sigma}} \quad \text{and} \quad K_2 \leq \frac{1}{\sqrt{\sigma}}$$

and (12) follows from Theorem 7, as similar to Lemma 6 we also have on Ω_K^c

$$e^{z(0)} \|\xi\| \leq e^{C\sqrt{\sigma}} \|\xi\| \leq R.$$

□

Acknowledgements

Supported by Bosch-foundation Nr. 32.5.8003.0010.0 (“Angewandte Mathematik”) and NSFC No. 10701072. Part of the work was done during a stay of W. Wang at the University of Augsburg.

References

- [1] L. Arnold, *Random Dynamical Systems*, Springer-Verlag, New York/Berlin, 1998.
- [2] D. Blömker, Amplitude equations for locally cubic non-autonomous nonlinearities, *SIAM J. on Appl. Dyn. Sys.* **2(2)**,(2003), 464–486.
- [3] D. Blömker & M. Hairer, Multiscale expansion of invariant measures for SPDEs, *Commun. in Math. Phys.* **251(3)**,(2004), 515–555.
- [4] D. Blömker, *Amplitude Equations for Stochastic Partial Differential Equations*, Vol. 3 of Interdisciplinary Mathematical Sciences, World Scientific Publishing, 2007.
- [5] D. Blömker, M. Hairer, & G. Pavliotis. Multiscale analysis for SPDEs with quadratic nonlinearities. *Nonlinearity*, **20**, (2007) 1721–1744.
- [6] D. Blömker, M. Hairer, & G. Pavliotis. Stochastic Swift-Hohenberg equation near a change of stability. pp. 27–37 in Proceedings of Equadiff 11, Proceedings of minisymposia and contributed talks, July 25–29, 2005, Bratislava, (Ed. M. Fila, A. Handlovicova, K. Mikula, M. Medved, P. Quittner und D. Sevcovic), 2007.
- [7] D. Blömker, F. Flandoli & M. Romito, Markovianity and ergodicity for a surface growth PDE, to appear in *Annals of Probability*
- [8] T. Caraballo, J. A. Langa & J.C.Robinson, Stability and random attractors for a reaction-diffusion equation with multiplicative noise, *Disc. and Cont. Dyn. Sys.* **6** (2000), 875–892.
- [9] T. Caraballo, J. A. Langa & J.C.Robinson, A stochastic pitchfork bifurcation in a reaction-diffusion equation, *Proc. Roy. Soc. London A* **457** (2001), 2041–2061.
- [10] I. D. Chueshov, J. Duan & B. Schmalfuss, Determining functionals for random partial differential equations, *NoDEA* **10** (2003), 431–454.
- [11] I. D.Chueshov, Approximate inertial manifolds of exponential order for semilinear parabolic equations subjected to additive white noise, *J. Dyn. and Diff. Equ.* **7(4)**(1995), 549–566.
- [12] H. Crauel & F. Flandoli, Attractor for random dynamical systems, *Prob. Theory and Relat. Fields.* **100** (1994), 365–393.

- [13] H. Crauel, A. Debussche & F. Flandoli, Random attractors, *J. Dyn. Diff. Equ.* **9(2)** (1997), 307–341.
- [14] G. Da Prato & J. Zabczyk, *Stochastic Equations in Infinite Dimensions*, Cambridge University Press, 1992.
- [15] J. Duan, K. Lu & B. Schmalfuß. Invariant manifolds for stochastic partial differential equations, *Ann. Prob.*, **31(4)** (2003), 2109–2135.
- [16] J. Duan, K. Lu & B. Schmalfuß. Smooth stable and unstable manifolds for stochastic partial differential equations, *J. Dyn. and Diff. Equ.*, **16(4)** (2004), 949–972.
- [17] J. Garcia-Ojalvo & J. M. Sancho, *Noise in Spatially Extended Systems*. Springer-Verlag, 1999.
- [18] Z. Huang & J. Yan, *Introduction to Infinite Dimensional Stochastic Analysis*. Science Press/Kluwer Academic Pub., Beijing/New York, 1997.
- [19] K. Lu & B. Schmalfuß, Invariant manifolds for stochastic wave equations, *J. Diff. Equa.* **236(2)** (2007), 460–492.
- [20] B. L. Rozovskii, *Stochastic Evolution Equations*. Kluwer Academic Publishers, Boston, 1990.
- [21] B. Schmalfuß, Backward cocycles and attractors of stochastic differential equations, In *International Seminar on Applied Mathematics-Nonlinear Dynamics: Attractor Approximation and Global Behaviour*, V. Reitmann, T. Riedrich & N. Kokscha (Eds.), 1992, 185–192.
- [22] R. Temam, *Infinite-dimensional dynamical systems in mechanics and physics*. Springer-Verlag, New York, second edition, 1997.
- [23] W. Wang & J. Duan, A dynamical approximation for stochastic partial differential equations, *J. Math. Phys.* **48**, (2007), 102701–14.
- [24] E. Waymire & J. Duan (Eds.), *Probability and Partial Differential Equations in Modern Applied Mathematics*. IMA Volume 140, Springer-Verlag, New York, 2005.
- [25] M. Yor, Some remarks about the joint law of Brownian motion and its supremum. Azema, J. (ed.) et al., *Seminaire de probabilites XXXI*. Berlin: Springer. *Lect. Notes Math.* **1655**, (1997), 306–314.