

Separability in domain semirings

Dexter Kozen, Bernhard Möller

Angaben zur Veröffentlichung / Publication details:

Kozen, Dexter, and Bernhard Möller. 2004. "Separability in domain semirings."
Augsburg: Institut für Informatik, Universität Augsburg.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under the following conditions:


Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publizieren/>



UNIVERSITÄT AUGSBURG




Separability in Domain Semirings

Dexter Kozen Bernhard Möller

Report 2004-16

Dezember 2004



INSTITUT FÜR INFORMATIK

D-86135 AUGSBURG

Copyright © Dexter Kozen Bernhard Möller
Institut für Informatik
Universität Augsburg
D-86135 Augsburg, Germany
<http://www.Informatik.Uni-Augsburg.DE>
— all rights reserved —

Separability in Domain Semirings

Dexter Kozen¹ Bernhard Möller²

¹ Department of Computer Science, Cornell University,
Upson Hall, Ithaca, NY 14853, USA

² Institut für Informatik, Universität Augsburg,
D-86135 Augsburg, Germany

Abstract. First, we show with two examples that in test semirings with an incomplete test algebra a domain operation may or may not exist. Second, we show that two notions of separability in test semirings coincide, respectively, with locality of composition and with extensionality of the diamond operators in domain semirings. We conclude with a brief comparison of dynamic algebras and modal Kleene algebras.

1 Basic Definitions

A *test semiring* [6] is a two-sorted structure $(K, \mathbf{test}(K))$, where K is an idempotent semiring and $\mathbf{test}(K) \subseteq K$ is a Boolean algebra embedded into K such that the operations of $\mathbf{test}(K)$ coincide with the restricted operations of K . In particular, $p \leq 1$ for all $p \in \mathbf{test}(K)$. In general, $\mathbf{test}(K)$ may be a proper subset of all elements below 1.

A *domain semiring* [1] is a structure (K, \ulcorner) , where K is an idempotent semiring such that the *domain operation* $\ulcorner: K \rightarrow \mathbf{test}(K)$ satisfies, for all $a, b \in K$ and $p \in \mathbf{test}(K)$,

$$a \leq \ulcorner a a, \tag{D1}$$

$$\ulcorner(pa) \leq p. \tag{D2}$$

The conjunction of (D1) and (D2) is equivalent to each of

$$\ulcorner a \leq p \Leftrightarrow a = pa, \tag{LLP}$$

$$\ulcorner a \leq p \Leftrightarrow \neg pa = 0, \tag{GLA}$$

which constitute elimination laws for domain. (LLP) says that $\ulcorner a$ is the least left preserver of a . (GLA) says that $\neg \ulcorner a$ is the greatest left

annihilator of a . Both properties obviously characterize domain in set-theoretic relations. An important consequence of the axioms is strictness of the domain operation:

$$a = 0 \Leftrightarrow \ulcorner a = 0 \text{ .} \tag{1}$$

Moreover, we have the following useful proof principle.

Lemma 1.1 (Indirect Domain Inequality)

$$\ulcorner a \leq \ulcorner b \Leftrightarrow (\forall p. pb = 0 \Rightarrow pa = 0) \text{ .}$$

Proof. $\ulcorner a \leq \ulcorner b$

$$\Leftrightarrow \{ \text{contraposition} \}$$

$$\neg \ulcorner b \leq \neg \ulcorner a$$

$$\Leftrightarrow \{ \text{indirect inequality} \}$$

$$\forall p. p \leq \neg \ulcorner b \Rightarrow p \leq \neg \ulcorner a$$

$$\Leftrightarrow \{ \text{by (GLA)} \}$$

$$\forall p. pb = 0 \Rightarrow pa = 0 \text{ .}$$

□

A domain semiring is called *modal* if it additionally satisfies

$$\ulcorner(a \ulcorner b) \leq \ulcorner(ab) \text{ .} \tag{DLoc}$$

In a modal semiring, domain is *local*:

$$\ulcorner(ab) = \ulcorner(a \ulcorner b) \text{ .}$$

Without (DLoc), only the inequality $\ulcorner(ab) \leq \ulcorner(a \ulcorner b)$ holds. The additional axiom (DLoc) guarantees that the domain of ab is independent from the inner structure and the “far end” of b ; information about the domain of b in interaction with a suffices. By this property the multimodal operators that can be defined in terms of domain become well-behaved under composition.

A *codomain* operation \lrcorner can easily be defined as a domain operation in the opposite semiring, where, as usual in algebra, opposition just

swaps the order of multiplication. We call a semiring K with local domain and codomain a *modal semiring*.

A *Kleene algebra* [5] is a structure $(K, *)$ such that K is an idempotent semiring and the *star* $*$ satisfies, for $a, b, c \in K$, the *unfold* and *induction laws*

$$\begin{aligned} 1 + aa^* &\leq a^*, & 1 + a^*a &\leq a^*, \\ b + ac \leq c &\Rightarrow a^*b \leq c, & b + ca \leq c &\Rightarrow ba^* \leq c. \end{aligned}$$

A Kleene algebra is **-continuous* if it satisfies

$$ab^*c = \sqcup \{ab^n c \mid n \in \mathbb{N}\}.$$

A *Kleene algebra with tests* (KAT) [6] is a test semiring $(K, \text{test}(K))$ such that K is a Kleene algebra. Finally, a *Kleene algebra with domain* (KAD) [1] is a Kleene algebra with tests that also provides a domain operation.

Below we examine in how far the existence of a domain operation depends on completeness of the test algebra and discuss the relation of the domain axioms to separability of elements by tests.

2 Completeness of the Test Algebra

Consider a test semiring $(K, \text{test}(K), +, 0, \cdot, 1)$. In the sequel, a, b, \dots range over K and p, q, \dots range over $\text{test}(K)$.

Define, for $a \in K$, the sets $\text{LP}(a), \text{LA}(a) \subseteq \text{test}(K)$ of *left-preservers* and *left-annihilators* of a by

$$\text{LP}(a) \stackrel{\text{def}}{=} \{p \mid a = pa\} \qquad \text{LA}(a) \stackrel{\text{def}}{=} \{p \mid pa = 0\}$$

Lemma 2.1 *1. $\text{LP}(a)$ is closed under upper bounds of arbitrary non-empty subsets and under infima of finite subsets. Hence, $1 \in \text{LP}(a)$. In particular, $\text{LP}(a)$ is a filter. Moreover,*

$$\text{LP}(a) \cap \text{LP}(b) \subseteq \text{LP}(a + b).$$

2. $\text{LA}(a)$ is closed under lower bounds of arbitrary non-empty subsets and under suprema of finite subsets. Hence, $0 \in \text{LA}(a)$. In particular, $\text{LA}(a)$ is an ideal. Moreover,

$$\text{LA}(a) \cap \text{LA}(b) = \text{LA}(a + b).$$

3. If K is a domain semiring then $\lceil a$ is the least element of $\text{LP}(a)$ and hence the infimum of $\text{LP}(a)$, whereas $\neg\lceil a$ is the greatest element of $\text{LA}(a)$ and hence the supremum of $\text{LA}(a)$.

This implies that a test semiring can be a domain semiring only if its test algebra has sufficiently many infima and suprema. Since infima are unique, we have

Corollary 2.2 *Every test semiring with a finite test algebra can be extended uniquely to a domain semiring.*

We now give an example of a test semiring that lacks infima and suprema to such an extent that it cannot be extended to a domain semiring.

Example 2.3 Consider an infinite set M and the test semiring $K = (\mathcal{P}(M), \mathcal{Q}, \cup, \emptyset, \cap, M)$ where

$$\mathcal{Q} \stackrel{\text{def}}{=} \{N \subseteq M \mid N \text{ finite or cofinite}\}.$$

Then $\mathcal{Q} = \text{test}(K)$ is a Boolean algebra that is incomplete.

For $A \in K$ and $Q \in \mathcal{Q}$ we have

$$Q \in \text{LP}(A) \Leftrightarrow A \subseteq Q, \quad Q \in \text{LA}(A) \Leftrightarrow Q \subseteq \bar{A}.$$

Let A now be infinite and coinfinite. Suppose $\text{LP}(A)$ has an infimum $Q \in \mathcal{Q}$. Then $A \subseteq Q$ but $A \neq Q$, since $A \notin \mathcal{Q}$. Let $x \in Q - A$. Now, Q must be cofinite, since $A \subseteq Q$. But then also $Q - \{x\}$ is cofinite with $Q - \{x\} \in \text{LP}(A)$ and $Q - \{x\} \subsetneq Q$, a contradiction. Hence

$\text{LP}(A)$ does not have an infimum in \mathcal{Q} . Symmetrically, $\text{LA}(\bar{A})$ does not have a supremum in \mathcal{Q} .

Thus, $\lceil A$ is undefined in K for infinite and coinfinite A . \square

Hence the question arises whether the test algebra needs to be complete to admit a domain operation. This is not the case, as the following example shows.

Example 2.4 Consider an infinite set M . Choose a finite subset $N \subseteq M$ and set

$$K \stackrel{\text{def}}{=} \{R \subseteq M \times M \mid R \cap I \text{ finite or cofinite} \wedge R - I \subseteq N \times N\}$$

where $I \stackrel{\text{def}}{=} \{(x, x) \mid x \in M\}$. Moreover, set

$$\text{test}(K) \stackrel{\text{def}}{=} \{R \mid R \subseteq I . R \text{ finite or cofinite}\}.$$

Then K is closed under \cup and $;$, so that $(K, \text{test}(K), \cup, \emptyset, ;, I)$ is a test semiring which under the standard definition

$$\ulcorner R \stackrel{\text{def}}{=} \{(x, x) \mid \exists y . (x, y) \in R\}$$

becomes a domain semiring. But $\text{test}(K)$ is incomplete.

K can even be extended into a KAD, since, for $R \in K$,

$$\begin{aligned} R^* &= ((R \cap I) \cup (R - I))^* = (R \cap I)^* ; ((R - I) ; (R \cap I)^*)^* \\ &= (R - I)^* = I \cup (R - I)^+ \in K . \end{aligned}$$

The third step uses that $R \cap I \subseteq I$ implies $(R \cap I)^* = I$. □

3 Separability, Locality and Extensionality

In [7] the following two separability properties are studied:

$$ab = 0 \Rightarrow \exists q . a = aq \wedge b = \neg qb , \quad (\text{Sep1})$$

$$a \not\leq b \Rightarrow \exists p, q . paq \neq 0 \wedge pbq = 0 . \quad (\text{Sep2})$$

Actually, both properties can be strengthened to equivalences. For (Sep1), from the assumption $a = aq \wedge b = \neg qb$ we get

$$ab = aq\neg qb = a0b = 0 .$$

So we will use

$$ab = 0 \Leftrightarrow \exists q . a = aq \wedge b = \neg qb . \quad (\text{Sep1}')$$

For (Sep2) we first note that, by contraposition, it is equivalent to

$$(\forall p, q . pbq = 0 \Rightarrow paq = 0) \Rightarrow a \leq b . \quad (\text{Sep2}')$$

Now from the assumptions $a \leq b$ and $pbq = 0$ we obtain $paq = 0$ by isotonicity of multiplication.

It turns out that, in a domain semiring, (Sep1) is equivalent to (DLoc) and (Sep2) is equivalent to

$$(\forall q . |a\rangle q \leq |b\rangle q) \Rightarrow a \leq b , \quad (\text{IDext})$$

where the forward modal diamond operator $|a\rangle$ is defined as

$$|a\rangle q \stackrel{\text{def}}{=} \ulcorner(aq) .$$

This operator is the same as $\langle a$ in Dynamic Logic (see Section 4); the notation $|a\rangle$ has been chosen, since in a test semiring with codomain one can analogously define a backward diamond operator $\langle a|$.

Property (IDext) means *extensional isotonicity*. It can be expressed more succinctly using the pointwise ordering on test transformers $f, g : \text{test}(K) \rightarrow \text{test}(K)$:

$$f \leq g \stackrel{\text{def}}{\Leftrightarrow} \forall q . f(q) \leq g(q) ;$$

(IDext) then becomes

$$|a\rangle \leq |b\rangle \Rightarrow a \leq b .$$

By the definition of the natural order on idempotent semirings it is equivalent to the property of extensionality:

$$(\forall q . |a\rangle q = |b\rangle q) \Rightarrow a = b , \quad (\text{Dext})$$

or, point-free,

$$|a\rangle = |b\rangle \Rightarrow a = b .$$

Property (Sep2) induces the following extensionality property.

Corollary 3.1 *In a domain semiring*

$$(\forall p, q. pbq = 0 \Leftrightarrow paq = 0) \Leftrightarrow |a\rangle = |b\rangle .$$

If the test semiring K also has a codomain operation, we refer to the analogues of the above locality and extensionality properties as *co-locality/co-extensionality* properties.

Now we make the connections between the various properties precise.

Lemma 3.2 *Consider a domain semiring K .*

1. $(Sep1) \Leftrightarrow (DLoc)$.
2. $(Sep2) \Leftrightarrow (IDext)$.

Proof. 1. We prove the equivalent statement $(Sep1') \Leftrightarrow (DLoc)$.
 (\Rightarrow)

$$\begin{aligned} & \ulcorner a \urcorner b \leq \ulcorner ab \urcorner \\ \Leftrightarrow & \quad \{ \text{by (GLA)} \} \\ & \neg \ulcorner ab \urcorner a \urcorner b = 0 \\ \Leftrightarrow & \quad \{ \text{by (Sep1')} \} \\ & \exists q. \neg \ulcorner ab \urcorner a = \neg \ulcorner ab \urcorner a q \wedge \urcorner b = \neg q \urcorner b \\ \Leftrightarrow & \quad \{ \text{Boolean algebra} \} \\ & \exists q. \neg \ulcorner ab \urcorner a = \neg \ulcorner ab \urcorner a q \wedge \urcorner b \leq \neg q \\ \Leftrightarrow & \quad \{ \text{by (LLP)} \} \\ & \exists q. \neg \ulcorner ab \urcorner a = \neg \ulcorner ab \urcorner a q \wedge b = \neg q b \\ \Leftrightarrow & \quad \{ \text{by (Sep1')} \} \\ & \neg \ulcorner ab \urcorner ab = 0 \\ \Leftrightarrow & \quad \{ \text{by (GLA)} \} \\ & \text{TRUE} . \end{aligned}$$

(\Leftarrow)

$$\begin{aligned} & ab = 0 \\ \Leftrightarrow & \quad \{ \text{strictness (1)} \} \end{aligned}$$

$$\begin{aligned}
& \ulcorner(ab) = 0 \\
\Leftrightarrow & \quad \{\{ \text{by (DLoc)} \} \\
& \ulcorner(a \ulcorner b) = 0 \\
\Leftrightarrow & \quad \{\{ \text{strictness (1)} \} \\
& a \ulcorner b = 0 \\
\Leftrightarrow & \quad \{\{ \text{by (GLA)} \} \\
& a = a \ulcorner \ulcorner b \\
\Leftrightarrow & \quad \{\{ \text{by(LLP)} \} \\
& a = a \ulcorner \ulcorner b \wedge b = \ulcorner b b \\
\Rightarrow & \quad \{\{ \text{setting } q = \ulcorner \ulcorner b \} \\
& (\text{Sep1}') .
\end{aligned}$$

2. We prove the equivalent statement $(\text{Sep2}') \Leftrightarrow (\text{IDext})$. It suffices to transform the premise of $(\text{Sep2}')$ into that of (IDext) :

$$\begin{aligned}
& \forall p, q . pbq = 0 \Rightarrow paq = 0 \\
\Leftrightarrow & \quad \{\{ \text{indirect domain inequality (Lemma 1.1)} \} \\
& \forall q . \ulcorner(aq) \leq \ulcorner(bq) \\
\Leftrightarrow & \quad \{\{ \text{definition of } \ulcorner \} \} \\
& \forall q . |a\rangle q \leq |b\rangle q .
\end{aligned}$$

□

The main application of this lemma is a representation theorem that follows immediately from Theorem 3.11 in [7].

Theorem 3.3 *Every *-continuous extensional KAD is isomorphic to a KAT of relations which, however, may be non-standard in that the star operation need not coincide with reflexive-transitive closure.*

The paper [7] also provides a sufficient condition for that algebra of relations to be standard.

Another application of the Lemma is the following. Since (Sep1) and (Sep2) are symmetric in the tests involved, we have

Corollary 3.4 *A semiring with domain and codomain is extensional/local iff it is co-extensional/co-local.*

The part on locality has already been shown in [1] using that in a semiring with domain and codomain one has

$$(DLoc) \Leftrightarrow (ab = 0 \Leftrightarrow a^\top \lrcorner b = 0).$$

4 Modal Kleene Algebra and Dynamic Algebra

Obviously the setting of modal Kleene algebra with its forward diamond operators is very similar to the one of dynamic algebra. Let us therefore briefly relate them and point out some of their differences. Roughly, a *dynamic algebra* is a structure $(K, B, \langle _ \rangle _)$ where K has operations $+$, \cdot and $*$, B is a Boolean algebra and $\langle _ \rangle _ : K \times B \rightarrow B$ is a scalar multiplication satisfying Segerberg’s induction axiom. In the original definition by Kozen [4] K was required to be a $*$ -continuous Kleene algebra. Later, Pratt [9] introduced a more liberal definition in which K was not required to satisfy any axioms at all.

The class **KAD** now blends Kleene algebras and dynamic algebras in a single-sorted structure. The forward diamond as defined above satisfies all axioms of dynamic algebra including the induction axiom (see [1] for details). Since the class **KAD** is based on **KAT**, it does not require $*$ -continuity; rather it uses Kozen’s later axiomatization [5] of $*$ as the least fixed point of an appropriate isotone function. Hence the class of dynamic algebras obtained via the diamond operator of **KAD** is larger than that of Kozen’s dynamic algebras.

Another related structure is that of Kleene modules [3, 8]; these are like dynamic algebras in the sense of Kozen, only with $*$ -continuity replaced by the least-fixpoint characterization of star. Further details on the precise relation between the various structures mentioned can be found in Section 5 of [2].

Acknowledgement We are most grateful to Peter Höfner for a number of comments, notably for his conjecture that Lemma 3.2.1 is independent of the existence of a codomain operation. Useful comments were also provided by Georg Struth.

References

1. J. Desharnais, B. Möller, G. Struth: Kleene algebra with domain. Institut für Informatik, Universität Augsburg, Report 2003-7, May 2003. Revised version to appear in *ACM Transactions on Computational Logic*.
2. J. Desharnais, B. Möller, G. Struth: Modal Kleene Algebra and Applications - A Survey. *Journal on Relational Methods in Computer Science (Electronic Journal - under <http://www.jormics.org>)*, 1:93-131, 2004.
3. T. Ehm, B. Möller, and G. Struth. Kleene modules. In R. Berghammer, B. Möller, and G. Struth, editors, *Relational and Kleene-Algebraic Methods in Computer Science*, volume 3051 of *LNCS*, pages 112–123. Springer, 2004.
4. D. Kozen. A representation theorem for *-free PDL. Technical Report RC7864, IBM, 1979.
5. D. Kozen. A completeness theorem for Kleene algebras and the algebra of regular events. *Information and Computation*, 110(2):366–390, 1994.
6. D. Kozen. Kleene algebra with tests. *Trans. Programming Languages and Systems*, 19(3):427–443, 1997.
7. D. Kozen: On the representation of Kleene algebras with tests. Computer Science Department, Cornell University, Technical Report 2003-1910, September 2003
8. Hans Leiß. Kleenean semimodules and linear languages. In Zoltán Ésik and Anna Ingólfssdóttir, editors, *FICS'02 Preliminary Proceedings*, number NS-02-2 in BRICS Notes Series, pages 51–53. Univ. of Aarhus, 2002.
9. V. R. Pratt. Dynamic algebras as a well-behaved fragment of relation algebras. In: C.H. Bergman, R.D. Maddux, and D.L. Pigozzi, editors, *Algebraic Logic and Universal Algebra in Computer Science*, volume 425 of *LNCS*, pages 77–110. Springer, 1990.