

## Separation of Quasiparticle and Phononic Heat Currents in $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$

B. Zeini,<sup>1</sup> A. Freimuth,<sup>1</sup> B. Büchner,<sup>1</sup> R. Gross,<sup>1</sup> A. P. Kampf,<sup>2</sup> M. Kläser,<sup>3</sup> and G. Müller-Vogt<sup>3</sup>

<sup>1</sup>*II. Physikalisches Institut, Universität zu Köln, 50937 Köln, Germany*

<sup>2</sup>*Theoretische Physik III, Universität Augsburg, 86135 Augsburg, Germany*

<sup>3</sup>*Kristall- und Materiallabor, Universität Karlsruhe, 76128 Karlsruhe, Germany*

(Received 9 February 1998)

Measurements of the transverse ( $k_{xy}$ ) and longitudinal ( $k_{xx}$ ) thermal conductivity in high magnetic fields are used to separate the quasiparticle thermal conductivity ( $k_{xx}^{\text{el}}$ ) of the  $\text{CuO}_2$ -planes from the phononic thermal conductivity in  $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ .  $k_{xx}^{\text{el}}$  is found to display a pronounced maximum below  $T_c$ . Our data analysis reveals distinct transport ( $\tau$ ) and Hall ( $\tau_H$ ) relaxation times below  $T_c$ : Whereas  $\tau$  is strongly enhanced,  $\tau_H$  follows the same temperature dependence as above  $T_c$ . [S0031-9007(99)08592-0]

PACS numbers: 74.25.Fy, 72.15.Jf, 74.72.Bk

The study of heat transport in the superconducting state is well known to provide valuable information on the quasiparticle (QP) excitations and their dynamics. Compared to other probes of the QP dynamics such as the microwave conductivity thermal transport has the advantage of probing only the QP response, since the superfluid does not carry heat. On the other hand, a major complication in the analysis of the thermal conductivity is often a substantial phononic contribution  $k_{xx}^{\text{ph}}$  to the heat current. Such a situation is realized in the high- $T_c$  superconductors (HTSC), where the heat current above  $T_c$  is known to be dominated by  $k_{xx}^{\text{ph}}$  [1]. Accordingly, the interpretation of experimental data is ambiguous: For example, the maximum in the temperature dependence of the thermal conductivity in the superconducting state of  $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$  (YBCO) has been attributed to both, a maximum of the electronic as well as the phononic contribution [1–4]. A clear separation of the phonon and QP heat currents is difficult and up to now an unsolved problem.

It has been pointed out in Refs. [5,6] that the transverse thermal conductivity  $k_{xy}$ —the thermal analogue of the Hall effect—is free of phonons, i.e.,  $k_{xy}$  is purely electronic.  $k_{xy}$  is related to the electronic thermal conductivity  $k_{xx}^{\text{el}}$  according to  $k_{xy} = k_{xx}^{\text{el}} \tan \alpha_R$  where  $\alpha_R$  is the thermal Hall angle. From the Wiedemann-Franz law [7] one expects that  $\alpha_R$  is equal to the electrical Hall angle  $\alpha_H$  as obtained from the electrical ( $\sigma_{xx}$ ) and Hall ( $\sigma_{xy}$ ) conductivities via  $\tan \alpha_H = \sigma_{xy}/\sigma_{xx}$ . In conventional metals  $\tan \alpha_H = \omega_c \tau$  where  $\omega_c = eB/m$  is the cyclotron frequency and  $\tau$  is the usual transport relaxation time. In contrast, in the normal state of the HTSC  $\tan \alpha_H$  is highly anomalous and has a temperature dependence which is *distinctly* different from  $\tau(T)$  [8–10]. Up to now no data on the QP Hall angle are available below  $T_c$ , but one may suspect that its behavior is anomalous as well. Therefore, in order to calculate  $k_{xx}^{\text{el}}$  from  $k_{xy}$  it is necessary to treat  $\tan \alpha_R = \omega_c \tau_R$  as an additional independent parameter which must be determined experimentally.

In this Letter we determine both  $k_{xx}^{\text{el}}$  and  $\tan \alpha_R$  below  $T_c$  in a single crystal of YBCO from combined

measurements of  $k_{xy}$  and  $k_{xx}$  in high magnetic fields. Our main results are as follows: (1) The electronic thermal conductivity of the  $\text{CuO}_2$ -planes shows a pronounced maximum below  $T_c$  which is strongly suppressed by a magnetic field. (2)  $\tan \alpha_R$ , as extracted from the thermal transport data below  $T_c$ , displays the same temperature and magnetic field dependence as  $\tan \alpha_H$  obtained from electrical transport data above  $T_c$ , and it passes smoothly through  $T_c$ . This shows that  $\tau_R \approx \tau_H$  and, remarkably, that  $\tau_H$  and  $\tau$  behave differently also below  $T_c$ .

The thermal conductivity tensor  $\underline{k}$  is the sum of an electronic and a phononic part  $\underline{k} = \underline{k}^{\text{el}} + \underline{k}^{\text{ph}}$ . It is defined via the heat current density  $\mathbf{j}_h = -\underline{k} \nabla T$  [11]. We assume that  $\underline{k}^{\text{ph}}$  remains diagonal even for  $\mathbf{B} \neq 0$ , i.e.,  $k_{xy}^{\text{ph}} = 0$ . In this case the transverse components of  $\underline{k}$  are purely electronic. The transverse thermal conductivity, also called the Righi-Leduc effect, is measured as follows: In a magnetic field  $\mathbf{B} = (0, 0, B)$  a temperature gradient  $\nabla_x T$  is applied in the  $x$  direction. Under the condition  $j_{h,y} = 0$  a transverse temperature gradient  $\nabla_y T$  is found in the  $y$  direction. Using the Onsager relations we find in this situation

$$j_{h,y} = k_{xy} \nabla_x T - k_{xx} \nabla_y T = 0, \quad (1)$$

with  $k_{xx} = k_{yy}$  for twinned crystals without in-plane anisotropy.  $k_{xy}$  can therefore be determined experimentally by measuring  $\nabla_x T$ ,  $\nabla_y T$ , and the *total* longitudinal thermal conductivity  $k_{xx}$ .

Our measurements were carried out at constant temperatures with the magnetic field applied perpendicular to the  $\text{CuO}_2$  planes. Typically, temperature gradients  $\nabla_x T$  of order 0.5 K/mm were applied using a small manganin heater mounted on top of the samples. The resulting transverse temperature gradients  $\nabla_y T$  of order  $10^{-3}$  K/mm in magnetic fields up to 14 T were measured with AuFe-Chromel thermocouples calibrated in the same field range [12]. To eliminate offset voltages due to misalignment of the thermocouple we have measured for both field directions  $\pm \mathbf{B}$  in order to determine the Righi-Leduc component of  $\nabla_y T$  which must be antisymmetric with respect to field reversal. We have measured

in two different modes: Either  $\mathbf{B}$  was reversed at fixed temperature or we have heated the sample to temperatures above  $T_c$  before the field was reversed. Because of vortex pinning effects this latter mode was used for all low temperature measurements. We note that sweeping the magnetic field at fixed temperatures results in a strong hysteresis of  $k_{xx}$  in almost the entire temperature range below  $T_c$ , similar as reported in Ref. [13];  $k_{xy}$ , which must be extracted from the asymmetry of the field dependence, can therefore not be determined by sweeping the magnetic field. We have tested our method by measurements on an insulator ( $k_{xy} = 0$ ) and on simple metals [14]. Details of our experimental setup will be described elsewhere. The results presented here have been obtained on a high quality twinned single crystal of YBCO with dimensions  $1.9 \text{ mm} \times 2 \text{ mm} \times 0.38 \text{ mm}$  and with a superconducting transition at  $T_c \approx 90.5 \text{ K}$ .

Representative experimental results are shown in Fig. 1.  $k_{xx}$  has a pronounced maximum at  $T_{\text{max}} < T_c$  which is strongly suppressed by the applied magnetic field. The absolute value of  $k_{xx}$  ( $\approx 10 \text{ W/Km}$  at  $T_c$ ), the relative upturn of  $k_{xx}$  in zero field as characterized by the ratio  $k_{xx}(T_{\text{max}})/k_{xx}(T_c) \approx 1.6$  for our sample, as well as the sensitivity of the maximum to magnetic fields are consistent with previous results [1–3]. The overall temperature dependence of  $k_{xy}$  is similar to that of  $k_{xx}$  but the maximum of  $k_{xy}$  occurs at higher temperatures and the relative change below  $T_c$  is larger in comparison




FIG. 1. Open symbols:  $k_{xx}$  (upper panel) and  $k_{xy}$  (lower panel) of  $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$  as a function of temperature  $T$  for various fixed magnetic fields as indicated in the figure. Full symbols:  $k_{xx}^{\text{rest}}$  as obtained from our data analysis (see text).

to  $k_{xx}$ . The absolute magnitude of  $k_{xy}$  is comparable to that reported previously in Ref. [6].

For our data analysis we assume that in YBCO three channels of heat conduction are present:

$$k_{xx} = k_{xx}^{\text{el}} + k_{xx}^{\text{ch}} + k_{xx}^{\text{ph}} = k_{xx}^{\text{el}} + k_{xx}^{\text{rest}}. \quad (2)$$

Here  $k_{xx}^{\text{el}}$  is the electronic contribution from the  $\text{CuO}_2$ -planes ( $\text{CuO}_2$  bilayers in YBCO) and  $k_{xx}^{\text{ph}}$  is that of the phonons.  $k_{xx}^{\text{ch}}$  describes a possible contribution from the  $\text{CuO}$  chains which are present in YBCO along the  $b$  direction of the orthorhombic structure. These chains are metallic for optimally doped samples and lead to a rather strong  $a$ - $b$  anisotropy in untwinned crystals ( $\sigma_{bb}/\sigma_{aa} \approx 2$ ) [8]. In a twinned crystal they should contribute to the electrical and the heat conduction on average;  $a$ - $b$  anisotropy is, of course, absent. In the subsequent data analysis  $k_{xx}^{\text{ch}}$  must be treated differently from  $k_{xx}^{\text{el}}$  since the  $\text{CuO}$ -chains as a quasi-one-dimensional channel for charge and heat transport should neither contribute to the transverse transport coefficients  $\sigma_{xy}$  and  $k_{xy}$  nor to the magnetic field dependence of the longitudinal ones.

We compare in Fig. 2  $\Delta k_{xx} = k_{xx}(B) - k_{xx}(0)$  and  $k_{xy}/B$ . Notably, these quantities have the same magnetic field dependence, i.e.,

$$\frac{\partial}{\partial B} \left( \frac{k_{xy}}{B} \right) \propto \frac{\partial k_{xx}}{\partial B}. \quad (3)$$

This observation provides the key to our data analysis. We define

$$\tan \alpha_R = \omega_c \tau_R = \frac{k_{xy}}{k_{xx}^{\text{el}}}, \quad (4)$$




FIG. 2.  $\Delta k_{xx} = k_{xx}(B) - k_{xx}(B = 0)$  ( $\bullet$ ) and  $k_{xy}/B$  ( $\circ$ ) versus magnetic field  $B$  at fixed temperatures given in the figure.

where  $\tau_R(B, T)$  is a “relaxation time” introduced to parametrize the field and temperature dependence of tan  $\alpha_R$ . Using Eqs. (2) and (4) we find

$$\frac{m}{e} \frac{\partial}{\partial B} \left( \frac{k_{xy}}{B} \right) = \tau_R \frac{\partial k_{xx}}{\partial B} + \left[ k_{xx}^{el} \frac{\partial \tau_R}{\partial B} - \tau_R \frac{\partial k_{xx}^{ph}}{\partial B} - \tau_R \frac{\partial k_{xx}^{ch}}{\partial B} \right]. \quad (5)$$

Apparently, our experimental results suggest that the term in brackets is zero. Since the three terms in brackets refer to three distinct channels of heat conduction this requires that  $\tau_R$ ,  $k_{xx}^{ph}$ , and  $k_{xx}^{ch}$  are separately field independent. Note that this is certainly reasonable in view of the origin of these contributions to the heat current.

Assuming thus that the three terms in brackets vanish  $\tau_R$  can be calculated from our data by comparing  $k_{xx}$  and  $k_{xy}/B$  at different magnetic fields according to  $e\tau_R/m = \Delta(k_{xy}/B)/\Delta k_{xx}$  [see Eq. (5)]. The result is shown in Fig. 3. Note that the values obtained for different magnetic fields coincide within the experimental accuracy consistent with the anticipated field independence of  $\tau_R$ .

As a check of our result for  $\tau_R$  we have also determined  $e\tau_H/m = \sigma_{xy}/B\sigma_{xx}$  for the same sample from measurements of  $\sigma_{xy}$  and  $\sigma_{xx}$  in the normal state [14]. These data as well as their extrapolation [15] to temperatures below  $T_c$  are also shown in Fig. 3. The extrapolated values for  $\tau_H^{-1}$  look very similar to  $\tau_R^{-1}$  regarding the temperature dependence (see inset of Fig. 3), but they appear to be systematically larger by roughly a factor of 2. However, note that  $\tau_R$  as extracted from the thermal transport data is clearly unaffected by the CuO chains and that this is also true for  $\sigma_{xy}$ . In contrast,  $\sigma_{xx}$  has a contribution from the CuO chains, i.e.,  $\sigma_{xx} = \sigma_{xx}^{pl} + \langle \sigma_{xx}^{ch} \rangle$ , where  $\sigma_{xx}^{pl}$  is the electrical conductivity of the CuO<sub>2</sub> planes and  $\langle \sigma_{xx}^{ch} \rangle$  is an average of the chain contribution appropriate for a twinned crystal. With  $\sigma_{xx}^{pl} \approx \langle \sigma_{xx}^{ch} \rangle$  [8] we conclude that  $e\tau_H/m = \sigma_{xy}/B\sigma_{xx}$  is underestimated by a factor of 2. Correcting the normal state data for this factor we find excellent agreement between  $\tau_H$  and  $\tau_R$ , i.e., our data tell  $\tau_R \approx \tau_H$ . This strongly supports the procedure of our data analysis.

Once  $\tau_R$  is known the remainder of our analysis is straightforward:  $k_{xx}^{el}(B \neq 0)$  follows from Eq. (4) using the data for  $k_{xy}(B)$ , and  $k_{xx}^{rest}$  is obtained subsequently from Eq. (2) for each field strength. As a test for internal consistency we have verified that  $k_{xx}^{rest}$  is indeed field independent. Finally,  $k_{xx}^{el}(B = 0)$  follows from Eq. (2) using  $k_{xx}^{rest}$  and the zero field data for  $k_{xx}$ . We have also determined  $k_{xx}^{el}(B = 0)$  directly from Eq. (4) using an extrapolation of  $B/k_{xy}$  to  $B = 0$  in good agreement with the results obtained from using  $k_{xx}^{rest}$  and Eq. (2).

In Fig. 4 we show  $k_{xx}^{el}(B)$  as obtained from our data analysis.  $k_{xx}^{el}$  represents the electronic thermal conductivity of the CuO<sub>2</sub>-planes in YBCO. Our results thus confirm explicitly that  $k_{xx}^{el}$  is strongly enhanced below  $T_c$ .  $k_{xx}^{el} \propto Tn_{QP}\tau$  implies that  $\tau$  is strongly enhanced below  $T_c$  overcompensating the decrease of the QP number density  $n_{QP}(T)$  with decreasing temperature. This confirms




FIG. 3.  $T < T_c$ :  $m/e\tau_R$  vs temperature  $T$  obtained from  $e\tau_R/m = \Delta(k_{xy}/B)/\Delta k_{xx}$ , where  $\Delta(k_{xy}/B) = k_{xy}(B_1)/B_1 - k_{xy}(B_2)/B_2$  and  $\Delta k_{xx} = k_{xx}(B_1) - k_{xx}(B_2)$ . Different symbols correspond to different values of  $B_1$  and  $B_2$ .  $T > T_c$ :  $m/e\tau_H$  obtained from  $\sigma_{xy}$  and  $\sigma_{xx}$ . Solid line: extrapolated normal state data (see text). Dashed line: extrapolated normal state data divided by a factor of 2. Inset: The same data on a double logarithmic scale. The dotted line corresponds to a  $T^2$  temperature dependence.

the results obtained for the QP relaxation time from the microwave conductivity [16].

We also find that  $k_{xx}^{el}$  is very sensitive to magnetic fields contrary to what is observed in the normal state, where the total thermal conductivity and thus  $k_{xx}^{el}$  is field independent. It is straightforward to attribute the field dependence below  $T_c$  to an additional scattering mechanism [17] characteristic for the superconducting state such as scattering of QPs on vortices [6,18,19]. Assuming the corresponding scattering rate  $\tau_v^{-1}$  to be proportional to the number of vortices  $n_v \propto B$  one expects for the total scattering rate that  $\tau^{-1} = \tau_{in}^{-1} + \tau_v^{-1} = \tau_{in}^{-1} + \alpha B$ . Here,  $\tau_{in}$  includes the same scattering processes as in the normal state, i.e., it has in general an elastic defect and an inelastic contribution; the latter collapses below  $T_c$ . Using  $k_{xy} = k_{xx}^{el} \omega_c \tau_R = L(T) T \sigma_{xx} \omega_c \tau_R$  where  $L$  is the Lorentz number and  $\sigma_{xx} = n_{QP}(T) e^2 \tau/m$  we find

$$\frac{B}{k_{xy}} = C(T) \tau^{-1} = C(T) (\tau_{in}^{-1} + \alpha B), \quad (6)$$

where  $C(T) = m^2/(LTn_{QP}e^2\tau_R)$  depends only on temperature. Thus a plot of  $B/k_{xy}$  vs  $B$  should yield a straight line. This is indeed the case as shown in Fig. 4. We note that by assuming  $\alpha$  to be temperature independent  $\tau_{in}(T)$  can be extracted from the slope and the intersection of the




FIG. 4. Left panel: Electronic thermal conductivity  $k_{xx}^{el}(B)$  of the  $\text{CuO}_2$  planes as a function of temperature for various magnetic fields as indicated in the figure. The zero field data have been obtained from  $k_{xx}^{el}(B=0) = k_{xx}(B=0) - k_{xx}^{\text{rest}}$  ( $\circ$ ) as well as from an extrapolation of  $B/k_{xy}$  to  $B=0$  ( $\bullet$ ). Right panels:  $B/k_{xy}$  as a function of magnetic field  $B$  at various fixed temperatures given in the figure.

$B/k_{xy}$  vs  $B$  curves (to within the constant factor  $\alpha$ ). We find that  $\tau_{\text{in}}$  increases strongly below  $T_c$  [14].

$k_{xx}^{\text{rest}}$  as obtained from our data analysis is shown in Fig. 1. Remarkably,  $k_{xx}^{\text{rest}}$  shows a pronounced maximum below  $T_c$ , too. This maximum may be due to the phononic contribution [1]. However, it may also arise from the chain contribution. The latter has previously been determined experimentally from the  $a$ - $b$  anisotropy of  $k_{xx}$  in detwinned single crystals of YBCO [2,20].  $k_{xx}^{\text{ch}}$  shows a pronounced maximum below  $T_c$  with an overall temperature dependence similar to that found here for  $k_{xx}^{\text{rest}}$  [20]. Furthermore, the field independence of  $k_{xx}^{\text{rest}}$  implies that the phononic contribution  $k_{xx}^{\text{ph}}$  is independent of  $B$ . Such a conclusion has recently been drawn also on the basis of low temperature results for the thermal conductivity in Bi-based HTSC [21].

Finally, our data show clearly that  $\tau_R \approx \tau_H$  and that—as in the normal state— $\tau_H$  and  $\tau$  behave differently also below  $T_c$ . In particular, whereas  $\tau$  is strongly enhanced below  $T_c$ ,  $\tau_H$  is unaffected by the superconducting transition and shows the same temperature dependence as above  $T_c$ . This finding should provide important information for the theoretical understanding of transport phenomena in the cuprates.

In summary, we have presented a separation of the QP and phononic contributions to the thermal conductivity below  $T_c$  in YBCO based on measurements of the longitudinal and transverse thermal conductivity in high magnetic fields. Our data analysis shows explicitly that the QP contribution to  $k_{xx}$  is strongly enhanced below  $T_c$  and that it is the QP contribution to the heat current which

is responsible for the magnetic field dependence of  $k_{xx}$ . We find that—as in the normal state—two relaxation times must be distinguished also below  $T_c$ : Whereas the QP relaxation time  $\tau$  is strongly enhanced below  $T_c$  and magnetic field dependent, the Hall relaxation time  $\tau_H$  remains independent of  $B$  below  $T_c$  and has the same temperature dependence as above  $T_c$ .

We are particularly grateful for stimulating discussions with W. Brenig, Ch. Bruder, M. Galffy, P. J. Hirschfeld, T. Kopp, D. Rainer, S. Uhlénbrück, and P. Wölfle. This work was supported by the Deutsche Forschungsgemeinschaft through SFB 341.

- [1] C. Uher, in *Physical Properties of High Temperature Superconductors*, edited by D.M. Ginsberg (World Scientific, Singapore, 1992), Vol. 3.
- [2] R. C. Yu *et al.*, Phys. Rev. Lett. **69**, 1431 (1992).
- [3] S. D. Peacor *et al.*, Phys. Rev. B **44**, 9508 (1991).
- [4] J. L. Cohn *et al.*, Phys. Rev. Lett. **71**, 1657 (1993).
- [5] A. Freimuth *et al.*, J. Low Temp. Phys. **95**, 383 (1994).
- [6] K. Krishana *et al.*, Phys. Rev. Lett. **75**, 3529 (1995).
- [7] The generalized Wiedemann-Franz law relates the electrical and thermal conductivity tensors,  $\underline{\sigma}$  and  $\underline{k}$ , respectively, according to  $\underline{k} = L T \underline{\sigma}$ .  $L$  is the Lorentz number.
- [8] Y. Iye, in *Physical Properties of High Temperature Superconductors*, edited by D.M. Ginsberg (World Scientific, Singapore, 1992), Vol. 3.
- [9] T. R. Chien *et al.*, Phys. Rev. Lett. **67**, 2088 (1991).
- [10] P. W. Anderson, Phys. Rev. Lett. **67**, 2092 (1991).
- [11] The heat current due to vortex motion is negligibly small [14].
- [12] The thermal conductance of the thermocouple was about 0.5% of that of the sample. The calibration was performed against an insulator above 30 K and against a second thermocouple of the same type placed in zero magnetic field below 30 K [see C. K. Chiang, Rev. Sci. Instrum. **45**, 985 (1974); and Ref. [14]].
- [13] H. Aubin *et al.*, Science **280**, 9a (1998).
- [14] B. Zeini, Ph.D. thesis, Universität zu Köln, (Shaker, 1997); B. Zeini *et al.* (to be published).
- [15] We have fitted the resistivity  $\rho$  and the inverse Hall coefficient  $R_H^{-1}$  to  $a + bT$  and to  $\tilde{a} + \tilde{b}T$ , respectively, where  $a$ ,  $b$ ,  $\tilde{a}$ , and  $\tilde{b}$  are constants.  $e\tau_R/m$  was then obtained from  $e\tau_H/m = \sigma_{xy}/B\sigma_{xx} \approx R_H/\rho$ .
- [16] D. A. Bonn and W. N. Hardy, in *Physical Properties of High Temperature Superconductors*, edited by D.M. Ginsberg (World Scientific, Singapore, 1996), Vol. 5.
- [17] A magnetic field dependence of  $k_{xx}^{el}$  may also result from the field dependence of  $n_{\text{QP}}$  in unconventional superconductors [G. E. Volovik, JETP Lett. **58**, 469 (1993)]. However,  $n_{\text{QP}}$  and thus  $k_{xx}^{el}$  should increase with  $B$  contrary to the behavior of  $k_{xx}$  found here.
- [18] M. B. Salamon *et al.*, J. Supercond. **8**, 449 (1995).
- [19] R. M. Cleary, Phys. Rev. **175**, 587 (1968).
- [20] R. Gagnon *et al.*, Phys. Rev. Lett. **78**, 1976 (1997).
- [21] K. Krishana *et al.*, Science **277**, 83 (1997).