

- [3] S.C. Brenner, L.-Y. Sung, H. Zhang, and Y. Zhang. *A quadratic C^0 interior penalty method for the displacement obstacle problem of clamped Kirchhoff plates*, preprint, 2011.
- [4] S.C. Brenner, L.-Y. Sung, and Y. Zhang. *Finite element methods for the displacement obstacle problem of clamped plates*, Math. Comp., (to appear).
- [5] H. R. Brezis and G. Stampacchia. *Sur la régularité de la solution d'inéquations elliptiques*, Bull. Soc. Math. France **96** (1968), 153–180.
- [6] F. Brezzi, W. Hager, and P.-A. Raviart. *Error estimates for the finite element solution of variational inequalities*, Numer. Math. **28** (1977), 431–443.
- [7] L.A. Caffarelli and A. Friedman. *The obstacle problem for the biharmonic operator*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **6** (1979), 151–184.
- [8] E. Casas. *Control of an elliptic problem with pointwise state control*, SIAM J. Control Optim. **24** (1986), 1309–1318.
- [9] R.S. Falk. *Error estimates for the approximation of a class of variational inequalities*, Math. Comp. **28** (1974), 963–971.
- [10] J. Frehse. *Zum Differenzierbarkeitsproblem bei Variationsungleichungen höherer Ordnung*, Abh. Math. Sem. Univ. Hamburg **36** (1971), 140–149.
- [11] J.-L. Lions and G. Stampacchia. *Variational inequalities*, Comm. Pure Appl. Math. **20** (1967), 493–519.
- [12] W. Liu, W. Gong and N. Yan. *A new finite element approximation of a state-constrained optimal control problem*, J. Comp. Math. **27** (2009), 97–114.

Comparison results for first-order FEMs

MIRA SCHEDENSACK

(joint work with Carsten Carstensen, Daniel Peterseim)

Various first-order finite element methods are known for the Poisson Model Problem (1) and for linear elasticity (2). The recent publications [1] started the comparison between some of these methods for the Poisson Model Problem, which is completed in this presentation and its underlying paper [3].

Given a bounded polygonal Lipschitz domain Ω in the plane and data $f \in L^2(\Omega)$, the Poisson model problem seeks the weak solution $u \in H^1(\Omega)$ of

$$(1) \quad -\Delta u = f \text{ in } \Omega \quad \text{and} \quad u = 0 \text{ on } \partial\Omega.$$

This presentation compares the error of three popular finite element methods (FEM) of Figure 1 for the numerical solution of (1), namely the conforming *Courant* FEM (CFEM) [4], the nonconforming *Crouzeix-Raviart* FEM (CR-NCFEM) [5], and the mixed *Raviart-Thomas* FEM (RT-MFEM) [7] with respective solutions u_C , u_{CR} , and (p_{RT}, u_{RT}) based on a shape-regular triangulation \mathcal{T} of Ω into triangles. The finite element space of CFEM reads $P_1(\Omega) \cap C_0(\Omega)$ for




FIGURE 1. CFEM (left), CR-NCFEM (middle), RT-MFEM (right).

$C_0(\Omega)$ the continuous functions with zero boundary conditions. The finite element

space of Crouzeix-Raviart $\text{CR}_0^1(\mathcal{T})$ consists of all piecewise affines which are continuous at the midpoints of interior edges and vanish at the midpoints of exterior edges. The Raviart-Thomas finite element space for the flux approximation reads $\text{RT}_0(\mathcal{T}) := \{p_{\text{RT}} \in P_1(\mathcal{T}, \mathbb{R}^2) \cap H(\text{div}, \Omega) \mid \forall T \in \mathcal{T} \exists a_T, b_T, c_T \in \mathbb{R} : p_{\text{RT}}|_T(x) = (a_T, b_T) + c_T x\}$.

The comparison is stated in terms of $A \lesssim B$ which abbreviates the existence of some constant C which only depends on the minimal angle in \mathcal{T} , but not on the domain Ω and not on the mesh-size $h_{\mathcal{T}}$, such that $A \leq CB$. The comparison includes data oscillations, namely $\text{osc}(f, \mathcal{T}) := \|h_{\mathcal{T}}(f - \Pi_0 f)\|$ where Π_0 denotes the L^2 orthogonal projection onto the piecewise constants.

The comparison result for CFEM, CR-NCFEM and RT-MFEM states that the errors of CFEM and CR-NCFEM are equivalent up to data oscillations, in the sense that

$$\|\nabla u - \nabla u_{\text{C}}\| \lesssim \|\nabla u - \nabla_{\text{NC}} u_{\text{CR}}\| \lesssim \|\nabla u - \nabla u_{\text{C}}\| + \text{osc}(f, \mathcal{T}).$$

The error of RT-MFEM is superior in the sense that

$$\|\nabla u - \nabla_{\text{NC}} u_{\text{CR}}\| \lesssim \|h_{\mathcal{T}} f\| + \|\nabla u - p_{\text{RT}}\| \lesssim \|\nabla u - \nabla_{\text{NC}} u_{\text{CR}}\| + \text{osc}(f, \mathcal{T}),$$

but the converse is false, i.e.,

$$\|\nabla u - \nabla_{\text{NC}} u_{\text{CR}}\| \not\lesssim \|\nabla u - p_{\text{RT}}\| + \text{osc}(f, \mathcal{T}).$$

The proof of the inequalities is an example of the medius analysis for it combines arguments of an *a priori* with those of an *a posteriori* error analysis. It is emphasised that no regularity assumption is made and the results hold for arbitrary coarse triangulations and not just in an asymptotic regime. The proof of the superiority of RT-MFEM considers a sequence of domains, on which the RT-MFEM has a steeper convergence rate than CR-NCFEM.

The results for the Poisson Model Problem can be generalised for the Navier-Lamé equations from linear elasticity, which seek $u \in H_0^1(\Omega; \mathbb{R}^2)$ with

$$(2) \quad f + 2\mu\Delta u + (\mu + \lambda)\nabla(\text{div } u) = 0 \quad \text{in } \Omega.$$

The compared FEMs are the conforming *Courant* FEM (CFEM) [2], the nonconforming *Kouhia-Stenberg* FEM (KS-NCFEM) [6], and the nonconforming *Crouzeix-Raviart* FEM (CR-NCFEM) [2] with respective solutions σ_{C} , σ_{KS} and σ_{CR} . The finite element space of KS-NCFEM reads $\text{KS} := (P_1(\mathcal{T}) \cap C_0(\Omega)) \times \text{CR}_0^1(\mathcal{T})$. The discretisation of CFEM and KS-NCFEM is based on the bilinear form

$$a(u_{\text{KS}}, v_{\text{KS}}) := \int_{\Omega} \varepsilon_{\text{NC}}(u_{\text{KS}}) : \mathbb{C} \varepsilon_{\text{NC}}(v_{\text{KS}}) \, dx,$$

while the discretisation of CR-NCFEM involves the bilinear form

$$a(u_{\text{CR}}, v_{\text{CR}}) := \int_{\Omega} (\mu D_{\text{NC}} u_{\text{CR}} : D_{\text{NC}} v_{\text{CR}} + (\mu + \lambda) \text{div}_{\text{NC}} u_{\text{CR}} \text{div}_{\text{NC}} v_{\text{CR}}) \, dx.$$

The comparison result for linear elasticity involves the Lamé modulus λ , which effects the locking and the \lesssim notation means, that, in addition, the underlying

constants do not depend on the Lamé modulus λ . Then

$$\|\sigma - \sigma_C\| \lesssim \lambda \|\sigma - \sigma_{KS}\| \lesssim \lambda (\|\sigma - \sigma_C\| + \text{osc}(f, \mathcal{T}))$$

and

$$\|\sigma - \sigma_{KS}\| + \text{osc}(f, \mathcal{T}) \approx \|\sigma - \sigma_{CR}\| + \text{osc}(f, \mathcal{T}).$$

REFERENCES

- [1] D. Braess, *An a posteriori error estimate and a comparison theorem for the nonconforming P_1 element*, *Calcolo* **46** (2009), 149–155.
- [2] S.C. Brenner, L.R. Scott, *The mathematical theory of finite element methods*, Texts in Applied Mathematics **15** (2008).
- [3] C. Carstensen, D. Peterseim, M. Schedensack, *Comparison Results of Three First-order Finite Element Methods for the Poisson Model Problem*, Preprint #831 (2011).
- [4] R. Courant, *On a method for the solution of boundary-value problems*, in: Theodore von Kármán Anniversary Volume (1941), 189–194.
- [5] M. Crouzeix, P.A. Raviart, *Conforming and nonconforming finite element methods for solving the stationary Stokes equations. I*, *Rev. Française Automat. Informat. Recherche Opérationnelle Sér. Rouge* **7** (1973), 33–75.
- [6] R. Kouhia, R. Stenberg, *A linear nonconforming finite element method for nearly incompressible elasticity and Stokes flow*, *Comput. Methods Appl. Mech. Engrg.* **124** (1995), 195–212.
- [7] P.A. Raviart, J.M. Thomas *A mixed finite element method for 2nd order elliptic problems*, in: *Mathematical aspects of finite element methods* (Proc. Conf., Consiglio Naz. delle Ricerche (C.N.R.), Rome, 1975) (1977), 292–315.

Quasi optimal adaptive pseudostress approximation of the Stokes equations

DIETMAR GALLISTL

(joint work with Carsten Carstensen, Mira Schedensack)

The pseudostress-velocity formulation [3, 5] of the stationary Stokes equations

$$(1) \quad -\Delta u + \nabla p = f \quad \text{and} \quad \text{div } u = 0 \quad \text{in } \Omega$$

with Dirichlet boundary conditions along the polygonal boundary $\partial\Omega$ allows the stresses-like variables σ in Raviart-Thomas mixed finite element spaces [2] $\text{RT}_k(\mathcal{T})$ with respect to a regular triangulation \mathcal{T} , and hence allows for higher flexibility in arbitrary polynomial degrees.

The weak form of problem (1) is formally equivalent and reads: Given $f \in L^2(\Omega; \mathbb{R}^2)$ and $g \in H^1(\Omega; \mathbb{R}^2) \cap H^1(\mathcal{E}(\partial\Omega); \mathbb{R}^2)$ with $\int_{\partial\Omega} g \cdot \nu \, ds = 0$ seek $\sigma \in H(\text{div}, \Omega; \mathbb{R}^{2 \times 2})/\mathbb{R}$ and $u \in L^2(\Omega; \mathbb{R}^2)$ such that

$$(2) \quad \begin{aligned} \int_{\Omega} \text{dev } \sigma : \tau \, dx + \int_{\Omega} u \cdot \text{div } \tau \, dx &= \int_{\partial\Omega} g \cdot \tau \, \nu \, ds, \\ \int_{\Omega} v \cdot \text{div } \sigma \, dx &= - \int_{\Omega} f \cdot v \, dx \end{aligned}$$