

Study concept and design: Foglia, Nolen, DeMauro, Das, Schmidt.
Acquisition, analysis, or interpretation of data: Foglia, Nolen, DeMauro, Bell, Stoll, Schmidt.
Drafting of the manuscript: Foglia, Nolen.
Critical revision of the manuscript for important intellectual content: Foglia, DeMauro, Das, Bell, Stoll, Schmidt.
Statistical analysis: Nolen, Das.
Obtained funding: Bell, Stoll.
Administrative, technical, or material support: Stoll.
Study supervision: Stoll, Schmidt.

Conflict of Interest Disclosures: The authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

Funding/Support: The National Institutes of Health and the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) provided grant support for the Neonatal Research Network's Generic Database Study via cooperative agreements.

Role of the Funder/Sponsor: The National Institutes of Health and the NICHD had no role in the design and conduct of the study; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Independent Statistical Analysis: Data collected at participating sites of the NICHD Neonatal Research Network were transmitted to RTI International, the data coordinating center for the network, which performed the storage, management, analysis, and interpretation of the data.

Group Information: Members of the NICHD Neonatal Research Network who contributed to this article: Rosemary D. Higgins, MD (NICHD, Bethesda, Maryland), Brenda B. Poindexter, MD, MS (Department of Pediatrics, Indiana University School of Medicine, Indianapolis), Seetha Shankaran, MD (Department of Pediatrics, Wayne State University, Detroit, Michigan), and Krisa P. Van Meurs, MD (Division of Neonatal and Developmental Medicine, Stanford University School of Medicine, Palo Alto, California).

Disclaimer: This article represents the views of the authors and does not necessarily represent the views of the NICHD.

Additional Information: The entire list of the members of the NICHD Neonatal Research Network appears in Shankaran et al.⁶

1. Rich W, Finer NN, Gantz MG, et al; SUPPORT and Generic Database Subcommittees of the Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Enrollment of extremely low birth weight infants in a clinical research study may not be representative. *Pediatrics*. 2012;129(3):480-484.
2. Schmidt B, Gillie P, Caco C, Roberts J, Roberts R. Do sick newborn infants benefit from participation in a randomized clinical trial? *J Pediatr*. 1999;134(2):151-155.
3. Rüegger CM, Kraus A, Koller B, et al. Randomized controlled trials in very preterm infants: does inclusion in the study result in any long-term benefit? *Neonatology*. 2014;106(2):114-119.
4. Vist GE, Bryant D, Somerville L, Birmingham T, Oxman AD. Outcomes of patients who participate in randomized controlled trials compared to similar patients receiving similar interventions who do not participate. *Cochrane Database Syst Rev*. 2008;(3):MR000009.
5. Fernandes N, Bryant D, Griffith L, et al. Outcomes for patients with the same disease treated inside and outside of randomized trials: a systematic review and meta-analysis. *CMAJ*. 2014;186(16):E596-E609.
6. Shankaran S, Laptook AR, Pappas A, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Effect of depth and duration of cooling on deaths in the NICU among neonates with hypoxic ischemic encephalopathy: a randomized clinical trial. *JAMA*. 2014;312(24):2629-2639.

COMMENT & RESPONSE

Presurgery Sedation and Patient Experience

To the Editor In a randomized clinical trial by Dr Maurice-Szamburski and colleagues,¹ premedication with oral lorazepam 2 hours prior to arriving in the operating room did not improve patient satisfaction after surgery and was associated with

prolonged time to extubation and decreased cognitive recovery compared with placebo or no medication. These results may not indicate that all sedative premedication is unwarranted but may suggest lorazepam is inadequate.

Maurice-Szamburski and colleagues explained why lorazepam was used instead of midazolam for their trial: "The choice of lorazepam was motivated by its use in the largest survey of sedative premedication published to date," referring to a study by Kain et al.² There is some discrepancy between this statement and the survey results, in which among adults, the most commonly used sedative premedication was midazolam (>75%), followed by diazepam (7%), and lorazepam (2%).² Seven years later, a follow-up survey study by Kain et al³ found that adult inpatients received midazolam most often (80%), followed by diazepam (3%), fentanyl (2%), and lorazepam (1%).

Lorazepam may have disadvantages compared with midazolam. Midazolam is a short-acting benzodiazepine with an elimination half-life of 2.0 hours to 2.5 hours, whereas lorazepam has a half-life of 10 hours to 20 hours. In a randomized clinical trial that compared midazolam with lorazepam for postoperative sedation, delays in emergence from sedation were observed in the lorazepam group.⁴

Tetsuji Fujita, MD

Author Affiliation: Jikei University School of Medicine, Tokyo, Japan.

Corresponding Author: Tetsuji Fujita, MD, Jikei University School of Medicine, 3-25-8 Nishishinbashi, Minato-ku, Tokyo, Minato-ku 105-8461, Japan (tetsu@jg8.so-net.ne.jp).

Conflict of Interest Disclosures: The author has completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

1. Maurice-Szamburski A, Auquier P, Viarre-Oreal V, et al; PremedX Study Investigators. Effect of sedative premedication on patient experience after general anesthesia: a randomized clinical trial. *JAMA*. 2015;313(9):916-925.
2. Kain ZN, Mayes LC, Bell C, Weisman S, Hofstadter MB, Rimar S. Premedication in the United States: a status report. *Anesth Analg*. 1997;84(2):427-432.
3. Kain ZN, Caldwell-Andrews AA, Krivutza DM, Weinberg ME, Wang SM, Gaal D. Trends in the practice of parental presence during induction of anesthesia and the use of preoperative sedative premedication in the United States, 1995-2002: results of a follow-up national survey. *Anesth Analg*. 2004;98(5):1252-1259.
4. Barr J, Zomorodi K, Bertaccini EJ, Shafer SL, Geller E. A double-blind, randomized comparison of IV lorazepam versus midazolam for sedation of ICU patients via a pharmacologic model. *Anesthesiology*. 2001;95(2):286-298.

To the Editor The key purpose of patient premedication with benzodiazepines is preoperative anxiolysis,¹ which allows for safer monitoring and induction of anesthesia. In contrast, the primary outcome parameter of the study by Dr Maurice-Szamburski and colleagues² was the patient's perception quantified with a patient satisfaction index (Evaluation du Vécu de l'Anesthésie Générale; EVAN-G)³ on the first postoperative day. From a clinical viewpoint, a patient's perception on the first postoperative day is not the key parameter to judge the value of premedication.

The more important outcomes were the secondary outcomes measuring preoperative and intraoperative conditions. Anxiety upon arrival in the operating room was significantly lower (visual analog scale scores for anxiety: 35 points for lorazepam group vs

44 points for placebo group; $P = .001$). However, the scores for quality of conditioning and well-being upon arrival in the operating room were not statistically significant. The present study was underpowered to shed light on these clinically important and desired effects of premedication.

In the lorazepam group, the duration of surgery was 11 minutes longer than in the placebo group and postoperative pain was significantly lower. The statistically significant secondary outcome of a prolonged time to extubation can be attributed to lorazepam only if dosages of sedatives and opioids were comparable between the groups, which were not presented in the article.

Because many patients do not want to remember anything about the operation, amnesia is an intended effect of premedication. The increased proportion of patients with amnesia (24% with lorazepam vs 6% with placebo) is a beneficial effect of lorazepam rather than a complication as the authors suggested.

Lorazepam produced clinically desired effects in the study by Maurice-Szamburski and colleagues. The data do not justify the conclusion of the authors that lorazepam lacks any benefit in routine use.

Axel R. Heller, MD, MBA
Thea Koch, MD

Author Affiliations: Department of Anesthesiology and Intensive Care Medicine, Universitätsklinikum Carl Gustav Carus Dresden, Dresden, Germany.

Corresponding Author: Axel R. Heller, MD, MBA, Department of Anesthesiology and Intensive Care Medicine, University Hospital Carl Gustav Carus, Fetscherstraße 74, 01307 Dresden, Germany (axel.heller@uniklinikum-dresden.de).

Conflict of Interest Disclosures: The authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

1. Bishop S, Reddy S. Premedication. *Anaesth Intensive Care Med*. 2010;11(10):407-409.
2. Maurice-Szamburski A, Auquier P, Viarre-Oreal V, et al; PremedX Study Investigators. Effect of sedative premedication on patient experience after general anesthesia: a randomized clinical trial. *JAMA*. 2015;313(9):916-925.
3. Auquier P, Pernoud N, Bruder N, et al. Development and validation of a perioperative satisfaction questionnaire. *Anesthesiology*. 2005;102(6):1116-1123.

In Reply If midazolam remains the most prescribed sedative premedication, lorazepam ranks third for outpatient surgery and fourth for inpatient surgery.¹ Midazolam and lorazepam both belong to the benzodiazepine class and share most of their indications. The PremedX study evaluated sedative premedication with lorazepam given orally 2 hours before surgery. Lorazepam was chosen over the more commonly used midazolam because, due to its shorter acting time, midazolam may not have been as potent as lorazepam upon arrival in the operating room.

In the study by Barr et al cited by Dr Fujita, these 2 drugs were the most frequently prescribed drugs for sedation. However, one should be careful about extrapolating results from this intensive care unit study with continuous administration of benzodiazepine and fentanyl up to 72 hours. Although delays in emergence were longer in the lorazepam group, the lengths of administration differed (mean [SD], 36.94 [30.90] hours with midazolam vs 15.02 [3.33] hours with lorazepam).²

In response to Drs Heller and Koch, we believe the purpose of premedication is not established. According to anesthesiologists, there are many different reasons for using sedative premedication.³ Because sedative premedication primarily concerns patients and their experience,⁴ the main outcome of the PremedX study was the patient experience and satisfaction assessed by a validated questionnaire, evaluating both the preoperative, perioperative, and postoperative periods.⁵ The EVAN-G mean global index showed no significant differences between the lorazepam, placebo, and no premedication groups in the whole population or in the subgroup of the most anxious patients, those expected to benefit the most from sedative premedication.

Even though anxiety upon operating room arrival was lower with lorazepam than with placebo, it was no different between lorazepam and no premedication (35 [95% CI, 32-38] vs 38 [95% CI, 35-41], respectively, $P = .41$) and was higher comparing placebo with no premedication (44 [95% CI, 40-47] vs 38 [95% CI, 35-41], respectively, $P = .05$), suggesting a nocebo effect of placebo. In our study, anesthesiologists were unable to distinguish between the lorazepam, placebo, and no premedication groups for quality of patient conditioning. Before invoking underpowering to explain the lack of efficacy, sedative premedication relevance should be questioned.

The randomization of the study was intended to eliminate other sources of variability, such as the dosing of sedatives and opioids. The types of drugs used for the anesthesia protocol were not different between groups.

Lorazepam was associated with a 10-point lower satisfaction score on the attention dimension of the EVAN-G scale. One explanation is that the amnesia from benzodiazepine led patients to forget the attention they received from caregivers, which would not necessarily be beneficial.

Given the results of the PremedX study, we continue to believe that routine sedative premedication with benzodiazepine should not be recommended.

Axel Maurice-Szamburski, MD

Author Affiliation: Service d'Anesthésie Réanimation, Hôpital de la Timone Adulte, Marseille, France.

Corresponding Author: Axel Maurice-Szamburski, MD, Service d'Anesthésie Réanimation, Hôpital de la Timone Adulte, 264 Rue Saint Pierre, 13385 Marseille CEDEX 05, France (amszamburski@gmail.com).

Conflict of Interest Disclosures: The author has completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and reported receiving a grant from the French Institutional Clinical Hospital Research Program, Ministry of Health.

1. Kain ZN, Caldwell-Andrews AA, Krivutza DM, Weinberg ME, Wang SM, Gaal D. Trends in the practice of parental presence during induction of anesthesia and the use of preoperative sedative premedication in the United States, 1995-2002: results of a follow-up national survey. *Anesth Analg*. 2004;98(5):1252-1259.
2. Barr J, Zomorodi K, Bertaccini EJ, Shafer SL, Geller E. A double-blind, randomized comparison of IV lorazepam versus midazolam for sedation of ICU patients via a pharmacologic model. *Anesthesiology*. 2001;95(2):286-298.
3. Mirakhor RK. Preanaesthetic medication: a survey of current usage. *J R Soc Med*. 1991;84(8):481-483.
4. Vetter TR, Ivankova NV, Pittet JF. Patient satisfaction with anesthesia: beauty is in the eye of the consumer. *Anesthesiology*. 2013;119(2):245-247.
5. Auquier P, Pernoud N, Bruder N, et al. Development and validation of a perioperative satisfaction questionnaire. *Anesthesiology*. 2005;102(6):1116-1123.