

The Finsler geometry of the rotating Kepler problem

Kai Cieliebak, Urs Frauenfelder, Otto van Koert

Angaben zur Veröffentlichung / Publication details:

Cieliebak, Kai, Urs Frauenfelder, and Otto van Koert. 2014. "The Finsler geometry of the rotating Kepler problem." *Publicationes Mathematicae Debrecen* 84 (3-4): 333-50. <https://doi.org/10.5486/pmd.2014.5368>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under the following conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publizieren>



The Finsler geometry of the rotating Kepler problem

By KAI CIELIEBAK (Augsburg), URS FRAUENFELDER (Seoul)
and OTTO VAN KOERT (Seoul)

Abstract. We investigate the Cartan and Finsler geometry of the rotating Kepler problem, a limit case of the restricted three body problem that arises if the mass of the one of the primaries goes to zero. We show that the Hamiltonian for the rotating Kepler problem can be regarded as the Legendre transform of a certain family of Finsler metrics on the two-sphere. For very negative energy levels, these Finsler metrics are close to the round metric, and the associated flag curvature is hence positive. On the other hand, we show that the flag curvature can become negative once the energy level becomes sufficiently high.

1. Introduction

The Kepler problem in a rotating frame arises as the limit of the restricted three body problem when the mass of one of the primaries goes to zero. Energy hypersurfaces of the Kepler problem as well as the restricted three body problem are noncompact. If one focuses on the bounded components of the energy hypersurfaces, i.e. one excludes the possibility of escape to infinity, then the only obstruction to compactness comes from collisions. However, it is well known in celestial mechanics that two body collisions can always be regularized. A particularly nice way to regularize the Kepler problem in an inertial frame was discovered by MOSER in [10]. By interchanging the roles of position and momenta Moser

Mathematics Subject Classification: 53C60, 70F07, 53Dxx.

Key words and phrases: Finsler geometry, Flag curvature, 3-body problem.

The second author was partially supported by the Basic Research fund 2010-0007669 funded by the Korean government, and the third author by the NRF Grant 2012011755 funded by the Korean government. The second and third author also hold joint appointments in the Research Institute of Mathematics, Seoul National University.

shows that the regularized planar Kepler problem is equivalent to the geodesic flow on the round two-sphere. Indeed, the point at infinity of the two-sphere corresponds to collision where the momenta explode. In [1] it was shown that for energies not too large the compact components of the regularized energy hypersurfaces for the planar restricted 3-body problem are starshaped. Here starshaped refers to the new cotangent bundle structure obtained after interchanging the roles of position and momenta; with respect to the old (q, p) -coordinates the Liouville vector field is given by $q \frac{\partial}{\partial q}$ and not by $p \frac{\partial}{\partial p}$. The challenging new question is now if starshaped can be replaced by the stronger condition of fiberwise convexity. If the energy hypersurfaces are fiberwise convex we obtain a Finsler metric after Legendre transformation and the integral curves of the restricted 3-body problem can be interpreted as geodesics for the Finsler metric. The relevance of this problem comes from its relation with the project initiated by Helmut Hofer to construct finite energy foliations for the restricted 3-body problem. Indeed, if the energy hypersurfaces are fiberwise convex then the Conley–Zehnder indices of the closed characteristics correspond to the Morse indices of the corresponding geodesics and hence are nonnegative. This idea can be used directly for the magnetic Hamiltonian of the unregularized problem. To treat collision orbits as well, we consider the regularized problem. Our first result is the following.

Theorem A. The bounded components of the regularized planar Kepler problem in a rotating frame are fiberwise convex.

In particular, the regularized Hamiltonian of the rotating Kepler problem is Legendre dual to a Finsler metric. The Legendre dual of a Finsler space was recognized by MIRON as a *Cartan space*, see [11] and the literature cited therein. Therefore, Theorem A provides the rotating Kepler problem with a Cartan structure. The sectional curvature of Riemannian geometry generalizes to the notion of flag curvature in Cartan, respectively Finsler geometry. In particular, for a Cartan surface the flag curvature is a function on the unit tangent bundle of the surface. This differs from the Riemannian case where the sectional curvature for a surface reduces to the Gaussian curvature and hence is given by a function on the surface itself. Since in our case the underlying surface is a sphere, the flag curvature is a function on $\mathbb{R}P^3$ regarded as the unit tangent bundle of the sphere. However, because the Kepler problem is invariant under rotations, the flag curvature is invariant under a circle action and quotienting out this circle action we again get a function on a sphere. However, be aware that this sphere does not coincide with the original one. The importance of the flag curvature comes from its relation with the Jacobi equation and hence with questions about

indices, see [3]. Our next result is rather surprising since by Moser's result in an inertial frame, the regularized Kepler problem is equivalent to the geodesic flow of the round two sphere whose curvature is hence always positive.

Observation B. *For the rotating Kepler problem there are regions in which the flag curvature is negative.*

Let us explain our motivation for studying this problem. As mentioned above, the rotating Kepler problem is a limiting case of the restricted 3-body problem. Conley first proposed to use hyperbolicity in the restricted 3-body problem in mission design, [6], [7], [8]. Suppose we have the following problem. Given a satellite on an orbit around the earth, is it possible to move this satellite to an orbit around the moon using the gravitational forces of the earth, moon and maybe the sun without spending much fuel. It was BELBRUNO who found the first realistic orbit using this idea, [4]. A variation of this orbit was successfully put into practice to save the first Japanese lunar mission, [5].

Conley's idea takes advantage of a hyperbolic orbit around the first Lagrange point, also known as a Lyapunov orbit. These Lyapunov orbits only appear for sufficiently high energy.

During a discussion with P. Albers, E. Belbruno, J. Fish and H. Hofer at the IAS in Princeton, the question arose whether (short) hyperbolic orbits can already arise below the first Lagrange point. For very small masses of the moon or for very negative energies it is known that this phenomenon never happens. Otherwise this question is completely open.

A possibility to locate the position of hyperbolic orbits in phase space would be to find regions of negative curvature of the Finsler metric. Indeed, [9] contains a short computation which illustrates the relation between the flag curvature and the Conley–Zehnder index. In particular, one can clearly see that if the flag curvature is negative along an entire orbit, then this orbit is hyperbolic. We therefore hope that the methods studied in this paper have some applications for mission design.

We conclude the introduction by pointing out that one could also try to understand the Conley–Zehnder indices by investigating whether there exists a convex embedding of a level set of the restricted 3-body problem into \mathbb{C}^2 . The latter question has been answered in the affirmative in [2] provided the energy levels are sufficiently far below the first Lagrange value.

2. A family of Minkowski metrics

For $p = (p_1, p_2) \in \mathbb{R}^2$ and $C > 0$ set $p^\perp = (p_2, -p_1)$ and consider the function $H_p : \mathbb{R}^2 - 0 \rightarrow \mathbb{R}$,

$$H_p(q) := \langle p^\perp, q \rangle - \frac{1}{|q|} + 2C.$$

Lemma 2.1. *Suppose $|p| < C^2$. Then the curve $H_p^{-1}(0)$ has two connected components: an unbounded one in the region $\{|q| > 1/C\}$, and a bounded component Σ_p in the region $\{|q| < 1/C\}$ which bounds a strictly convex domain containing the origin.*

PROOF. Given $q \neq 0$, let us determine all $\lambda \in \mathbb{R} - 0$ such that $H_p(\lambda^{-1}q) = 0$, i.e.

$$|\lambda|\lambda - 2C|q|\lambda - |q|\langle p^\perp, q \rangle = 0.$$

Suppose without loss of generality that $\langle p^\perp, q \rangle \geq 0$. Then the equation has the 3 solutions

$$\lambda_0 = C|q| \left(1 + \sqrt{1 + \frac{\langle p^\perp, q \rangle}{|q|C^2}} \right) > 0, \quad \lambda_\pm = -C|q| \left(1 \pm \sqrt{1 - \frac{\langle p^\perp, q \rangle}{|q|C^2}} \right) \leq 0.$$

By hypothesis we have

$$0 \leq x = \frac{\langle p^\perp, q \rangle}{|q|C^2} < 1,$$

and thus $1 - \sqrt{1-x} \leq 1 - (1-x) < 1$. We see that $|\lambda_0|, |\lambda_+| > C|q|$ and $|\lambda_-| < C|q|$. Thus $\lambda_0^{-1}q$ and $\lambda_+^{-1}q$ belong to the bounded component contained in $\{|q| < 1/C\}$ and $\lambda_-^{-1}q$ to the unbounded component contained in $\{|q| > 1/C\}$.

For convexity we compute the gradient and Hessian

$$\nabla H_p(q) = \begin{pmatrix} p_2 + \frac{q_1}{|q|^3} \\ -p_1 + \frac{q_2}{|q|^3} \end{pmatrix}, \quad \text{Hess } H_p(q) = \frac{1}{|q|^5} \begin{pmatrix} |q|^2 - 3q_1^2 & -3q_1q_2 \\ -3q_1q_2 & |q|^2 - 3q_2^2 \end{pmatrix}.$$

A tangent vector to $H_p^{-1}(0)$ is given by

$$v := -(\nabla H_p(q))^\perp = \begin{pmatrix} p_1 - \frac{q_2}{|q|^3} \\ p_2 + \frac{q_1}{|q|^3} \end{pmatrix}.$$

A short computation yields

$$\begin{aligned} |q|^5 \langle v, \text{Hess } H_p(q)v \rangle &= (|q|^2 - 3q_1^2) \left(p_1 - \frac{q_2}{|q|^3} \right)^2 - 6q_1q_2 \left(p_2 + \frac{q_1}{|q|^3} \right) \left(p_1 - \frac{q_2}{|q|^3} \right) \\ &\quad + (|q|^2 - 3q_2^2) \left(p_2 + \frac{q_1}{|q|^3} \right)^2 = |q|^{-2} \left(|p|^2|q|^4 + 2|q|\langle p^\perp, q \rangle + 1 - 3|q|^2\langle p, q \rangle^2 \right) \end{aligned}$$

To show positivity of the right hand side, we write $\langle p^\perp, q \rangle = |p| |q| \cos t$, $\langle p, q \rangle = |p| |q| \sin t$. Then the right hand side equals $|q|^{-2} f(t)$ with the periodic function

$$f(t) := a^2 + 2a \cos t + 1 - 3a^2 \sin^2 t, \quad a := |p| |q|^2.$$

The derivative $f'(t) = -2a \sin t(1 + 3a \cos t)$ vanishes iff $\sin t = 0$ or $\cos t = -1/(3a)$. In the case $\sin t = 0$ we have $f(t) = (1 - a)^2 > 0$, since $0 < a < 1$ due to the conditions $|p| < C^2$ and $|q| < 1/C$. In the case $\cos t = -1/(3a)$ we have

$$f(t) = a^2 - \frac{2a}{3a} + 1 - 3a^2 \left(1 - \frac{1}{(3a)^2}\right) = 2 \left(\frac{1}{3} - a^2\right).$$

The equation $H_p(q) = 0$ yields

$$\frac{1}{|q|} - 2C = \langle p^\perp, q \rangle = |p| |q| \cos t = -\frac{|p| |q|}{3a} = -\frac{1}{3|q|},$$

thus $|q| = 2/(3C)$. Using this and $|p| < C^2$, we estimate

$$a = |p| |q|^2 = \frac{4|p|}{9C^2} < \frac{4}{9}.$$

Hence $a^2 < 1/3$, which shows $f(t) > 0$ and concludes the proof of the lemma. \square

The convex curve Σ_p endows $T_p^* \mathbb{R}^2$ with the structure of a Cartan space with fundamental function $F_p^* : \mathbb{R}^2 \rightarrow \mathbb{R}_{\geq 0}$ in the sense of [11]: $F_p^*(q)$ is the unique $\lambda > 0$ such that $\lambda^{-1}q \in \Sigma_p$. This λ has been computed in the preceding proof to be

$$F_p^*(q) = C|q| \left(1 + \sqrt{1 + \frac{\langle p^\perp, q \rangle}{|q|C^2}}\right).$$

This Cartan structure is non-reversible (i.e. $F_p^*(-q) \neq F_p^*(q)$) for $p \neq 0$ and does not appear to belong to one of the standard classes (Randers, Berwald, ...).

3. Kepler problem in a rotating frame

Consider the Kepler problem in a frame that is rotating around the origin with angular velocity $a > 0$. This system is again Hamiltonian with Hamiltonian

$$H(p, q) = a(p_2q_1 - p_1q_2) - \frac{1}{|q|} + \frac{|p|^2}{2} = \frac{1}{2}((p_1 - aq_2)^2 + (p_2 + aq_1)^2) + U(q),$$

with the potential

$$U(q) = -\frac{1}{|q|} - \frac{a^2}{2}|q|^2.$$

The function $U(r) = -\frac{1}{r} - \frac{a^2}{2}r^2$ attains its maximum $-\frac{3}{2}a^{2/3}$ at the point $r = a^{-2/3}$. So the energy level

$$\{H(p, q) = -c\}, \quad c > \frac{3}{2}a^{2/3}$$

has an unbounded component and a bounded component Σ on which we have

$$|q| \leq a^{-2/3}.$$

The function

$$H_p(q) = H(p, q) + c$$

is of the type considered above with p replaced by ap and $2C = |p|^2/2 + c$. Let us verify the hypothesis of Lemma 2.1,

$$a|p| < C^2 = \left(\frac{|p|^2}{4} + \frac{c}{2}\right)^2.$$

Using $c > \frac{3}{2}a^{2/3}$ and setting $x := |p|$, it suffices to show

$$16ax \leq (x^2 + 3a^{2/3})^2 = x^4 + 6x^2a^{2/3} + 9a^{4/3}.$$

Replacing x by $xa^{1/3}$, this is equivalent to non-negativity of the function

$$g(x) = x^4 + 6x^2 - 16x + 9.$$

Since g attains its minimum 0 at $x = 1$, the hypotheses of the lemma are satisfied. Hence the level set Σ is convex in q for each fixed p . Adding $p = \infty$, Σ gives a fiberwise convex hypersurface in T^*S^2 , where the coordinate on S^2 is p and the fiber coordinate is q . Let $F^* : T^*S^2 \rightarrow \mathbb{R}$ be the fundamental function of the corresponding Cartan space, which we computed above to be

$$F^*(p, q) = \frac{1}{4}(|p|^2 + 2c)|q| \left(1 + \sqrt{1 + \frac{16a\langle p^\perp, q \rangle}{|q|(|p|^2 + 2c)^2}}\right).$$

The fiberwise Legendre transform of F^* (via $\frac{1}{2}F^{*2}$) yields a Finsler metric $F : TS^2 \rightarrow \mathbb{R}$ whose geodesic flow equals the Hamiltonian flow of H on the energy level Σ .

Note that H , and thus also F , is invariant under the rotation of S^2 around the vertical axis.

4. Polar coordinates

To take advantage of the symmetry of F^* it is useful to rewrite it in polar coordinates (x, y) defined by

$$p_1 = x \cos y, \quad p_2 = x \sin y.$$

The dual coordinates (r, t) on the cotangent bundle are determined by

$$r dx + t dy = q_1 dp_1 + q_2 dp_2.$$

In particular,

$$q_1 = (\cos y)r - \frac{\sin y}{x}t, \quad q_2 = (\sin y)r + \frac{\cos y}{x}t$$

and hence

$$|q|^2 = r^2 + \frac{1}{x^2}t^2, \quad \langle p^\perp, q \rangle = -t, \quad |p|^2 = x^2.$$

Hence in polar coordinates the fundamental function reads

$$F^*(x, y, r, t) = \frac{1}{4}(x^2 + 2c)\sqrt{r^2 + \frac{1}{x^2}t^2} \left(1 + \sqrt{1 - \frac{16at}{\sqrt{r^2 + \frac{1}{x^2}t^2}(x^2 + 2c)^2}} \right).$$

Note that F^* does not explicitly depend on y .

4.1. A one-parameter family of Cartan metrics. The above formula for the Cartan metrics F^* describes a two-parameter family with parameters c and a . However, each such Cartan metric is a constant multiple of one with a -parameter equal to 1.

Proposition 4.1. $F_{c,a}^* = a^{1/3}F_{ca^{-2/3},1}^*$.

PROOF. Use the coordinate transformation $x = a^{1/3}x'$, $y = y'$. On the cotangent fibers, this coordinate transformation induces $r = a^{-1/3}r'$ and $t = t'$. We get

$$F_{c,a}^*(x, y, r, t) = \frac{1}{4}(a^{2/3}x'^2 + 2c)\sqrt{r'^2 a^{-2/3} + \frac{1}{x'^2 a^{2/3}}t'^2} \cdot \left(1 + \sqrt{1 - \frac{16at'}{\sqrt{r'^2 a^{-2/3} + \frac{1}{x'^2 a^{2/3}}t'^2}(a^{2/3}x'^2 + 2c)^2}} \right)$$

$$\begin{aligned}
&= a^{1/3} \frac{1}{4} (x'^2 + 2ca^{-2/3}) \sqrt{r'^2 + \frac{1}{x'^2} t'^2} \\
&\quad \cdot \left(1 + \sqrt{1 - \frac{16t'}{\sqrt{r'^2 + \frac{1}{x'^2} t'^2} (x'^2 + 2ca^{-2/3})^2}} \right) \\
&= a^{1/3} F_{ca^{-2/3}, 1}^*(x', y', r', t'). \quad \square
\end{aligned}$$

We shall henceforth put $a = 1$.

5. Spray coefficients

An elegant way of computing the flag curvature of a Finsler surface is via spray coefficients, see [3, Section 12.5]. Since we are on the Cartan side we first have to Legendre transform the fundamental function F^* , in order to apply the formulas in [3] directly. However, we do not know an explicit description of the Legendre transform of F^* . We therefore derive in this section a formula to obtain the spray coefficients of a Cartan surface directly from the fundamental function. This allows us in turn to implement a MAPLE algorithm for computing the flag curvature of a Cartan surface. We abbreviate

$$\mathcal{L}^* = \frac{1}{2} F^{*2}$$

and denote by \mathcal{L} the Legendre dual of \mathcal{L}^* . The coordinates on the tangent bundle are given by

$$\begin{aligned}
u(x, y, r, t) &= \mathcal{L}_r^*(x, y, r, t) \\
v(x, y, r, t) &= \mathcal{L}_t^*(x, y, r, t).
\end{aligned}$$

Here a subscript denotes differentiation with respect to the corresponding variable. We first recall some useful facts of the Legendre transformation for which we refer to [3, Section 14.8] or [11]. Since \mathcal{L}^* and \mathcal{L} are 2-homogeneous they numerically coincide

$$\mathcal{L}^*(x, y, r, t) = \mathcal{L}(x, y, u(x, y, r, t), v(x, y, r, t)). \quad (1)$$

The cometric coefficients are defined by

$$g^{11} = \mathcal{L}_{rr}^*, \quad g^{12} = g^{21} = \mathcal{L}_{tr}^*, \quad g^{22} = \mathcal{L}_{tt}^*. \quad (2)$$

The metric coefficients inverse to the cometric coefficients can be obtained by

$$g_{11} = \mathcal{L}_{uu}, \quad g_{12} = g_{21} = \mathcal{L}_{uv}, \quad g_{22} = \mathcal{L}_{vv}.$$

Moreover, since the Legendre transform is involutive, one has the identities

$$r = \mathcal{L}_u, \quad t = \mathcal{L}_v.$$

We further abbreviate the determinant of the cometric by

$$g = g^{11}g^{22} - (g^{12})^2.$$

Differentiating (1) we obtain

$$\frac{\partial \mathcal{L}^*}{\partial x} = \frac{d\mathcal{L}}{dx} = \frac{\partial \mathcal{L}}{\partial x} + \frac{\partial \mathcal{L}}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial \mathcal{L}}{\partial v} \frac{\partial v}{\partial x}$$

and therefore

$$\mathcal{L}_x = \mathcal{L}_x^* - \mathcal{L}_u \mathcal{L}_{rx}^* - \mathcal{L}_v \mathcal{L}_{tx}^*.$$

Similarly

$$\mathcal{L}_y = \mathcal{L}_y^* - \mathcal{L}_u \mathcal{L}_{ry}^* - \mathcal{L}_v \mathcal{L}_{ty}^*.$$

From the identity

$$0 = \frac{\partial r}{\partial x} = \frac{\partial \mathcal{L}_u}{\partial x} + \frac{\partial \mathcal{L}_u}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial \mathcal{L}_u}{\partial v} \frac{\partial v}{\partial x}$$

one obtains

$$\mathcal{L}_{ux} = -\mathcal{L}_{uu} \mathcal{L}_{rx}^* - \mathcal{L}_{uv} \mathcal{L}_{tx}^* = -g_{11} \mathcal{L}_{rx}^* - g_{12} \mathcal{L}_{tx}^* = \frac{1}{g} \left(-g^{22} \mathcal{L}_{rx}^* + g^{12} \mathcal{L}_{tx}^* \right)$$

In a similar vein, one derives the formulas

$$\mathcal{L}_{uy} = -\mathcal{L}_{uu} \mathcal{L}_{ry}^* - \mathcal{L}_{uv} \mathcal{L}_{ty}^* = \frac{1}{g} \left(-g^{22} \mathcal{L}_{ry}^* + g^{12} \mathcal{L}_{ty}^* \right)$$

$$\mathcal{L}_{vx} = -\mathcal{L}_{vu} \mathcal{L}_{rx}^* - \mathcal{L}_{vv} \mathcal{L}_{tx}^* = \frac{1}{g} \left(g^{12} \mathcal{L}_{rx}^* - g^{11} \mathcal{L}_{tx}^* \right)$$

$$\mathcal{L}_{vy} = -\mathcal{L}_{vu} \mathcal{L}_{ry}^* - \mathcal{L}_{vv} \mathcal{L}_{ty}^* = \frac{1}{g} \left(g^{12} \mathcal{L}_{ry}^* - g^{11} \mathcal{L}_{ty}^* \right).$$

Plugging these identities into the formulas for the spray coefficients G , H (this H is not the Hamiltonian!) in [3, p 330] we get

$$\begin{aligned} 2G &= g\mathcal{L}_{vv}\mathcal{L}_x - g\mathcal{L}_{vx}\mathcal{L}_v - g\mathcal{L}_{uv}\mathcal{L}_y + g\mathcal{L}_{uy}\mathcal{L}_v \\ &= g^{11}(\mathcal{L}_x^* - \mathcal{L}_u\mathcal{L}_{rx}^* - \mathcal{L}_v\mathcal{L}_{tx}^*) - (g^{12}\mathcal{L}_{rx}^* - g^{11}\mathcal{L}_{tx}^*)\mathcal{L}_v \\ &\quad + g^{12}(\mathcal{L}_y^* - \mathcal{L}_u\mathcal{L}_{ry}^* - \mathcal{L}_v\mathcal{L}_{ty}^*) + (-g^{22}\mathcal{L}_{ry}^* + g^{12}\mathcal{L}_{ty}^*)\mathcal{L}_v \\ &= (g^{11}\mathcal{L}_x^* + g^{12}\mathcal{L}_y^*) - (g^{11}\mathcal{L}_{rx}^* + g^{12}\mathcal{L}_{ry}^*)\mathcal{L}_u \end{aligned}$$

$$\begin{aligned}
 & + (g^{11}\mathcal{L}_{tx}^* - g^{12}\mathcal{L}_{rx}^* - g^{11}\mathcal{L}_{tx}^* - g^{12}\mathcal{L}_{ty}^* - g^{22}\mathcal{L}_{ry}^* + g^{12}\mathcal{L}_{ty}^*)\mathcal{L}_v \\
 & = (g^{11}\mathcal{L}_x^* + g^{12}\mathcal{L}_y^*) - (g^{11}\mathcal{L}_{rx}^* + g^{12}\mathcal{L}_{ry}^*)\mathcal{L}_u - (g^{12}\mathcal{L}_{rx}^* + g^{22}\mathcal{L}_{ry}^*)\mathcal{L}_v \\
 2H & = g\mathcal{L}_{uu}\mathcal{L}_y - g\mathcal{L}_{uy}\mathcal{L}_u - g\mathcal{L}_{vu}\mathcal{L}_x + g\mathcal{L}_{vx}\mathcal{L}_u \\
 & = g^{22}(\mathcal{L}_y^* - \mathcal{L}_u\mathcal{L}_{ry}^* - \mathcal{L}_v\mathcal{L}_{ty}^*) - (-g^{22}\mathcal{L}_{ry}^* + g^{12}\mathcal{L}_{ty}^*)\mathcal{L}_u \\
 & \quad + g^{12}(\mathcal{L}_x^* - \mathcal{L}_u\mathcal{L}_{rx}^* - \mathcal{L}_v\mathcal{L}_{tx}^*) + (g^{12}\mathcal{L}_{rx}^* - g^{11}\mathcal{L}_{tx}^*)\mathcal{L}_u \\
 & = (g^{12}\mathcal{L}_x^* + g^{22}\mathcal{L}_y^*) - (g^{12}\mathcal{L}_{tx}^* + g^{22}\mathcal{L}_{ty}^*)\mathcal{L}_v \\
 & \quad + (g^{22}\mathcal{L}_{ry}^* - g^{12}\mathcal{L}_{ty}^* - g^{22}\mathcal{L}_{ry}^* - g^{12}\mathcal{L}_{rx}^* + g^{12}\mathcal{L}_{rx}^* - g^{11}\mathcal{L}_{tx}^*)\mathcal{L}_u \\
 & = (g^{12}\mathcal{L}_x^* + g^{22}\mathcal{L}_y^*) - (g^{12}\mathcal{L}_{tx}^* + g^{22}\mathcal{L}_{ty}^*)\mathcal{L}_v - (g^{11}\mathcal{L}_{tx}^* + g^{12}\mathcal{L}_{ty}^*)\mathcal{L}_u
 \end{aligned}$$

5.1. A formula for the flag curvature following Bao–Chern–Shen. We shall use formula (12.5.14) from [3] to compute the flag curvature. This formula is given by

$$K = \frac{(G_{xv} - G_{yu})v + 2GG_{uu} + 2HG_{uv} - G_uG_u - G_vH_u}{v\mathcal{L}_v}. \tag{3}$$

All these terms can be rewritten entirely into terms that exist on the Cartan side. In particular, we shall use the above formulas for G and H in terms of g^{ij} and \mathcal{L}^* and their derivatives.

As before, we define the Hamiltonian or Legendre dual of \mathcal{L} by

$$\mathcal{L}^* = \frac{1}{2}F^{*2}$$

The cometric can be obtained via Equation (2). As the Hamiltonian is given in terms of the base coordinates x, y and cotangent fiber coordinates r, t , we need to apply the chain rule to compute (3), which is given in terms of x, y and the tangent fiber coordinates u, v . We need to use the following relations,

$$\frac{\partial}{\partial u} = \frac{\partial r}{\partial u} \frac{\partial}{\partial r} + \frac{\partial t}{\partial u} \frac{\partial}{\partial t} = g_{11} \frac{\partial}{\partial r} + g_{21} \frac{\partial}{\partial t}.$$

Similarly,

$$\frac{\partial}{\partial v} = g_{12} \frac{\partial}{\partial r} + g_{22} \frac{\partial}{\partial t}.$$

Note that the metric can be expressed in terms of the cometric and its determinant. The dependence on the base coordinate is a little tricky, since the independent variables are now x, y, u, v , whereas the Hamiltonian is given in terms x, y, r, t . This means that the (co)-metric comes in. Let us write out the required expression. We shall use the notation

$$\left(\frac{\partial G}{\partial x} \right)_{y,u,v}$$

to indicate the derivative of G keeping y, u, v fixed. We obtain

$$\begin{aligned} \left(\frac{\partial G}{\partial x}\right)_{y,u,v} &= \left(\frac{\partial G}{\partial x}\right)_{y,r,t} + \left(\frac{\partial G}{\partial r}\right)_{x,y,t} \left(\frac{\partial r}{\partial x}\right)_{y,u,v} + \left(\frac{\partial G}{\partial t}\right)_{x,y,r} \left(\frac{\partial t}{\partial x}\right)_{y,u,v} \\ &= \left(\frac{\partial G}{\partial x}\right)_{y,r,t} + \left(\frac{\partial G}{\partial r}\right)_{x,y,t} \left(\left(\frac{\partial g_{11}}{\partial x}\right)_{y,u,v} u + \left(\frac{\partial g_{12}}{\partial x}\right)_{y,u,v} v \right) \\ &\quad + \left(\frac{\partial G}{\partial t}\right)_{x,y,r} \left(\left(\frac{\partial g_{21}}{\partial x}\right)_{y,u,v} u + \left(\frac{\partial g_{22}}{\partial x}\right)_{y,u,v} v \right). \end{aligned}$$

Similarly, we have

$$\begin{aligned} \left(\frac{\partial G}{\partial y}\right)_{x,u,v} &= \left(\frac{\partial G}{\partial y}\right)_{x,r,t} + \left(\frac{\partial G}{\partial r}\right)_{x,y,t} \left(\left(\frac{\partial g_{11}}{\partial y}\right)_{x,u,v} u + \left(\frac{\partial g_{12}}{\partial y}\right)_{x,u,v} v \right) \\ &\quad + \left(\frac{\partial G}{\partial t}\right)_{x,y,r} \left(\left(\frac{\partial g_{21}}{\partial y}\right)_{x,u,v} u + \left(\frac{\partial g_{22}}{\partial y}\right)_{x,u,v} v \right). \end{aligned}$$

In the above we have used

$$r = g_{11}u + g_{12}v \text{ and } t = g_{21}u + g_{22}v.$$

Using the cometric g^{ij} , we can conversely express u and v in terms of r and t . Hence we can translate all derivatives in the expression for the flag curvature in terms of quantities on the Cartan side.

6. Computing the flag curvature of the rotating Kepler problem

We now come to the computer implementation of the curvature computation. The MAPLE-program is included in the appendix. Although the program itself is straightforward with the above formulas, there are some practical issues involved which we shall briefly discuss.

Even though our computer setup was sufficient to compute the full expression for the flag curvature of the Cartan metric $F_{c,1}^*$ as a function of x, r, t , the resulting expression was too unwieldy to simplify to reasonable proportions. However, it is possible to evaluate and simplify the curvature exactly in specific points, for instance if the r coordinate is 0. The simplified expression for the curvature along this ray is still very long, see Equation (4).

In order to make these evaluations work, one only needs to realize that it is beneficial for a computer to quickly evaluate the results as soon as further differentiation is not needed. This is the reason for the many substitutions in the MAPLE program.

6.1. Analytical and numerical results. We fix the fiber coordinate $(r, t) = (0, x)$ and compute the curvature as a function of x , which is the coordinate on the base sphere S^2 . The MAPLE program yields the following expression for the flag curvature. Write

$$\alpha := \sqrt{x^4 + 4x^2c + 4c^2 - 16x}$$

to simplify the resulting expression.

$$\begin{aligned} & K(F_{c,1}^*)(x, 0, 0, x) \\ &= \frac{2}{(x^2 + 2c + \alpha)(x^2\alpha + 2c\alpha + x^4 + 4x^2c + 4c^2 - 8x)(x^4 + 4x^2c + 4c^2 - 16x)^2} \\ & \cdot (5824x^2c^4 - 5888x^3c^5 - 3840x^2c - 2240c^6x - 6320x^5c^4 - 384x^2\alpha + 1120x^6c^5 \\ & + 2x^{14}c + 28x^{12}c^2 - 6528x^5c + 256c^8 - 864\alpha c^5x - 1872\alpha c^4x^3 + 896x^2c^7 \\ & - 1296x^7 + 204x^{10} - 768c^5 - 9x^{13} + 2096x^8c - 160x^{11}c - 1060c^2x^9 \\ & - 3520c^3x^7 + 11520x^4c^3 + 3840c^3x + 7584x^6c^2 - 5952x^3c^2 - 648x^7\alpha c^2 \\ & - 126x^9\alpha c - 1120x^3\alpha c + 2448x^4\alpha c^2 + 1152\alpha c^2x + 1920x^4 + 168x^{10}c^3 \\ & + 1344x^4c^6 + 560x^8c^4 + 1032x^6\alpha c - 1584x^5\alpha c^3 + 1632\alpha c^3x^2 + 128\alpha c^7 \\ & + 384\alpha c^6x^2 - 9x^{11}\alpha + 2x^{12}\alpha c + 132x^8\alpha + 320x^6\alpha c^4 + 120x^8\alpha c^3 + 24x^{10}\alpha c^2 \\ & + 480\alpha c^5x^4 - 528\alpha x^5 - 384\alpha c^4) \end{aligned} \quad (4)$$

By plotting graphs we see that the curvature can become negative for values of c close to $\frac{3}{2}$. See the arxiv version of our paper for such a graph, [arXiv:1110.1021](https://arxiv.org/abs/1110.1021)

Alternatively, we can evaluate the expression for the flag curvature at suitable points. For instance, putting $\alpha = 1$, we find

$$K(F_{1.55,1}^*)(1/2, 0, 0, 1/2) \sim -2.85 < 0$$

This gives a proof of Observation B. We remind the reader that the Finsler metric of the rotating Kepler problem is close to the round metric for large values of c , and hence the flag curvature is then strictly positive.

Finally we consider the flag curvature as a function on the unit cotangent bundle. Because of the S^1 -symmetry, it suffices to use the coordinates x, ϕ , where the cotangent fiber coordinates are given by

$$(r, t) = (\sin \phi, \cos \phi).$$

The full expression for the curvature for $r \neq 0$ is considerably more complicated

than Equation (4), so we shall only present numerical results for this general case.

Using MAPLE we made a graph of the flag curvature by evaluating the curvature on a 256×256 lattice. See the arxiv version of our paper for this graph. On this lattice the minimal value of the flag curvature for $c = 1.55$ was -5.55 , and the maximal value 15.21.

7. Appendix

Here is a general program to (numerically) compute the flag curvature of a Cartan metric.

```

> K:=proc(F,x0,y0,r0,t0)
> local L,Lx,Ly,Lr,Lt,Lxt,Lyr,Lrr,Lrt,Ltt,D,
>       G,H,Gr,Gt,Gar,Gat,Grr,Grt,Gxt,Gyr,Hr,K,
>       Garre,Garte,Grre,Grte,Gxte,Gyre,Hre,Gre,Gte,
>       Ht,Hrr,Hrt,Htt,detg,gi,g,gie,ge,
>       Lsx,Lsy,Lsq1x,Lsq1y,Lsq2x,Lsq2y:
>
> ## Hamiltonian given
> H:=F^2/2:
>
> Hr:=diff(H,r):
> Ht:=diff(H,t):
> Hrr:=diff(Hr,r):
> Hrt:=diff(Hr,t):
> Htt:=diff(Ht,t):
>
> gi[1,1]:=Hrr: gi[1,2]:=Hrt: gi[2,1]:=Hrt: gi[2,2]:=Htt:
> detg:=gi[1,1]*gi[2,2]-gi[1,2]^2:
>
> ##obtain Finsler 'metric' g_{ij}: note that the dependence on
> ##the fiber coordinates is not correct, since we haven't
> ##performed a Legendre transformation
> g[1,1]:=1/detg*gi[2,2]: g[1,2]:=-1/detg*gi[1,2]:
> g[2,1]:=-1/detg*gi[2,1]: g[2,2]:=1/detg*gi[1,1]:
>
> Lr:=r:
> Lt:=t:
>
> Lsx:=diff(H,x): Lsy:=diff(H,y):
> Lsq1x:=diff(Hr,x): Lsq1y:=diff(Hr,y):
> Lsq2x:=diff(Ht,x): Lsq2y:=diff(Ht,y):

```

```

>
> ##Formulas from Section 5
> G:= 1/2*( gi[1,1]*Lsx+gi[1,2]*Lsy-Lr*(gi[1,1]*Lsq1x
>         +gi[1,2]*Lsq1y)-Lt*(gi[1,2]*Lsq1x+gi[2,2]*Lsq1y) ):
> H:=1/2*(gi[1,2]*Lsx+gi[2,2]*Lsy-Lt*(gi[1,2]*Lsq2x
>         +gi[2,2]*Lsq2y)-Lr*(gi[1,1]*Lsq2x+gi[1,2]*Lsq2y) ):
>
> ##avoid double computations
> Gar:=diff(G,r): Gat:=diff(G,t):
>
> Gr:=Gar*g[1,1]+Gat*g[2,1]:
> Gt:=Gar*g[1,2]+Gat*g[2,2]:
>
> ##evaluate quickly
>
> ge[1,1]:=subs(x=x0,y=y0,r=r0,t=t0,g[1,1]):
> ge[1,2]:=subs(x=x0,y=y0,r=r0,t=t0,g[1,2]):
> ge[2,1]:=subs(x=x0,y=y0,r=r0,t=t0,g[2,1]):
> ge[2,2]:=subs(x=x0,y=y0,r=r0,t=t0,g[2,2]):
> gie[1,1]:=subs(x=x0,y=y0,r=r0,t=t0,gi[1,1]):
> gie[1,2]:=subs(x=x0,y=y0,r=r0,t=t0,gi[1,2]):
> gie[2,1]:=subs(x=x0,y=y0,r=r0,t=t0,gi[2,1]):
> gie[2,2]:=subs(x=x0,y=y0,r=r0,t=t0,gi[2,2]):
>
> ##avoid double computations
> Garre:=subs(x=x0,y=y0,r=r0,t=t0,diff(Gr,r) ):
> Garte:=subs(x=x0,y=y0,r=r0,t=t0, diff(Gr,t) ):
>
> ##evaluate when no more differentiation is needed.
> Grre:=Garre*ge[1,1]+Garte*ge[2,1] :
> Grte:=Garre*ge[1,2]+Garte*ge[2,2] :
>
>
> Gxte:=subs(x=x0,y=y0,r=r0,t=t0, diff(Gt,x) )
>       +subs(x=x0,y=y0,r=r0,t=t0, diff(Gt,r) )
>       *subs(x=x0,y=y0,r=r0,t=t0,( diff(g[1,1],x)
>       *(gie[1,1]*r+gie[1,2]*t)+diff(g[1,2],x)*(gie[2,1]*r
>       +gie[2,2]*t) ) ) +subs(x=x0,y=y0,r=r0,t=t0, diff(Gt,t) )
>       *subs(x=x0,y=y0,r=r0,t=t0,( diff(g[2,1],x)
>       *(gie[1,1]*r+gie[1,2]*t)+diff(g[2,2],x)*(gie[2,1]*r
>       +gie[2,2]*t) ) ):
>
> Gyre:=subs(x=x0,y=y0,r=r0,t=t0,diff(Gr,y) )

```

```

> +subs(x=x0,y=y0,r=r0,t=t0,diff(Gr,r) )
> *subs(x=x0,y=y0,r=r0,t=t0,( diff(g[1,1],y)
> *(gie[1,1]*r+gie[1,2]*t)+diff(g[1,2],y)*(gie[2,1]*r
> +gie[2,2]*t) ) )+subs(x=x0,y=y0,r=r0,t=t0,diff(Gr,t) )
> *subs(x=x0,y=y0,r=r0,t=t0,( diff(g[2,1],y)
> *(gie[1,1]*r+gie[1,2]*t)+diff(g[2,2],y)*(gie[2,1]*r
> +gie[2,2]*t) ) ) :
>
> Hre:=subs(x=x0,y=y0,r=r0,t=t0,diff(H,r) )*ge[1,1]
> +subs(x=x0,y=y0,r=r0,t=t0,diff(H,t) )*ge[2,1] :
>
>
> Gre:=subs(x=x0,y=y0,r=r0,t=t0,Gr):
> Gte:=subs(x=x0,y=y0,r=r0,t=t0,Gt):
>
> K:= subs(x=x0,y=y0,r=r0,t=t0,
> ( (Gxte-Gyre)*(gie[2,1]*r+gie[2,2]*t)+2*G*Grre+2*H*Grte
> -Gre*Gre-Gte*Hre )/( (gie[2,1]*r+gie[2,2]*t)*Lt ) ):
>
>
> end proc;

> a:=1:c:=1.55:x0:=0.75: y0:=0: r0:=0: t0:=1:
> F:=1/4*(x^2+2*c)*sqrt(r^2+x^(-2)*t^2)*
> ( 1+1*sqrt(1-16*a*(t)/sqrt(r^2+x^(-2)*t^2))/(x^2+2*c)^2) );

> K(F,x0,y0,r0,t0);

```

The following program is optimized for the rotationally symmetric Cartan metric associated with the rotating Kepler Hamiltonian. Here we use the same notation as in Section 4.

```

> K:=proc(F)
> local H,Hr,Ht,Hrr,Hrt,Htt,gi,detg,g,
> Lr,Lt,Lsx,Lsq1x,Lsq2x,
> G,Gr,Gt,Gar,Gat,Grre,Gtt,Gre,Gte,Hre,x0,r0,t0,
> Garre,Garte,Grte,Gxte,
> gie,detge,ge,Lsq1xe,Lsq2xe:
>
> ## Hamiltonian given
> H:=F^2/2:
>
> Hr:=diff(H,r):
> Ht:=diff(H,t):
> Hrr:=diff(Hr,r):

```

```

> Hrt:=diff(Hr,t):
> Htt:=diff(Ht,t):
>
> gi[1,1]:=Hrr: gi[1,2]:=Hrt: gi[2,1]:=Hrt: gi[2,2]:=Htt:
>
> detg:=gi[1,1]*gi[2,2]-gi[1,2]^2:
>
> ##obtain Finsler 'metric' g_{ij}: note that the dependence on
> ##the fiber coordinates is not correct, since we haven't
> ##performed a Legendre transformation
> g[1,1]:=1/detg*gi[2,2]: g[1,2]:=-1/detg*gi[1,2]:
> g[2,1]:=-1/detg*gi[2,1]: g[2,2]:=1/detg*gi[1,1]:
>
>
> Lr:=r:
> Lt:=t:
>
> Lsx:=diff(H,x):
> Lsq1x:=diff(Hr,x):
> Lsq2x:=diff(Ht,x):
>
> ##Formulas from Section 5
> G:= 1/2*( gi[1,1]*Lsx-Lr*(gi[1,1]*Lsq1x)-Lt*(gi[1,2]*Lsq1x) ):
> H:=1/2*(gi[1,2]*Lsx-Lt*(gi[1,2]*Lsq2x)-Lr*(gi[1,1]*Lsq2x) ):
>
> ##avoid double computations
> Gar:=diff(G,r): Gat:=diff(G,t):
>
> Gr:=Gar*g[1,1]+Gat*g[2,1]:
> Gt:=Gar*g[1,2]+Gat*g[2,2]:
>
> ##evaluate quickly
> x0:=x: r0:=0: t0:=x:
> ge[1,1]:=subs(x=x0,r=r0,t=t0,g[1,1]):
> ge[1,2]:=subs(x=x0,r=r0,t=t0,g[1,2]):
> ge[2,1]:=subs(x=x0,r=r0,t=t0,g[2,1]):
> ge[2,2]:=subs(x=x0,r=r0,t=t0,g[2,2]):
> gie[1,1]:=subs(x=x0,r=r0,t=t0,gi[1,1]):
> gie[1,2]:=subs(x=x0,r=r0,t=t0,gi[1,2]):
> gie[2,1]:=subs(x=x0,r=r0,t=t0,gi[2,1]):
> gie[2,2]:=subs(x=x0,r=r0,t=t0,gi[2,2]):
> detge:=subs(x=x0,r=r0,t=t0,detg):
> Lsq1xe:=subs(x=x0,r=r0,t=t0,Lsq1x):

```

```

> Lsq2xe:=subs(x=x0,r=r0,t=t0,Lsq2x):
>
> ##avoid double computations
> Garre:=subs(x=x0,r=r0,t=t0,diff(Gr,r) ):
> Garte:=subs(x=x0,r=r0,t=t0, diff(Gr,t) ):
>
> ##evaluate when no more differentiation is needed.
> Grre:=Garre*ge[1,1]+Garte*ge[2,1] :
> Grte:=Garre*ge[1,2]+Garte*ge[2,2] :
>
> Gxte:=subs(x=x0,r=r0,t=t0, diff(Gt,x) )
>       +subs(x=x0,r=r0,t=t0, diff(Gt,r) )*(-ge[1,1]*Lsq1xe
>       -ge[1,2]*Lsq2xe)+subs(x=x0,r=r0,t=t0, diff(Gt,t) )
>       *(-ge[1,2]*Lsq1xe-ge[2,2]*Lsq2xe):
> Hre:=subs(x=x0,r=r0,t=t0,diff(H,r) )*ge[1,1]
>       +subs(x=x0,r=r0,t=t0,diff(H,t) )*ge[2,1] :
>
> Gre:=subs(x=x0,r=r0,t=t0,Gr):
> Gte:=subs(x=x0,r=r0,t=t0,Gt):
>
> K:= subs(x=x0,r=r0,t=t0,
>       ( (Gxte)*(gie[2,1]*r+gie[2,2]*t)+2*G*Grre+2*H*Grte
>       -Gre*Gre-Gte*Hre )/( (gie[2,1]*r+gie[2,2]*t)*Lt) ):
>
> end proc;

> F:=1/4*(x^2+2*c)*sqrt(r^2+x^(-2)*t^2)*
>       ( 1+1*sqrt(1-16*a*(t)/sqrt(r^2+x^(-2)*t^2)/(x^2+2*c)^2));

> GK:=K(F);
> curv:=simplify(GK,assume=positive);
> plot({subs(a=1,c=2,curv)},x=-10..10);

```

References

- [1] P. ALBERS, U. FRAUENFELDER, O. VAN KOERT and G. PATERNAIN, The contact geometry of the restricted 3-body problem, *Comm. Pure Appl. Math.* **65** (2012), 229–263.
- [2] P. ALBERS, J. FISH, U. FRAUENFELDER, H. HOFER and O. VAN KOERT, Global surfaces of section in the planar restricted 3-body problem, *Arch. Ration. Mech. Anal.* **204** (2012), 273–284.
- [3] D. BAO, S.-S. CHERN and Z. SHEN, An Introduction to Riemann–Finsler Geometry, *Springer*, 2000.

- [4] E. BELBRUNO, Lunar capture orbits, a method of constructing earth-moon trajectories and the lunar gas mission, Proceedings of AIAA/DGGLR/JSASS Inter. Elec. Prop. Conf. 87-1054, 1989.
- [5] E. BELBRUNO and J. MILLER, A ballistic lunar capture trajectory for the Japanese spacecraft Hiten, Technical Report JPL-IOM 312/90.4-1731-EAB, Jet Propulsion Laboratory, 1990.
- [6] C. CONLEY, Low energy transit orbits in the restricted three-body problem, *SIAM J. Appl. Math.* **16** (1968), 732-746.
- [7] C. CONLEY, Topological dynamics (Sympos., Colorado State Univ., Ft. Collins, Colo., 1967), *Benjamin, New York*, 1968.
- [8] C. CONLEY, On the ultimate behavior of orbits with respect to an unstable critical point I, oscillating, asymptotic and capture orbits, *J. Diff. Equ.* **5** (1969), 136-158.
- [9] A. HARRIS and G. PATERNAIN, Dynamically convex Finsler metrics and J -holomorphic embedding of asymptotic cylinders, *Ann. Global Anal. Geom.* **34** (2008), 115-134.
- [10] J. MOSER, Regularization of Kepler's problem and the averaging method on a manifold, *Comm. Pure Appl. Math.* **23** (1970), 609-636.
- [11] R. MIRON, D. HRIMIUC, H. SHIMADA and S. SABAU, The geometry of Hamilton and Lagrange spaces, Fundamental Theories of Physics 118, *Kluwer Academic Publishers Group, Dordrecht*, 2001.

KAI CIELIEBAK
 MATHEMATISCHES INSTITUT
 ZIMMER L1 - 2009
 UNIVERSITÄT AUGSBURG
 UNIVERSITÄTSSTRASSE 14
 86159 AUGSBURG
 GERMANY
E-mail: kai.cieliebak@math.uni-augsburg.de

URS FRAUENFELDER
 DEPARTMENT OF MATHEMATICS AND
 RESEARCH INSTITUTE OF MATHEMATICS
 SEOUL NATIONAL UNIVERSITY
 BUILDING 27, ROOM 403
 SAN 56-1, SILLIM-DONG
 GWANAK-GU, SEOUL
 POSTAL CODE 151-747
 SOUTH KOREA
E-mail: frauenf@snu.ac.kr

OTTO VAN KOERT
 DEPARTMENT OF MATHEMATICS AND
 RESEARCH INSTITUTE OF MATHEMATICS
 SEOUL NATIONAL UNIVERSITY
 BUILDING 27, ROOM 402
 SAN 56-1, SILLIM-DONG
 GWANAK-GU, SEOUL
 POSTAL CODE 151-747
 SOUTH KOREA
E-mail: okoert@snu.ac.kr