

The restricted three body problem and  
holomorphic curves

Urs Frauenfelder, Otto van Koert

September 18, 2019



# Contents

<b>1</b>	<b>Introduction</b>	<b>5</b>
1.1	The Birkhoff conjecture . . . . .	5
1.2	The power of holomorphic curves . . . . .	6
1.3	Systolic inequalities and symplectic embeddings . . . . .	8
1.4	Beyond the Birkhoff conjecture . . . . .	10



# Chapter 1

## Introduction

### 1.1 The Birkhoff conjecture

The study of the restricted three body problem has a long history. Nevertheless many questions still remain to be solved. We quote from page 328 of Birkhoff's inspiring essay [38]

“This state of affairs seems to me to make it probable that the restricted problem of three bodies admit of reduction to the transformation of a discoid into itself as long as there is a closed oval of zero velocity about J(upiter), and that in consequence there always exists at least one direct periodic orbit of simple type.”

Translated into modern language Birkhoff asks if below the first critical energy value in each bounded component of the restricted three body problem there exists a disk-like global surface of section. The Poincaré return map associated to the disk-like global surface of section then has by Brouwer's translation theorem at least one fixed point. These fixed points give rise to direct periodic orbits.

What do we know a century after Birkhoff published his essay about his question? Unfortunately to the authors' knowledge we still cannot answer the question by Birkhoff with hundred percent certainty in the affirmative. Nevertheless the modern methods of symplectic geometry make it quite likely that the answer to this century old question can be found in the near future. The purpose of these notes is to make young ambitious researchers familiar with the main players of Birkhoff's question, namely the restricted three body problem and disk-like global surfaces of section and to introduce them to the modern techniques of symplectic geometry which reduce Birkhoff's question to questions about symplectic embeddings and systolic geometry.

Through the introduction of holomorphic curves into symplectic topology Gromov revolutionized the subject [104]. Hofer [119] and Hofer, Wysocki, and Zehnder [121, 123, 127] later extended the theory of holomorphic curves to the

case of symplectizations of contact manifolds and discovered in [125] that disk-like global surfaces of section can be constructed with the help of holomorphic curves. The theory of Hofer, Wysocki, and Zehnder was later refined by Siefring with the discovery of his intersection number and by Hryniewicz with the introduction of fast finite energy planes. These theories allow us to find sufficient and necessary conditions for a periodic orbit to bound a disk-like global surface of section. In [9] the moon was “contacted”, i.e., it was shown that for energies below and slightly above the first critical value the bounded components of the energy hypersurface of the regularized restricted three body problem are of contact type. This result enables us to apply holomorphic curve techniques to the restricted three body problem.

The restricted three body problem studies the dynamics of a massless body attracted by two massive bodies according to Newton’s law of gravitation. For example the “massless body” could be imagined as a satellite and the two massive bodies are the earth and the moon. Another option is to think of the “massless body” as the moon and the two massive bodies are the sun and the earth. Or one could think of the “massless body” as the planet Tatooine attracted by the two stars Tatoo I and Tatoo II as in the Star Wars saga. In fact when the planet Tatooine in the Star Wars saga first appeared nobody knew if such planets exist in reality. Due to amazing progress in the search for exoplanets in the last decades we now know that such worlds with two suns exist, see for example [73].

## 1.2 The power of holomorphic curves

The question about the existence of a global surface of section is a question about all orbits not just periodic ones. However, holomorphic curves seem to confirm Poincaré’s fantastic insight [204] that periodic orbits in some sense are the “skeleton” of the dynamics. The power of the technology of holomorphic curves lies in the fact that they reduce the question about existence of global surfaces of section to questions about periodic orbits.

Not every periodic orbit can bind a disk-like global surface of section. Indeed, if a periodic orbit binds a disk-like global surface of section, it necessarily has to satisfy some topological conditions. The mere fact that it is the boundary of an embedded disk implies that it is unknotted. Another obstruction is that it necessarily has to be linked to every other periodic orbit. A more subtle obstruction which comes from contact topology is that its self-linking number has to be  $-1$ . Now suppose that the dynamics can be interpreted as the Reeb flow of a starshaped hypersurface  $\Sigma \subset \mathbb{C}^2$ , which we know to hold true in the case of interest for Birkhoff’s conjecture [9]. Then results of Hryniewicz and Salomão [134, 135] show that these necessary conditions are basically sufficient as well.

The idea behind this result which goes back to the ground breaking work of Hofer, Wysocki, and Zehnder [125] is the following. One considers finite energy planes in the symplectization  $\Sigma \times \mathbb{R}$ . These are finite energy solutions of

a holomorphic curve equation for maps from the plane to the symplectization. Note that  $\Sigma$  has three dimensions, therefore the dimension of  $\Sigma \times \mathbb{R}$  is four, while holomorphic planes are two-dimensional. A pair of two-dimensional objects in a four dimensional space generically intersects in isolated points. For holomorphic curves the local intersection index is positive. On the other hand to define an intersection number for holomorphic planes that is invariant under homotopies is not an easy task, because a plane is noncompact and one has to worry about intersections at infinity. In his celebrated approach to the Weinstein conjecture [119] Hofer already noted that the finiteness of energy for a plane guarantees that asymptotically the projection of a finite energy plane to  $\Sigma$  converges to a periodic Reeb orbit. If one fixes the asymptotics Siefring [219] managed to define an intersection number for finite energy planes which has all the properties one would expect for an intersection number from the closed case. In particular, it is homotopy invariant and moreover, in view of positivity of intersection for holomorphic curves, its vanishing guarantees that two curves with different image do not intersect at all. Even more is true. If Siefring's intersection number for two finite energy planes vanishes, then even their projections to  $\Sigma$  do not intersect unless the images coincide.

One now fixes a periodic Reeb orbit and considers the moduli space of finite energy planes asymptotic to this fixed periodic orbit. There are various symmetries on this moduli space. First there are the reparametrizations of maps from the plane to  $\Sigma \times \mathbb{R}$  and moreover, there is an  $\mathbb{R}$ -action which comes from the obvious  $\mathbb{R}$ -action on the second factor of the target. If one considers the moduli space modulo these symmetries, the dimension of the space becomes two less than the Conley-Zehnder index of the asymptotic Reeb orbit. In particular, if the Conley-Zehnder index is three, the dimension of the moduli space is one. Moreover, due to some automatic transversality miracle it can be shown in this case that the moduli space is actually a manifold, so that it is a disjoint union of circles and intervals. In case the Conley-Zehnder index is bigger than three Hryniewicz [132] found a way to associate a subspace to the space of finite energy planes which after dividing out the symmetries becomes a one-dimensional manifold. Namely, he discovered the notion of "fast finite energy" planes, namely finite energy planes which have a fast exponential asymptotic decay, where the decay rate is chosen in such a way to guarantee that the projection of the finite energy plane to  $\Sigma$  is an immersion transverse to the Reeb vector field.

Now one considers the moduli space modulo symmetries of fast finite energy planes to a fixed asymptotic Reeb orbit  $\gamma$  with vanishing Siefring self-intersection number. By the properties of Siefring's intersection number the projections of these fast finite energy planes to  $\Sigma$  do not intersect and therefore build a local foliation of  $\Sigma$ . To get a global foliation one has to understand the compactness properties of this moduli space. A compactification of the moduli space of finite energy planes is provided by the SFT-compactness theorem [42]. In the case of fast finite energy planes with vanishing Siefring self-intersection number SFT-compactness implies that if the moduli space is noncompact a sequence of fast finite energy planes has to converge to a negatively punctured fast finite energy plane and at these punctures the projection of the punctured fast finite energy

plane converges to periodic Reeb orbits which are unlinked to the Reeb orbit  $\gamma$ . In particular, if  $\gamma$  is linked to every other periodic Reeb orbit this scenario cannot occur and therefore the moduli space of fast finite energy planes has to be compact. In case it is not empty, we get an open book decomposition of  $\Sigma$  whose binding is the periodic Reeb orbit  $\gamma$ . Each page of this open book corresponds to the projection of a fast finite energy plane to  $\Sigma$ . It turns out that each page is a global surface of section for the Reeb flow.

### 1.3 Systolic inequalities and symplectic embeddings

As we have seen in the previous section in order to prove compactness of the moduli space of fast finite energy planes of vanishing Siefring self-intersection number asymptotic to the Reeb orbit  $\gamma$  one needs to make sure that  $\gamma$  is linked to every other periodic Reeb orbit. However, looking at the SFT-compactness theorem more closely reveals that if the moduli space is noncompact and therefore it contains a negatively punctured fast finite energy plane in its closure, then the periodic orbits at the negative punctures satisfy additional conditions

- (i) The periods of the periodic orbits at the negative punctures are less than the period of  $\gamma$ .
- (ii) At least one periodic orbit at a negative puncture orbit has Conley-Zehnder index less than or equal to 2.

In particular, we deduce from (i) that if  $\gamma$  has the shortest period among all periodic orbits, compactness is guaranteed. The periodic orbit of smallest period is referred to as the *systole*. Hence the Birkhoff conjecture prompts the following question, which if answered positively, actually would imply the Birkhoff conjecture with the help of holomorphic curve techniques.

**Question:** *Does the retrograde periodic orbit represent the systole?*

We point out, that we only expect a positive answer to the question for energy values below the first critical one. Indeed, above the first critical value a periodic orbit bifurcates out of the critical point known as Lyapunov orbit, which at least for energies slightly above the first critical value represents the systole.

In Riemannian geometry the systole is just the shortest geodesic and there is a huge literature on the subject, see for example [36]. The introduction of systolic question to the contact world is rather new and started with the work of Álvarez Paiva and Balacheff [13]. How much the Riemannian world and the contact world can differ was recently realized through the discoveries of Abbondandolo, Bramham, Hryniewicz, and Salomão [2]. The work of Lee [157] represents first steps to approach the systole in Hill's lunar problem via Symplectic homology.

In view of (ii) a different approach to the Birkhoff conjecture is to rule out periodic orbits of Conley-Zehnder index 2. The following definition is due to Hofer, Wysocki, and Zehnder [125]

**Definition 1.3.1** *A starshaped hypersurface in  $\Sigma \subset \mathbb{C}^2$  is called dynamically convex if the Conley-Zehnder indices of all periodic Reeb orbits are at least 3.*

Armed with this notion we can now ask

**Question:** *Is the restricted three body problem dynamically convex?*

Again this question only applies to energies below the first critical value. Apart from the fact that the dynamics of the restricted three body problem just below the first critical value can be interpreted as the Reeb flow of a starshaped hypersurface [9] the culprit is again the Lyapunov orbit which bifurcates out of the critical point. This periodic orbit has Conley-Zehnder index 2!

Only in rare cases dynamical convexity can be checked directly. Indeed, first finding all periodic orbits and then compute their Conley-Zehnder indices is in general not doable. However, Hofer, Wysocki, and Zehnder [125] found a much more tractable criterion.

**Theorem [Hofer, Wysocki, Zehnder]:** *A strictly convex hypersurface  $\Sigma \subset \mathbb{C}^2$  is dynamically convex.*

It is interesting to note that while dynamical convexity is a symplectic notion, the concept of convexity is not. Indeed, it can well happen that a non-convex starshaped hypersurface admits a different symplectic embedding which is convex. On the other hand the theorem of Hofer, Wysocki, and Zehnder tells us that a starshaped hypersurface in the complex plane which has a periodic orbit of Conley-Zehnder index less than three does not admit a convex symplectic embedding at all. To our knowledge the following problem is still open.

**Question:** *Does there exist a dynamically convex starshaped hypersurface  $\Sigma \subset \mathbb{C}^2$  which does not admit a convex symplectic embedding at all?*

The question about symplectic embeddings was one of the driving forces for the development of symplectic topology. Starting with Gromov's celebrated non-squeezing theorem [104] the question which symplectic manifolds can be symplectically embedded into each other is a highly active area of research, see for example [213, 214] for a survey.

Moreover, the question about convex symplectic embeddings and the systole are not unrelated. Indeed, it is known that in the convex case the Hofer-Zehnder capacity coincides with the systole [130], and obtaining estimates for the systole in the convex case is strongly related to the famous Viterbo conjecture [236, 198].

What do we know about dynamical convexity for the restricted three body problem. For energies below the first critical value the Levi-Civita regularization provides us with an embedding into  $\mathbb{C}^2$  and it was shown in [9] that the

corresponding hypersurface is starshaped. In view of the Theorem of Hofer, Wysocki, and Zehnder the natural question is, if this embedding is maybe even convex. In fact this is sometimes true. It was shown in [6] that for small mass ratios around the small mass the embedding is convex and therefore dynamical convexity holds. On the other hand this fails around the heavy body, even in the limit when the mass of the light body is zero. This limit is the rotating Kepler problem, namely the Kepler problem in rotating coordinates. In fact it was noted in [7] that the Levi-Civita embedding of the rotating Kepler problem is not convex. On the other hand in the same paper, the Conley-Zehnder indices of all periodic orbits were computed and the outcome was that the rotating Kepler problem is dynamically convex. The question arose if the rotating Kepler problem might provide an example of a dynamical convex starshaped hypersurface in  $\mathbb{C}^2$  which does not admit a convex symplectic embedding. This turned out to be wrong as well. In fact in [93] a convex embedding for the rotating Kepler problem was found which used a combination of the Ligon-Schaaf and Levi-Civita embedding.


## 1.4 Beyond the Birkhoff conjecture

Birkhoff's assumption that there is a closed oval of zero velocity around the massive bodies precludes a satellite from traveling from the earth to the moon. This assumption of Birkhoff is equivalent to the satellite having an energy below the first critical value. Although we basically concentrate us in this monograph on this case, our motivation is actually to go above the first critical value. The binding orbit of the global surface of section that Birkhoff has in mind is the retrograde periodic orbit. Now it is impossible that the retrograde periodic orbit binds a global surface of section above the first critical point. The reason is that from the critical point a new periodic orbit bifurcates which is known as the Lyapunov orbit. The retrograde periodic orbit and the Lyapunov orbit are unlinked and therefore the retrograde orbit cannot bind a global surface of section anymore. Here is what we expect to happen. The holomorphic curves giving rise to global surfaces of section below the first critical value break precisely at the Lyapunov orbit and give rise to a finite energy foliation similar as in [129]. Interesting research in this direction was carried out in [83, 202]. If this picture is correct, then it follows from the theory developed in [129] that the stable and unstable manifold of the Lyapunov orbit intersect. For small mass ratios this phenomenon was first noted by Conley [59] and McGehee [173], and this fact turned out to have fantastic applications to space mission design.

The traditional way from the earth to the moon uses the Hohmann transfer [131]. This transfer uses two engine impulses; one at the beginning to bring the spacecraft to the transfer orbit and one at the end to stop. This method only takes advantage of the dynamics of the two body problem which have been well known since the times of Kepler. The Hohmann transfer was for example successfully applied for the Apollo Moon landings. It is fast, but one of the drawbacks is that it uses a lot of fuel. Not just at the beginning but also at the

end of the journey to stop. Especially the fuel one needs to stop the rocket has to be carried during its whole journey which is very expensive. There is also some danger. Namely if the stopping does not work out properly the rocket either crashes into the moon or just flies by. It was Conley [58, 59, 60] who first propagated the idea to use the dynamics of the restricted three body problem to find low energy transit orbits to the moon. In this spirit the spacecraft ISEE-3 (International Sun-Earth Explorer-3) was brought in 1978 to a halo orbit at a Lagrange point of the earth-sun system using three-body trajectories [79, 80]. It was Belbruno who found the first realistic low energy transit orbit from an orbit around the earth to an orbit around the moon [31, 32]. Although many people at first were skeptical about these new methods, the rescue of the Japanese Hiten mission [34] impressively proved that chaotic motion can be applied to real mission design. In January 1990 the Japanese launched their first lunar mission. There were two robotic spacecrafts involved in this mission: MUSES-A (later on renamed Hiten) and MUSES-B (later on renamed Hagoromo). MUSES-B was supposed to go to the moon while MUSES-A was to stay on an orbit around the earth as a communication relay with MUSES-B. Unfortunately, contact with MUSES-B was lost and only MUSES-A remained. Since MUSES-A was never supposed to go to the moon it had only little fuel, too little to travel to the moon on a standard route. Nevertheless, on October 2, 1991, Hiten successfully arrived at the moon after a long travel which took advantage of chaotic motion in the earth-moon-satellite system as well as in the sun-earth-satellite system. The many ups and downs in this thrilling story can be nicely read in Belbruno's book *Fly me to the moon* [33]. While the traditional space mission design is referred to as *patched conics*, namely patched two body problems whose solutions are given by conic sections, the new approach to space mission design pioneered by Belbruno can be thought of as *patched restricted three body problems*. We refer the reader to the beautiful book by Koon, Lo, Marsden, and Ross [151] for more information on this new paradigm in space mission design.

*Acknowledgements:* Part of these notes are extracted from lectures the first named author held at the University of Augsburg in the winter term 2014/15 and the two authors held at the CIMPA-NIMS Research school on dynamical systems which took place in Daejeon 25.5.–3.6.2015. We would like to thank heartily all the participants of these lectures. You greatly helped that these notes took shape. We sincerely thank CIMPA and NIMS for making this research school possible. We immensely profited from discussions with many people, in particular, Peter Albers, Edward Belbruno, Alain Chenciner, Hagyun Choi, Kai Cieliebak, Etienne Ghys, Joel Fish, Helmut Hofer, Umberto Hryniewicz, Jungsoo Kang, Seongchan Kim, Junyoung Lee, Pedro Salomão, Felix Schlenk, Martin Schwingenheuer, Richard Siefring, Francesco Topputo, Lei Zhao.



# Bibliography

- [1] C. Abbas, H. Hofer, *Holomorphic Curves and Global Questions in Contact Geometry*, book in preparation.
- [2] A. Abbondandolo, B. Bramham, U. Hryniewicz, P. Salomão, *Sharp systolic inequalities for Reeb flows on the three-sphere*, arXiv:1504.05258
- [3] R. Abraham, J. Marsden, *Foundations of Mechanics*, 2nd ed., Addison-Wesley, Reading (1978).
- [4] R. Abraham, J. Marsden, T. Ratiu, *Manifolds, tensor analysis, and applications*, 2nd ed., Applied Mathematical Sciences **75**, Springer (1988).
- [5] B. Aebischer, M. Borer, M. Kalin, C. Leuenberger, *Symplectic Geometry*, Progress in Mathematics **124**, Birkhäuser, Basel (1994).
- [6] P. Albers, J. Fish, U. Frauenfelder, H. Hofer, O. van Koert, *Global surfaces of section in the planar restricted 3-body problem*, Arch. Ration. Mech. Anal. **204** (2012), no. 1, 273–284.
- [7] P. Albers, J. Fish, U. Frauenfelder, O. van Koert, *The Conley-Zehnder indices of the rotating Kepler problem*, Math. Proc. Cambridge Phil. Soc. **154**, no. 2 (2013), 243–260.
- [8] P. Albers, U. Frauenfelder, *Rabinowitz Floer homology: A survey*, in Global Differential Geometry, Springer Proc. in Math. (2012), 437–461.
- [9] P. Albers, U. Frauenfelder, O. van Koert, G. Paternain, *Contact geometry of the restricted three-body problem*, Comm. Pure Appl. Math. **65** (2012), no. 2, 229–263.
- [10] P. Albers, H. Hofer, *On the Weinstein conjecture in higher dimensions*, Comm. Math. Helv. (2009), 429–436.
- [11] A. Albouy, *There is a projective dynamics*, EMS Newsletter **89** (2013), 37–43.
- [12] J. Alexander, *A lemma on systems of knotted curves*, Proc. Nat. Acad. Sci. U.S.A. **9** (1923), 93–95.

- [13] J. Álvarez Paiva, F. Balacheff, *Contact geometry and isosystolic inequalities*, *Geom. Funct. Anal.* **24** (2014), 648–669.
- [14] P. Appell, *De l'homographie en mécanique*, *American Journal of Mathematics*, **12** (1890), 103–114.
- [15] R. Arenstorf, *Periodic solutions of the restricted three-body problem representing analytic continuations of Keplerian elliptic motions*, *Am. J. Math.* **85** (1963), 27–35.
- [16] V. Arnold, *On a characteristic class entering into conditions of quantization*, *Func. Anal.* **1** (1967), 1–8.
- [17] V. Arnold, *Dynamical Systems V: Bifurcation Theory and Catastrophe Theory*, *Encyclopaedia of Mathematical Sciences* **5**, Springer Berlin Heidelberg (1994).
- [18] V. Arnold, *Mathematical methods of classical mechanics*, *Graduate texts in mathematics* **60**, New York, Springer (2000).
- [19] V. Arnold, A. Givental, *Symplectic topology*, In *Dynamical Systems IV*. Springer-Verlag, Berlin (1990).
- [20] V. Arnold, V. Kozlov, A. Neishtadt, *Mathematical Aspects of Classical and Celestial Mechanics*, 3. edition, Springer-Verlag Berlin Heidelberg (2006).
- [21] M. Atiyah, *Convexity and commuting Hamiltonians*, *Bull. London Math. Soc.* **14** (1982), 1–15.
- [22] H. Bacry, H. Ruegg, J. Souriau, *Dynamical Groups and Spherical Potentials in Classical Mechanics*, *Commun. math. Phys.* **3** (1966), 323–333.
- [23] V. Bangert, Y. Long, *The existence of two closed geodesics on every Finsler 2-sphere*, *Math. Ann.* **346** (2010), 335–366.
- [24] V. Bargmann, *Zur Theorie des Wasserstoffatoms: Bemerkungen zur gleichnamigen Arbeit von V. Fock*, *Zeitschrift für Physik* **99** (1936), 576–582.
- [25] R. Barrar, *Existence of periodic orbits of the second kind in the restricted problem of three bodies* *Astron. J.* **70** (1965), 3–5.
- [26] R. Barrar, *Periodic orbits of the second kind*, *Indiana Univ. Math. J.* **22** (1972), 33–41.
- [27] J. Barrow-Green, *Poincaré and the Three Body Problem*, AMS-LMS (1997).
- [28] A. Bathkin, N. Bathkina, *Hénon's generating solutions and the structure of periodic orbits families of the restricted three-body problem*, arXiv:1411.4933
- [29] E. Belbruno, *Two body motion under the inverse square central force and equivalent geodesic flows*, *Celest. Mech.* **15** (1977), 467–476.

- [30] E. Belbruno, *Regularizations and geodesic flows*, Lecture notes in Pure and Applied Mathematics **80** (1981), 1–11.
- [31] E. Belbruno, *Lunar capture orbits, a method of constructing earth-moon trajectories and the lunar gas mission*, Proceedings of AIAA/DGGLR/JSASS Inter. Elec. Propl. Conf., number 87-1054 (1987).
- [32] E. Belbruno, *Capture Dynamics and Chaotic Motions in Celestial Mechanics*, Princeton University Press (2004).
- [33] E. Belbruno, *Fly me to the moon*, Princeton University Press: Princeton (2007).
- [34] E. Belbruno, J. Miller, *A ballistic lunar capture trajectory for the Japanese spacecraft Hiten*, Technical Report JPL-IOM 312/90.4-1731-EAB, Jet Propulsion Laboratory (1990).
- [35] J. van den Berg, F. Pasquotto, C. Vandervorst, *Closed characteristics on non-compact hypersurfaces in  $\mathbb{R}^{2n}$* , Math. Ann. **343** (2009), 247–284.
- [36] M. Berger, *A Panoramic View of Riemannian Geometry*, Springer-Verlag Berlin Heidelberg (2003).
- [37] G. Birkhoff, *Proof of Poincaré's geometric theorem*, Trans. Amer. Math. Soc. **14** (1913), 14–22.
- [38] G. Birkhoff, *The restricted problem of three bodies*, Rend. Circ. Matem. Palermo **39** (1915), 265–334.
- [39] A. Bolsinov, A. Fomenko, *Integrable Hamiltonian systems. Geometry, topology, classification* Translated from the 1999 Russian original. Chapman & Hall/CRC, Boca Raton, FL, 2004. xvi+730 pp. ISBN: 0-415-29805-9
- [40] R. Bott, L. Tu, *Differential forms in algebraic topology*, Graduate Texts in Mathematics, 82. Springer-Verlag, New York-Berlin, 1982. xiv+331 pp. ISBN: 0-387-90613-4
- [41] F. Bourgeois, *A Morse-Bott approach to contact homology*, Ph.D. thesis, Stanford University (2002).
- [42] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki, E. Zehnder, *Compactness results in Symplectic Field theory*, Geom. and Top. **7** (2003), 799–889.
- [43] T. Bröcker, K. Jänich, *Introduction to Differential Topology*, Cambridge University Press (1982).
- [44] R. Broucke, *Periodic orbits in the restricted three-body theorem with Earth-Moon masses*, Pasadena, Jet Propulsion Laboratory, California Institute of Technology (1968).

- [45] L. Brouwer, *Beweis des ebenen Translationssatzes*, Math. Ann. **72** (1912), 37–54.
- [46] L. Brouwer, *Über die periodischen Transformationen der Kugel*, Math. Ann. **80** (1919), 39–41.
- [47] E. Brown, *An Introductory Treatise on the lunar Theory*, Cambridge University Press (1896).
- [48] A. Bruno, *The restricted 3-body problem: Plane periodic orbits*, De Gruyter Expositions in Mathematics **17**, Walter de Gruyter, Berlin, New York (1994).
- [49] A. Cannas da Silva, *Lectures on Symplectic geometry* 2nd printing, Lecture Notes in Mathematics **1764**, Springer (2008).
- [50] C. Charlier, *Die Mechanik des Himmels*, Leipzig, Veit (1902).
- [51] A. Chenciner, R. Montgomery, *A remarkable periodic solution of the three-body problem in the case of equal masses*, Ann. of Math. (2) **152** (2000), no. 3, 881–901.
- [52] A. Chenciner (2007), *Three body problem*, Scholarpedia, 2(10):2111.
- [53] A. Chenciner, *Poincaré and the Three-Body Problem*, Poincaré 1912–2012, Séminaire Poincaré XVI (2012), 45–133.
- [54] K. Cieliebak, U. Frauenfelder, *A Floer homology for exact contact embeddings*, Pacific J. Math. **239** (2009), no. 2, 251–316.
- [55] K. Cieliebak, U. Frauenfelder, O. van Koert, *The Finsler geometry of the rotating Kepler problem*. Publ. Math. Debrecen **84** (2014), no. 3-4, 333–350.
- [56] C. Conley, *On Some New Long Periodic Solutions of the Plane Restricted Three Body Problem*, Comm. Pure Appl. Math. **16** (1963), 449–467.
- [57] C. Conley, *The Retrograde Circular Solutions of the Restricted Three-Body Problem Via a Submanifold Convex to the Flow*, SIAM J. Appl. Math. **16**, no. 3, (1968), 620–625.
- [58] C. Conley, *Low energy transit orbits in the restricted three-body problem*, SIAM J. Appl. Math. **16** (1968), 732–746.
- [59] C. Conley, *Twist mappings, linking, analyticity, and periodic solutions which pass close to an unstable periodic solution*, Topological dynamics (Sympos., Colorado State Univ., Ft. Collins, Colo., 1967), Benjamin, New York (1968), 129–153.
- [60] C. Conley, *On the ultimate behavior of orbits with respect to an unstable critical point I, oscillating, asymptotic and capture orbits*, J. Differential Equations **5** (1969), 136–158.

- [61] C. Conley, E. Zehnder, *The Birkhoff-Lewis fixed point theorem and a conjecture of V.I. Arnold*, Invent. Math. **73** (1983), 33–49.
- [62] C. Conley-Zehnder, *Morse-type index theory for flows and periodic solutions of Hamiltonian equations*, Commun. Pure Appl. Math. **37** (1984), 207–253.
- [63] A. Constantin, B. Kolev, *The theorem of K er ekjart  on periodic homeomorphisms of the disc and the sphere*, Enseign. Math. (2) **40** (1994), no. 3–4, 193–204.
- [64] D. Cristofaro-Gardiner, M. Hutchings, *From one Reeb orbit to two*, arXiv:1202.4839
- [65] R. Cushman, L. Bates, *Global aspects of classical integrable systems* Second edition. Birkh user/Springer, Basel, 2015. xx+477 pp. ISBN: 978-3-0348-0917-7
- [66] R. Cushman, J. Duistermaat, *A Characterization of the Ligon-Schaaf Regularization Map*, Comm. Pure Appl. Math. **50** (1997), 773–787.
- [67] G. Darwin, *Periodic orbits*, Acta Math. **21**, (1897), 99–242.
- [68] G. Darwin, *On certain families of periodic orbits*, Monthly Notices Roy. Astron. Soc. **70**, (1909). 108–143.
- [69] C. Delaunay, *Nouvelle th orie analytique du mouvement de la lune*, Comptes Rendus **23**, 968–970.
- [70] C. Delaunay, *Th orie du mouvement de la lune I*, M moire de l’Academie des Sciences **28**, 1–883.
- [71] C. Delaunay, *Th orie du mouvement de la lune II*, M moire de l’Academie des Sciences **29**, 1–931.
- [72] T. Delzant, *Hamiltoniens p riodiques et image convexe de l’application moment*, Bull. Soc. Math. France **116** (1988), 315–339.
- [73] L. Doyle, W. Welsh, *Worlds with Two Suns*, Scientific American **309**(5), (2013), 40–47.
- [74] J. Duistermaat, *On the Morse index in Variational Calculus*, Adv. Math. **21** (1976), 173–195.
- [75] C. Ehresmann, *Les connexions infinit simales dans un espace fibr  diff rentiable*, Colloque de Topologie, Bruxelles (1950), 29–55.
- [76] Y. Eliashberg, A. Givental, H. Hofer, *Introduction to symplectic field theory*, Geom. Funct. Anal. **10** (2000), 560–673.
- [77] L. Euler, *Un corps  tant attir  en raison d ciproque quarr e des distances vers deux points fixes donn s*, M moires de l’Acad. de Berlin (1760), 228–249.

- [78] L. Euler, *De motu corporis ad duo centra virium fixa attracti*, Novi Commentarii Academiae Scientiarum Imperialis Petropolitanae **10** (1766), 207–242, **11** (1767), 152–184.
- [79] R. Farquhar, *The control and use of libration-point satellites*, Technical Report TR R-346, NASA (1970).
- [80] R. Farquhar, D. Muhonen, C. Newman, H. Heuberger, *Trajectories and orbital maneuvers for the first libration-point satellite*, J. Guid. and Control **3** (1980), 549–554.
- [81] A. Fathi, *An orbit closing proof of Brouwer’s lemma on translation arcs*, L’enseignement Math. **33** (1987), 315–322.
- [82] S. Ferraz-Mello (2009), *Celestial mechanics*, Scholarpedia, 4(1):4416.
- [83] J. Fish, R. Siefring, *Connected sums and finite energy foliations I: Contact connected sums*, arXiv:1311.4221
- [84] A. Floer, *Morse theory for Lagrangian intersections*, J. Differential Geom. **28** (1988), 513–547.
- [85] A. Floer, H. Hofer, D. Salamon, *Transversality in elliptic Morse theory for the symplectic action*, Duke Math. Journal, **80** (1996), 251–292.
- [86] V. Fock, *Zur Theorie des Wasserstoffatoms*, Zeitschrift für Physik **98** (1935), 145–154.
- [87] A. Fomenko, *Symplectic geometry*, Advanced studies in contemporary mathematics **5**, New York, Gordon and Breach (1988).
- [88] J. Franks, *A new proof of the Brouwer plane translation theorem*, Erg. Th. and Dyn. Syst., **12** (1992), 217–226.
- [89] J. Franks, *Geodesics on  $S^2$  and periodic points of annulus homeomorphisms*, Invent. Math., **108** (1992), 403–418.
- [90] J. Franks, *Area preserving homeomorphisms of open surfaces of genus zero*, New York Jour. Math. **2** (1996), 1–19.
- [91] U. Frauenfelder, *Dihedral homology and the moon*, J. Fixed Point Theory Appl. **14** (2013), 55–69.
- [92] U. Frauenfelder, J. Kang, *Real holomorphic curves and invariant global surfaces of section*, Proc. Lond. Math. Soc. (3) **112** (2016), no. 3, 477–511.
- [93] U. Frauenfelder, O. van Koert, L. Zhao, *A convex embedding for the rotating Kepler problem*, arXiv:1605.06981
- [94] D. Fuks, *The Maslov-Arnold characteristic classes*, Dokl. Akad. Nauk SSSR **178** (1968), 303–306.

- [95] N. Gavrilov, A. Shilnikov, *Example of a blue sky catastrophe*, AMS Transl. Series II, v.200, 99–105, 2000.
- [96] H. Geiges, *An introduction to contact topology*, Cambridge Studies in Adv. Math. **109**, Cambridge Univ. Press (2008).
- [97] H. Geiges, *The Geometry of Celestial Mechanics*, book in progress.
- [98] J. Ginsburg, D. Smith, *A History of Mathematics in America before 1900*, The Mathematical Association of America (Carus Mathematical Monograph Number 5).
- [99] V. Ginzburg, V. Guillemin, Y. Karshon, *Moment Maps, Cobordism, and Hamiltonian Group Actions*, Math. Surv. and Monographs, **98**. Amer. Math. Soc.
- [100] V. Ginzburg, B. Gürel, *A  $C^2$ -smooth counterexample to the Hamiltonian Seifert conjecture in  $\mathbb{R}^4$* , Ann. of Math. (2) **158** (2003), no.3, 953–976.
- [101] H. Goldstein, *Prehistory of the Runge-Lenz vector*, Amer. Jour. Phys. **43** (1975), no. 8, 737–738.
- [102] H. Goldstein, *More on the prehistory of the Runge-Lenz vector*, Amer. Jour. Phys. **44** (1976), no. 11, 1123–1124.
- [103] E. Goursat, *Sur les transformations isogonales en Mécanique*, Comptes Rendus des Séances de l'Académie des Sciences, Paris, **108**, (1889), 446–448.
- [104] M. Gromov, *Pseudo Holomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307–347.
- [105] V. Guillemin, S. Sternberg, *Convexity Properties of the Moment Mapping*, Inv. Math. **67** (1982), 491–513.
- [106] L. Guillou, *Théorème de translation plane de Brouwer et généralisations du théorème de Poincaré-Birkhoff*, Topology **33**(2) (1994), 331–351.
- [107] G. Györgyi, *Kepler's equation, Fock variables, Bacry's generators and Dirac brackets*, Nuovo Cimento, 53A (1968), 717–735.
- [108] Y. Hagihara, *Celestial mechanics*, **1-2**, MIT Press, Cambridge, Massachusetts (1970-1972); **3-5**, Japan Society for the Promotion of Science, Tokyo, (1974-1976).
- [109] G. Hall, K. Meyer, *Introduction to Hamiltonian Dynamical Systems and the N-Body Problem*, Applied Mathematical Sciences **90**, Springer (1992).
- [110] A. Hatcher, *Algebraic Topology*, Cambridge University Press (2001).
- [111] G. Heckman, T. de Laat, *On the Regularization of the Kepler Problem*, Jour. Symp. Geom. **10**, vol. 3 (2012), 463–473.

- [112] G. Heckman, L. Zhao, *Angular Momenta of Relative Equilibrium Motions and Real Moment Map Geometry*, *Invent. Math.* **205** (2016), no. 3, 671–691.
- [113] M. Hénon, *Numerical exploration of the restricted problem. V. Hill's case: Periodic orbits and their stability*, *Astron. Astrophys.* **1**, 223–238.
- [114] M. Hénon, *Generating Families in the Restricted Three-Body Problem*, Springer (1997).
- [115] G. Hill, *On the part of the Motion of the Lunar Perigee which is a Function of the Mean Motions of the Sun and the Moon*, John Wilson & Son, Cambridge, Massachusetts (1877), Reprinted in *Acta* **8**, (1886), 1–36.
- [116] G. Hill, *Researches in the lunar theory*, *Amer. J. Math.* **1** (1878), 5–26, 129–147, 245–260.
- [117] A. Hildebrandt, *On the problem of two fixed centres and certain of its generalizations*, *Amer. J. Math.* **33**, no. 1/4 (1911), 337–362.
- [118] L. Hörmander, *Fourier integral operators I*, *Acta Math.* **127** (1971), 79–183.
- [119] H. Hofer, *Pseudoholomorphic curves in symplectisations with application to the Weinstein conjecture in dimension three*, *Invent. Math.* **114** (1993), 515–563.
- [120] H. Hofer, K. Wysocki, E. Zehnder, *A characterisation of the tight three-sphere*, *Duke Math. J.* **81** (1995), 159–226.
- [121] H. Hofer, K. Wysocki, E. Zehnder, *Properties of pseudo-holomorphic curves in symplectizations. II. Embedding controls and algebraic invariants*, *Geom. Funct. Anal.* **5** (1995), no. 2, 270–328.
- [122] H. Hofer, K. Wysocki, E. Zehnder, *Unknotted periodic orbits for Reeb flows on the three-sphere*, *Topol. Methods Nonlinear Anal.* **7**, (1996), no. 2, 219–244.
- [123] H. Hofer, K. Wysocki, E. Zehnder, *Properties of pseudo-holomorphic curves in symplectizations. I. Asymptotics*, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **13** (1996), no. 3, 337–379.
- [124] H. Hofer, K. Wysocki, E. Zehnder, *Correction to: "A characterisation of the tight three sphere"*, *Duke Math. J.* **89** (1997), no. 3, 603–617.
- [125] H. Hofer, K. Wysocki, E. Zehnder, *The dynamics on a strictly convex energy surface in  $\mathbb{R}^4$* , *Ann. Math.*, **148** (1998) 197–289.
- [126] H. Hofer, K. Wysocki, E. Zehnder, *A characterization of the tight 3-sphere. II*, *Comm. Pure Appl. Math.* **52**, no. 9 (1999), 1139–1177.

- [127] H. Hofer, K. Wysocki, E. Zehnder, *Properties of pseudo-holomorphic curves in symplectizations. III. Fredholm theory*, Topics in nonlinear analysis, Progr. Nonlinear Differential Equations Appl., **35**, Birkhäuser, Basel (1999), 381–475.
- [128] H. Hofer, K. Wysocki, E. Zehnder, *Pseudoholomorphic curves and dynamics in three dimensions*, Handbook of dynamical systems, Vol. 1A, North-Holland, Amsterdam (2002), 1129–1188.
- [129] H. Hofer, K. Wysocki, E. Zehnder, *Finite energy foliations of tight three-spheres and Hamiltonian dynamics*, Ann. Math. **157** (2003), 125–257.
- [130] H. Hofer, E. Zehnder, *Symplectic invariants and Hamiltonian dynamics*, Reprint of the 1994 edition. Modern Birkhäuser Classics. Birkhäuser Verlag, Basel, 2011. xiv+341 pp. ISBN: 978-3-0348-0103-4
- [131] W. Hohmann, *Die Erreichbarkeit der Himmelskörper*, Oldenbourg, Munich (1925).
- [132] U. Hryniewicz, *Fast finite-energy planes in symplectizations and applications*, Trans. Amer. Math. Soc. **364** (2012), 1859–1931.
- [133] U. Hryniewicz, *Systems of global surfaces of section for dynamically convex Reeb flows on the 3-sphere*, J. Symplectic Geom. **12** (2014), no. 4, 791–862.
- [134] U. Hryniewicz, P. Salomão, *On the existence of disk-like global sections for Reeb flows on the tight 3-sphere*, Duke Math. J. **160**, no. 3, (2011), 415–465.
- [135] U. Hryniewicz, P. Salomão, *Elliptic bindings for dynamically convex Reeb flows on the real projective three-space*, Calc. Var. Partial Differential Equations **55** (2016), no. 2, Art. 43, 57 pp.
- [136] L. Hulthén, *Über die quantenmechanische Herleitung der Balmerterme*, Zeitschrift für Physik **86** (1933), 21–23.
- [137] C. Hummel, *Gromov’s compactness theorem for pseudo-holomorphic curves*, Progress in Mathematics, 151. Birkhäuser Verlag, Basel, 1997. viii+131 pp. ISBN: 3-7643-5735-5
- [138] M. Hutchings, *An index inequality for embedded pseudoholomorphic curves in symplectizations*, J. Eur. Math. Soc. (JEMS) **4** (2002), 313–361.
- [139] C. Jacobi, *Vorlesungen über Dynamik*, Berlin, Reimer (1866).
- [140] P. Kahn, *Pseudohomology and Homology*, math.AT/0111223.
- [141] J. Kang, *Some remarks on symmetric periodic orbits in the restricted three-body problem*, Disc. Cont., Dyn. Sys. (A), **34**, no. 12 (2014), 5229–5245.
- [142] J. Kang, *On reversible maps and symmetric periodic points*, arXiv:14103997

- [143] T. Kato, *On the convergence of the perturbation method I*, Progr., Theor. Phys. **4** (1949), 514–523.
- [144] T. Kato, *On the convergence of the perturbation method II*, Progr., Theor. Phys. **5** (1950), 207–212.
- [145] T. Kato, *Perturbation theory for Linear Operators*, Springer, Grundlehren edition (1976).
- [146] B. von K erekjart , * ber die periodischen Transformationen der Kreisscheibe und der Kugelfl che*, Math. Ann. **80**, (1919–1920), 36–38.
- [147] W. Killing, *Die Mechanik in den Nicht-Euklidischen Raumformen*, Jour. reine und angewandte Math. **98** (1885), 1–48.
- [148] D. Kim, *Planar Circular Restricted Three Body Problem*, Master thesis, Seoul National University (2011).
- [149] S. Kim, *Hamiltonian mechanics and Symmetries*, Master thesis, Seoul National University (2014).
- [150] S. Kobayashi, K. Nomizu, *Foundations of Differential Geometry I & II*, Wiley Classics Library (1996).
- [151] W. Koon, M. Lo, J. Marsden, S. Ross, *Dynamical Systems, the Three-Body Problem and Space Mission Design*, Marsden Books (2011).
- [152] V. Kozlov, A. Harin, *Kepler’s problem in constant curvature spaces*, Celestial Mech. Dyn. Astron. **54**, no. 4 (1992), 393–399.
- [153] M. Kriener, *An intersection formula for finite energy half cylinders*, PhD thesis, ETH Zurich (1998).
- [154] M. Kummer, *On the stability of Hill’s Solutions of the Plane Restricted Three Body Problem*, Amer. J. Math. **101**(6), (1979), 1333–1354.
- [155] P. Kustaanheimo, E. Stiefel, *Perturbation theory of Kepler motion based on spinor regularization*, J. Reine Angew. Math. **218**, (1965), 204–219.
- [156] J. Lagrange, *Recherches sur la mouvement d’un corps qui est attir  vers deux centres fixes*, Miscellanea Taurinensia **4** (1766–69), 67–121, M canique Analytique, 2nd edition, Paris (1811), 108–121.
- [157] J. Lee, *Fiberwise Convexity of Hill’s lunar problem*, J. Topol. Anal. **9** (2017), no. 4, 571–630.
- [158] J. Lee, *Spectrum estimates of Hill’s lunar problem*, arXiv:1510.07248
- [159] T. Levi-Civita, *Sur la r gularisation du probl me des trois corps*, Acta Math. **42** (1920), 99–144.

- [160] P. Libermann, C. Marle, *Symplectic geometry and analytical mechanics*, Reidel, Dordrecht (1987).
- [161] R. Lickorish, *An introduction to knot theory*, Graduate texts in mathematics **175**, New York, Springer (1997).
- [162] T. Ligon, M. Schaaf, *On the Global Symmetry of the Classical Kepler Problem*, Reports on Math. Phys. **9** (1976), 281–300.
- [163] G. Lion, M. Vergne, *The Weil representation, Maslov index and Theta Series*, Progress in Mathematics **6**, Birkhäuser (1980).
- [164] J. Llibre, L. Roberto, *On the periodic orbits and the integrability of the regularized Hill lunar problem*, J. Math. Phys. **52**(8), 082701, 8 pp. (2011).
- [165] G. Lodder, *Dihedral homology and the free loop space*, Proc. London Math. Soc. (3) **60** (1990), 201–224.
- [166] R. Lohner, *Computation of guaranteed enclosures for the solutions of ordinary initial and boundary value problems*, Computational ordinary differential equations (London, 1989), 425–435, Inst. Math. Appl. Conf. Ser. New Ser., 39, Oxford Univ. Press, New York, 1992.
- [167] Y. Long, *Index theory for symplectic paths with applications*, Progress in Mathematics, 207. Birkhäuser Verlag, Basel, 2002. xxiv+380 pp. ISBN: 3-7643-6647-8
- [168] Y. Long, D. Zhang, C. Zhu, *Multiple brake orbits in bounded convex symmetric domains*, Adv. Math. **203** (2006), 568–635.
- [169] J. Marsden, T. Ratiu, *Introduction to Mechanics and Symmetry*, 2nd edition, Text in Appl. Math. **17**, Springer Verlag (1999).
- [170] T. Matukuma, *On the periodic orbits in Hill's case*, Proc. Imp. Acad. Japan **6** (1930), 6–8, **8** (1932), 147–150, **9** (1933), 364–366.
- [171] D. McDuff, D. Salamon, *Introduction to Symplectic Topology* 2nd edition, Oxford University Press (1998).
- [172] D. McDuff, D. Salamon, *J-holomorphic Curves and Symplectic Topology* 2nd edition, Amer. Math. Soc., Providence, RI (2012).
- [173] R. McGehee, *Some homoclinic orbits for the restricted three-body problem*, Thesis (Ph.D.)—The University of Wisconsin - Madison. 1969. 63 pp.
- [174] W. Merry, *On the Rabinowitz Floer homology of twisted cotangent bundles*, Calc. Var. Partial Differential Equations **42** (2011), no. 3-4, 355–404.
- [175] K. Meyer, D. Schmidt, *Hill's lunar Equations and the Three-Body Problem*, J. Diff. Eq. **44**, (1982), 263–272.

- [176] J. Milnor, *Topology from the Differential Viewpoint*, The University Press of Virginia (1965).
- [177] J. Milnor, *On the geometry of the Kepler problem*, Amer. Math. Monthly **90**(6), (1983), 353–365.
- [178] H. Mineur, *Sur les systèmes mécaniques admettant  $n$  intégrales premières uniformes et l'extension à ces systèmes de la méthode de quantification de Sommerfeld*, C. R. Acad. Sci., Paris **200** (1935), 1571–1573.
- [179] H. Mineur, *Sur les systèmes mécanique dans lesquels figurent des paramètres fonctions du temps. Étude des systèmes admettant  $n$  intégrales premières uniformes en involution. Extension à ces systèmes des conditions de quantification de Bohr-Sommerfeld*, Journal de l'École Polytechnique, Série III, 143ème année (1937), 173–191 and 237–270.
- [180] R. Moeckel, *Lectures on central configurations*, downloaded from [www-users.math.umn.edu/~rmoeckel/notes/CentralConfigurations](http://www-users.math.umn.edu/~rmoeckel/notes/CentralConfigurations)
- [181] C. Moore, *Braids in Classical Gravity*, Physical Review Letters **70**, 3675–3679 (1993).
- [182] R. Moore, R. Kearfott, M. Cloud, *Introduction to interval analysis*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2009. xii+223 pp. ISBN: 978-0-898716-69-6
- [183] E. Mora, *Pseudoholomorphic cylinders in symplectisations*, Doctoral dissertation, New York University (2003).
- [184] M. Morse, *Calculus of variations in the large*, American Mathematical Society Colloquium Publication **18**, Providence, RI (1934).
- [185] J. Moser, *Regularization of Kepler's problem and the averaging method on a manifold*, Comm. Pure Appl. Math. **23** (1970), 609–636.
- [186] J. Moser, *Periodic orbits near an equilibrium and a theorem by Alan Weinstein*, Comm. Pure Appl. Math. **29** (1976), 724–747.
- [187] J. Moser, C. Siegel, *Lectures on celestial mechanics*, Die Grundlehren der mathematischen Wissenschaften **187**, Berlin, Springer (1971).
- [188] J. Moser, E. Zehnder, *Notes on dynamical systems*, Courant lecture notes in mathematics **12**, New York, NY, Courant Inst. of Math. Sciences (2005).
- [189] F. Moulton, *Periodic Orbits*, Carnegie Inst. of Washington, Washington, D.C. (1920).
- [190] D. Mumford, J. Fogarty, F. Kirwan, *Geometric invariant theory*, vol. **34** of *Ergebnisse der Mathematic und ihrer Grenzgebiete*, Springer, Berlin, 3. edition (1994).

- [191] W. Neutsch, K. Scherer, *Celestial mechanics*, Mannheim, BI-Wiss.-Verl. (1992).
- [192] P. Newstead, *Geometric invariant theory*, In *Moduli spaces and vector bundles*, vol. **359** of London Math. Soc. Lecture Note Ser., (2009), 99–127.
- [193] D. Ó Mathúna, *Integrable systems in celestial mechanics*, Progress in mathematical physics **51**, Boston u. a., Birkhäuser (2008).
- [194] J. Ortega, T. Ratiu, *Momentum maps and Hamiltonian reduction*, Progress in Math. **222**, Birkhäuser, Boston (2004).
- [195] L. O’shea, R. Sjamaar, *Moment maps and Riemannian symmetric pairs*, Math. Ann. **31** (2000), 415–457.
- [196] Y. Osipov, *Geometrical interpretation of Kepler’s problem*, *Uspehi Mat. Nauk*, **27**(2) (1972), 161 (In Russian).
- [197] Y. Osipov, *The Kepler problem and geodesic flows in spaces of constant curvature*, *Celest. Mech.* **16** (1977), 191–208.
- [198] Y. Ostrover, *When symplectic topology meets Banach space geometry*, Proceedings of the ICM 2014, vol 2, 959–981.
- [199] J. Palis, C. Pugh, *Fifty problems in dynamical systems*, in *Dynamical Systems*, Lect. Notes Math. **468** (1975), 345–353.
- [200] W. Pauli, *Über das Modell des Wasserstoffmolekülions*, *Annalen der Physik* **68** (1922), 177–240.
- [201] W. Pauli, *Über das Wasserstoffspektrum vom Standpunkt der neuen Quantenmechanik*, *Zeitschrift für Physik* **36** (1926), 336–363.
- [202] N. de Paulo, P. Salomão, *Systems of transversal sections near critical energy levels of Hamiltonian systems in  $\mathbb{R}^4$* , arXiv:1310.8464
- [203] H. Plummer, *On periodic orbits in the neighbourhood of centres of libration*, *Monthly Notices Roy. Astron. Soc.* **62**, 6 (1901).
- [204] H. Poincaré, *Les Méthodes Nouvelles de la Mécanique Céleste I-III*, Gauthiers-Villars, Paris (1899).
- [205] H. Poincaré, *Sur les lignes géodésiques des surfaces convexes*, *Transaction Amer. Math. Soc.* **6** (1905), 237–274.
- [206] H. Poincaré, *Sur un théorème de géométrie*, *Rend. Circ. Matem. Palermo* **33** (1912), 375–407.
- [207] C. Pugh, C. Robinson, *The  $C^1$  closing lemma, including Hamiltonians*, *Ergodic Theory Dynam. Systems* **3** (1983), no. 2, 261–313.

- [208] P. Rabinowitz, *Periodic solutions of Hamiltonian systems*, Comm. Pure Appl. Math. **31** (1978), 157–184.
- [209] A. Robadey, *Différentes modalités de travail sur le général dans les recherches de Poincaré sur les systèmes dynamiques*, Thèse de Doctorat, Université Paris 7 Denis Diderot (2006).
- [210] J. Robbin, D. Salamon, *The Maslov index for paths*, Topology **32** (1993), 827–844.
- [211] J. Robbin, D. Salamon, *The spectral flow and the Maslov index*, Bull. L.M.S. **27** (1995), 1–33.
- [212] C. Robinson, *A global approximation theorem for Hamiltonian systems*, Proc. Symp. Pure Math. **XIV**, Global Analysis, AMS (1970), 233–244.
- [213] F. Schlenk, *Embedding problems in symplectic geometry*, de Gruyter Expositions in Mathematics **40**, Walter de Gruyter, Berlin (2005).
- [214] F. Schlenk, *Symplectic embedding problems, old and new*, preprint.
- [215] M. Schwarz, *Cohomology operations from  $S^1$ -Cobordism in Floer Homology*, Ph.D thesis, ETH Zürich (1996).
- [216] M. Schwarz, *Equivalences for Morse Homology*, Geometry and Topology in Dynamics, Contemp. Math. **246**, AMS, Providence, RI (1999), 197–216.
- [217] H. Seifert, *Periodische Bewegungen mechanischer Systeme*, Math. Z. **51** (1948), 197–216.
- [218] R. Siefring, *Relative asymptotic behavior of pseudoholomorphic half-cylinders*, Comm. Pure Appl. Math. **61** (2008), no. 12, 1631–1684.
- [219] R. Siefring, *Intersection theory of punctured pseudoholomorphic curves*, Geom. Topol. **15** (2011), no. 4, 2351–2457.
- [220] R. Siefring, *Finite-energy pseudoholomorphic planes with multiple asymptotic limits*, arXiv:1607.00324
- [221] C. Simó, T.J. Stuchi, *Central stable/unstable manifolds and the destruction of KAM tori in the planar Hill problem* Phys. D **140** (2000), no. 1-2, 1–32.
- [222] C. Simó, *New families of solutions in  $N$ -body problems*, (English summary) European Congress of Mathematics, Vol. I (Barcelona, 2000), 101–115, Progr. Math., 201, Birkhäuser, Basel, 2001.
- [223] J. Souriau, *Quantification géométrique. Application.*, Ann. Inst. H. Poincaré **6**, 311–341.
- [224] J. Souriau, *Structures des Systèmes Dynamiques*, Dunod, Paris (1970).

- [225] J. Souriau, *Sur la variété de Képler*, in *Convegno die Geometria Simpletica e Fisica Matematica*, Ist. Naz. di Alta Matematica, Roma 1973, Symposia Mathematica, Vol. XIV, Academic Press, London (1974).
- [226] J. Souriau, *Géométrie globale du problème à deux corps*, Atti Accad. Sci. Torino CI. Sci. Fis. Mat. Natur. **117**(1) (1983), 369–418.
- [227] S. Suhr, K. Zehmisch, *Linking and closed orbits*, Abh. Math. Semin. Univ. Hambg. **86** (2016), no. 1, 133–150.
- [228] F. Spirig, J. Waldvogel, *Chaotic motion in Hill's lunar problem*, From Newton to chaos, 217–230, NATO Adv. Sci. Inst. Ser. B Phys. **336**, Plenum, New York (1995).
- [229] E. Strömberg, *Connaissance actuelle des orbites dans le problème des trois corps*, Bull. Astron. **9** (1935), 87–130.
- [230] V. Szebehely, *Theory of Orbits - The Restricted Problem of Three Bodies*, Academic Press, New York (1967).
- [231] B. Szökefalvi-Nagy, *Spektraldarstellungen linearer Transformationen des Hilbertschen Raumes*, Ergebnisse der Mathematik und ihrer Grenzgebiete. Berlin: Springer (1942).
- [232] C. Taubes, *The Seiberg-Witten equations and the Weinstein conjecture*, Geom. Topol. **11** (2007), 2117–2202.
- [233] J. Thorpe, *Elementary Topics in Differential Geometry*, Undergraduate texts in mathematics, New York, Springer (1979).
- [234] W. Tucker, *Validated numerics. A short introduction to rigorous computations*. Princeton University Press, Princeton, NJ, 2011. xii+138 pp. ISBN: 978-0-691-14781-9
- [235] H. Varvoglis, C. Vozikis, K. Wodnar, *The two fixed centers: An exceptional integrable system*, Celestial Mech. Dyn. Astronomy **89** (2004), 343–356.
- [236] C. Viterbo, *Metric and isoperimetric problems in symplectic geometry*, J. Amer. Math. Soc. **13**, no. 2, (2000), 411-431.
- [237] T. Vozmischeva, *Integrable Problems of Celestial Mechanics in Spaces of Constant Curvature*, Kluwer Academic Publishers, Dordrecht (2003).
- [238] A. Weinstein, *Symplectic manifolds and their lagrangian submanifolds*, Adv. in Math. **6** (1971), 329–346.
- [239] A. Weinstein, *Lectures on Symplectic manifolds*, CBMS Conference Series **29**, American Mathematical Society, Providence, RI (1977).
- [240] A. Weinstein, *On the hypothesis of Rabinowitz periodic orbit theorems*, J. Diff. Eq. **33** (1979), 353–358.

- [241] A. Wintner, *Über die Konvergenzfragen der Mondtheorie*, Math. Z. **30** (1929), no. 1, 211–227.
- [242] A. Wintner, *The Analytical Foundations of Celestial Mechanics*, Princeton University Press (1941).
- [243] Z. Xia, *Melnikov Method and Transversal Homoclinic Points in the Restricted Three-Body Problem*, Jour. Diff. Eq. **96** (1992), 170–184.
- [244] A. Zinger, *Pseudocycles and integral homology*, Trans. Amer. Math. Soc. **360** (2008), no. 5, 2741–2765.