


Complexes of Ni(II) and Zn(II) as models for hydrolytic enzymes

Hendrik Schwöppe, Dirk Volkmer, Bernt Krebs

Anorganisch-Chemisches Institut, Universität Münster, Wilhelm-Klemm-Strasse 8, 48149 Münster, Germany
schwoph@nwz.uni-muenster.de



Hydrolytic enzymes play an important role in nature. In order to mimic the active sites of that type of enzymes we synthesized complexes of Zn(II) and Ni(II) with the new ligand 1-[N,N-bis(benzimidazolylmethyl)]amino-2,3-dihydroxypropane (bdapoh). The complexes were characterized by X-ray crystallography and tested for their hydrolytic activity towards 4-nitrophenylacetate (npa).

Crystals of $[\text{Zn}_2(\text{bdapo})_2 \text{C}_6\text{H}_5\text{CO}_2](\text{NO}_3)_2 \cdot 2\text{CH}_3\text{OH}$ (**1**) were obtained from $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$, $\text{Na}(\text{C}_6\text{H}_5\text{CO}_2)$ and bdapoh in methanol. $[\text{Ni}_3(\text{bdapo})_2(\text{CH}_3\text{COO})_2(\text{ClO}_4)_2] \cdot \text{C}_4\text{H}_{10}\text{O} \cdot 6\text{CH}_3\text{OH}$ (**2**) was obtained as dark green crystals from $\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$, $\text{Ni}(\text{CH}_3\text{COO})_2 \cdot 4\text{H}_2\text{O}$ and bdapoh in methanolic solution.

Kinetic measurements were carried out in buffered ethanol-water mixtures 1:1 at constant pH. We used the biological buffers mops (pH < 7) and hepes (pH > 7). The hydrolysis rate of npa was measured by monitoring the increase of the absorption at 400 nm of the released 4-nitrophenolate. The activity of the complex

decreases from Zn^{2+} ($k = 1.69 \cdot 10^{-3} \text{ mol}^{-1} \text{ s}^{-1}$) to Ni^{2+} ($k = 0.66 \cdot 10^{-3} \text{ mol}^{-1} \text{ s}^{-1}$). At higher pH the hydrolytic activity of the complexes increases. Using low temperature $^1\text{H-NMR}$ we determined an equilibrium of mono- and dinuclear species in solution for (**1**). The free enthalpy of activation for the dinuclear complex is -91.2 kJ/mol.