

Turbulent cascades in foreign exchange markets

S. Ghashghaie*, W. Breymann†, J. Peinke‡, P. Talkner§ & Y. Dodge||

* Fürstensteinerstrasse 4, 4053 Basel, Switzerland

† Institute für Physik der Universität Basel, 4056 Basel, Switzerland

‡ Experimentalphysik II, Universität Bayreuth, 95440 Bayreuth, Germany

§ Paul Scherrer Institut, 5232 Villigen, Switzerland

|| Groupe de Statistiques, Université de Neuchâtel, 2000 Neuchâtel, Switzerland

THE availability of high-frequency data for financial markets has made it possible to study market dynamics on timescales of less than a day¹. For foreign exchange (FX) rates Müller *et al.*² have shown that there is a net flow of information from long to short timescales: the behaviour of long-term traders (who watch the markets only from time to time) influences the behaviour of short-term traders (who watch the markets continuously). Motivated by this hierarchical feature, we have studied FX market

dynamics in more detail, and report here an analogy between these dynamics and hydrodynamic turbulence³⁻⁸. Specifically, the relationship between the probability density of FX price changes (Δx) and the time delay (Δt) (Fig. 1a) is much the same as the relationship between the probability density of the velocity differences (Δv) of two points in a turbulent flow and their spatial separation Δr (Fig. 1b). Guided by this similarity we claim that there is an information cascade in FX market dynamics that corresponds to the energy cascade in hydrodynamic turbulence. On the basis of this analogy we can now rationalize the statistics of FX price differences at different time delays, which is important for, for example, option pricing. The analogy also provides a conceptual framework for understanding the short-term dynamics of speculative markets.

A flow of energy from large to small scales is one of the main characteristics of fully developed homogeneous isotropic turbulence in three spatial dimensions²⁴. It provides a mechanism for dissipating large amounts of energy in a viscous fluid. Energy is pumped into the system at large scales of the order of, say, metres (by a moving car or a flying aeroplane) or kilometres (by meteorological events), transferred to smaller scales through a hierarchy of eddies of decreasing sizes, and dissipated at the smallest scale—of the order of millimetres in the above examples. This cascade of kinetic energy extending over several orders of magnitude generates a scaling behaviour of the eddies and manifests itself in a scaling of the moments $\langle (\Delta v)^n \rangle$ of Δv as $(\Delta r)^{\zeta_n}$ (refs 5,6). Here the angle brackets $\langle \rangle$ denote the mean value of the enclosed quantity and Δv is the difference of the velocity component in the direction of the spatial separation of length Δr . Under the assumption that the eddies of each size are space-filling and that the downward energy flow is homogeneous, $\zeta_n = n/3$ (ref. 8). The probability densities $P_{\Delta r}(\Delta v)$ are then scale invariant. This means that if the velocity differences are normalized by their respective standard deviation, the resulting standardized probability densities do not depend on Δr . But for $n > 3$, experimentally determined values of ζ_n follow a concave curve definitely below the ($\zeta_n = n/3$)-line (Fig. 2b). The dependence of the standardized probability density on Δr also provides evidence that eddies of a given size are not space-filling but rather fluctuating in space and time in a typical intermittent way (Fig. 1b).

Our analyses of FX markets are based on a data set provided by Olsen and Associates containing all worldwide 1,472,241 bid-ask quotes for US dollar-German mark exchange rates which have emanated from the interbank Reuters network from 1 October 1992 until 30 September 1993. From these data we have determined the probability densities of price changes $P_{\Delta t}(\Delta x)$ with time delays Δt varying from five minutes up to approximately two days, which are displayed in Fig. 1a. In comparison, Fig. 1b shows the analogous turbulent probability densities $P_{\Delta r}(\Delta v)$, which exhibit the same characteristic features. Using the probability density $P_{\Delta t}(\Delta x)$, the information acquired by observing the market after a time Δt can be quantified as $I(\Delta t) = -\int P_{\Delta t}(\Delta x) \log P_{\Delta t}(\Delta x) d(\Delta x)$. It turns out that the dependence of this information on Δt is directly related to the scaling of the variance of Δx with Δt . In turbulence, on the other hand,

the variance of the velocity differences at a distance Δr is proportional to the mean energy which is contained in an eddy of size Δr . This further supports the proposed analogy between energy and information.

Given the analogy between turbulence and FX market dynamics, we expect the moments of FX price changes to scale with the time delay as $\langle (\Delta x)^n \rangle \propto (\Delta t)^{\zeta_n}$. Scaling has already been reported for the mean absolute values of FX returns in ref. 9 and by Evertsz in ref. 1, and for the second moments of the variations of the Standard & Poor's 500 economic index in ref. 10. For FX




FIG. 1 a, Data points: standardized probability density $P_{\Delta t}(\Delta x)$ of price changes $\Delta x = x(t) - x(t + \Delta t)$ for time delays $\Delta t = 640$ s, 5,120 s, 40,960 s, 163,840 s (from top to bottom). The middle prices $x(t) = (x_{\text{bid}}(t) + x_{\text{ask}}(t))/2$ have been used (data provided by Olsen & Associates (see text)). The probability density has been calculated in a similar way as ref. 10. Full lines: results of (least-squares) fits carried out according to ref. 7; $\lambda^2 = 0.25, 0.23, 0.13, 0.06$ (from top to bottom). For better visibility, the curves have been vertically shifted with respect to each other. Note the systematic change in shape of the densities. b, Data points: standardized probability density $P_{\Delta r}(\Delta v)$ of velocity differences $P_{\Delta r}(\Delta v)$ for a turbulent flow with $\Delta r = 3.3\eta, 18.5\eta, 138\eta, 325\eta$ (data taken from ref. 13, $R_i = 598$. Here η is the Kolmogorov scale, where viscous dissipation occurs. Full lines: results of (least-squares) fits carried out according to ref. 7; $\lambda^2 = 0.19, 0.10, 0.04, 0.01$.

market data, it has also been observed that the kurtosis, which is the ratio of the fourth moment and the squared variance, decreases with increasing time delay, that is, the tails of the probability density lose weight^{9,11}. Figure 2a shows the higher-order moments of the FX price changes. They scale for time delays Δt varying from about five minutes up to several hours. As in turbulence, the scaling exponents ξ_n depend on the order of the moments in a nonlinear way. Moreover, the ξ_n of the FX data are close to the ζ_n of turbulent data (Fig. 2b). This is also in agreement with the observation that the shapes of the turbulent and FX probability densities, $P_{\Delta r}(\Delta v)$ and $P_{\Delta r}(\Delta x)$, depend on their respective scale parameters in a similar way: both show a decrease of kurtosis with increasing scale parameter¹² (Fig. 1a and 1b).

Even though the probability densities $P_{\Delta r}(\Delta x)$ and $P_{\Delta r}(\Delta v)$ are in principle determined by their moments, this is of little help in practice because errors drastically increase when calculating higher-order moments from observed data. In turbulence, for different experimental situations, the change in shape of $P_{\Delta r}(\Delta v)$ has been successfully parametrized using a method motivated by the energy cascade^{7,13,14}. The standardized probability density is approximated by a superposition of gaussian densities with log-normally distributed variances. The variance λ^2 of this log-normal distribution is a measurable form parameter containing information on the energy cascade¹⁵. It is the only parameter that must be




FIG. 2 a, Scaling of moments $\langle (\Delta x)^n \rangle$ with $n = 2, 4$ and 6 . b, n -dependence of the scaling factors ξ_n and ζ_n for the n th moments of the probability densities of FX price changes (squares) and hydrodynamic velocity differences taken from ref. 22 (crosses) and ref. 23 (triangles). Note the same qualitative deviation of all curves from a straight line.

TABLE 1 Correspondence between fully developed three-dimensional turbulence and FX markets

Hydrodynamic turbulence	FX markets
Energy	Information
Spatial distance	Time delay
Laminar periods interrupted by turbulent bursts (intermittency)	Clusters of low and high volatility
Energy cascade in space hierarchy $\langle (\Delta v)^n \rangle \propto (\Delta r)^{\xi_n}$	Information cascade in time hierarchy $\langle (\Delta x)^n \rangle \propto (\Delta t)^{\xi_n}$

adjusted to the data. Applying this method to the FX market data yields a surprisingly good fit (Fig. 1a). Although the data in the centre dominate the fit, the agreement is reasonable also as regards the tails of the probability densities as long as the time delay is shorter than about two days. This is in contrast to fits with Levy distributions which significantly deviate in the tails¹⁰. The parameter λ^2 , which measures the spread of the variances of the superimposed gaussians (that is, the variance of the log-normal distribution), decreases linearly with $\log \Delta t$ (Fig. 3). This result further confirms the similarity of the statistical behaviour of FX markets with the classical picture of turbulence as given by Kolmogorov⁵⁻⁷. In particular, it verifies the scaling of the moments, $\langle (\Delta x)^n \rangle \propto (\Delta t)^{\xi_n}$, in an independent way.

An important aspect of turbulent flows is their intermittent behaviour, that is, the typical occurrence of laminar periods which are interrupted by turbulent bursts. In FX markets this corresponds to clusters of high and low volatility¹², which give rise to relatively high values of the probability densities $P_{\Delta r}(\Delta x)$ both in the centre and the tails. This particular aspect is well reproduced for time delays smaller than two days by the proposed log-normal superpositions of gaussians. However, long interruptions of the trading process (particularly during weekends) affect the probability densities and lead to deviations from the expected shape. Hence, we conclude that the range which is governed by the information cascade has an upper limit. We note that in turbulence also there is an upper length scale, where the energy cascade sets in and beyond which scaling fails.

Different types of models have been proposed to describe the statistical characteristics of price differences of financial indices as, for example, the behaviour of the volatility. Prominent approaches are models using mixtures of distributions¹⁶⁻¹⁸ and ARCH/GARCH-type models^{19,20} (see, for example, the overview in ref. 21). But these studies do not address the scaling behaviour of the probability distributions of FX price changes.

It is unlikely that there is a set of a few partial differential equations (like the Navier Stokes equations in hydrodynamics) which might serve as a model of FX market dynamics. More striking is the similarity of the two phenomena, which is accounted for by the existence of a cascade in both cases. This has prompted us to introduce concepts of turbulence in the description of FX




FIG. 3 Dependence of the form parameter λ^2 on Δt (in seconds).

markets, in particular that of an information cascade. Table 1 summarizes the corresponding notions. How far this analogy goes still has to be shown. In any case, we have reason to believe that the qualitative picture of turbulence that has developed during the past 70 years will help our understanding of the apparently remote field of financial markets.

1. Proc. 1st Int. Conf. on High Frequency Data in Finance (Olsen & Associates, Zürich, 1995).
2. Müller, U. A. et al. *J. empirical Fin.* (in the press).
3. Landau, L. D. & Lifshitz, E. M. *Fluid Mechanics* 2nd edn (Pergamon, Oxford, 1987).
4. Monin, A. S. & Yaglom, A. M. *Statistical Fluid Mechanics* Vols 1 & 2 (ed. Lumely, J.) (MIT Press, Cambridge, MA, 1971 & 1975).
5. Kolmogorov, A. N. *J. Fluid Mech.* **13**, 82–85 (1962).
6. Obukhov, A. M. *J. Fluid Mech.* **13**, 77–81 (1962).
7. Castaing, B., Gagne, Y. & Hopfinger, E. *Physica* **D46**, 177–200 (1990).
8. Kolmogorov, A. N. *Dokl. Akad. Nauk. SSSR* **30**, 301–305 (1941).
9. Müller, U. A. et al. *J. Banking Fin.* **14**, 1189–1208 (1990).
10. Mantegna, R. N. & Stanley, H. E. *Nature* **376**, 46–49 (1995).
11. Baillie, R. T. & Bollerslev, T. *J. Business econ. Statist.* **7**, 297–305 (1989); *Rev. econ. Stud.* **58**, 565–585 (1991).
12. Vassilicos, J. C. *Nature* **374**, 408–409 (1995).
13. Chabaud, B. et al. *Phys. Rev. Lett.* **73**, 3227–3230 (1994).
14. Peinke, J. et al. in *Fractals in the Natural and Applied Sciences* Vol. A41, (ed. Novak, M. M.) 295–304 (Elsevier Science Amsterdam, 1994).
15. Naert, A. et al. *J. Phys. II Fr.* **4**, 215–224 (1994).
16. Granger, C. W. & Orr, D. J. *J. Am. statist. Ass.* **67**, 275–285 (1972).
17. Clark, P. K. *Econometrica* **41**, 135–155 (1973).
18. Kon, S. J. *J. Fin.* **39**, 147–165 (1984).
19. Engle, R. F. *Econometrica* **50**, 987–1007 (1982).
20. Bollerslev, T., Chous, R. Y. & Kroner, K. F. *J. Econometrics* **52**, 5–59 (1992).
21. Taylor, S. J. *Math. Fin.* **4**, 183–204 (1994).
22. Stolovitzky, G., Sreenivasan, K. R. & Juneja, A. *Phys. Rev.* **E48**, 3217–3220 (1993).
23. Anselmet, F., Gagne, Y., Hopfinger, E. J. & Antonia, R. A. *J. Fluid Mech.* **140**, 63–89 (1984).
24. Richardson, L. F. *Weather Prediction by Numerical Process* 66 (Cambridge Univ. Press, 1922).

ACKNOWLEDGEMENTS. Some of this work was done at the Institute for Theoretical Physics of the Eötvös University in Budapest; S.G. and W.B. express their thanks for the hospitality. The authors thank C. Beck, B. Castaing, R. Friedrich, H. Gersbach, G. Györgyi, O. Korn and T. Tel for discussions, and C. Jeeroburkhan for reading the manuscript. We also thank T. Gehrig for valuable remarks about the market behaviour. W.B. acknowledges partial support of the Swiss National Science Foundation. J.P. acknowledges support by a Heisenberg fellowship of the Deutsche Forschungsgemeinschaft. We also thank Olsen & Associates for providing the FX data.

CORRESPONDENCE should be addressed to W.B. (e-mail: breymann@ubaclu.unibas.ch).