

Homogenization of linearized elasticity in a two-component medium with slip displacement conditions

Tanja Lochner^{a,*}, Malte A. Peter^{a,b}

^a*Institute of Mathematics, University of Augsburg, 86135 Augsburg, Germany*

^b*Augsburg Centre for Innovative Technologies, University of Augsburg, 86135 Augsburg, Germany*

Abstract

We consider the periodic homogenization of the linearized elasticity problem with slip displacement conditions for a two-scale composite of two solids. Such interface jumps in displacement arise e.g. in contact problems with imperfect bonding. Motivated by applications in carbon-fibre-reinforced concretes, we aim to investigate the impact of the length of the carbon fibres on the effective properties, which is why we assume one of the materials to be connected, whereas the other one is either connected or disconnected. The method of periodic unfolding is applied, deriving some new compactness results in passing, to determine the macroscopic limit problems rigorously, which show significant differences for the two cases.

Keywords: periodic homogenization, linear elasticity, imperfect interface, slip displacement

1. Introduction

In a two-scale composite of two solids, we consider the upscaling in the context of periodic homogenization of the linear elasticity problem with slip displacement conditions in normal and tangential direction at the internal interface, and Dirichlet and Neumann boundary conditions at separate parts of the outer boundary. While one of the components is assumed globally connected, we consider two distinct cases for the other component: globally connected or disconnected.

The problem is motivated by recently proposed concretes reinforced with short carbon fibres, cf. [1], [2], [3]. Similarly to carbon-fibre-reinforced ceramics, the bonding of the fibres to the concrete matrix is rather weak, so that it cannot be modelled as a perfect bond. Hence, we apply general linear slip displacement conditions. Moreover, the carbon fibres in this new material occur as single fibres and they are rather short (of the order of 1 cm in length). Therefore, the question arises what connectivity should be assumed in the homogenization setting. We investigate the impact of this choice by considering the two different connectivities alluded to above as extreme cases.

Related to our choice of internal interface condition, there are several authors studying homogenization problems with jumps on the interface and focus on the disconnectedness of one of the subdomains. In [4], [5] and [6], thermal diffusion with homogeneous Dirichlet boundary conditions

*Corresponding author

Email addresses: tanja.lochner@math.uni-augsburg.de (Tanja Lochner),
malte.peter@math.uni-augsburg.de (Malte A. Peter)

on the outer boundary is considered. To derive the homogenized solution in the first paper, Tartar's method of oscillating test functions [7] was applied and, in the second and third, the periodic unfolding method [8] was used. Moreover, in this scalar diffusion setting, scaling of the interfacial term was considered systematically in [9] in terms of two-scale convergence [10, 11]. Nonlinear variants of these problems were considered more recently by [12], [13] and [14]. In [15], the (vector-valued) elasticity problem with prescribed interface jumps in displacements and tractions and Neumann condition on a part of the external boundary was studied. A related vector-valued problem with perfect bonding was recently considered by [16].

As mentioned above, the connectivity of the domain is known to have a significant impact. This is highlighted in [9] in terms of two-scale-convergence compactness results. We refer to [4], [5], [12] and [6] for the impact in terms of the upscaled system of equations.

Although the periodic homogenization of linearized elasticity is well-known for standard (internal and external) boundary conditions in the literature, cf. e.g. [17] and [18], difficulties arise due to the fact that the slip displacement conditions only yield Robin-type interface conditions, which complicates the a-priori estimates of the symmetric gradient. In the disconnected case, we can resolve this issue by choosing an appropriate solution space and using standard extension operators. In the connected case, we use the recently developed extension operators from [19] to handle the homogeneous Dirichlet boundary conditions at the external boundary.

The paper is organized as follows. In section 2, we state the problem for a composite with periodic microstructure, for which we prove the existence and uniqueness of the solution in the connected and disconnected case in section 3. In section 4, we derive the homogenized problems in both cases, where general compactness results are proven first, which may be of general interest independent of the particular boundary-value problem considered here. A comparison and discussion of the limit problems is given in section 5.

2. Statement of the problem for a composite with periodic microstructure

We consider a composite of two materials with a periodic microstructure and distinguish two cases. In the disconnected case, one material is connected, whereas the other one is not. On the other hand, in the connected case, the composite consists of two globally connected materials. We choose a bounded connected Lipschitz-domain $\Omega \subset \mathbb{R}^3$, which can be represented as a union of axis-parallel cuboids and which has corner coordinates in \mathbb{Q}^3 .

In order to describe the periodic microstructure, we define the reference cell $Y = [0, 1]^3 \subset \mathbb{R}^3$ and two subsets $Y_0, Y_1 \subset Y$ such that $Y_0^\circ \cap Y_1^\circ = \emptyset$, $\Sigma_Y := \overline{Y_0} \cap \overline{Y_1}$ Lipschitz-continuous and $Y = Y_0 \cup \Sigma_Y \cup Y_1$. In the disconnected case, we assume that $Y_0 \subset\subset Y$, i.e. $\partial Y_0 \cap \partial Y = \emptyset$, and, in the connected one, we prescribe that the boundary of Y_0 resp. Y_1 on opposite faces is the same and if we extend Y 1-periodically, this splits the space into two connected sets. We define the following sets:

- $\Omega_\alpha^\varepsilon := \bigcup_{\xi \in \Lambda^\varepsilon} \varepsilon(Y_\alpha + \xi)$ is the Y -periodically extended domain Y_α scaled with ε , $\alpha = 0, 1$,
- $\Sigma^\varepsilon := \bigcup_{\xi \in \Lambda^\varepsilon} \varepsilon(\Sigma_Y + \xi)$ is the Y -periodically extended interface Σ_Y scaled with ε ,
- $\Omega^\varepsilon = \Omega_0^\varepsilon \cup \Omega_1^\varepsilon \cup \Sigma^\varepsilon$ is the union of the scaled cells,

where $\Lambda^\varepsilon := \{\xi \in \mathbb{Z}^3 : \varepsilon(Y + \xi) \subset \Omega\}$. The idea is that Ω_0^ε represents one material and Ω_1^ε the other one. We split the external boundary into two parts $\partial\Omega = \Gamma_1 \cup \Gamma_2$, where Γ_1 and Γ_2 are disjoint sets

and Γ_1 has positive measure. In both cases, we only admit the scaling factor ε in such a way that $\varepsilon^{-1}\Omega$ can be represented as a finite union of axis-parallel cuboids with corner coordinates in \mathbb{Z}^3 . Thus, we can guarantee that the domain Ω can completely be filled up with scaled reference cells. This condition can be relaxed but we assume it in what follows to avoid well-known technicalities induced by otherwise non-matching boundaries of Ω and its ε -periodic approximation. Due to the choice of ε , there holds $\Omega^\varepsilon = \Omega$, $\Gamma_1 = \Gamma_1 \cap \partial\Omega^\varepsilon$ and $\Gamma_2 = \Gamma_2 \cap \partial\Omega^\varepsilon$. Let n be the normal to Σ^ε with orientation from Ω_0^ε to Ω_1^ε and ν the outward-pointing normal to Γ_2 .

We consider the linear elasticity equation with slip displacement conditions on the interface of the materials. We let $u_\varepsilon: \Omega^\varepsilon \rightarrow \mathbb{R}^3$ be the displacement field. The stress tensor $\sigma^\varepsilon = (\sigma_{ij}^\varepsilon)_{1 \leq i, j \leq 3}$ is defined by

$$\sigma_{ij}^\varepsilon := \sum_{k, l=1}^3 a_{ijkl}^\varepsilon e_{kl}(u^\varepsilon) := \sum_{k, l=1}^3 a_{ijkl}^\varepsilon \frac{1}{2} (\partial_k u_l^\varepsilon + \partial_l u_k^\varepsilon),$$

where the symmetric gradient $e(u^\varepsilon) = (e_{kl}(u^\varepsilon))_{1 \leq k, l \leq 3}$ is the linear strain tensor and $A^\varepsilon(x) = (a_{ijkl}^\varepsilon(x))_{1 \leq i, j, k, h \leq 3}$ is a tensor of fourth order, which describes the properties of the materials of the solid. In order to formulate the slip displacement conditions, we introduce some additional notation:

- $u_n^\varepsilon := u^\varepsilon \cdot n$ resp. $u_{\tau^i}^\varepsilon := u^\varepsilon \cdot \tau^i$, $i = 1, 2$, the projection of the displacement field in normal resp. tangential direction of the interface,
- $\sigma_n^\varepsilon := (\sigma^\varepsilon \cdot n) \cdot n$ resp. $\sigma_{\tau^i}^\varepsilon := (\sigma^\varepsilon \cdot n) \cdot \tau^i$, $i = 1, 2$, the projection of the normal stress in normal resp. tangential direction of the interface,
- $[\varphi]_{\Sigma^\varepsilon} := (\varphi_1 - \varphi_0)|_{\Sigma^\varepsilon}$ the jump on the interface,

where $\varphi_\alpha := \varphi|_{\Omega_\alpha^\varepsilon}$ is the restriction of φ to $\Omega_\alpha^\varepsilon$ and $\varphi_\alpha|_{\Sigma^\varepsilon}$ is the trace of φ_α (we write only φ_α if it is clear) for $\alpha = 0, 1$.

With this notation, the linear elasticity problem is given by

$$\begin{cases} -\nabla \cdot \sigma^\varepsilon = f^\varepsilon & \text{in } \Omega_0^\varepsilon \cup \Omega_1^\varepsilon, \\ u^\varepsilon = 0 & \text{on } \Gamma_1, \\ \sigma^\varepsilon \cdot \nu = g & \text{on } \Gamma_2, \end{cases} \quad (1)$$

with the slip displacement condition on the interface Σ^ε

$$\begin{cases} \varepsilon [u_n^\varepsilon]_{\Sigma^\varepsilon} = \frac{1}{K_N} \sigma_n^{\Sigma^\varepsilon}, \\ \varepsilon [u_{\tau^i}^\varepsilon]_{\Sigma^\varepsilon} = \frac{1}{K_T} \sigma_{\tau^i}^{\Sigma^\varepsilon}, \quad i = 1, 2, \\ [\sigma_n^\varepsilon]_{\Sigma^\varepsilon} = 0, \\ [\sigma_{\tau^i}^\varepsilon]_{\Sigma^\varepsilon} = 0, \quad i = 1, 2, \end{cases} \quad (2)$$

where the constants $K_N, K_T > 0$ are the normal and tangential stiffness, respectively, and $\sigma^{\Sigma^\varepsilon}$ is the stress tensor of the interface. We refer to [20] for more details on the slip displacement conditions.

For simplification, we choose $A^\varepsilon \in M(\alpha, \beta, Y)$ (see Definition 2.1 below) with

$$A^\varepsilon(x) = (a_{ijkh}^\varepsilon(x))_{1 \leq i,j,k,h \leq 3} := \left(a_{ijkh} \left(\frac{x}{\varepsilon} \right) \right)_{1 \leq i,j,k,h \leq 3} = A \left(\frac{x}{\varepsilon} \right),$$

where $A(\cdot) = (a_{ijkh}(\cdot))_{1 \leq i,j,k,h \leq 3} \in M(\alpha, \beta, Y)$ and all components a_{ijkh} are Y -periodic for all $i, j, k, h \in \{1, 2, 3\}$. Under additional assumptions we can assume a more general tensor A dependent on both the microscopic and the macroscopic coordinate, see Remark 4.15 and Remark 4.21.

Definition 2.1. Let $\alpha, \beta \in \mathbb{R}$ with $0 < \alpha < \beta$ and let \mathcal{O} be an open set in \mathbb{R}^3 . We denote by $M(\alpha, \beta, \mathcal{O})$ the set of all tensors $B = (b_{ijkh})_{1 \leq i,j,k,h \leq 3}$ such that

- (i) $b_{ijkh} \in L^\infty(\mathcal{O})$ for all $i, j, k, h \in \{1, 2, 3\}$
- (ii) $b_{ijkh} = b_{jikh} = b_{khij}$ for all $i, j, k, h \in \{1, 2, 3\}$
- (iii) $\alpha|m|^2 \leq Bmm$ for all symmetric matrices m
- (iv) $|B(x)m| \leq \beta|m|$ for all matrices m

a.e. in \mathcal{O} , where

$$\begin{cases} Bm := \left((Bm)_{ij} \right)_{1 \leq i,j \leq 3} = \left(\left(\sum_{k,h=1}^3 B_{ijkh} m_{kh} \right)_{ij} \right)_{1 \leq i,j \leq 3}, \\ Bm\tilde{m} := \sum_{i,j,k,h=1}^3 b_{ijkh} m_{ij} \tilde{m}_{kh}, \\ |m| := \left(\sum_{i,j=1}^3 m_{ij}^2 \right)^{\frac{1}{2}}, \end{cases}$$

for quadratic matrices $m = (m_{ij})_{1 \leq i,j \leq 3}$ and $\tilde{m} = (\tilde{m}_{ij})_{1 \leq i,j \leq 3}$.

We test problem (1) with some sufficiently smooth function φ , use integration by parts, the boundary conditions, the fact that σ^ε is symmetric and the jump of the normal stress on Σ^ε is zero, meaning that $\sigma_0^\varepsilon = \sigma_1^\varepsilon$ on Σ^ε and thus $\sigma_0^\varepsilon, \sigma_1^\varepsilon$ coincide with $\sigma^{\Sigma^\varepsilon}$ on Σ^ε , to obtain the weak formulation

$$\begin{aligned} & \int_{\Omega_0^\varepsilon} A^\varepsilon e(u_0^\varepsilon) e(\varphi_0) dx + \int_{\Omega_1^\varepsilon} A^\varepsilon e(u_1^\varepsilon) e(\varphi_1) dx \\ & + \varepsilon \int_{\Sigma^\varepsilon} \left(K_N [u_n^\varepsilon]_{\Sigma^\varepsilon} n + K_T \sum_{i=1}^2 [u_{\tau_i}^\varepsilon]_{\Sigma^\varepsilon} \tau^i \right) \cdot (\varphi_1 - \varphi_0) dS(x) \\ & = \int_{\Omega_0^\varepsilon} f^\varepsilon \cdot \varphi_0 dx + \int_{\Omega_1^\varepsilon} f^\varepsilon \cdot \varphi_1 dx + \int_{\Gamma_2 \cap \partial \Omega_0^\varepsilon} g \cdot \varphi_0 dS(x) + \int_{\Gamma_2 \cap \partial \Omega_1^\varepsilon} g \cdot \varphi_1 dS(x). \end{aligned} \quad (3)$$

Note that we have not yet introduced the function spaces, which complete the weak formulation. This is done in the next section and it will turn out that different function spaces are required depending on the connectedness of Ω_0^ε . Throughout the paper, we denote by C a constant independent of ε whose value may change from line to line and we write $\mathcal{D}(\Omega) := \mathcal{C}_0^\infty(\Omega)$ and $\mathcal{D}(\bar{\Omega}) := \mathcal{C}^\infty(\Omega)$.

3. Existence of solutions

Before we can apply homogenization methods to obtain the upscaled problem, we have to justify the existence, uniqueness and uniform boundedness of the solution. In subsection 3.1, we introduce the complete weak form and we prove the result for the disconnected case and in subsection 3.2 for the connected case.

3.1. Existence result for disconnected case

First of all, we notice that $\Gamma_2 \cap \partial\Omega_0^\varepsilon = \emptyset$ and $\Gamma_2 \cap \partial\Omega_1^\varepsilon = \Gamma_2$ in the disconnected case, so the third term on the right-hand side in (3) disappears. Since infinitesimal rotations do not induce forces, we assume that there are no rotations, meaning that the curl, $\nabla \times \cdot$, has to be zero. Therefore, we define the solution space for the disconnected case as

$$\mathcal{W}_d(\Omega^\varepsilon) = \{u \in [L^2(\Omega^\varepsilon)]^3 : u_0 \in [H^1(\Omega_0^\varepsilon)]^3, u_1 \in [H^1(\Omega_1^\varepsilon, \Gamma_1)]^3, \nabla \times u_0 = 0 \text{ in } \Omega_0^\varepsilon\},$$

endowed with the norm

$$\|u\|_{\mathcal{W}_d(\Omega^\varepsilon)}^2 := \|e(u_0)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}}^2 + \|e(u_1)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 + \varepsilon \| [u]_{\Sigma^\varepsilon} \|_{[L^2(\Sigma^\varepsilon)]^3}^2$$

for all $u \in \mathcal{W}_d(\Omega^\varepsilon)$, whereby $H^1(\mathcal{O}, \tilde{\Gamma}) := \{u \in H^1(\mathcal{O}) : u = 0 \text{ on } \tilde{\Gamma}\}$ for some open set \mathcal{O} with $\tilde{\Gamma} \subset \partial\mathcal{O}$. In what follows, we often use the fact that

$$\nabla u_0 = \frac{1}{2} (\nabla u_0 + (\nabla u_0)^T) + \frac{1}{2} (\nabla u_0 - (\nabla u_0)^T) = \frac{1}{2} (\nabla u_0 + (\nabla u_0)^T) = e(u_0)$$

for all $u \in \mathcal{W}_d(\Omega^\varepsilon)$. Before we prove that $\mathcal{W}_d(\Omega^\varepsilon)$ is a Banach space, we gather some estimates.

Lemma 3.1. *For every $v \in \mathcal{W}_d(\Omega^\varepsilon)$, there holds*

$$\begin{aligned} \text{(i)} \quad & \varepsilon \|v_1\|_{[L^2(\Sigma^\varepsilon)]^3}^2 \leq C \left(\|v_1\|_{[L^2(\Omega_1^\varepsilon)]^3}^2 + \varepsilon^2 \|\nabla v_1\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 \right) \\ \text{(ii)} \quad & \|v_0\|_{[L^2(\Omega_0^\varepsilon)]^3}^2 \leq C \left(\varepsilon^2 \|e(v_0)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}}^2 + \varepsilon \|v_0\|_{[L^2(\Sigma^\varepsilon)]^3}^2 \right) \end{aligned}$$

for constants $C > 0$ independent of ε .

Proof. Both estimates follow by scaling and summation from standard results for the trace operator (for (i)) and Korn's inequality from [21, Corollary 5.8] (for (ii)). \square

Theorem 3.2. *($\mathcal{W}_d(\Omega^\varepsilon), \|u\|_{\mathcal{W}_d(\Omega^\varepsilon)}$) defines a Banach space.*

Proof. Using the fact that $u_1 = 0$ on Γ_1 for $u \in \mathcal{W}_d(\Omega^\varepsilon)$, it is easily proven that $\|\cdot\|_{\mathcal{W}_d(\Omega^\varepsilon)}^2$ defines a norm on $\mathcal{W}_d(\Omega^\varepsilon)$. Let $\{u^{(k)}\}_k$ be a Cauchy sequence in $\mathcal{W}_d(\Omega^\varepsilon)$. With the help of [17, Theorem 4.2], we can construct a global extension operator on domains with holes, which do not intersect the boundary of the reference cell. More precisely, there exists a linear extension operator $P_\varepsilon : [H^1(\Omega_1^\varepsilon)]^3 \rightarrow [H^1(\Omega)]^3$ such that

$$\begin{aligned} \|P_\varepsilon v\|_{[H^1(\Omega)]^3} &\leq c_1 \|v\|_{[H^1(\Omega_1^\varepsilon)]^3}, \\ \|P_\varepsilon v\|_{[L^2(\Omega)]^3} + \|e(P_\varepsilon v)\|_{[L^2(\Omega)]^{3 \times 3}} &\leq c_2 \left(\|v\|_{[L^2(\Omega_1^\varepsilon)]^3} + \|e(v)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}} \right), \\ \|e(P_\varepsilon v)\|_{[L^2(\Omega)]^{3 \times 3}} &\leq c_3 \|e(v)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}} \end{aligned} \tag{4}$$

for all $v \in [H^1(\Omega_1^\varepsilon)]^3$, where the constants c_1, c_2, c_3 do not depend on ε . Korn's inequality for functions with zero trace on part of the boundary yields

$$\|u_1\|_{[H^1(\Omega_1^\varepsilon)]^3}^2 \leq \|P_\varepsilon u_1\|_{[H^1(\Omega)]^3}^2 \leq C \|e(P_\varepsilon u_1)\|_{[L^2(\Omega)]^{3 \times 3}}^2 \leq C \|e(u_1)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2. \tag{5}$$

Using this inequality twice and Lemma 3.1 (i) and (ii), we get for all $u \in \mathcal{W}_d(\Omega^\varepsilon)$

$$\begin{aligned} & \|u_0\|_{[H^1(\Omega_0^\varepsilon)]^3}^2 + \|u_1\|_{[H^1(\Omega_1^\varepsilon)]^3}^2 \\ & \leq C \left((1 + \varepsilon^2) \|e(u_0)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}}^2 + \varepsilon \| [u]_{\Sigma^\varepsilon} \|_{[L^2(\Sigma^\varepsilon)]^3}^2 + \varepsilon \|u_1\|_{[L^2(\Sigma^\varepsilon)]^3}^2 + \|e(u_1)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 \right) \\ & \leq C \|u\|_{\mathcal{W}_d(\Omega^\varepsilon)}^2. \end{aligned} \quad (6)$$

Therefore, $\{u_0^{(k)}\}_k$ and $\{u_1^{(k)}\}_k$ are Cauchy sequences in the Banach spaces $H^1(\Omega_0^\varepsilon)$ resp. $H^1(\Omega_1^\varepsilon)$. Thus there exists $u_0 \in H^1(\Omega_0^\varepsilon)$ and $u_1 \in H^1(\Omega_1^\varepsilon)$ with $u_0^{(k)} \rightarrow u_0$ and $u_1^{(k)} \rightarrow u_1$ for $k \rightarrow \infty$. It remains to show that $u = (u_0, u_1) \in \mathcal{W}_d(\Omega^\varepsilon)$. Due to the continuity of the trace operator, u_1 is equal to zero on Γ_2 . Using the convergence of $\{u_0^{(k)}\}_k$ and the fact that $\nabla u_0^{(k)} = e(u_0^{(k)})$, we get

$$\partial_{x_j}(u_0)_i = \lim_{k \rightarrow \infty} \partial_{x_j} \left(u_0^{(k)} \right)_i = \lim_{k \rightarrow \infty} \frac{1}{2} \left(\partial_{x_j} \left(u_0^{(k)} \right)_i + \partial_{x_i} \left(u_0^{(k)} \right)_j \right) = \frac{1}{2} (\partial_{x_j}(u_0)_i + \partial_{x_i}(u_0)_j)$$

for all $i, j \in \{1, 2, 3\}$, which shows that $\nabla \times u_0 = 0$. \square

Theorem 3.3. *Let $f^\varepsilon \in [L^2(\Omega^\varepsilon)]^3$, $g \in [L^2(\Gamma_2)]^3$. Then, there exists a unique weak solution $u \in \mathcal{W}_d(\Omega^\varepsilon)$ of (3) for all admissible $0 < \varepsilon \leq 1$.*

Proof. Our aim is to apply the Lax–Milgram Theorem. Let $\varepsilon > 0$. We denote the left-hand side of (3) as a mapping $a: \mathcal{W}_d(\Omega^\varepsilon) \times \mathcal{W}_d(\Omega^\varepsilon) \rightarrow \mathbb{R}$,

$$\begin{aligned} a(u, v) &= \int_{\Omega_0^\varepsilon} A^\varepsilon e(u_0) e(v_0) dx + \int_{\Omega_1^\varepsilon} A^\varepsilon e(u_1) e(v_1) dx \\ &+ \varepsilon \int_{\Sigma^\varepsilon} \left(K_N [u_n]_{\Sigma^\varepsilon} n + K_T \sum_{i=1}^2 [u_{\tau^i}]_{\Sigma^\varepsilon} \tau^i \right) \cdot (v_1 - v_0) dS(x) \end{aligned}$$

and the right-hand side of (3) as a mapping $F: \mathcal{W}_d(\Omega^\varepsilon) \rightarrow \mathbb{R}$,

$$F(v) = \int_{\Omega^\varepsilon} f^\varepsilon \cdot v dx + \int_{\Gamma_2} g \cdot v_1 dS(x).$$

First of all, we prove that a is a continuous coercive bilinear form. The linearity in both components is clear. It is easy to compute that a is coercive using the properties of A^ε and the splitting of u_0 and u_1 in normal and tangential part. The continuity can be shown with the help of the boundedness of A^ε and Hölder's inequality. It remains to prove that F is linear and continuous. The linearity is clear. Let $v \in \mathcal{W}_d(\Omega^\varepsilon)$. Using the same estimate as in (6) for v_0 and (5) for v_1 , we obtain

$$\|v_0\|_{[L^2(\Omega_0^\varepsilon)]^3}^2 \leq C \|v\|_{\mathcal{W}_d(\Omega^\varepsilon)}^2, \quad (7)$$

$$\|v_1\|_{[L^2(\Omega_1^\varepsilon)]^3}^2 \leq C \|e(v_1)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 \leq C \|v\|_{\mathcal{W}_d(\Omega^\varepsilon)}. \quad (8)$$

Thus, Hölder's inequality leads to

$$|F(v)| \leq C \|f^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^3} \|v\|_{\mathcal{W}_d(\Omega^\varepsilon)} + \|g\|_{[L^2(\Gamma_2)]^3} \|v_1\|_{[L^2(\Gamma_2)]^3}. \quad (9)$$

It remains to estimate $\|v_1\|_{[L^2(\Gamma_2)]^3}^2$ in (9) by the norm $\|v\|_{\mathcal{W}_d(\Omega^\varepsilon)}$. We use the extension operator P_ε from (4) together with the trace operator and (5) to receive

$$\|v_1\|_{[L^2(\Gamma_2)]^3}^2 \leq C\|P_\varepsilon v_1\|_{[H^1(\Omega)]^3}^2 \leq C\|v_1\|_{[H^1(\Omega_1^\varepsilon)]^3}^2 \leq C\|e(v_1)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 \leq C\|v\|_{\mathcal{W}_d(\Omega^\varepsilon)}^2.$$

We receive

$$|F(v)| \leq C \left(\|f^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^3} + \|g\|_{[L^2(\Gamma_2)]^3} \right) \|v\|_{\mathcal{W}_d(\Omega^\varepsilon)}.$$

Therefore, all assumptions of the Lax–Milgram theorem are fulfilled and we get the existence and uniqueness of the solution. \square

In [4] and [22], a similar approach to the proof of existence of solutions was chosen in the scalar case, where Poincaré’s inequality instead of Korn’s inequality is used. The following two theorems show the uniform boundedness of the weak solutions with respect to two different norms.

Theorem 3.4. *Let $u^\varepsilon \in \mathcal{W}_d(\Omega^\varepsilon)$ be the weak solution of (3) and $\|f^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^3} \leq \tilde{C}$ for an ε -independent constant \tilde{C} . Then, there exists an ε -independent constant C with*

$$\|u^\varepsilon\|_{\mathcal{W}_d(\Omega^\varepsilon)} \leq C$$

for all admissible $0 < \varepsilon \leq 1$.

Proof. We use the estimates in the proof of Theorem 3.3 with $v = u^\varepsilon$ to get

$$\min\{\alpha, K_N, K_T\} \|u^\varepsilon\|_{\mathcal{W}_d(\Omega^\varepsilon)}^2 \leq a(u^\varepsilon, u^\varepsilon) = F(u^\varepsilon) \leq C \left(\|f^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^3} + \|g\|_{[L^2(\Gamma_2)]^3} \right) \|u^\varepsilon\|_{\mathcal{W}_d(\Omega^\varepsilon)},$$

which proves that $\|u^\varepsilon\|_{\mathcal{W}_d(\Omega^\varepsilon)} \leq \tilde{C}$ for a constant $\tilde{C} > 0$ independent of ε . \square

Theorem 3.5. *With the same assumption as in Theorem 3.4, there holds*

$$\|u^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^3}^2 + \|e(u_0^\varepsilon)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}}^2 + \|e(u_1^\varepsilon)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 \leq \tilde{C}$$

for a constant $\tilde{C} > 0$ independent of ε .

Proof. If we choose $v = u^\varepsilon$ in the estimates (7) and (8) and note that the constants there are independent of ε we get the desired result. \square

3.2. Existence result for connected case

In the connected case, we do not have to restrict the solution space as much as in the disconnected case. Therefore, we define the solution space for the connected case as

$$\mathcal{W}_c(\Omega^\varepsilon) = \{u \in [L^2(\Omega^\varepsilon)]^3 : u_0 \in [H^1(\Omega_0^\varepsilon, \Gamma_1 \cap \partial\Omega_0^\varepsilon)]^3, u_1 \in [H^1(\Omega_1^\varepsilon, \Gamma_1 \cap \partial\Omega_1^\varepsilon)]^3\}.$$

Using [19, Theorem 4.4], which states Korn’s inequality for a constant independent of ε , the space $\mathcal{W}_c(\Omega^\varepsilon)$ endowed with the norm

$$\|u\|_{\mathcal{W}_c(\Omega^\varepsilon)}^2 = \|e(u_0)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}}^2 + \|e(u_1)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 + \varepsilon \| [u]_{\Sigma^\varepsilon} \|_{[L^2(\Sigma^\varepsilon)]^3}^2$$

becomes a Banach space. Note that the norms on $\mathcal{W}_d(\Omega_\varepsilon)$ and $\mathcal{W}_c(\Omega_\varepsilon)$ are identical.

Theorem 3.6. *Let $f^\varepsilon \in [L^2(\Omega^\varepsilon)]^3$, $g \in [L^2(\Gamma_2)]^3$. Then there exists a unique solution $u^\varepsilon \in \mathcal{W}_c(\Omega^\varepsilon)$ of (3) for all admissible $0 < \varepsilon \leq 1$.*

Proof. We prove this statement via Lax–Milgram theorem like in Theorem 3.3. Let $\varepsilon > 0$ and the mapping a be defined as in Theorem 3.3. The same proof shows that a is a coercive continuous bilinear form. The right-hand side of (3) is slightly different from before, so we define the mapping $F: \mathcal{W}_c(\Omega^\varepsilon) \rightarrow \mathbb{R}$,

$$F(v) = \int_{\Omega^\varepsilon} f^\varepsilon \cdot v \, dx + \int_{\Gamma_2 \cap \partial\Omega_0^\varepsilon} g \cdot v_0 \, dS(x) + \int_{\Gamma_2 \cap \partial\Omega_1^\varepsilon} g \cdot v_1 \, dS(x). \quad (10)$$

We have to prove that $F \in \mathcal{W}_c(\Omega^\varepsilon)'$, i.e. F is linear and continuous. The linearity is clear. Let $v \in \mathcal{W}_c(\Omega^\varepsilon)$. Using Hölder's inequality and Korn's inequality ([19, Theorem 4.4] applied on Ω_0^ε and Ω_1^ε), we obtain

$$\left| \int_{\Omega^\varepsilon} f^\varepsilon \cdot v \, dx \right| \leq C \|f^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^3} \left(\|e(v_0)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}}^2 + \|e(v_1)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 \right)^{\frac{1}{2}}.$$

Let $\alpha = 0$ or 1 . We apply Hölder's inequality to get

$$\left| \int_{\Gamma_2 \cap \partial\Omega_\alpha^\varepsilon} g \cdot v_\alpha \, dS(x) \right| \leq \|g\|_{[L^2(\Gamma_2)]^3} \|v_\alpha\|_{[L^2(\Gamma_2 \cap \partial\Omega_\alpha^\varepsilon)]^3}.$$

In order to estimate the boundary terms, we use the extension operator L_α^ε from $\Omega_\alpha^\varepsilon$ to Ω defined in [19, Theorem 3.4]. Therefore, we can apply the trace operator on $L_\alpha^\varepsilon v_\alpha$, use the estimates of the trace operator and the Korn's inequality ([19, Theorem 4.4]) to get the desired estimate

$$\|v_\alpha\|_{[L^2(\Gamma_2 \cap \partial\Omega_\alpha^\varepsilon)]^3} \leq C \|L_\alpha^\varepsilon(v_\alpha)\|_{[H^1(\Omega)]^3} \leq C \|v_\alpha\|_{[H^1(\Omega_\alpha^\varepsilon)]^3} \leq C \|e(v_\alpha)\|_{[L^2(\Omega_\alpha^\varepsilon)]^{3 \times 3}}.$$

Summarizing the previous estimates, we get the continuity of the mapping F

$$|F(v)| \leq C \left(\|f^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^{3 \times 3}} + \|g\|_{[L^2(\Gamma_2)]^3} \right) \|v\|_{\mathcal{W}_c(\Omega^\varepsilon)}.$$

Lax–Milgram Theorem guarantees the existence and uniqueness of the solution. \square

Theorem 3.7. *Let $u^\varepsilon \in \mathcal{W}_c(\Omega^\varepsilon)$ be the weak solutions of (3) and $\|f^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^3} \leq \tilde{C}$ for an ε -independent constant \tilde{C} . Then, exists a constant C independent of ε such that*

$$\|u^\varepsilon\|_{\mathcal{W}_c(\Omega^\varepsilon)} \leq C$$

for all admissible $0 < \varepsilon \leq 1$.

Proof. The proof is analogous to the proof of Theorem 3.4. \square

4. Periodic homogenization

This section is devoted to the rigorous upscaling of the problem in the disconnected (subsection 4.2) and connected case (subsection 4.3). We mainly use the periodic unfolding method, which is briefly introduced together with some standard results for convenience first (subsection 4.1).

4.1. Introduction to periodic unfolding method

The periodic unfolding method was formally introduced by Doina Cioranescu, Alain Damlamian und George Griso in 2002. The main idea is to write the function, which is defined on Ω^ε , as a function of micro- and macro-scale, which makes it easier to work with perforated domains which change with ε . We refer to [8], [23] and [5] for the definitions and theorems in this section and further results. The first paper considers a fixed domain, the second a perforated domain with connected or disconnected holes and the third a domain consisting of two components.

To define the periodic unfolding operator $\mathcal{T}_i^\varepsilon, i = 0$ or 1 , we need another representation of $x \in \Omega$. Let $x \in \mathbb{R}^3$. We denote by $[x]$ the unique linear combination of the integers $\xi_j \in \mathbb{Z}$ and the periodicity vectors $b_j \in \mathbb{R}^3$, which are in our case the unit vectors e_j for $j = 1, 2, 3$, i.e. $[x] = \sum_{j=1}^3 \xi_j b_j$, such that $\{x\} := x - [x] \in Y$. Therefore we can rewrite $x \in \mathbb{R}^3$ via

$$x = \varepsilon \left(\left[\frac{x}{\varepsilon} \right] + \left\{ \frac{x}{\varepsilon} \right\} \right).$$

Definition 4.1. *Let $i = 0$ or 1 . For a Lebesgue-measurable function ϕ on Ω_i^ε , the periodic unfolding operator $\mathcal{T}_i^\varepsilon$ is defined as*

$$\mathcal{T}_i^\varepsilon(\phi)(x, y) := \begin{cases} \phi \left(\varepsilon \left[\frac{x}{\varepsilon} \right] + \varepsilon y \right) & \text{for a.e. } (x, y) \in \hat{\Omega}^\varepsilon \times Y_i, \\ 0 & \text{for a.e. } (x, y) \in \Pi^\varepsilon \times Y_i, \end{cases}$$

where $\hat{\Omega}^\varepsilon = \bigcup_{\xi \in \Lambda^\varepsilon} \varepsilon(\xi + Y)$ and $\Pi^\varepsilon = \Omega \setminus \hat{\Omega}^\varepsilon$.

Due to our assumption, there holds $\hat{\Omega}^\varepsilon = \Omega^\varepsilon$ and $\Pi^\varepsilon = \emptyset$ for admissible ε . Note that the definition of the periodic unfolding operator does not require Ω to be representable as a union of axis-parallel cuboids. We denote by \tilde{f} the extension of the function f to Ω by zero and by $\mathcal{M}_\mathcal{O}(\phi) = \frac{1}{|\mathcal{O}|} \int_{\mathcal{O}} \phi dy$ the mean value of the function ϕ over the domain \mathcal{O} , where \mathcal{O} is a set with finite measure. We summarize a number of well-known results on periodic unfolding in our notation for future reference.

Proposition 4.2. *The operator $\mathcal{T}_i^\varepsilon: L^2(\Omega_i^\varepsilon) \rightarrow L^2(\Omega \times Y_i), i = 0, 1$, is linear and continuous and*

- (i) $\mathcal{T}_i^\varepsilon(vw) = \mathcal{T}_i^\varepsilon(v)\mathcal{T}_i^\varepsilon(w)$ for all Lebesgue-measurable functions v, w in Ω_i^ε ,
- (ii) $\frac{1}{|Y|} \int_{\Omega \times Y_i} \mathcal{T}_i^\varepsilon(\phi)(x, y) dx dy = \int_{\hat{\Omega}_i^\varepsilon} \phi(x) dx = \int_{\Omega_i^\varepsilon} \phi(x) dx - \int_{\Pi_i^\varepsilon} \phi(x) dx$ for all $\phi \in L^2(\Omega_i^\varepsilon)$,
- (iii) $\|\mathcal{T}_i^\varepsilon(\phi)\|_{L^2(\Omega \times Y_i)} \leq |Y|^{\frac{1}{2}} \|\phi\|_{L^2(\Omega_i^\varepsilon)}$ for all $\phi \in L^2(\Omega_i^\varepsilon)$,
- (iv) $\mathcal{T}_i^\varepsilon(\phi) \rightarrow \phi$ strongly in $L^2(\Omega \times Y_i)$ for all $\phi \in L^2(\Omega)$,
- (v) if $\{\phi^\varepsilon\}$ is a sequence in $L^2(\Omega)$ with $\phi^\varepsilon \rightarrow \phi$ strongly in $L^2(\Omega)$, then $\mathcal{T}_i^\varepsilon(\phi^\varepsilon) \rightarrow \phi$ strongly in $L^2(\Omega \times Y_i)$,
- (vi) if $\phi \in L^2(Y_i)$ Y -periodic and $\phi^\varepsilon(x) = \phi\left(\frac{x}{\varepsilon}\right)$, then $\mathcal{T}_i^\varepsilon(\phi^\varepsilon) \rightarrow \phi$ strongly in $L^2(\Omega \times Y_i)$,
- (vii) if $\phi^\varepsilon \in L^2(\Omega_i^\varepsilon)$ with $\|\phi^\varepsilon\|_{L^2(\Omega_i^\varepsilon)} \leq C$ and $\mathcal{T}_i^\varepsilon(\phi^\varepsilon) \rightharpoonup \phi$ weakly in $L^2(\Omega \times Y_i)$, then $\tilde{\phi}^\varepsilon \rightharpoonup \frac{|Y_i|}{|Y|} \mathcal{M}_{Y_i}(\phi)$ weakly in $L^2(\Omega)$,
- (viii) if $\phi \in H^1(\Omega_i^\varepsilon)$, then $\nabla_y [\mathcal{T}_i^\varepsilon(\phi)] = \varepsilon \mathcal{T}_i^\varepsilon(\nabla \phi)$ and $\mathcal{T}_i^\varepsilon(\phi) \in L^2(\Omega, H^1(Y_i))$.

The next result shows the equivalence between classic two-scale convergence of a sequence [10, 11] and the weak convergence of the unfolded sequence.

Theorem 4.3. *Let $\{\phi^\varepsilon\}$ be a bounded sequence in $L^2(\Omega)$. The following assertions are equivalent:*

- (i) $\{\mathcal{T}^\varepsilon(\phi^\varepsilon)\}$ converges weakly to ϕ in $L^2(\Omega \times Y)$,
- (ii) $\{\phi^\varepsilon\}$ two-scale converges to ϕ , which we denote by $\phi^\varepsilon \xrightarrow{2} \phi$ for convenience.

Due to the equivalence, we often use assertions (i) and (ii) of Theorem 4.3 synonymously.

Remark 4.4. *The results are still true for vector- and matrix-valued functions, if we use the period unfolding operator for every component and the standard inner product for vectors and the Frobenius inner product for matrices.*

4.2. Homogenization in the disconnected case

In section 4.2.1, we prove some general compactness results via the periodic unfolding method, which we apply in section 4.2.2 to derive the homogenized problem for the disconnected case.

4.2.1. Compactness results for the disconnected case

In the first theorem, we compute the limit of a sequence $\{u_1^\varepsilon\}$ in $[H^1(\Omega_1^\varepsilon, \Gamma_1)]^3$, for which we have to define the Hilbert space

$$[L^2(\Omega, H_{\text{per},0}^1(Y_1))]^3 := \{u \in [L^2(\Omega, H_{\text{per}}^1(Y_1))]^3 : \mathcal{M}_{Y_1}(u) = 0\}.$$

Theorem 4.5. *Let $\{u_1^\varepsilon\}$ be a sequence in $[H^1(\Omega_1^\varepsilon, \Gamma_1)]^3$ with*

$$\|u_1^\varepsilon\|_{[L^2(\Omega_1^\varepsilon)]^3} + \|e(u_1^\varepsilon)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}} \leq C$$

for a constant C independent of ε . Then, there exist a subsequence (again denoted by ε), $u_1 \in [H^1(\Omega, \Gamma_1)]^3$ and $\hat{u}_1 \in [L^2(\Omega, H_{\text{per},0}^1(Y_1))]^3$ such that

$$\begin{aligned} \mathcal{T}_1^\varepsilon(u_1^\varepsilon) &\rightharpoonup u_1 \text{ weakly in } [L^2(\Omega, H^1(Y_1))]^3, \\ \mathcal{T}_1^\varepsilon(e(u_1^\varepsilon)) &\rightharpoonup e(u_1) + e_y(\hat{u}_1) \text{ weakly in } [L^2(\Omega \times Y_1)]^{3 \times 3}. \end{aligned}$$

Proof. The sequence $\{u_1^\varepsilon\}$ is bounded in $[H^1(\Omega_1^\varepsilon)]^3$ because, using (5), we can estimate

$$\|u_1^\varepsilon\|_{[L^2(\Omega_1^\varepsilon)]^3}^2 + \|\nabla u_1^\varepsilon\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 \leq C \|e(u_1^\varepsilon)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 \leq C.$$

Since the domain Ω_1^ε is connected, we can apply [24, Theorem 4.43] to get, up to a subsequence,

$$\begin{aligned} \mathcal{T}_1^\varepsilon(u_1^\varepsilon) &\rightharpoonup u_1 \text{ weakly in } [L^2(\Omega, H^1(Y_1))]^3, \\ \mathcal{T}_1^\varepsilon(\nabla u_1^\varepsilon) &\rightharpoonup \nabla u_1 + \nabla_y \hat{u}_1 \text{ weakly in } [L^2(\Omega \times Y_1)]^{3 \times 3} \end{aligned}$$

for some $u_1 \in [H^1(\Omega, \Gamma_1)]^3$ and $\hat{u}_1 \in [L^2(\Omega, H_{\text{per},0}^1(Y_1))]^3$. The definition of the linear strain tensor $e(u_1^\varepsilon) = \frac{1}{2}(\nabla u_1^\varepsilon + (\nabla u_1^\varepsilon)^T)$ directly yields

$$\mathcal{T}_1^\varepsilon(e(u_1^\varepsilon)) \rightharpoonup e(u_1) + e_y(\hat{u}_1) \text{ weakly in } [L^2(\Omega \times Y_1)]^{3 \times 3}.$$

□

In the rest of the section, we want to derive the limit of a sequence $\{u_0^\varepsilon\}$ in the Hilbert space

$$\left[\tilde{H}^1(\Omega_0^\varepsilon)\right]^3 := \{u \in [H^1(\Omega_0^\varepsilon)]^3 : \nabla \times u = 0 \text{ a.e. in } \Omega_0^\varepsilon\}.$$

The next lemma states the connection between the curl and symmetric gradient of some function ϕ and of the unfolded function $\mathcal{T}^\varepsilon(\phi)$.

Lemma 4.6. *Let $i = 0$ or 1 and $\phi \in [H^1(\Omega_0^\varepsilon)]^3$. Then, there holds*

- (i) $\nabla_y \times \mathcal{T}_i^\varepsilon(\phi) = \varepsilon \mathcal{T}_i^\varepsilon(\nabla \times \phi)$,
- (ii) $e_y(\mathcal{T}_i^\varepsilon(\phi)) = \varepsilon \mathcal{T}_i^\varepsilon(e(\phi))$.

Proof. These are simple computations using the linearity of $\mathcal{T}_i^\varepsilon$. □

The strategy of the proof of Theorem 4.7 follows [5, Theorem 2.18].

Theorem 4.7. *Let $\{u_0^\varepsilon\}$ be a sequence in $[\tilde{H}^1(\Omega_0^\varepsilon)]^3$ with*

$$\|u_0^\varepsilon\|_{[L^2(\Omega_0^\varepsilon)]^3} + \|e(u_0^\varepsilon)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}} \leq C \quad (11)$$

for a constant C independent of ε . Then, there exists a subsequence (again denoted by ε) and $u_0 \in [L^2(\Omega)]^3$ such that

$$\begin{aligned} \mathcal{T}_0^\varepsilon(u_0^\varepsilon) &\rightharpoonup u_0 \text{ weakly in } [L^2(\Omega, H^1(Y_0))]^3, \\ \varepsilon \mathcal{T}_0^\varepsilon(\nabla u_0^\varepsilon) &\rightarrow 0 \text{ strongly in } [L^2(\Omega \times Y_0)]^{3 \times 3}. \end{aligned}$$

Proof. Let $\{u_0^\varepsilon\}$ be a sequence in $[\tilde{H}^1(\Omega_0^\varepsilon)]^3$, for which (11) holds. Then, there holds due to Proposition 4.2 (iii)

$$\begin{aligned} \|\mathcal{T}_0^\varepsilon(u_0^\varepsilon)\|_{[L^2(\Omega \times Y_0)]^3} &\leq |Y|^{\frac{1}{2}} \|u_0^\varepsilon\|_{[L^2(\Omega_0^\varepsilon)]^3} \leq C, \\ \|\nabla_y \mathcal{T}_0^\varepsilon(u_0^\varepsilon)\|_{[L^2(\Omega \times Y_0)]^{3 \times 3}} &= \varepsilon \|\mathcal{T}_0^\varepsilon(e(u_0^\varepsilon))\|_{[L^2(\Omega \times Y_0)]^{3 \times 3}} \leq \varepsilon |Y|^{\frac{1}{2}} \|e(u_0^\varepsilon)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}} \leq \varepsilon C. \end{aligned}$$

Thus, $\mathcal{T}_0^\varepsilon(u_0^\varepsilon)$ is bounded in $[L^2(\Omega, H^1(Y_0))]^3$. Since $[L^2(\Omega, H^1(Y_0))]^3$ is a Hilbert space, there exists a subsequence (again denoted by ε) and $u_0 \in [L^2(\Omega, H^1(Y_0))]^3$ such that

$$\begin{aligned} \mathcal{T}_0^\varepsilon(u_0^\varepsilon) &\rightharpoonup u_0 \text{ weakly in } [L^2(\Omega, H^1(Y_0))]^3, \\ \nabla_y(\mathcal{T}_0^\varepsilon(u_0^\varepsilon)) &\rightarrow 0 \text{ strongly in } [L^2(\Omega \times Y_0)]^{3 \times 3}, \end{aligned}$$

which yield

$$\nabla_y u_0 = 0 \text{ a.e. in } \Omega \times Y_0.$$

This proves that u_0 is independent of y . □

We can even derive from $\nabla \times u_0^\varepsilon = 0$ that the limit satisfies $\nabla \times u_0 = 0$.

Theorem 4.8. Let $\{u_0^\varepsilon\}$ be a bounded sequence in $[\tilde{H}^1(\Omega_0^\varepsilon)]^3$ with

$$\mathcal{T}_0^\varepsilon(u_0^\varepsilon) \rightharpoonup u_0 \text{ weakly in } [L^2(\Omega, H^1(Y_0))]^3$$

for some $u_0 \in [L^2(\Omega)]^3$. Then, the weak limit satisfies $\nabla \times u_0 = 0$.

Proof. Let u_0^ε be as in the assumption and $\varphi(x, y) = (\psi_i(x)\eta_i(y))_{1 \leq i \leq 3} \in [\mathcal{D}(\Omega) \otimes H^1(Y_0)]^3$. We write $\varphi^\varepsilon := (\psi_i \eta_i^\varepsilon)_{1 \leq i \leq 3} := \varphi(x, \frac{x}{\varepsilon})$. Since

$$\mathcal{T}_0^\varepsilon(\varphi_i^\varepsilon) = \mathcal{T}_0^\varepsilon(\psi_i) \mathcal{T}_0^\varepsilon(\eta_i^\varepsilon) = \mathcal{T}_0^\varepsilon(\psi_i) \eta_i \longrightarrow \psi_i \eta_i \text{ strongly in } L^2(\Omega \times Y_0) \quad (12)$$

there holds

$$\int_{\Omega_0^\varepsilon} u_0^\varepsilon \cdot \varphi^\varepsilon dx = \int_{\Omega \times Y_0} \mathcal{T}_0^\varepsilon(u_0^\varepsilon) \cdot (\mathcal{T}_0^\varepsilon(\psi_i) \mathcal{T}_0^\varepsilon(\eta_i))_{1 \leq i \leq 3} dy dx \longrightarrow \int_{\Omega \times Y_0} u_0 \cdot \varphi dy dx. \quad (13)$$

Clearly, $\varphi^\varepsilon \in [H^1(\Omega_0^\varepsilon)]^3$. Thus, the trace of $\mathcal{T}_0^\varepsilon(\varphi^\varepsilon)$ with respect to ∂Y_0 exists for all $\varepsilon > 0$ and we can apply the trace theorem to calculate

$$\|\mathcal{T}_0^\varepsilon(\varphi^\varepsilon) - \varphi\|_{[L^2(\Omega \times \partial Y_0)]^3} \leq C \left(\|\mathcal{T}_0^\varepsilon(\varphi^\varepsilon) - \varphi\|_{[L^2(\Omega \times Y_0)]^3} + \|\nabla_y(\mathcal{T}_0^\varepsilon(\varphi^\varepsilon) - \varphi)\|_{[L^2(\Omega \times Y_0)]^{3 \times 3}} \right).$$

The convergences (12) and

$$\nabla_y \mathcal{T}_0^\varepsilon(\varphi_i^\varepsilon) = \varepsilon \mathcal{T}_0^\varepsilon(\nabla \psi_i) \eta_i + \mathcal{T}_0^\varepsilon(\psi_i) \nabla_y \eta_i \longrightarrow \psi_i \nabla_y \eta_i \text{ strongly in } [L^2(\Omega \times Y_0)]^3$$

yield

$$\|\mathcal{T}_0^\varepsilon(\varphi^\varepsilon) - \varphi\|_{[L^2(\Omega \times \partial Y_0)]^3} \rightarrow 0.$$

Since $u_0^\varepsilon \in [H^1(\Omega_0^\varepsilon)]^3$, the traces exist and we can compute, using the convergences from above,

$$\begin{aligned} \int_{\partial \Omega_0^\varepsilon} (u_0^\varepsilon \times \varphi^\varepsilon) \cdot n \left(\frac{x}{\varepsilon} \right) dS(x) &= \int_{\Omega \times \partial Y_0} (\mathcal{T}_0^\varepsilon(u_0^\varepsilon) \times \mathcal{T}_0^\varepsilon(\varphi^\varepsilon)) \cdot n(y) dS(y) dx \\ &\longrightarrow \int_{\Omega \times \partial Y_0} (u_0 \times \varphi) \cdot n dS(y) dx \end{aligned} \quad (14)$$

Now, for test functions with $\nabla_y \times \varphi = 0$, there holds, using the identity for the divergence of a cross product and the weak convergence of u_0^ε ,

$$\begin{aligned} 0 &= \int_{\Omega_0^\varepsilon} \nabla \times u_0^\varepsilon \cdot \varphi^\varepsilon dx \\ &= - \int_{\partial \Omega_0^\varepsilon} (u_0^\varepsilon \times \varphi^\varepsilon) \cdot n dS(x) + \int_{\Omega_0^\varepsilon} u_0^\varepsilon \cdot \nabla_x \times \varphi \left(x, \frac{x}{\varepsilon} \right) dx \\ &\stackrel{(13)}{\underset{(14)}{=}} - \int_{\Omega \times \partial Y_0} (u_0 \times \varphi) \cdot n dS(y) dx + \int_{\Omega \times Y_0} u_0 \cdot \nabla_x \times \varphi dy dx \\ &= - \int_{\Omega \times Y_0} u_0 \cdot \nabla_y \times \varphi dy dx + \int_{\partial \Omega \times Y_0} (u_0 \times \varphi) \cdot n dy dS(x) + \int_{\Omega \times Y_0} \nabla_x \times u_0 \cdot \varphi dy dx \\ &= \int_{\Omega} \nabla \times u_0(x) \cdot \int_{Y_0} \varphi(x, y) dy dx. \end{aligned} \quad (15)$$

If we choose $\psi_1 = \psi_2 = \psi_3 = \Psi$ for some $\Psi \in \mathcal{D}(\Omega)$, $\eta_2 = \eta_3 = 0$ and $\eta_1(y) = ay_1$, where $a \in \mathbb{R}^3$, we receive with (15)

$$0 = \int_{\Omega} \nabla \times u_0(x) \int_{Y_0} \varphi(x, y) dy dx = C \int_{\Omega} (\nabla \times u_0)_1 \Psi dx,$$

which shows, due the fundamental lemma of the calculus of variations, that $(\nabla \times u_0)_1 = 0$. Using similar functions for the other components, we get $\nabla \times u_0 = 0$. \square

In the next theorem, we compute the limit of the symmetric gradient of u_0^ε , where the strategy of the proof follows that of [5, Theorem 2.19]. For convenience, we define the Hilbert space

$$\left[L^2(\Omega, \tilde{H}^1(Y_0)) \right]^3 := \left\{ v \in \left[L^2(\Omega, H^1(Y_0)) \right]^3 : \nabla_y \times v = 0 \text{ a.e. in } \Omega \times Y_0 \right\}.$$

Theorem 4.9. *Let $\{u_0^\varepsilon\}$ be a sequence in $\left[\tilde{H}^1(\Omega_0^\varepsilon) \right]^3$ with*

$$\|u_0^\varepsilon\|_{[L^2(\Omega_0^\varepsilon)]^3} + \|e(u_0^\varepsilon)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}} \leq C \quad (16)$$

for a constant C independent of ε . Then, there exists a function $\hat{u}_0 \in \left[L^2(\Omega, \tilde{H}^1(Y_0)) \right]^3$ such that up to a subsequence

$$\begin{aligned} Z_0^\varepsilon &:= \frac{1}{\varepsilon} [\mathcal{T}_0^\varepsilon(u_0^\varepsilon) - \mathcal{M}_{Y_0}(\mathcal{T}_0^\varepsilon(u_0^\varepsilon))] \rightharpoonup \hat{u}_0 \text{ weakly in } \left[L^2(\Omega, \tilde{H}^1(Y_0)) \right]^3, \\ \mathcal{T}_0^\varepsilon(e(u_0^\varepsilon)) &\rightharpoonup e_y(\hat{u}_0) \text{ weakly in } [L^2(\Omega \times Y_0)]^{3 \times 3}. \end{aligned}$$

Furthermore, there holds $\mathcal{M}_{Y_0}(\hat{u}_0) = 0$.

Proof. Let $\{u_0^\varepsilon\}$ be a bounded sequence in $\left[\tilde{H}^1(\Omega_0^\varepsilon) \right]^3$. If we apply the Poincaré–Wirtinger inequality in $H^1(Y_0)$ and Proposition 4.2 (iii), we receive

$$\|Z_0^\varepsilon\|_{[L^2(\Omega \times Y_0)]^3} \leq \frac{C}{\varepsilon} \|\nabla_y \mathcal{T}_0^\varepsilon(u_0^\varepsilon)\|_{[L^2(\Omega \times Y_0)]^{3 \times 3}} \leq C|Y|^{\frac{1}{2}} \|e(u_0^\varepsilon)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}} \leq C.$$

With Lemma 4.6 and $\nabla \times u_0^\varepsilon = 0$ a.e. in Ω_0^ε , we get that

$$\nabla_y \times Z_0^\varepsilon = \frac{1}{\varepsilon} \nabla_y \times \mathcal{T}_0^\varepsilon(u_0^\varepsilon) = \mathcal{T}_0^\varepsilon(\nabla \times u_0^\varepsilon) = \mathcal{T}_0^\varepsilon(0) = 0$$

a.e. in $\Omega \times Y_0$. Furthermore, there holds

$$\|e_y(Z_0^\varepsilon)\|_{[L^2(\Omega \times Y_0)]^{3 \times 3}} = \frac{1}{\varepsilon} \|e_y(\mathcal{T}_0^\varepsilon(u_0^\varepsilon))\|_{[L^2(\Omega \times Y_0)]^{3 \times 3}} = \|\mathcal{T}_0^\varepsilon(e(u_0^\varepsilon))\|_{[L^2(\Omega \times Y_0)]^{3 \times 3}} \leq C.$$

Thus, the sequence $\{Z_0^\varepsilon\}$ is bounded in the Hilbert space $\left[L^2(\Omega, \tilde{H}^1(Y_0)) \right]^3$ and, therefore, there exists a function $\hat{u}_0 \in \left[L^2(\Omega, \tilde{H}^1(Y_0)) \right]^3$ such that up to a subsequence

$$Z_0^\varepsilon \rightharpoonup \hat{u}_0 \text{ weakly in } \left[L^2(\Omega, \tilde{H}^1(Y_0)) \right]^3.$$

So

$$\mathcal{T}_0^\varepsilon(e(u_0^\varepsilon)) = e_y(Z_0^\varepsilon) \rightharpoonup e_y(\hat{u}_0) \text{ weakly in } [L^2(\Omega \times Y_0)]^{3 \times 3}$$

and since

$$\mathcal{M}_{Y_0}(Z_0^\varepsilon) = \frac{1}{\varepsilon} [\mathcal{M}_{Y_0}(\mathcal{T}_0^\varepsilon(u_0^\varepsilon)) - \mathcal{M}_{Y_0}(\mathcal{T}_0^\varepsilon(u_0^\varepsilon))] = 0$$

for all ε , we receive $\mathcal{M}_{Y_0}(\hat{u}_0) = 0$. \square

Due to Proposition 4.2 (viii), there exists the trace of the unfolding operator with respect to y and we can prove the following result.

Theorem 4.10. *Let $u^\varepsilon, \varphi \in \mathcal{W}_d(\Omega^\varepsilon)$. Then, there holds*

$$\frac{1}{\varepsilon|Y|} \int_{\Omega} \int_{\Sigma_Y} (\mathcal{T}_1^\varepsilon(u_1^\varepsilon) - \mathcal{T}_0^\varepsilon(u_0^\varepsilon)) \cdot (\mathcal{T}_1^\varepsilon(\varphi_1) - \mathcal{T}_0^\varepsilon(\varphi_0)) \, dS(y) \, dx = \int_{\Sigma^\varepsilon} (u_1^\varepsilon - u_0^\varepsilon) \cdot (\varphi_1 - \varphi_0) \, dS(x).$$

Proof. The proof is similar to that of [5, Lemma 2.14]. \square

A direct consequence is the following corollary.

Corollary 4.11. *Let $u^\varepsilon \in \mathcal{W}_d(\Omega^\varepsilon)$. Then, there holds*

$$\frac{1}{\varepsilon|Y|} \int_{\Omega} \int_{\Sigma_Y} |\mathcal{T}_1^\varepsilon(u_1^\varepsilon) - \mathcal{T}_0^\varepsilon(u_0^\varepsilon)|^2 \, dS(y) \, dx = \int_{\Sigma^\varepsilon} |u_1^\varepsilon - u_0^\varepsilon|^2 \, dS(x).$$

Proof. The statement follows directly from Theorem 4.10, if we choose $\varphi = u^\varepsilon$. \square

4.2.2. Passage to the limit for the disconnected case

In this section, we apply the results proven in the section before to derive the homogenized problem. We define the Banach spaces

$$\left[L^2(\Omega, \tilde{H}_0^1(Y_0)) \right]^3 := \left\{ u \in [L^2(\Omega, H^1(Y_0))]^3 : \nabla_y \times u = 0 \text{ a.e. in } \Omega \times Y_0, \mathcal{M}_{Y_0}(u) = 0 \right\}$$

endowed with the norm $\left(\int_{\Omega} \|e_y(u)\|_{[L^2(Y_0)]^{3 \times 3}}^2 \, dx \right)^{\frac{1}{2}}$ and

$$\mathcal{Z}(\Omega, Y_1, \Gamma_1) := [H^1(\Omega, \Gamma_1)]^3 \times [L^2(\Omega, H_{\text{per},0}^1(Y_1))]^3 \times [L_{\text{curl}}^2(\Omega)]^3,$$

whereby $[L_{\text{curl}}^2(\Omega)]^3 := \{u \in [L^2(\Omega)]^3 : \nabla \times u = 0 \text{ a.e. in } \Omega\}$, endowed with the norm

$$\|u\|_{\mathcal{Z}(\Omega, Y_1, \Gamma_1)}^2 = \|e(u_1)\|_{[L^2(\Omega)]^{3 \times 3}}^2 + \int_{\Omega} \|e_y(\hat{u}_1)\|_{[L^2(Y_1)]^{3 \times 3}}^2 \, dx + \|u_0\|_{[L^2(\Omega)]^3}^2$$

for all $u = (u_1, \hat{u}_1, u_0) \in \mathcal{Z}(\Omega, Y_1, \Gamma_1)$. Having introduced this notation, we can formulate the limit problem. Afterwards, we rewrite the limit problem with the help of auxiliary cell problems and show that the solution is unique.

Theorem 4.12. Let $\{u^\varepsilon\}$ be a sequence of weak solutions of the problems (3) with $u^\varepsilon \in \mathcal{W}(\Omega^\varepsilon)$ and $\{f^\varepsilon\} \subset [L^2(\Omega^\varepsilon)]^3$ such that

$$\mathcal{T}_\alpha^\varepsilon(f^\varepsilon) \rightharpoonup f|_{\Omega \times Y_\alpha} \text{ in } [L^2(\Omega \times Y_\alpha)]^3$$

for some $f \in [L^2(\Omega \times Y)]^3$ and $\alpha = 0$ or 1 . Then, there exists $u = (u_1, \hat{u}_1, u_0) \in \mathcal{Z}(\Omega, Y_1, \Gamma_1)$ and $\hat{u}_0 \in [L^2(\Omega, \tilde{H}_0^1(Y_0))]^3$ such that

$$\begin{cases} \mathcal{T}_1^\varepsilon(u_1^\varepsilon) \rightharpoonup u_1 \text{ weakly in } [L^2(\Omega, H^1(Y_1))]^3, \\ \mathcal{T}_1^\varepsilon(e(u_1^\varepsilon)) \rightharpoonup e(u_1) + e_y(\hat{u}_1) \text{ weakly in } [L^2(\Omega \times Y_1)]^{3 \times 3}, \\ \mathcal{T}_0^\varepsilon(u_0^\varepsilon) \rightharpoonup u_0 \text{ weakly in } [L^2(\Omega, \tilde{H}^1(Y_0))]^3, \\ \mathcal{T}_0^\varepsilon(e(u_0^\varepsilon)) \rightharpoonup e_y(\hat{u}_0) \text{ weakly in } [L^2(\Omega \times Y_0)]^{3 \times 3}, \end{cases} \quad (17)$$

where u is the solution of the problem

$$\begin{aligned} & \int_\Omega \int_{Y_1} A(y)(e(u_1) + e_y(\hat{u}_1))(e(v_1) + e_y(\hat{v}_1)) \, dy \, dx \\ & + \int_\Omega \int_{\Sigma_Y} \left(K_N [u_1 \cdot n - u_0 \cdot n] n + K_T \sum_{i=1}^2 [u_1 \cdot \tau^i - u_0 \cdot \tau^i] \tau^i \right) \cdot (v_1 - v_0) \, dS(y) \, dx \\ & = \int_\Omega \int_{Y_1} f \, dy \cdot v_1 \, dx + \int_\Omega \int_{Y_0} f \, dy \cdot v_0 \, dx + \int_{\Gamma_2} g \cdot v_1 \, dS(x) \end{aligned} \quad (18)$$

for all $v = (v_1, \hat{v}_1, v_0) \in \mathcal{Z}(\Omega, Y_1, \Gamma_1)$.

Proof. Let $\{u^\varepsilon\}$ be a sequence of weak solutions of problem (3) with $u^\varepsilon \in \mathcal{W}_d(\Omega^\varepsilon)$. We apply Theorem 3.4 and Theorem 3.5 to get

$$\|u^\varepsilon\|_{[L^2(\Omega^\varepsilon)]^3}^2 + \|e(u_0^\varepsilon)\|_{[L^2(\Omega_0^\varepsilon)]^{3 \times 3}}^2 + \|e(u_1^\varepsilon)\|_{[L^2(\Omega_1^\varepsilon)]^{3 \times 3}}^2 < C.$$

Using the theorems from section 4.2.1, the convergences (17) follow directly. We rewrite the weak formulation of problem (3) using Theorem 4.10 and Proposition 4.2 (i) and (ii) to get the unfolded problem, whereas the integral over $\Gamma_2 \cap \partial\Omega_0^\varepsilon$ vanishes since $\Gamma_2 \cap \partial\Omega_0^\varepsilon = \emptyset$,

$$\begin{aligned} & \int_\Omega \int_{Y_1} \mathcal{T}_1^\varepsilon(A^\varepsilon) \mathcal{T}_1^\varepsilon(e(u_1^\varepsilon)) \mathcal{T}_1^\varepsilon(e(\varphi_1)) \, dy \, dx + \int_\Omega \int_{Y_0} \mathcal{T}_0^\varepsilon(A^\varepsilon) \mathcal{T}_0^\varepsilon(e(u_0^\varepsilon)) \mathcal{T}_0^\varepsilon(e(\varphi_0)) \, dy \, dx \\ & + \int_\Omega \int_{\Sigma_Y} (K_N [\mathcal{T}_1^\varepsilon(u_1^\varepsilon \cdot n) - \mathcal{T}_0^\varepsilon(u_0^\varepsilon \cdot n)] n) \cdot (\mathcal{T}_1^\varepsilon(\varphi_1) - \mathcal{T}_0^\varepsilon(\varphi_0)) \, dS(y) \, dx \\ & + \int_\Omega \int_{\Sigma_Y} \left(K_T \sum_{i=1}^2 [\mathcal{T}_1^\varepsilon(u_1^\varepsilon \cdot \tau^i) - \mathcal{T}_0^\varepsilon(u_0^\varepsilon \cdot \tau^i)] \tau^i \right) \cdot (\mathcal{T}_1^\varepsilon(\varphi_1) - \mathcal{T}_0^\varepsilon(\varphi_0)) \, dS(y) \, dx \\ & = \int_\Omega \int_{Y_1} \mathcal{T}_1^\varepsilon(f^\varepsilon) \cdot \mathcal{T}_1^\varepsilon(\varphi_1) \, dy \, dx + \int_\Omega \int_{Y_0} \mathcal{T}_0^\varepsilon(f^\varepsilon) \cdot \mathcal{T}_0^\varepsilon(\varphi_0) \, dy \, dx + \int_{\Gamma_2} g \cdot \varphi_1 \, dS(x). \end{aligned} \quad (19)$$

We choose the test functions

$$\varphi_0 = \varphi_0^\varepsilon := v_0 + \varepsilon \left(w_i^0 \widehat{\psi}_i^0 \right)_{1 \leq i \leq 3} \quad \text{and} \quad \varphi_1 = \varphi_1^\varepsilon := v_1 + \varepsilon \left(w_i^1 \widehat{\psi}_i^1 \right)_{1 \leq i \leq 3}$$

with $v_1 \in [\mathcal{D}(\overline{\Omega}, \Gamma_1)]^3 := \{\phi \in [\mathcal{D}(\overline{\Omega})]^3 : v \text{ is equal to } 0 \text{ in a neighbourhood of } \Gamma_1\}$, $w^0, w^1, v_0 \in [\mathcal{D}(\Omega)]^3$, $\widehat{\psi}^0(x) = \psi^0\left(\frac{x}{\varepsilon}\right)$ with $\psi^0 \in [H^1(Y_0)]^3$ (Y-periodically extended to \mathbb{R}^3), $\widehat{\psi}^1(x) = \psi^1\left(\frac{x}{\varepsilon}\right)$ with $\psi^1 \in [H_{\text{per}}^1(Y_1)]^3$ such that $\nabla \times \varphi_0^\varepsilon = 0$. Then, $\varphi^\varepsilon \in \mathcal{W}_d(\Omega^\varepsilon)$,

$$\mathcal{T}_0^\varepsilon(\varphi_0^\varepsilon) \in [L^2(\Omega, \tilde{H}^1(Y_0))]^3 \text{ and } \mathcal{T}_1^\varepsilon(\varphi_1^\varepsilon) \in [L^2(\Omega, H^1(Y_1))]^3.$$

Let $\alpha = 0$ or 1 . Since $\mathcal{T}_\alpha^\varepsilon(w_i^\alpha)\mathcal{T}_\alpha^\varepsilon(\widehat{\psi}_i^\alpha) \rightarrow w_i^\alpha\psi_i^\alpha$ strongly in $L^2(\Omega \times Y_\alpha)$ for all $i = 1, 2, 3$, there holds

$$\mathcal{T}_\alpha^\varepsilon(\varphi_\alpha^\varepsilon) \rightarrow v_\alpha \text{ strongly in } [L^2(\Omega \times Y_\alpha)]^3.$$

Every component of the symmetric gradient of $\varphi_\alpha^\varepsilon$ is of the form

$$e_{kl}(\varphi_\alpha^\varepsilon) = e_{kl}(v_\alpha) + \frac{1}{2} \left[\varepsilon \partial_{x_k} w_l^\alpha(x) \psi_l^\alpha\left(\frac{x}{\varepsilon}\right) + w_l^\alpha(x) \partial_{y_k} \psi_l^\alpha\left(\frac{x}{\varepsilon}\right) \right. \\ \left. + \varepsilon \partial_{x_l} w_k^\alpha(x) \psi_k^\alpha\left(\frac{x}{\varepsilon}\right) + w_k^\alpha(x) \partial_{y_l} \psi_k^\alpha\left(\frac{x}{\varepsilon}\right) \right].$$

If we apply the periodic unfolding operator to $e_{kl}(\varphi_\alpha^\varepsilon)$, we get

$$\mathcal{T}_\alpha^\varepsilon(e_{kl}(\varphi_\alpha^\varepsilon)) \rightarrow e_{kl}(v_\alpha) + \frac{1}{2} [w_l^\alpha(x) \partial_{y_k} \psi_l^\alpha(y) + w_k^\alpha(x) \partial_{y_l} \psi_k^\alpha(y)] = e_{kl}(v_\alpha) + (e_y(\hat{v}_\alpha))_{kl}$$

strongly in $L^2(\Omega \times Y_\alpha)$ with $\hat{v}_\alpha = (w_i^\alpha(x)\psi_i^\alpha(y))_{1 \leq i \leq 3}$. So

$$\mathcal{T}_\alpha^\varepsilon(e(\varphi_\alpha^\varepsilon)) \rightarrow e(v_\alpha) + e_y(\hat{v}_\alpha) \text{ strongly in } [L^2(\Omega \times Y_\alpha)]^{3 \times 3}.$$

We estimate the interface term with respect to the normal direction using Hölder's inequality, Corollary 4.11 and boundedness of the solution

$$\int_\Omega \int_{\Sigma_Y} (K_N [\mathcal{T}_1^\varepsilon(u_1^\varepsilon \cdot n) - \mathcal{T}_0^\varepsilon(u_0^\varepsilon \cdot n)] n) \cdot \mathcal{T}_0^\varepsilon \left(\varepsilon \left(w_i^\alpha \widehat{\psi}_i^\alpha \right)_{1 \leq i \leq 3} \right) dS(y) dx \\ \leq C \varepsilon^{\frac{1}{2}} \|u_1^\varepsilon - u_0^\varepsilon\|_{L^2(\Sigma^\varepsilon)} \varepsilon \|(\mathcal{T}_0^\varepsilon(w_i^\alpha)\psi_i^\alpha)_{1 \leq i \leq 3}\|_{[L^2(\Omega \times \Sigma_Y)]^3} \leq C \varepsilon.$$

Therefore, this integral converges to zero. The analogous result holds for the terms with τ_i , $i = 1, 2$ instead of n . Since

$$\mathcal{T}_0^\varepsilon(v_0) \rightarrow v_0 \text{ in } [L^2(\Omega \times Y_0)]^3 \text{ and } \mathcal{T}_0^\varepsilon(e(v_0)) \rightarrow e(v_0) \text{ in } [L^2(\Omega \times Y_0)]^{3 \times 3}$$

we know that $e_y(\mathcal{T}_0^\varepsilon(v_0)) \rightarrow 0$ strongly in $[L^2(\Omega \times Y_0)]^{3 \times 3}$ and thus

$$\mathcal{T}_0^\varepsilon(v_0) \rightarrow v_0 \text{ strongly in } [L^2(\Omega, \tilde{H}^1(Y_0))]^3.$$

This yields the convergence of the traces of $\mathcal{T}_0^\varepsilon(v_0)$ with respect to y . Similarly, we receive for v_1

$$\mathcal{T}_1^\varepsilon(v_1) \rightarrow v_1 \text{ strongly in } [L^2(\Omega, H^1(Y_1))]^3.$$

If we plug in the test function in (19) and pass to the limit, we get

$$\begin{aligned}
& \int_{\Omega} \int_{Y_1} A(y)(e(u_1) + e_y(\hat{u}_1))(e(v_1) + e_y(\hat{v}_1)) \, dy dx + \int_{\Omega} \int_{Y_0} A(y)e_y(\hat{u}_0)(e(v_0) + e_y(\hat{v}_0)) \, dy dx \\
& + \int_{\Omega} \int_{\Sigma_Y} \left(K_N [u_1 \cdot n - u_0 \cdot n] n + K_T \sum_{i=1}^2 [u_1 \cdot \tau^i - u_0 \cdot \tau^i] \tau^i \right) \cdot (v_1 - v_0) \, dS(y) \, dx \\
& = \int_{\Omega} \int_{Y_1} f \, dy \cdot v_1 \, dx + \int_{\Omega} \int_{Y_0} f \, dy \cdot v_0 \, dx + \int_{\Gamma_2} g \cdot v_1 \, dS(x). \tag{20}
\end{aligned}$$

Since $\mathcal{D}(\bar{\Omega}, \Gamma_1)$ is dense in $H^1(\Omega, \Gamma_1)$ and $\mathcal{D}(\Omega) \times H_{\text{per}}^1(Y_1)$ is dense in $L^2(\Omega, H_{\text{per}}^1(Y_1))$, the homogenized problem is true for all $v_1 \in [H^1(\Omega, \Gamma_1)]^3$ and $\hat{v}_1 \in [L^2(\Omega, H_{\text{per},0}^1(Y_1))]^3$.

To improve this result, we choose as a test function

$$\varphi_1 = 0 \quad \text{and} \quad \varphi_0 = \varepsilon e(v_0(x)) \cdot \psi\left(\frac{x}{\varepsilon}\right) = \varepsilon e(v_0(x)) \cdot \frac{x}{\varepsilon}$$

with $v_0 \in [\mathcal{D}(\Omega)]^3$ with $\nabla \times v_0 = 0$ and $\psi(y) = y \in [H^1(Y_0)]^3$ Y -periodically extended to \mathbb{R}^3 . Using the fact that $\nabla \times v_0 = 0$, it is easy to compute that $\varphi^\varepsilon \in \mathcal{W}_d(\Omega^\varepsilon)$. Therefore, we can rewrite (19)

$$\begin{aligned}
& \int_{\Omega} \int_{Y_0} \mathcal{T}_0^\varepsilon(A^\varepsilon) \mathcal{T}_0^\varepsilon(e(u_0^\varepsilon)) \mathcal{T}_0^\varepsilon(e(\varphi_0)) \, dy dx \\
& = \int_{\Omega} \int_{\Sigma_Y} (K_N [\mathcal{T}_1^\varepsilon(u_1^\varepsilon \cdot n) - \mathcal{T}_0^\varepsilon(u_0^\varepsilon \cdot n)] n) \cdot \mathcal{T}_0^\varepsilon(\varphi_0) \, dS(y) \, dx \\
& + \int_{\Omega} \int_{\Sigma_Y} \left(K_T \sum_{i=1}^2 [\mathcal{T}_1^\varepsilon(u_1^\varepsilon \cdot \tau^i) - \mathcal{T}_0^\varepsilon(u_0^\varepsilon \cdot \tau^i)] \tau^i \right) \cdot \mathcal{T}_0^\varepsilon(\varphi_0) \, dS(y) \, dx + \int_{\Omega} \int_{Y_0} \mathcal{T}_0^\varepsilon(f^\varepsilon) \cdot \mathcal{T}_0^\varepsilon(\varphi_0) \, dy dx
\end{aligned}$$

With similar argumentation as above, the right-hand side disappears for $\varepsilon \rightarrow 0$ and

$$\mathcal{T}_0^\varepsilon \left(\varepsilon e_{kl} \left(e(v_0(x)) \frac{x}{\varepsilon} \right) \right) \rightarrow e_{kl}(v_0)(x) = (e_y(e(v_0(x))y))_{kl}.$$

Thus, we obtain for all $v_0 \in [\mathcal{D}(\Omega)]$ with $\nabla \times v_0 = 0$

$$\int_{\Omega} \int_{Y_0} A(y)e_y(\hat{u}_0)e(v_0) \, dy dx = 0.$$

If we choose $v_1 = \hat{v}_1 = v_0 = 0$ in (20) there holds

$$\int_{\Omega} \int_{Y_0} A(y)e_y(\hat{u}_0)e_y(\hat{v}_0) \, dy dx = 0.$$

Thus, we obtain for all $v_1 \in [H^1(\Omega, \Gamma_1)]^3$, $\hat{v}_1 \in [L^2(\Omega, H_{\text{per},0}^1(Y_1))]^3$, $v_0 \in [\mathcal{D}(\Omega)]^3$ with $\nabla \times v_0 = 0$

$$\begin{aligned}
& \int_{\Omega} \int_{Y_1} A(y)(e(u_1) + e_y(\hat{u}_1))(e(v_1) + e_y(\hat{v}_1)) \, dy dx \\
& + \int_{\Omega} \int_{\Sigma_Y} \left(K_N [u_1 \cdot n - u_0 \cdot n] n + K_T \sum_{i=1}^2 [u_1 \cdot \tau^i - u_0 \cdot \tau^i] \tau^i \right) \cdot (v_1 - v_0) \, dS(y) \, dx \\
& = \int_{\Omega} \int_{Y_1} f \, dy \cdot v_1 \, dx + \int_{\Omega} \int_{Y_0} f \, dy \cdot v_0 \, dx + \int_{\Gamma_2} g \cdot v_1 \, dS(x).
\end{aligned}$$

Since $\{v \in [\mathcal{D}(\Omega)]^3 : \nabla \times v = 0 \text{ a.e. in } \Omega\}$ is dense in $L^2_{\text{curl}}(\Omega)$, we get the desired result. Using that the solution is unique, which we prove below, all the convergences above hold true for the whole sequence. \square

In the next theorem, we rewrite the homogenized problem from Theorem 4.12 as a macroscopic problem with an auxiliary cell problem.

Theorem 4.13. *Let $\{u^\varepsilon\}$ be as in Theorem 4.12. We can reformulate the homogenized problem (18) as follows: Find $u_1 \in [H^1(\Omega, \Gamma_1)]^3$, $u_0 \in [L^2_{\text{curl}}(\Omega)]^3$ such that*

$$\begin{aligned} & \int_{\Omega} A_1^{\text{hom}} e(u_1) e(v_1) \, dx \\ & + \int_{\Omega} \int_{\Sigma_Y} \left(K_N (u_1 \cdot n - u_0 \cdot n) n + K_T \sum_{i=1}^2 (u_1 \cdot \tau^i - u_0 \cdot \tau^i) \tau^i \right) \cdot (v_1 - v_0) \, dS(y) \, dx \\ & = \int_{\Omega} \int_{Y_0} f \, dy \cdot v_0 \, dx + \int_{\Omega} \int_{Y_1} f \, dy \cdot v_1 \, dx + \int_{\Gamma_2} g \cdot v_1 \, dS(x) \end{aligned} \quad (21)$$

where

$$(A_1^{\text{hom}})_{ijkh} = \int_{Y_1} a_{ijkh}(y) - \sum_{l,m=1}^3 a_{ijlm} \left(e_y(\chi_1^{kh}) \right)_{lm} \, dy$$

and $\chi_1^{lm} \in [H^1_{\text{per},0}(Y_1)]^3$, $l, m \in \{1, 2, 3\}$, is the unique solution of

$$\begin{cases} \left(-\sum_{j=1}^3 \frac{\partial}{\partial y_j} \left[(Ae_y(\chi_1^{lm}))_{ij} - a_{ijlm} \right] \right)_{1 \leq i \leq 3} = 0 & \text{in } Y_1, \\ \left(-\sum_{j=1}^3 \left[(Ae_y(\chi_1^{lm}))_{ij} - a_{ijlm} \right] n_j \right)_{1 \leq i \leq 3} = 0 & \text{on } \Sigma_Y. \end{cases} \quad (22)$$

Proof. This result follows by standard arguments. We refer to [19] to see the steps of the proof in detail. \square

Similar to [17, Theorem II.1.1], it can be proven that there exist $\alpha^{\text{hom}}, \beta^{\text{hom}} \in \mathbb{R}$ with $0 < \alpha^{\text{hom}} < \beta^{\text{hom}}$ such that $A_1^{\text{hom}} \in M(\alpha^{\text{hom}}, \beta^{\text{hom}}, \Omega)$. With this fact we are able to prove the uniqueness of the solution of (21).

Theorem 4.14. *The solutions $u_1 \in [H^1(\Omega, \Gamma_1)]^3$ and $u_0 \in [L^2_{\text{curl}}(\Omega)]^3$ of the macroscopic problem (21) are unique.*

Proof. We assume that there exist two solutions $u = (u_1, u_0), w = (w_1, w_0) \in [H^1(\Omega, \Gamma_1)]^3 \times [L^2_{\text{curl}}(\Omega)]^3$ of the problem (21). If we choose as test function $v_1 = u_1 - w_1$ and $v_0 = u_0 - w_0$, we can estimate, using that A_1^{hom} is coercive and Korn's inequality for functions which are zero on part of the boundary,

$$\begin{aligned} 0 & = \int_{\Omega} A_1^{\text{hom}} e(u_1 - w_1) e(u_1 - w_1) \, dx + \int_{\Omega} \int_{\Sigma_Y} \left(K_N [u_1 - u_0 - w_1 + w_0] \cdot nn \right. \\ & \quad \left. + K_T \sum_{i=1}^2 [u_1 - u_0 - w_1 + w_0] \cdot \tau^i \tau^i \right) \cdot (u_1 - w_1 - u_0 + w_0) \, dS(y) \, dx \\ & \geq \alpha^{\text{hom}} C \|u_1 - w_1\|_{[H^1(\Omega)]^3}^2 + \min\{K_N, K_T\} |\Sigma_Y| \|u_1 - u_0 - w_1 + w_0\|_{[L^2(\Omega)]^3}^2. \end{aligned}$$

Thus, $u_1 = w_1$. Using this and $\|u_1 - u_0 - w_1 + w_0\|_{[L^2(\Omega)]^3}^2 = 0$, we receive $u_0 = w_0$, which proves the uniqueness of the solution. \square

Remark 4.15. We can generalize the result using $A^\varepsilon = A(\cdot, \frac{\cdot}{\varepsilon}) \in M(\alpha, \beta, \Omega)$ instead of $A^\varepsilon = A(\frac{\cdot}{\varepsilon}) \in M(\alpha, \beta, Y)$ and under the additional assumption that $\mathcal{T}_1^\varepsilon(A^\varepsilon) \rightarrow B_1$ a.e. in $\Omega \times Y_1$ and $\mathcal{T}_0^\varepsilon(A^\varepsilon) \rightarrow B_0$ a.e. in $\Omega \times Y_0$. Then, $B_1 \in M(\alpha, \beta, \Omega \times Y_1)$ and $B_0 \in M(\alpha, \beta, \Omega \times Y_0)$ and the homogenized problem is: Find the unique solutions $u_1 \in [H^1(\Omega, \Gamma_1)]^3$, $u_0 \in [L^2_{\text{curl}}(\Omega)]^3$ such that

$$\begin{aligned} & \int_{\Omega} B_1^{\text{hom}} e(u_1) e(v_1) \, dx + \int_{\Omega} \int_{\Sigma_Y} \left(K_N(u_1 - u_0) \cdot nn + K_T \sum_{i=1}^2 (u_1 - u_0) \cdot \tau^i \tau^i \right) \cdot (v_1 - v_0) \, dS(y) \, dx \\ &= \int_{\Omega} \int_{Y_0} f \, dy \cdot v_0 \, dx + \int_{\Omega} \int_{Y_1} f \, dy \cdot v_1 \, dx + \int_{\Gamma_2} g \cdot v_1 \, dS(x) \end{aligned}$$

with

$$(B_1^{\text{hom}})_{ijkh}(x) = \int_{Y_1} (B_1)_{ijkh}(x, y) - \sum_{l,m=1}^3 (B_1)_{ijlm}(x, y) \left(e_y(\chi_1^{kh}) \right)_{lm}(x, y) \, dy$$

and $\chi_1^{lm} \in [L^2(\Omega, H^1_{\text{per},0}(Y_1))]^3$, $l, m \in \{1, 2, 3\}$ is the unique solution of

$$\begin{cases} \left(-\sum_{j=1}^3 \frac{\partial}{\partial y_j} \left[(B_1(\cdot, y) e_y(\chi_1^{lm})(\cdot, y))_{ij} - (B_1)_{ijlm}(\cdot, y) \right] \right)_{1 \leq i \leq 3} = 0 & \text{in } Y_1, \\ \left(-\sum_{j=1}^3 \left[(B_1(\cdot, y) e_y(\chi_1^{lm})(\cdot, y))_{ij} - (B_1)_{ijlm}(\cdot, y) \right] \right)_{1 \leq i \leq 3} \cdot n = 0 & \text{on } \Sigma_Y \end{cases}$$

a.e. in Ω .

4.3. Homogenization in the connected case

In section 4.3.1, we prove a convergence result, which we apply in section 4.3.2 to derive the homogenized problem for the connected case. In order to handle the different boundary conditions at the exterior boundary, we use a special extension operator established in [19] in the context of two-scale convergence. For some general introduction to the two-scale convergence method, we refer to [25], and we note that the two-scale convergence and weak convergence of the unfolded sequence are equivalent (cf. Theorem 4.3).

4.3.1. Compactness result for the connected case

The following theorem gives us some information about the two-scale limit of the solutions u_α^ε , $\alpha \in \{0, 1\}$.

Theorem 4.16. For $\alpha \in \{0, 1\}$, let $\{u_\alpha^\varepsilon\}$ be a sequence in $[H^1(\Omega_\alpha^\varepsilon)]^3$ with $u_\alpha^\varepsilon = 0$ on $\Gamma_1 \cap \partial\Omega_\alpha^\varepsilon$ and

$$\|e(u_\alpha^\varepsilon)\|_{[L^2(\Omega_\alpha^\varepsilon)]^{3 \times 3}} \leq C$$

for a constant C independent of ε . Then, there exists, up to a subsequence, $u_\alpha \in [H^1(\Omega, \Gamma_1)]^3$ and $\hat{u}_\alpha \in [L^2(\Omega, H^1_{\text{per}}(Y)/\mathbb{R})]^3$, such that

$$\begin{aligned} \mathcal{T}^\varepsilon(\tilde{u}_\alpha^\varepsilon) &\rightharpoonup u_\alpha(x) \text{ weakly in } [L^2(\Omega \times Y)]^3, \\ \mathcal{T}^\varepsilon(e(\tilde{u}_\alpha^\varepsilon)) &\rightharpoonup e(u_\alpha)(x) + e_y(\hat{u}_\alpha)(x, y) \text{ weakly in } [L^2(\Omega \times Y)]^{3 \times 3}, \end{aligned}$$

where $\tilde{\cdot}$ is the extension to Ω defined in [19, Theorem 3.5] and \mathcal{T}^ε is the unfolding operator from [23, Definition 1.2].

Proof. Let $\{u_1^\varepsilon\}$ be a sequence in $[H^1(\Omega_\alpha^\varepsilon)]^3$ with the properties of the assumption. Using Korn's inequality ([19, Theorem 4.4]), there holds

$$\|u_\alpha^\varepsilon\|_{[L^2(\Omega_\alpha^\varepsilon)]^3} + \|\nabla u_\alpha^\varepsilon\|_{[L^2(\Omega_\alpha^\varepsilon)]^{3 \times 3}} \leq C \|e(u_\alpha^\varepsilon)\|_{[L^2(\Omega_\alpha^\varepsilon)]^{3 \times 3}} \leq C.$$

Therefore, $\{u_\alpha^\varepsilon\}$ is a bounded sequence in $[H^1(\Omega_\alpha^\varepsilon)]^3$ with $u_\alpha^\varepsilon = 0$ in $\partial\Omega_\alpha^\varepsilon \cap \Gamma_1$. The extension operator defined in [19, Theorem 3.5] guarantees that $\{\tilde{u}_\alpha^\varepsilon\}$ is a bounded sequence in $[H^1(\Omega, \Gamma_1)]^3$. Since $[H^1(\Omega, \Gamma_1)]^3$ is a Hilbert space there exists a $u_\alpha \in [H^1(\Omega, \Gamma_1)]^3$ with

$$\tilde{u}_\alpha^\varepsilon \rightharpoonup u_\alpha \text{ weakly in } [H^1(\Omega, \Gamma_1)]^3.$$

Therefore, we can apply [25, Theorem 20] and Theorem 4.3 to get for some $\hat{u}_\alpha \in [L^2(\Omega, H_{\text{per}}^1(Y)/\mathbb{R})]^3$

$$\mathcal{T}^\varepsilon(\tilde{u}_\alpha^\varepsilon) \rightharpoonup u_\alpha(x) \text{ weakly in } [L^2(\Omega \times Y)]^3$$

and

$$\mathcal{T}^\varepsilon(\nabla \tilde{u}_\alpha^\varepsilon) \rightharpoonup \nabla u_\alpha(x) + \nabla_y \hat{u}_\alpha(x, y) \text{ weakly in } [L^2(\Omega \times Y)]^{3 \times 3}.$$

The definition of the linear strain tensor $e(\tilde{u}_\alpha^\varepsilon) = \frac{1}{2}(\nabla \tilde{u}_\alpha^\varepsilon + (\nabla \tilde{u}_\alpha^\varepsilon)^T)$ directly yields

$$\mathcal{T}^\varepsilon(e(\tilde{u}_\alpha^\varepsilon)) \rightharpoonup e(u_\alpha)(x) + e_y(\hat{u}_\alpha)(x, y) \text{ weakly in } [L^2(\Omega \times Y)]^{3 \times 3}.$$

□

4.3.2. Passage to the limit in the connected case

We endow the space

$$\mathcal{V}(\Omega, Y, \Gamma_1) := [H^1(\Omega, \Gamma_1)]^3 \times [L^2(\Omega, H_{\text{per},0}^1(Y_0))]^3 \times [H^1(\Omega, \Gamma_1)]^3 \times [L^2(\Omega, H_{\text{per},0}^1(Y_1))]^3$$

with the norm (due to Korn's inequality for functions with zero trace on part of the boundary and for periodic functions)

$$\|u\|_{\mathcal{V}(\Omega, Y, \Gamma_1)}^2 = \|e(u_0)\|_{[L^2(\Omega)]^{3 \times 3}}^2 + \|e_y(\hat{u}_0)\|_{[L^2(\Omega \times Y_0)]^{3 \times 3}}^2 + \|u_1\|_{[L^2(\Omega)]^3}^2 + \|e_y(\hat{u}_1)\|_{[L^2(\Omega \times Y_1)]^{3 \times 3}}^2$$

for all $u = (u_0, \hat{u}_0, u_1, \hat{u}_1) \in \mathcal{V}(\Omega, Y, \Gamma_1)$.

Theorem 4.17. *Let $\{u^\varepsilon\}$ be a sequence of weak solutions of the problem (3) with $u^\varepsilon \in \mathcal{W}_c(\Omega^\varepsilon)$ and $\{f^\varepsilon\} \subset [L^2(\Omega^\varepsilon)]^3$ such that*

$$\mathcal{T}^\varepsilon(f^\varepsilon) \rightharpoonup f \text{ weakly in } [L^2(\Omega \times Y)]^3$$

for some $f \in [L^2(\Omega \times Y)]^3$. Let $\alpha = 0$ or 1 . Then, there exist functions $u_\alpha \in [H^1(\Omega, \Gamma_1)]^3$ and $\hat{u}_\alpha \in [L^2(\Omega, H_{\text{per}}^1(Y)/\mathbb{R})]^3$ such that

$$\begin{cases} \mathcal{T}^\varepsilon(\tilde{u}_\alpha^\varepsilon) \rightharpoonup u_\alpha(x) \text{ weakly in } [L^2(\Omega \times Y)]^3, \\ \mathcal{T}^\varepsilon(e(\tilde{u}_\alpha^\varepsilon)) \rightharpoonup e(u_\alpha)(x) + e_y(\hat{u}_\alpha)(x, y) \text{ weakly in } [L^2(\Omega \times Y)]^{3 \times 3}, \end{cases} \quad (23)$$

where $\tilde{\cdot}$ is the extension to Ω defined in [19, Theorem 3.5] and \mathcal{T}^ε is the unfolding operator from [23, Definition 1.2]. Furthermore, the restriction $u = (u_0, \hat{u}_0|_{\Omega \times Y_0}, u_1, \hat{u}_1|_{\Omega \times Y_1}) \in \mathcal{V}(\Omega, Y, \Gamma_1)$ is the solution of the problem

$$\begin{aligned} & \int_{\Omega} \int_{Y_0} A(e(u_0) + e_y(\hat{u}_0))(e(v_0) + e_y(\hat{v}_0)) \, dy dx + \int_{\Omega} \int_{Y_1} A(e(u_1) + e_y(\hat{u}_1))(e(v_1) + e_y(\hat{v}_1)) \, dy dx \\ & + \int_{\Omega} \int_{\Sigma_Y} \left(K_N(u_1 \cdot n - u_0 \cdot n)n + K_T \sum_{i=1}^2 (u_1 \cdot \tau^i - u_0 \cdot \tau^i)\tau^i \right) \cdot (v_1 - v_0) \, dS(y) dx \quad (24) \\ & = \int_{\Omega} \int_{Y_0} f \, dy \cdot v_0 \, dx + \int_{\Omega} \int_{Y_1} f \, dy \cdot v_1 \, dx + \int_{\Gamma_2} g \cdot h_0 v_0 \, dS(x) + \int_{\Gamma_2} g \cdot h_1 v_1 \, dS(x) \end{aligned}$$

for all $v \in \mathcal{V}(\Omega, Y, \Gamma_1)$.

Proof. Let $\{u^\varepsilon\}$ be the sequence of the solutions of problem (3). We have already proven in Theorem 3.6 and Theorem 3.7 the existence, uniqueness and uniform boundedness of this sequence. So we can apply Theorem 4.16, which proves the convergences in (23). We choose some test functions φ^ε of the form

$$\varphi_0^\varepsilon := v_0 + \varepsilon w_0^\varepsilon \quad \text{and} \quad \varphi_1^\varepsilon := v_1 + \varepsilon w_1^\varepsilon$$

with $v_0, v_1 \in [\mathcal{D}(\bar{\Omega}, \Gamma_1)]^3 := \{\phi \in [\mathcal{D}(\bar{\Omega})]^3 : v \text{ is equal to 0 in a neighbourhood of } \Gamma_1\}$, $w_0^\varepsilon(x) := w_0(x, \frac{x}{\varepsilon})$ and $w_1^\varepsilon(x) := w_1(x, \frac{x}{\varepsilon})$ with $w_0, w_1 \in [\mathcal{D}(\Omega, C_{\text{per}}^\infty(Y))]^3$. Therefore, $\varphi_0^\varepsilon|_{\Gamma_1} = 0$, $\varphi_1^\varepsilon|_{\Gamma_1} = 0$, $\varphi_0^\varepsilon \in [H^1(\Omega_0^\varepsilon)]^3$ and $\varphi_1^\varepsilon \in [H^1(\Omega_1^\varepsilon)]^3$ and the weak formulation is

$$\begin{aligned} & \int_{\Omega_0^\varepsilon} A^\varepsilon e(u_0^\varepsilon) e(\varphi_0^\varepsilon) dx + \int_{\Omega_1^\varepsilon} A^\varepsilon e(u_1^\varepsilon) e(\varphi_1^\varepsilon) dx \\ & + \varepsilon \int_{\Sigma^\varepsilon} \left(K_N [u_n^\varepsilon]_{\Sigma^\varepsilon} n + K_T \sum_{i=1}^2 [u_{\tau^i}^\varepsilon]_{\Sigma^\varepsilon} \tau^i \right) \cdot (\varphi_1^\varepsilon - \varphi_0^\varepsilon) \, dS(x) \\ & = \int_{\Omega_0^\varepsilon} f^\varepsilon \cdot \varphi_0^\varepsilon \, dx + \int_{\Omega_1^\varepsilon} f^\varepsilon \cdot \varphi_1^\varepsilon \, dx + \int_{\Gamma_2 \cap \partial \Omega_0^\varepsilon} g \cdot \varphi_0^\varepsilon \, dS(x) + \int_{\Gamma_2 \cap \partial \Omega_1^\varepsilon} g \cdot \varphi_1^\varepsilon \, dS(x). \end{aligned}$$

Let $\alpha = 0$ or 1 . Using the fact that

$$e(\varphi_\alpha^\varepsilon)(x) = e(v_\alpha)(x) + \varepsilon e(w_\alpha) \left(x, \frac{x}{\varepsilon} \right) + e_y(w_\alpha) \left(x, \frac{x}{\varepsilon} \right),$$

and similar argumentation as in the disconnected case, we receive

$$\mathcal{T}^\varepsilon(e(\varphi_\alpha^\varepsilon)) \rightarrow e(v_\alpha) + e_y(w_\alpha) \text{ strongly in } [L^2(\Omega \times Y)]^{3 \times 3}.$$

Using the periodic unfolding operator, the following convergence holds

$$\begin{aligned} \int_{\Omega_\alpha^\varepsilon} A^\varepsilon e(u_\alpha^\varepsilon) e(\varphi_\alpha^\varepsilon) \, dx &= \int_{\Omega} \int_{Y_\alpha} \mathcal{T}^\varepsilon(A^\varepsilon) \mathcal{T}^\varepsilon(e(\tilde{u}_\alpha^\varepsilon)) \mathcal{T}^\varepsilon(e(\varphi_\alpha^\varepsilon)) \, dx \\ &\longrightarrow \int_{\Omega} \int_{Y_\alpha} A(e(u_\alpha) + e_y(\hat{u}_\alpha))(e(v_\alpha) + e_y(w_\alpha)) \, dy dx. \end{aligned}$$

We receive the convergence of the interface term from Lemma 4.18 and the fact that the terms with w_1^ε and w_0^ε vanish. Moreover,

$$\int_{\Omega_\alpha^\varepsilon} f^\varepsilon \cdot \varphi_\alpha^\varepsilon \, dx = \int_{\Omega} \int_{Y_\alpha} \mathcal{T}^\varepsilon(f^\varepsilon) \cdot \mathcal{T}^\varepsilon(v_\alpha) \, dx + \varepsilon \int_{\Omega} \int_{Y_\alpha} \mathcal{T}^\varepsilon(f^\varepsilon) \cdot \mathcal{T}^\varepsilon(w_\alpha^\varepsilon) \, dx \longrightarrow \int_{\Omega} \int_{Y_\alpha} f \, dy \cdot v_\alpha \, dx.$$

For the proof of the convergence of the boundary integral we refer to section 7.6.2 in [19]

$$\int_{\Gamma_2 \cap \partial\Omega_\alpha^\varepsilon} g \cdot \varphi_\alpha^\varepsilon \, dS(x) \longrightarrow \int_{\Gamma_2} g \cdot h_\alpha v_\alpha \, dS(x)$$

for some $h_\alpha \in L^2(\Gamma_2)$. Under further assumption on the boundary $\partial\Omega$ and the exterior boundaries of Y_0 and Y_1 we can formulate the limit functions h_0 and h_1 explicitly (see [19, Theorem 7.17]). Using the convergence results above and the density of $[\mathcal{D}(\bar{\Omega}, \Gamma_1)]^3$ in $[H^1(\Omega, \Gamma_1)]^3$ (see [26, Theorem 3.1]) and $[\mathcal{D}(\Omega, C_\infty^\infty(Y))]^3$ in $[L^2(\Omega, H_{\text{per}}^1(Y))]^3$, we get the desired homogenized problem (24) up to uniqueness. To ensure the uniqueness, we can restrict the space $[L^2(\Omega, H_{\text{per}}^1(Y)/\mathbb{R})]^3$ to $[L^2(\Omega, H_{\text{per},0}^1(Y_0))]^3$ resp. $[L^2(\Omega, H_{\text{per},0}^1(Y_1))]^3$ for the functions \hat{u}_0, \hat{u}_1 and the test functions in the limit problem, which does not change the limit problem. Since the solution is unique, which we prove below, all the convergences above hold true for the whole sequence. \square

We want to prove the limit result for the interface term used in the proof of the last theorem.

Lemma 4.18. *Let $\{u^\varepsilon\}$ be the sequence as in Theorem 4.17. Then, there holds up to a subsequence for all $v_0, v_1 \in [\mathcal{D}(\bar{\Omega}, \Gamma_1)]^3$*

$$\begin{aligned} & \varepsilon \int_{\Sigma^\varepsilon} \left(K_N [\tilde{u}_1^\varepsilon \cdot n - \tilde{u}_0^\varepsilon \cdot n] n + K_T \sum_{i=1}^2 [\tilde{u}_1^\varepsilon \cdot \tau^i - \tilde{u}_0^\varepsilon \cdot \tau^i] \tau^i \right) \cdot (v_1 - v_0) \, dS(x) \\ & \xrightarrow{\varepsilon \rightarrow 0} \int_{\Omega} \int_{\Sigma_Y} \left(K_N [u_1 \cdot n - u_0 \cdot n] n + K_T \sum_{i=1}^2 [u_1 \cdot \tau^i - u_0 \cdot \tau^i] \tau^i \right) \cdot (v_1 - v_0) \, dS(y) \, dx, \end{aligned} \quad (25)$$

whereby only n and τ^i , $i = 1, 2$, depend on y .

Proof. If we apply the boundary unfolding operator $\mathcal{T}_b^\varepsilon$ defined in [23, Definition 3.1] we get

$$\varepsilon K_N \int_{\Sigma^\varepsilon} \tilde{u}_1^\varepsilon \cdot n v_1 \cdot n \, dS(x) = \sum_{i,j=1}^3 \frac{K_N}{|Y|} \int_{\Omega \times \Sigma_Y} \mathcal{T}^\varepsilon(\tilde{u}_1^\varepsilon)_i \mathcal{T}_b^\varepsilon(n_i) \mathcal{T}_b^\varepsilon(n_j) \mathcal{T}^\varepsilon(v_1)_j \, dS(y) \, dx.$$

Using the fact that $\mathcal{T}_b^\varepsilon(n)(x, y) = n(y)$ for all $(x, y) \in \Omega \times \Sigma_Y$ and Proposition 4.2 applied to v_1 , we can conclude that

$$\varepsilon K_N \int_{\Sigma^\varepsilon} \tilde{u}_1^\varepsilon \cdot n v_1 \cdot n \, dS(x) \longrightarrow K_N \int_{\Omega} \int_{\Sigma_Y} u_1 \cdot n n \cdot v_1 \, dS(y) \, dx.$$

Following the same steps of the proof for all other summands of (25), we get the desired result. \square

We want to split the problem (24) into a micro- and macroscopic problem.

Theorem 4.19. *Let $\{u^\varepsilon\}$ be as in Theorem 4.17. We can reformulate the homogenized problem (24): Find $u_0, u_1 \in [H^1(\Omega, \Gamma_1)]^3$ with*

$$\begin{aligned} & \int_{\Omega} A_0^{\text{hom}} e(u_0) e(v_0) \, dx + \int_{\Omega} A_1^{\text{hom}} e(u_1) e(v_1) \, dx \\ & + \int_{\Omega} \int_{\Sigma_Y} \left(K_N (u_1 \cdot n - u_0 \cdot n) n + K_T \sum_{i=1}^2 (u_1 \cdot \tau^i - u_0 \cdot \tau^i) \tau^i \right) \cdot (v_1 - v_0) \, dS(y) \, dx \quad (26) \\ & = \int_{\Omega} \int_{Y_0} f \, dy \cdot v_0 \, dx + \int_{\Omega} \int_{Y_1} f \, dy \cdot v_1 \, dx + \int_{\Gamma_2} g \cdot h_0 v_0 \, dS(x) + \int_{\Gamma_2} g \cdot h_1 v_1 \, dS(x) \end{aligned}$$

whereby

$$(A_\alpha^{\text{hom}})_{ijkh} = \int_{Y_\alpha} a_{ijkh}(y) - \sum_{l,m=1}^3 a_{ijlm} \left(e_y(\chi_\alpha^{kh}) \right)_{lm} \, dy$$

and $\chi_\alpha^{lm} \in [H^1(Y_\alpha)]^3$, $l, m \in \{1, 2, 3\}$ is the unique solution of

$$\begin{cases} \left(-\sum_{j=1}^3 \frac{\partial}{\partial y_j} \left[(Ae_y(\chi_\alpha^{lm}))_{ij} - a_{ijlm} \right] \right)_{1 \leq i \leq 3} = 0 & \text{in } Y_\alpha, \\ \left(-\sum_{j=1}^3 \left[(Ae_y(\chi_\alpha^{lm}))_{ij} - a_{ijlm} \right] n_j \right)_{1 \leq i \leq 3} = 0 & \text{on } \Sigma_Y, \\ \chi_\alpha^{lm} \text{ is } Y\text{-periodic with } \mathcal{M}_{Y_\alpha}(\chi_\alpha^{lm}) = 0. \end{cases}$$

for $\alpha \in \{0, 1\}$.

Proof. This result can be shown in the same way as in the proof of Theorem 4.13. \square

Similar to [17, Theorem II.1.1], it can be proven that there exist $\alpha_0^{\text{hom}}, \beta_0^{\text{hom}}, \alpha_1^{\text{hom}}, \beta_1^{\text{hom}} \in \mathbb{R}$ with $0 < \alpha_0^{\text{hom}} < \beta_0^{\text{hom}}, 0 < \alpha_1^{\text{hom}} < \beta_1^{\text{hom}}$ such that $A_0^{\text{hom}} \in M(\alpha_0^{\text{hom}}, \beta_0^{\text{hom}}, \Omega)$, $A_1^{\text{hom}} \in M(\alpha_1^{\text{hom}}, \beta_1^{\text{hom}}, \Omega)$. With this fact we are able to prove the uniqueness of the solution of (26).

Theorem 4.20. *There exist unique solutions $u_0, u_1 \in [H^1(\Omega, \Gamma_1)]^3$ of the macroscopic problem (26).*

Proof. The proof is analogous to that of Theorem 4.14. \square

Remark 4.21. *As in the disconnected case, we can generalize the result under the assumptions that $A^\varepsilon = A(\cdot, \frac{\cdot}{\varepsilon}) \in M(\alpha, \beta, \Omega)$ and $\mathcal{T}_1^\varepsilon(A^\varepsilon) \rightarrow B_1$ a.e. in $\Omega \times Y_1$ and $\mathcal{T}_0^\varepsilon(A^\varepsilon) \rightarrow B_0$ a.e. in $\Omega \times Y_0$. The homogenized problem is the same as above but the cell problems and the definition of the homogenized tensors A_0^{hom} and A_1^{hom} need to be adjusted in an analogous fashion as in Remark 4.15.*

5. Comparison of the homogenized problem in the disconnected and connected case and discussion

In this section, we state and compare the strong formulations of the upscaled problems in the disconnected and connected case and discuss their differences.

In the disconnected case, we only know so far that the solution u_0 is unique if $\nabla \times u_0 = 0$ (see Theorem 4.14). Thus, we have to prove that a general solution already satisfies the curl-free condition, which is physically expected owing to the coupling of u_0 and u_1 through the slip

displacement conditions. (The disconnected components cannot undergo rigid-body motions due to the bond to the connected part.) By using the unique decomposition of functions $v \in L^2(\Omega)$, namely $v = \nabla p + \nabla \times w$ for some functions p, w (for more details we refer to [27]), the interface term yields the desired result. In the disconnected case (cf. eq. (21)), the strong formulation is given by: Find u_1, u_0 such that

$$\begin{aligned} -\nabla \cdot \left(A_1^{\text{hom}} e(u_1) \right) + \int_{\Sigma_Y} K_N(u_1 - u_0) \cdot nn + K_T \sum_{i=1}^2 (u_1 - u_0) \cdot \tau^i \tau^i \, dS(y) &= \int_{Y_1} f \, dy \quad \text{in } \Omega, \\ -\int_{\Sigma_Y} K_N(u_1 - u_0) \cdot nn + K_T \sum_{i=1}^2 (u_1 - u_0) \cdot \tau^i \tau^i \, dS(y) &= \int_{Y_0} f \, dy \quad \text{in } \Omega, \\ A_1^{\text{hom}} e(u_1) \cdot n &= g \quad \text{on } \Gamma_2, \\ u_1 &= 0 \quad \text{on } \Gamma_1, \end{aligned}$$

while in the connected case (cf. eq. (26)), it is: Find u_0, u_1 such that

$$\begin{aligned} -\nabla \cdot \left(A_1^{\text{hom}} e(u_1) \right) + \int_{\Sigma_Y} K_N(u_1 - u_0) \cdot nn + K_T \sum_{i=1}^2 (u_1 - u_0) \cdot \tau^i \tau^i \, dS(y) &= \int_{Y_1} f \, dy \quad \text{in } \Omega, \\ -\nabla \cdot \left(A_0^{\text{hom}} e(u_0) \right) - \int_{\Sigma_Y} K_N(u_1 - u_0) \cdot nn + K_T \sum_{i=1}^2 (u_1 - u_0) \cdot \tau^i \tau^i \, dS(y) &= \int_{Y_0} f \, dy \quad \text{in } \Omega, \\ A_0^{\text{hom}} e(u_0) \cdot n &= gh_0 \quad \text{on } \Gamma_2, \\ A_1^{\text{hom}} e(u_1) \cdot n &= gh_1 \quad \text{on } \Gamma_2, \\ u_1 &= 0 \quad \text{on } \Gamma_1, \\ u_0 &= 0 \quad \text{on } \Gamma_1. \end{aligned}$$

Apart from the different boundary conditions at the exterior boundary, the main difference is that, in the disconnected case, there is no contribution of any homogenized tensor A_0^{hom} . This means that no stress is transmitted globally by the material occupying the disconnected domain. Moreover, adding the two equations stated in Ω for the disconnected case makes the interfacial terms disappear and leaves only the total volumetric forcing given by f in the homogenized elasticity equation for u_1 . Nevertheless, as f is a force density w.r.t. volume, the density of the material does play a role. Similarly, inertial effects are expected to have an impact in the associated time-dependent case.

In terms of the application to concretes reinforced with short carbon fibres, it is unlikely that the disconnected case is the correct model. However, it may be anticipated that modelling the carbon fibres as connected in only one spatial dimension but not in the other two might lead to a mixture of the two cases, which may lead to appropriate models if realistic unit cells are used. This, together with simulation results comparing to experimental data, is planned as future work.

Acknowledgements

This work was supported by the German Research Foundation (DFG) through a project within the Special Priority Program (SPP) 2020.

References

- [1] M. Hambach, H. Möller, T. Neumann, D. Volkmer, Portland cement paste with aligned carbon fibers exhibiting exceptionally high flexural strength (> 100 MPa), *Cement and Concrete Research* 89 (2016) 80–86.
- [2] P. Lauff, M. Raith, C. Grosse, M. Rutzen, D. Volkmer, L. Reischmann, U. Weiß, M. A. Peter, O. Fischer, Investigation of localized damage indicators of a short-fibre reinforced high performance concrete under dynamic and flexural load, in: W. Derkowski et al. (Ed.), *Concrete: Innovations in Materials, Design and Structures - Proceedings of the fib Symposium 2019, Kraków, Poland, 27-29 May 2019*, pp. 1952–1959.
- [3] M. Rutzen, D. Volkmer, U. Weiß, L. Reischmann, M. A. Peter, P. Lauff, O. Fischer, M. Raith, C. Große, Microstructural analysis of crack growth caused by static and dynamic loads in a carbon fiber reinforced cement past, in: A. Zingoni (Ed.), *Advances in Engineering Materials, Structures and Systems: Innovations, Mechanics and Applications*, 2019, pp. 617–622.
- [4] P. Donato, S. Monsurrò, Homogenization of two heat conductors with an interfacial contact barrier, *Analysis and Applications* 02 (03) (2004) 247–273.
- [5] P. Donato, K. H. Le Nguyen, R. Tardieu, The periodic unfolding method for a class of imperfect transmission problems., *J. Math. Sci., New York* 176 (6) (2011) 891–927.
- [6] R. Bunoiu, C. Timofte, Upscaling of a diffusion problem with interfacial flux jump leading to a modified barenblatt model, *ZAMM - Journal of Applied Mathematics and Mechanics / Zeitschrift für Angewandte Mathematik und Mechanik* 99 (2) (2018) .
- [7] L. Tartar, Quelques remarques sur l’homogénéisation, functional analysis and numerical analysis, *Proc. Japan-France Seminar 1976* (1978) 468–482.
- [8] D. Cioranescu, A. Dalmlamian, G. Griso, The periodic unfolding method in homogenization., *SIAM J. Math. Anal.* 40 (4) (2008) 1585–1620.
- [9] M. A. Peter, M. Böhm, Different choices of scaling in homogenization of diffusion and interfacial exchange in a porous medium, *Mathematical Methods in the Applied Sciences* 31 (2008) 1257–1282.
- [10] G. Nguetseng, A general convergence result for a functional related to the theory of homogenization, *SIAM Journal on Mathematical Analysis* 20 (3) (1989) 608–629.
- [11] G. Allaire, Homogenization and two-scale convergence., *SIAM J. Math. Anal.* 23 (6) (1992) 1482–1518.
- [12] P. Donato, K. H. Le Nguyen, Homogenization of diffusion problems with a nonlinear interfacial resistance, *Nonlinear Differ. Equ. Appl.* 22 (2015) 1345–1380.
- [13] K. H. Le Nguyen, Homogenization of of heat transfer process in composite materials, *J. Elliptic Parabolic Equations* 1 (2015) 175–188.
- [14] I. Graf, M. A. Peter, J. Sneyd, Homogenization of a nonlinear multiscale model of calcium dynamics in biological cells, *Journal of Mathematical Analysis and Applications* 419 (1) (2014) 28–47.
- [15] J. Orlik, Two-scale homogenization in transmission problems of elasticity with interface jumps, *Applicable Analysis* 91 (7) (2011) 1299–1319.
- [16] M. Ptashnyk, B. Seguin, Homogenization of a system of elastic and reaction-diffusion equations modelling plant cell wall biomechanics, *ESAIM: Mathematical Modelling and Numerical Analysis* 50 (2) (2016) 593–631.
- [17] O. Oleinik, A. Shamaev, G. Yosifian, *Mathematical problems in elasticity and homogenization*, North-Holland, Amsterdam, London, New York, Tokyo, 1992.
- [18] D. Cioranescu, P. Donato, *An Introduction to Homogenization*, Oxford Univ. Press, Oxford [u.a], 1999.
- [19] M. Höpker, Extension operators for sobolev spaces on periodic domains, their applications, and homogenization of a phase field model for phase transitions in porous media, PhD thesis, University of Bremen (2016).
- [20] B. Lombard, J. Piraux, Numerical modeling of elastic waves across imperfect contacts., *SIAM J. Sci. Comput.* 28 (1) (2006) 172–205.
- [21] G. Alessandrini, A. Morassi, E. Rosset, The linear constraints in Poincaré and Korn type inequalities., *Forum Math.* 20 (3) (2008) 557–569.
- [22] S. Monsurrò, Homogenization of a two-component composite with interfacial thermal barrier, *Adv. in Math. Sci. and Appl.* 13 (01) (2003) 43–63.
- [23] D. Cioranescu, A. Dalmlamian, P. Donato, G. Griso, R. Zaki, The periodic unfolding method in domains with holes., *SIAM J. Math. Anal.* 44 (2) (2012) 718–760.
- [24] D. Cioranescu, A. Dalmlamian, G. Griso, *The periodic unfolding method*, Springer, Singapore, 2018.
- [25] D. Lukkassen, G. Nguetseng, P. Wall, Two-scale convergence, *Int. J. of Pure and Appl. Math.* 2 (1) (2002) 35–86.
- [26] J.-M. E. Bernard, Density results in Sobolev spaces whose elements vanish on a part of the boundary., *Chin. Ann. Math., Ser. B* 32 (6) (2011) 823–846.

- [27] R. Dautray, J.-L. Lions, *Mathematical Analysis and Numerical Methods for Science and Technology*, Springer, Berlin, Heidelberg, 2000.