

Orchestrating cells on a chip employing standing surface acoustic waves towards neural networks [Abstract]

Manuel S. Brugger, S. Grundeen, L. Theogarajan, Achim Wixforth, Christoph Westerhausen

Angaben zur Veröffentlichung / Publication details:

Brugger, Manuel S., S. Grundeen, L. Theogarajan, Achim Wixforth, and Christoph Westerhausen. 2017. "Orchestrating cells on a chip employing standing surface acoustic waves towards neural networks [Abstract]." *European Biophysics Journal* 46 (S1): S103. <https://doi.org/10.1007/s00249-017-1222-x>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under the following conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publizieren>



O-174 (P-930)**Orchestrating cells on a chip employing standing surface acoustic waves towards neural networks**

M. S. Brugger¹, S. Grunden², L. Theogarajan², A. Wixforth^{1,3}, C. Westerhausen^{1,3}

¹Chair for Experimental Physics 1, University of Augsburg, 86159 Augsburg, Germany; ²Department of Electrical and Computer Engineering, UCSB, CA, 93106, USA; ³Nanosystems Initiative Munich, 80799 Munich, Germany

We propose the implementation of a new lab-on-a-chip based system for the controlled growth and formation of complex neural networks on a semiconductor chip. By combining microfluidic techniques with surface acoustic waves (SAW), we can create and stimulate simple life-on-a-chip systems. For this purpose, we constructed a chip consisting of a piezoelectric LiNbO₃ substrate and four interdigital transducers for the excitation of SAW to form standing waves with according nodes and antinodes in a checkerboard pattern. The anticipated formation of the pressure node lattice has been visualized using atomic force microscopy. By adding a PDMS-microchannel, this equidistant and regular patterning lattice allows us to simultaneously control the position of objects in a liquid environment in space and time. The possibility and accuracy to pattern cell-sized single objects on these chips were validated by patterning small beads of different sizes. Ensuring the conditions for cell growth, we successfully demonstrate single cell alignment, their adhesion and growth within the well-defined pressure nodes on the chip. Finally, we verified the biocompatibility of SAW for primary neural cells. This gives us confidence that it will allow us to apply our technique to single neurons in the future.