

Regional policies targeting residential solid fuel and agricultural emissions can improve air quality and public health in the Greater Bay Area and across China

Luke Conibear, Carly L. Reddington, Ben J. Silver, Christoph Knote, Stephen R. Arnold, Dominick V. Spracklen

Angaben zur Veröffentlichung / Publication details:

Conibear, Luke, Carly L. Reddington, Ben J. Silver, Christoph Knote, Stephen R. Arnold, and Dominick V. Spracklen. 2021. "Regional policies targeting residential solid fuel and agricultural emissions can improve air quality and public health in the Greater Bay Area and across China." *GeoHealth* 5 (4): e2020GH000341. <https://doi.org/10.1029/2020gh000341>.

Special Section:

Atmospheric PM_{2.5} in China: Indoor, Outdoor, and Health Effects

Key Points:

- Ambient fine particulate matter exposure inside the Greater Bay Area is strongly controlled by emissions outside the Greater Bay Area
- Residential solid fuel and agricultural emissions lack effective controls that could improve air quality and public health across China
- Improving particulate air quality inside the Greater Bay Area will require regional cooperation inside and outside the Greater Bay Area

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

L. Conibear,
L.A.Conibear@leeds.ac.uk

Citation:

Conibear, L., Reddington, C. L., Silver, B. J., Knote, C., Arnold, S. R., & Spracklen, D. V. (2021). Regional policies targeting residential solid fuel and agricultural emissions can improve air quality and public health in the greater bay area and across China. *GeoHealth*, 5, e2020GH000341. <https://doi.org/10.1029/2020GH000341>

Received 22 OCT 2020

Accepted 1 FEB 2021

Author Contributions:

Conceptualization: Luke Conibear, Carly L. Reddington, Stephen R. Arnold, Dominick V. Spracklen

Data curation: Luke Conibear, Carly L. Reddington, Ben J. Silver

© 2021. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Regional Policies Targeting Residential Solid Fuel and Agricultural Emissions Can Improve Air Quality and Public Health in the Greater Bay Area and Across China

Luke Conibear¹ , Carly L. Reddington¹, Ben J. Silver¹ , Christoph Knote², Stephen R. Arnold¹ , and Dominick V. Spracklen¹

¹Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK,

²Faculty of Medicine, University of Augsburg, Germany

Abstract Air pollution exposure is a leading public health problem in China. The majority of the total air pollution disease burden is from fine particulate matter (PM_{2.5}) exposure, with smaller contributions from ozone (O₃) exposure. Recent emission reductions have reduced PM_{2.5} exposure. However, levels of exposure and the associated risk remain high, some pollutant emissions have increased, and some sectors lack effective emission control measures. We quantified the potential impacts of relevant policy scenarios on ambient air quality and public health across China. We show that PM_{2.5} exposure inside the Greater Bay Area (GBA) is strongly controlled by emissions outside the GBA. We find that reductions in residential solid fuel use and agricultural fertilizer emissions result in the greatest reductions in PM_{2.5} exposure and the largest health benefits. A 50% transition from residential solid fuel use to liquefied petroleum gas outside the GBA reduced PM_{2.5} exposure by 15% in China and 3% within the GBA, and avoided 191,400 premature deaths each year across China. Reducing agricultural fertilizer emissions of ammonia by 30% outside the GBA reduced PM_{2.5} exposure by 4% in China and 3% in the GBA, avoiding 56,500 annual premature deaths across China. Our simulations suggest that reducing residential solid fuel or industrial emissions will reduce both PM_{2.5} and O₃ exposure, whereas other policies may increase O₃ exposure. Improving particulate air quality inside the GBA will require consideration of residential solid fuel and agricultural sectors, which currently lack targeted policies, and regional cooperation both inside and outside the GBA.

1. Introduction

Air pollution exposure was the fourth leading risk factor to the disease burden in China in 2017, associated with 7.5% (95% uncertainty interval, UI: 6.5–8.6) of the healthy life lost (GBD 2017 Risk Factor Collaborators, 2018; Yin et al., 2020). The majority of this disease burden was attributed to ambient fine particulate matter (PM_{2.5}) exposure (71%), with contributions from household PM_{2.5} (23%) and ambient ozone (O₃, 6%) exposure (GBD 2017 Risk Factor Collaborators, 2018). The health effects incurred by air pollution in the Pearl River Delta (PRD) in South China have been estimated to result in economic losses equivalent of up to 2.3% of regional gross domestic product (GDP, D. Huang et al., 2012; Lu et al., 2016). In the absence of air pollution controls, GDP losses at the provincial level may exceed 3% by 2030 (Xie et al., 2016). This highlights the need for carefully designed policies that will reduce air pollution.

There was a 5% reduction in the loss of healthy life associated with air pollution exposure from 2013 to 2017 (8.0%, 95UI: 6.9 to 9.0, GBD 2017 Risk Factor Collaborators, 2018). This improvement in public health was primarily attributed to reduced PM_{2.5} exposure, resulting from emission reductions of the 2013–2017 Air Pollution Prevention and Control Action Plan (APPCAP, Cheng et al., 2019; Ding et al., 2019; Guo et al., 2018; J. Huang et al., 2018; X. Jiang et al., 2015; Li et al., 2019a Ministry of Environmental Protection of China, 2013; Silver et al., 2020a; B. Zheng et al., 2018; Y. Zheng et al., 2017b). For example, a key region targeted by the APPCAP was the PRD which achieved a 28% reduction in ambient PM_{2.5} concentrations (from 47 μg m⁻³ to 34 μg m⁻³), exceeding the targeted 15% reduction (China Ministry of Ecological Environment, 2013, 2017). This reduction in ambient PM_{2.5} concentrations was primarily due to emission reductions in the industrial and power generation sectors (Ding et al., 2019; B. Zheng et al., 2018; Q. Zhang et al., 2019).

Formal analysis: Luke Conibear, Carly L. Reddington

Investigation: Luke Conibear, Carly L. Reddington

Methodology: Luke Conibear, Carly L. Reddington

Project Administration: Stephen R. Arnold, Dominick V. Spracklen

Resources: Stephen R. Arnold, Dominick V. Spracklen

Software: Luke Conibear, Christoph Knote

Validation: Luke Conibear

Visualization: Luke Conibear

Writing – original draft: Luke Conibear

Writing – review & editing: Luke Conibear, Carly L. Reddington, Ben J. Silver, Christoph Knote, Stephen R. Arnold, Dominick V. Spracklen

Previous work on air pollution in the PRD has found that emissions outside the PRD contributed 53% and local emissions contributed 47% to ambient PM_{2.5} concentrations inside the PRD for 2010 (Hou et al., 2019). The contribution of emissions outside the PRD to ambient PM_{2.5} concentrations inside the PRD was highest in winter and autumn, likely due to the strong prevailing north-easterly wind (Hou et al., 2019). X. Jiang et al., (2015) found that ambient PM_{2.5} concentrations across the PRD in 2017 were 20% primary (organic and black carbon), 45% secondary (sulfate, nitrate, and ammonium), with 35% from other components. Fang et al., (2019) found that air quality in the Guangdong-Hong Kong-Macau Greater Bay Area (GBA) over 2015–2017 was worst in Foshan, Guangzhou, and Dongguan, likely due to both large local emissions and accumulation of secondary pollution. Ambient PM_{2.5} concentrations were lower in cities near the Pearl River estuary, such as Hong Kong and Shenzhen, likely due to more effective dilution by the cleaner surrounding marine air (Fang et al., 2019). Industrial emissions dominated the contribution to ambient PM_{2.5} concentrations in the PRD in 2010 (Reddington et al., 2019). Emissions across the PRD have changed in recent years. From 2010 to 2015, sulfur dioxide (SO₂) emissions reduced by 43%, PM_{2.5} emissions reduced by 28%, and nitrogen oxides (NO_x) emissions reduced by 13%, likely due to control measures on industry and energy generation (Bian et al., 2019; B. Zheng et al., 2018). In contrast, emissions of volatile organic compounds (VOCs) increased by 11% in the PRD over 2010–2015 (Bian et al., 2019; B. Zheng et al., 2018). Much of the PRD is VOC-limited, especially in autumn and winter, suggesting that increasing VOC emissions and decreasing NO_x emissions may increase O₃ concentrations (Jin & Holloway, 2015; Ou et al., 2016; Wang et al., 2017). Emissions of ammonia (NH₃), dominated by the agricultural sector, stayed approximately the same (Bian et al., 2019). This highlights the difficulty in simultaneously reducing local PM_{2.5} and O₃ concentrations (H. Liu et al., 2013). Indeed, long-term monitoring studies show PM_{2.5} concentrations across the PRD and GBA have declined in recent years, whereas O₃ concentrations have increased (Silver et al., 2020a; Silver et al., 2018). Despite improvements, PM_{2.5} concentrations across the GBA are still damaging to public health. Epidemiological studies of air pollution exposure in the GBA find associations with mortality and morbidity (Hedley et al., 2002; Peters et al., 1996; Y. Tao et al., 2012; Z. Zhang et al., 2014), in addition to the epidemiological studies across the whole of China (Baumgartner et al., 2014; R. Chen et al., 2019; Clark et al., 2019; Ebenstein et al., 2017; T. Li et al., 2018; F. Liang et al., 2019a; L. Liang et al., 2019b; Rich et al., 2015; Y. Tao et al., 2012; Tian et al., 2019; Wu et al., 2018; Yan et al., 2019; Yin et al., 2017).

Despite these recent improvements in air quality, ambient PM_{2.5} concentrations across China still largely exceed the national standard (35 μg m⁻³) and ambient O₃ concentrations increased (Silver et al., 2018; Silver et al., 2020b). There is a lack of effective emission control measures for the residential and agriculture sectors, which are both key contributors to the associated disease burden (Cai et al., 2018; Reddington et al., 2019; H. Zheng et al., 2019). The national 2018–2020 3-year plan requires emission reductions of 15% in SO₂, 15% in NO_x, and 10% in VOCs. The 3-year plan also requires that all cities that exceeded the annual-mean ambient PM_{2.5} concentration standard of 35 μg m⁻³ in 2015 to achieve 18% reductions by 2020 (Ministry of Environmental Protection of China, 2018).

We used a regional chemical transport model to explore how potential air quality policy scenarios can address these issues surrounding ambient air pollution exposure and the loss of healthy life across China, with a specific focus on the GBA. We explored the impacts of a range of potential policies and contrasted emission sectors where existing policies are already being implemented (such as land transport and industry) to sectors lacking effective policies (such as residential and agriculture).

2. Methods

2.1. Model

Simulations were conducted using the Weather Research and Forecasting model online-coupled with Chemistry (WRFChem) version 3.7.1 (Grell et al., 2005; Skamarock et al., 2008). Detailed information on the model setup is provided in Table S1 and in our previous work (Reddington et al., 2019; Silver et al., 2020a). The model domain covered China at 30 km (~0.3°) horizontal resolution with a 10 km (~0.1°) nest over the GBA (Figure S1). The GBA is an urban area surrounded by rural areas, and includes Dongguan, Foshan, Guangzhou, Huizhou, Jiangmen, Shenzhen, Zhaoqing, Zhongshan, Zhuhai, Hong Kong, and Macao.

Anthropogenic emissions for China were provided by the Multi-resolution Emission Inventory for China (MEIC) emission inventory at $0.25 \times 0.25^\circ$ horizontal resolution (M. Li et al., 2017; MEIC Research Group & Tsinghua University, 2019; B. Zheng et al., 2018). Emissions were for black carbon, organic carbon, $PM_{2.5}$, coarse particulate matter (PM_{10}), carbon monoxide, NH_3 , NO_x , SO_2 , and non-methane VOCs. VOCs were speciated according to the Model for Ozone and Related Chemical Tracers (MOZART, Emmons et al., 2010). Anthropogenic emissions of methane inside China, and all anthropogenic emissions outside of China, were from the Emission Database for Global Atmospheric Research with Task Force on Hemispheric Transport of Air Pollution (EDGAR-HTAP) version 2.2 for 2010 at $0.1 \times 0.1^\circ$ horizontal resolution (Janssens-Maenhout et al., 2015). Sectoral emissions were provided for land transport, industry, residential energy use, power generation, shipping, aircraft, and agriculture. A diurnal cycle was applied to the anthropogenic emissions (Qi et al., 2017; B. Zheng et al., 2017a).

Open biomass burning emissions were from the Fire Inventory from National Center for Atmospheric Research (FINN) version 1.5 (Wiedinmyer et al., 2011), with emissions distributed evenly throughout the boundary layer. Biogenic emissions were calculated online using the Model of Emissions of Gases and Aerosol from Nature (MEGAN, Guenther et al., 2006). Dust emissions were calculated online using Global Ozone Chemistry Aerosol Radiation and Transport (GOCART) with Air Force Weather Agency modifications (Legrand et al., 2019).

Gas phase chemistry was simulated using the extended MOZART scheme (Emmons et al., 2010; A. Hodzic & Jimenez, 2011; Knote et al., 2014). Aerosol physics and chemistry were simulated using the updated Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) scheme, with aqueous chemistry and four sectional discrete size bins; $0.039\text{--}0.156\ \mu\text{m}$, $0.156\text{--}0.625\ \mu\text{m}$, $0.625\text{--}2.5\ \mu\text{m}$, and $2.5\text{--}10\ \mu\text{m}$ (Alma Hodzic & Knote, 2014; Zaveri et al., 2008). The secondary organic aerosol formation was based on an updated volatility basis set mechanism (Knote et al., 2015).

Microphysics were simulated using the Morrison two-moment scheme (Morrison et al., 2009). Chemical initial- and boundary-conditions were taken from the MOZART/Goddard Earth Observing System Model (National Center for Atmospheric Research 2016). Meteorological initial- and boundary-conditions were taken from the European Center for Medium-Range Weather Forecasts Re-Analysis (ERA)-Interim global product (Dee et al., 2011), on a N256 ($\sim 35\ \text{km}$ at the equator) grid at the surface, on a N128 ($\sim 70\ \text{km}$ at the equator) grid above the surface, and updated every 6 h. WRF meteorology was nudged to these fields above the boundary layer.

2.2. Simulations

We performed two sets of simulations. The first set of simulations were to explore the relative roles of emissions inside and outside the GBA on ambient air pollution within the GBA. The second set of simulations explored the impacts of specific policy scenarios implemented inside and outside the GBA on ambient air quality and human health inside the GBA and across China.

The first set of simulations were for the months of April 2014, July 2014, October 2014, and January 2015, using anthropogenic emissions from MEIC for the corresponding years (2014 and 2015) with offline-nesting. The outer domain of each simulation was spun-up for 1 month, and the inner domain was spun-up for 3 days. A control simulation was performed with no changes in anthropogenic emissions. Simulations were implemented with 15% reductions in all anthropogenic emissions inside and outside the GBA. The 15% emission reduction was chosen as it is the average emission reduction target for Hong Kong and the PRD region in 2020 relative to 2015 across SO_2 , NO_x , PM_{10} , and VOC emissions (Hong Kong Environment Bureau & Ministry of Environmental Protection of China, 2019).

The second set of simulations were for the year of 2015, each with a 1-month spin-up, using anthropogenic emissions from MEIC for 2015 with online-nesting. A control (CTL) scenario was simulated where anthropogenic emissions were kept at 2015 levels. Six simulations were performed to represent relevant air quality policy scenarios (Table 1).

Two scenarios focused on the industrial sector, as this is a key source of VOC emissions in China, which are important precursors to ambient O_3 concentrations. We reduced industrial VOC emissions by 10% in

Table 1
Air Quality Policy Scenarios

Scenario	Description
CTL	A control scenario for 2015 in China
RES	Reduce residential emissions by 50% outside the GBA to approximate a 50% transition to liquefied petroleum gas
IND-GBA	Reduce industrial volatile organic compound emissions by 10% inside the GBA to approximate the attainment of this aim from the 2018–2020 3-year plan
IND-CHN	Same as for IND-GBA, though applied to the whole of China
TRA-GBA	Reduce land transport nitrogen oxides emissions by 80% inside the GBA to approximate the aim of the 2018–2020 3-year plan to enhance land transport standards in the Pearl River Delta through transitioning away from diesel
TRA-CHN	Same as for TRA-GBA, though applied to the whole of China
AGR	Reduce agricultural ammonia emissions by 30% outside the GBA to approximate the attainment of this aim from the 2018–2020 3-year plan

Note. Emission changes were applied to the residential (RES), industrial (IND), land transport (TRA), and agricultural (AGR) sectors, relative to a control (CTL), either inside the Guangdong-Hong Kong-Macau Greater Bay Area (GBA) or across China (CHN).

accordance with the aims of the 2018–2020 3-year plan, separately inside (IND-GBA) and outside the GBA (IND-CHN).

Two scenarios focused on the land transport sector, as this is a key source of NO_x emissions in China, which are another important precursor to ambient O₃ concentrations. The 2018–2020 3-year plan requires the PRD to enhance land transport standards and transition away from diesel to decrease NO_x emissions. We reduced land transport NO_x emissions by 80% to approximate diesel vehicles implementing the China six emission standard (X. Liang et al., 2019c). We applied this separately inside (TRA-GBA) and outside the GBA (TRA-CHN).

One scenario focused on the residential sector, as this is a leading contributing sector to air quality degradation across China and lacks effective emission controls (Reddington et al., 2019; Zhao et al., 2018; H. Zheng et al., 2019). The 2018–2020 3-year plan introduced specific policies for the residential sector in Beijing-Tianjin-Hebei and the surrounding areas in winter to achieve 70% clean heating by 2021 under the Clean Heating Plan (Ministry of Environmental Protection of China, 2017; National Development and Reform Commission of China, 2017). However, there are no specific policies for reducing the use of solid fuels for residential cooking and heating in South China. Solid fuels accounted for 80% of the total fuel consumption by energy in rural households in the PRD in 2017 (X. Jiang et al., 2015). We simulated a 50% transition from solid fuels to liquefied petroleum gas (LPG) outside the GBA by reducing residential emissions from solid fuel use by 50% (RES) as being representative of a partial transition under continued fuel stacking (Barrington-Leigh et al., 2019; Carter et al., 2019; S. Tao et al., 2018; Zhu et al., 2018).

One scenario focused on the agricultural sector, as this important sector has previously lacked effective emission controls (Cai et al., 2018; H. Zheng et al., 2019). The 2018–2020 3-year plan required agricultural NH₃ emissions to reduce by 40% in Beijing-Tianjin-Hebei and Yangtze River Delta by 2020. Previous work has suggested national use of fertilizer could be reduced by 30% whilst maintaining crop yields (X. Chen et al., 2014; X. Liu et al., 2016). Here we simulated the more conservative 30% reduction in agricultural NH₃ emissions nationally outside the GBA (AGR).

The agricultural and residential scenarios are intended to be approximations of expanding realistic policies at the regional scale. They are not intended to be simulations of the exact policies under the more spatially limited measures in the 2018–2020 3-year plan and clean heating plan, respectively.

2.3. Model Evaluation

We compared simulated ambient PM_{2.5} (Figure 1) and O₃ concentrations (Figure 2) against hourly measured data from over 1,600 sites across China, Macao, Hong Kong, and Taiwan as detailed in Silver et al. (2018).




Figure 1. Evaluation of modeled ambient fine particulate matter ($\text{PM}_{2.5}$) concentrations against measurements for 2015. (a) Simulated (background, combined domain) and measured (circles) annual-mean $\text{PM}_{2.5}$ concentrations. (b) Simulated (combined domain) versus measured annual-mean $\text{PM}_{2.5}$ concentrations across China and inside the Guangdong-Hong Kong-Macau Greater Bay Area (GBA). Evaluation metrics were the normalized mean bias factor (NMBF) and the normalized mean absolute error factor (NMAEF). Across China, the NMBF was 0.30 and the NMAEF was 0.43. Inside the GBA, the NMBF was 0.13 and the NMAEF was 0.13. Dotted lines show the 1:1, 2:1, and 1:2 ratios.

The same model setup was evaluated in our previous work (Reddington et al., 2019; Silver et al., 2020a). For the first set of simulations in 2014, we evaluated $\text{PM}_{2.5}$ concentrations from the nested domain over the GBA. For the second set of simulations in 2015, we evaluated a combined domain that overlaid the nested domain on top of the parent domain. The normalized mean bias factor (NMBF) and the normalized mean absolute error factor (NMAEF) were used to evaluate the model (S. Yu et al., 2006).

For the first set of simulations, the model slightly overestimated $\text{PM}_{2.5}$ concentrations in October 2014 (NMBF = 0.09 and NMAEF = 0.13), July 2014 (NMBF = 0.24 and NMAEF = 0.28), and January 2015 (NMBF = 0.26 and NMAEF = 0.26). The model slightly underestimated $\text{PM}_{2.5}$ concentrations in April 2014 (NMBF = -0.24 and NMAEF = 0.28).




Figure 2. Evaluation of modeled ambient ozone (O_3) concentrations against measurements for 2015. (a) Simulated (background, combined domain) and measured (circles) annual-mean O_3 concentrations. (b) Simulated (combined domain) versus measured annual-mean O_3 concentrations across China and inside the Guangdong-Hong Kong-Macau Greater Bay Area (GBA). Evaluation metrics were the normalized mean bias factor (NMBF) and the normalized mean absolute error factor (NMAEF). Across China, the NMBF was -0.27 and the NMAEF was 0.31. Inside the GBA, the NMBF was -0.25 and the NMAEF was 0.26. Dotted lines show the 1:1, 2:1, and 1:2 ratios.

For the second set of simulations, the model slightly overestimated PM_{2.5} concentrations (NMBF = 0.30 and NMAEF = 0.43, Figure 1) and slightly underestimated O₃ concentrations (NMBF = -0.27 and NMAEF = 0.31, Figure 2) across China. The model biases were smaller over the GBA for both PM_{2.5} (NMBF = 0.13 and NMAEF = 0.13) and O₃ (NMBF = -0.25 and NMAEF = 0.26) concentrations.

2.4. Health Impact Assessment

The health impact assessment used the second set of simulations for 2015 to estimate the disease burden attributable to air pollution exposure using population attributable fractions (PAF) of relative risk (RR). Intervention-driven variations in exposure were used to predict associated variations in outcome.

The exposure to PM_{2.5} (z) per grid cell was relative to the counterfactual exposure level of 2.4 $\mu\text{g m}^{-3}$ (cf) where no excess risk was assumed (Equation 1). The RR for a specific exposure and population age group was estimated through the Global Exposure Mortality Model (GEMM, Burnett et al., 2018). The RR was a function of the parameters θ , α , μ , and ν (Equation 2, Table S2). We used the GEMM for non-accidental mortality (non-communicable disease, NCD, plus lower respiratory infections, LRI), using parameters that included the China cohort, with age-specific modifiers for adults over 25 years of age in 5-year intervals. The GEMM functions have mean, lower, and upper uncertainty intervals. The PAF was estimated as a function of the RR and the population count (P) (Equation 3)

$$z = \max(0, PM_{2.5} - cf) \quad (1)$$

$$RR(z, age) = e^{\left\{ \theta \frac{\log\left(1 + \frac{z}{\alpha}\right)}{1 + e^{\left(\frac{\mu - z}{\nu}\right)}} \right\}} \quad (2)$$

$$PAF = P \times \left(1 - \frac{1}{RR(z, age)} \right) \quad (3)$$

The health impact assessment for O₃ exposure followed the methodology of the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) for 2017 (GBD 2017 Risk Factor Collaborators, 2018). The exposure to O₃ (z) per grid cell was calculated as the change in maximum 6-monthly-mean daily-maximum 8-hour O₃ concentrations (6mDM8h), relative to the counterfactual exposure level of 35.7 ppb (cf) where no excess risk was assumed (Equation 4, Turner et al., 2016). The 6mDM8h was calculated by first quantifying 24 separate 8-h rolling mean O₃ concentrations, then finding the maximum of these each day, then creating 12 separate 6-monthly means to account for seasonal variations, then finding the maximum of these over the year. The PAF was a function of the hazard ratio (HR), which was 1.06 (95UI: 1.02 to 1.10) for chronic obstructive pulmonary disease (COPD), based on data from five epidemiological cohorts (Equation 5, GBD 2017 Risk Factor Collaborators 2018).

$$z = \max(0, O_3 - cf) \quad (4)$$

$$PAF = P \times \left(1 - e^{-z \frac{\log HR}{10}} \right) \quad (5)$$

Premature mortality (MORT), years of life lost (YLL), and years lived with disability (YLD) per exposure, health outcome, age bracket, and grid cell were estimated as a function of the PAF and the corresponding baseline mortality and morbidity rate (I_{MORT} , I_{YLL} , and I_{YLD}) following Equations 6–8, respectively. Disability-adjusted life years (DALYs), i.e. the total loss of healthy life, were estimated as the total of YLL and YLD following Equation 9. The rates of MORT, YLL, YLD, and DALYs were calculated per 100,000 population:

$$MORT = PAF \times I_{MORT} \quad (6)$$




Figure 3. The impact of 15% emission reductions outside and inside the Guangdong-Hong Kong-Macau Greater Bay Area (GBA) on monthly-mean (January, April, July, and October) ambient fine particulate matter ($PM_{2.5}$) exposure within the GBA. Simulated changes in $PM_{2.5}$ exposure are from the nested domain at 10 km over the GBA.

$$YLL = PAF \times I_{YLL} \quad (7)$$

$$YLD = PAF \times I_{YLD} \quad (8)$$

$$DALYs = YLL + YLD \quad (9)$$

The United Nations adjusted population count dataset for 2015 at $0.05 \times 0.05^\circ$ resolution was obtained from the Gridded Population of the World, Version 4 (GPWv4) (Center for International Earth Science Information Network & NASA Socioeconomic Data and Applications Center, 2016). Population age composition was taken from the GBD2017 for 2015 for adults 25–80 years in 5-year intervals, and for 80 years plus (Global Burden of Disease Study 2017, 2018). Cause-specific (NCD, LRI, and COPD) baseline mortality and morbidity rates were taken from the GBD2017 for 2015 for MORT, YLL, and YLD for each age bracket (Institute for Health Metrics and Evaluation, 2020).

Shapefiles were used to aggregate results at the country, province, and prefecture level (Hijmans et al., 2016). Regional groupings were also applied as follows (Figure S2): North China (Beijing, Tianjin, Hebei, Shanxi, and Inner Mongolia), North East China (Liaoning, Jilin, and Heilongjiang), East China (Shanghai, Jiangsu, Zhejiang, Anhui, Fujian, Jiangxi, and Shandong), South Central China (Henan, Hubei, Hunan, Guangdong, Guangxi, Hainan, Hong Kong, and Macau) including the GBA, South West China (Chongqing, Sichuan, Guizhou, Yunnan, and Tibet), and North West China (Shaanxi, Gansu, Qinghai, Ningxia, and Xinjiang), and the GBA individually.

Uncertainty intervals at the 95% confidence level (95UI) were estimated through using the derived uncertainty intervals from the exposure–outcome associations, baseline mortality and morbidity rates, and population age fractions. Health impact assessments of the disease burden associated with air pollution exposure have many uncertainties, as detailed in Nethery & Dominici, (2019).

3. Results

3.1. Contribution of Emissions outside the GBA to Ambient Air Quality inside the GBA

Figure 3 compares the impacts of 15% emission reductions inside and outside the GBA on ambient $PM_{2.5}$ exposure inside the GBA. Emission reductions outside the GBA produce larger reductions in $PM_{2.5}$ exposure for January (8%) and October (6%) than emission reductions inside the GBA (4% for January and 5% for October). In contrast, emission reductions outside the GBA produce smaller reductions in $PM_{2.5}$ exposure for April (3%) and July (2%) than emission reductions inside the GBA (5% for April and 8% for July). These results demonstrate the importance of pollution transport into the GBA in autumn and winter when ambient $PM_{2.5}$ concentrations are generally higher than in spring and summer.

3.2. Current Disease Burden Associated with Ambient Air Pollution Exposure

Figure 4 shows the simulated disease burden due to ambient $PM_{2.5}$ and O_3 exposure. The simulated population-weighted annual-mean $PM_{2.5}$ exposure is $72.8 \mu g m^{-3}$ across China and $39.6 \mu g m^{-3}$ inside the GBA. The population-weighted 6mDM8h O_3 exposure is 63.5 ppb across China and 61.3 ppb inside the GBA. There are 2,779,000 (95UI: 2,701,000–2,865,000) premature deaths per year associated with ambient $PM_{2.5}$ exposure in China, of which 105,000 (95UI: 102,000–108,000) premature deaths are inside the GBA. The $PM_{2.5}$ disease burden is particularly large across East and South China. There are 123,000 (95UI: 86,000–171,000) premature deaths per year associated with ambient O_3 exposure in China, of which 5,700 (95UI: 4,000–7,900) premature deaths are inside the GBA. The O_3 disease burden is largest in South China. The DALYs rate from $PM_{2.5}$ exposure is 4,476 (95UI: 3,947–5,084) per 100,000 population across China, and 4,887 (95UI: 4,309–5,551) per 100,000 population inside the GBA. The DALYs rate from O_3 exposure is 186 (95UI:




Figure 4. The disease burden associated with ambient air pollution exposure from the control scenario. Rate of disability-adjusted life years (DALYs) per 100,000 population from (a) ambient fine particulate matter ($PM_{2.5}$) exposure in China, (b) ambient $PM_{2.5}$ exposure in the Guangdong–Hong Kong–Macau Greater Bay Area (GBA), (c) ambient ozone (O_3) exposure in China, and (d) ambient O_3 exposure in the GBA.


122–267) per 100,000 population across China, and 182 (95UI: 120–262) per 100,000 population inside the GBA.

Our disease burden estimates are comparable to previous work. Our calculated disease burden associated with ambient $PM_{2.5}$ exposure in China is 13% higher than that from Burnett et al. (2018), due to our higher estimate of $PM_{2.5}$ exposure. Our calculated disease burden associated with ambient O_3 exposure in China is 27% lower than that from the GBD 2017 Risk Factor Collaborators, (2018), due to our lower estimate of O_3 exposure. Combined with our evaluation of $PM_{2.5}$ and O_3 concentrations, this demonstrates that our modeling framework is suitable for the assessment of emission scenarios on air quality and public health.

3.3. Impacts of Policy Scenarios on Ambient Air Quality and Public Health in China

Figure 5 summarizes the impacts of each scenario on ambient air quality and public health. The results per region are in Table S3. Ambient air quality is reported as the population-weighted annual-mean $PM_{2.5}$ exposure and the population-weighted 6mDM8h O_3 exposure.

The residential scenario of a 50% transition from residential solid fuel use to LPG outside the GBA result in the largest reductions across China in both $PM_{2.5}$ exposure (1.4 – $14.3 \mu g m^{-3}$, 3–18%) and O_3 exposure (0.3 – 1.8 ppb, 0–2%). The reductions in $PM_{2.5}$ exposure are largest in North China ($14.3 \mu g m^{-3}$, 17%) and exceed $10 \mu g m^{-3}$ in South West, South Central, East, and North East China (Figure S3). The reductions in O_3 exposure are largest in South West (1.8 ppb, 2%) and South Central China (1.0 ppb, 1%, Figure S4). The residential scenario resulted in the largest public health benefits across China. The reduction in $PM_{2.5}$ exposure avoids 188,200 (95UI: 182,900–194,000) premature deaths (277 DALYs per 100,000, 95UI: 245–315), of which 29% are in East China and 24% are in South Central China (Figure S5). The reduction in O_3 exposure avoids 3,200 (95UI: 2,300–4,400) premature deaths (4 DALYs per 100,000, 95UI: 2–5), of which 66% are in South Central and South West China (Figure S6). The reduction of residential solid fuel use emissions



outside the GBA avoids 2,400 (95UI: 2,200–2,500) premature deaths (116 DALYs per 100,000, 95UI: 101–133) each year inside the GBA.

The next largest public health benefits result from the agricultural scenario of a 30% reduction in NH_3 emissions from fertilizer use outside the GBA. Under the agricultural scenario, $\text{PM}_{2.5}$ exposure decreases by 1.2–3.8 $\mu\text{g m}^{-3}$ (3–5%) across China. The reductions in $\text{PM}_{2.5}$ exposure are largest in North (3.8 $\mu\text{g m}^{-3}$, 4%), South Central (3.7 $\mu\text{g m}^{-3}$, 5%), and South West China (3.5 $\mu\text{g m}^{-3}$, 5%). However, the agricultural scenario increases O_3 exposure by 0.1–0.7 ppb (0–1%), mainly across North and South West China (both 0.7 ppb, 1%). Despite this increase in O_3 exposure, there are 56,500 (95UI: 55,500–57,600) fewer premature deaths (98 DALYs per 100,000, 95UI: 87–111) across China, as the O_3 disease burden is ~5% of the $\text{PM}_{2.5}$ disease burden. The reduction of agricultural fertilizer emissions outside the GBA avoids 1,900 (95UI: 1,800–1,900) premature deaths (93 DALYs per 100,000, 95UI: 82–105) each year inside the GBA.

The regional land transport scenario of an 80% reduction in NO_x emissions across China reduces $\text{PM}_{2.5}$ exposure by 0.3–2.8 $\mu\text{g m}^{-3}$ (1–4%) across China. The reductions in $\text{PM}_{2.5}$ exposure are largest in South Central and South West China (both 2.8 $\mu\text{g m}^{-3}$, 4%). However, under the regional land transport scenario, O_3 exposure increases by 1.4 ppb (2%) across China. O_3 exposure increases in East, North, South Central, and North East China (0.6–3.5 ppb, 1–6%), while O_3 exposure decreases in South West and North West China (1.0–1.6 ppb, 2%). Reducing NO_x emissions decreases O_3 exposure in the rural areas of South West China, as they are considered to be NO_x -limited (Jin & Holloway, 2015; Wang et al., 2017). In contrast, reducing NO_x emissions increases O_3 exposure in the urban areas of East China, as they are considered to be VOC-limited (Jin & Holloway, 2015; Wang et al., 2017). Despite the dichotomy in exposure change, there are 27,200 (95UI: 24,200–29,900) fewer premature deaths (72 DALYs per 100,000, 95UI: 62–84) across China, as the decrease in the $\text{PM}_{2.5}$ disease burden outweighs the increase in the O_3 disease burden. The regional reduction in land transport NO_x emissions avoids 200 (95UI: 100–300) premature deaths (17 DALYs per 100,000, 95UI: 17–19) each year inside the GBA.

The regional industrial scenario of a 10% reduction in VOC emissions across China reduces both $\text{PM}_{2.5}$ exposure (0.4–1.1 $\mu\text{g m}^{-3}$, 1%) and O_3 exposure (0.3–0.9 ppb, 0–1%) across China. The largest reduction in both $\text{PM}_{2.5}$ and O_3 exposure are in East, North, and South Central China. Reducing VOC emissions in urban areas across China reduces O_3 concentrations, as these areas are considered to be VOC-limited (Jin & Holloway, 2015; Wang et al., 2017). These exposure reductions avoid 14,300 (95UI: 13,900–14,800) premature deaths (18 DALYs per 100,000, 95UI: 16–20) from $\text{PM}_{2.5}$ exposure and avoid 2,800 (95UI: 2,000–4,400) premature deaths (1 DALYs per 100,000, 95UI: 1–2) from O_3 exposure. Within these public health benefits, there are 900 (95UI: 900–1,000) fewer premature deaths (42 DALYs per 100,000, 95UI: 35–48) each year inside the GBA.

The local industrial and land transport scenarios to the GBA produce minor changes in both $\text{PM}_{2.5}$ and O_3 exposure.

4. Discussion

Our scenarios inside the GBA focused on industrial VOC emissions and land transport NO_x emissions, as these are key local sources of precursor emissions to O_3 concentrations where existing policies are being implemented. Our scenarios outside the GBA focused on residential emissions and agricultural NH_3 emissions, as these key regional sources lack effective policies except in specific regions. A comparative air quality and health impact assessment of these scenarios across different regions in China enabled the impacts from emission changes inside the GBA to be contrasted against emission changes outside the GBA.

Figure 5. The impacts of policy scenarios on air quality and public health per region in China. Columns are population-weighted annual-mean ambient fine particulate matter ($\text{PM}_{2.5}$) exposure, population-weighted maximum 6-monthly-mean daily-maximum 8-h ambient ozone (O_3) exposure, and the annual-sum of premature mortality (MORT) associated with $\text{PM}_{2.5}$ and O_3 exposure. Rows are for China, North China, North East China, East China, South Central China, South West China, North West China, and the Guangdong-Hong Kong-Macau Greater Bay Area (GBA, see Figure S2 for more details). Scenarios are for the residential (RES), industrial within the GBA (IND-GBA), industrial over China (IND-CHN), land transport within the GBA (TRA-GBA), land transport over China (TRA-CHN), and agriculture (AGR), with the percentage change relative to the control (see Table 1 for more details).

The importance of pollution transport into the GBA in winter was also seen by Hou et al. (2019), who found that pollution outside the PRD contributed 53% of the annual-mean ambient $PM_{2.5}$ concentrations inside the PRD, with contributions highest in winter. The larger contribution of pollution transport in winter is due to the strong prevailing North Easterly winds, which bring in pollution from the highly polluted regions outside the GBA (Hou et al., 2019).

Our work further confirms the large contribution of residential solid fuels to particulate air quality across China and suggests policies to reduce emissions from this sector would reduce $PM_{2.5}$ and O_3 exposure. Previous policies that promoted residential solid fuel use had large impacts on air quality and public health in China. For example, the China's Huai River policy provided heavily subsidized coal for household heating to the north of the Huai River but not to the south from 1950 to 1980, and resulted in ambient PM_{10} exposure increasing by $41.7 \mu\text{g m}^{-3}$ in the north which reduced life expectancy by 3.1 years (Ebenstein et al., 2017; Yuyu Chen et al., 2013). Residential solid fuel use emissions from cooking did decrease from 1992 to 2015 in China (Aunan & Wang, 2014; Du et al., 2018; S. Tao et al., 2018; Yilin Chen et al., 2016), leading to large public health benefits (Zhao et al., 2018; H. Zheng et al., 2019). However, these improvements were primarily due to urbanization and increased incomes rather than specific policies (Zhao et al., 2018). New policies focusing on residential solid fuels are required because residential solid fuels are still widely used for heating (S. Tao et al., 2018; Zhu et al., 2018), urbanization and income growth are projected to slow (L. Jiang & O'Neill, 2017), and stacking clean fuels with solid fuels is still persistent (Barrington-Leigh et al., 2019; Zhu et al., 2018).

The 2018–2020 3-year plan introduced the first specific policies for residential heating in Beijing–Tianjin–Hebei and the surrounding areas. This policy is estimated to reduce $PM_{2.5}$ emissions from household fuels by 15%–41%, reduce total $PM_{2.5}$ exposure by 10–44%, and avoid 32,000–56,000 premature deaths (J. Liu et al., 2019a; Meng et al., 2019; Qin et al., 2017; Zhao et al., 2018). This exposure change is of a similar magnitude to our 17% reduction in $PM_{2.5}$ exposure in North China under the residential scenario, where 28,900 (95UI: 28,100–29,800) premature deaths were avoided. There are currently no specific policies for tackling residential cooking and heating from solid fuels in South China, despite this source making a large contribution to current emissions. For example, solid fuels account for 80% of the total fuel consumption in terms of energy generated within rural households in the PRD and are widely used in South West China (X. Jiang et al., 2015; S. Tao et al., 2018; Zhu et al., 2018). Our residential scenario reduced $PM_{2.5}$ exposure by $10.3 \mu\text{g m}^{-3}$ in South Central China and by $11.9 \mu\text{g m}^{-3}$ in South West China, avoiding 73,800 (95UI: 71,700–76,200) premature deaths across these regions. We found that air quality inside the GBA can be improved from reductions in residential emissions outside the GBA, as pollution transport into the GBA is important. These reductions in residential emissions may be realized in North China from current policies, however, solid fuel use remains high in many rural areas of South China (Yun et al., 2020), where there are currently no specific policies to reduce residential emissions.

Reducing residential solid fuel use will also lead to further public health benefits from reductions in household $PM_{2.5}$ exposure (Aunan et al., 2018; Chan et al., 2019; Kim et al., 2016; Li et al., 2019b; S. Li et al., 2019d; Vermeulen et al., 2019; K. Yu et al., 2018; Zhao et al., 2018). A reduction in residential solid fuel use may also reduce greenhouse gas emissions (Rive & Aunan, 2010; Smith & Haigler, 2008), depending on the replacement fuel (Qin et al., 2017; Yang & Jackson, 2013). The transition from residential solid fuel use to LPG may change VOC emissions and therefore O_3 formation, though these changes are not accounted for here (Conibear et al., 2020; Lyu et al., 2017). The reduction in agricultural NH_3 emissions could increase acid rain (M. Liu et al., 2019b), and may therefore be more suitable for areas with little acid rain and large $PM_{2.5}$ concentrations rather than areas with existing acid rain issues.

We found that reducing NO_x emissions from land transport by 80% across China avoided 33,000 (95UI: 32,100–34,000) premature deaths from $PM_{2.5}$ exposure. Anenberg et al., (2017) estimated that the China 6 emission standard would avoid 69,500 premature deaths from $PM_{2.5}$ exposure. These larger estimates from Anenberg et al. (2017) were for 2040 and included increased car ownership, population growth, and aging. A transition away from diesel, such as the China six emission standard, will also likely lead to reductions in primary particulate emissions from land transport, that are not accounted for here.

The GBA aims that by 2025 there will be a further 30% reduction in annual-mean ambient $PM_{2.5}$ concentrations to reach $25 \mu\text{g m}^{-3}$, and that daily-maximum 8-hour O_3 concentrations are under 75 ppb (Hong Kong Environment Bureau & Ministry of Environmental Protection of China, 2019; Tsinghua University, 2017; United Nations Environment Programme, 2019). If we adjust our simulated $PM_{2.5}$ exposure in the GBA by the positive bias (13%), then our control $PM_{2.5}$ exposure is just under the annual standard ($35 \mu\text{g m}^{-3}$) at $34 \mu\text{g m}^{-3}$. If we adjust our simulated O_3 exposure in the GBA by the negative bias (25%), then our control O_3 (6mDM8h) exposure is 77 ppb. Using these bias-adjusted exposures for the different scenarios, we find that no single scenario achieves either of these 2025 goals in the GBA. Furthermore, although the residential scenario provided large public health benefits by avoiding 191,400 (95UI: 185,200–198,400) premature deaths each year across China, this represents only 7% of the current disease burden from ambient air pollution exposure. This means that even under this residential scenario, more than 2,500,000 premature deaths would still occur every year due to ambient air pollution exposure. This highlights that further strong emission reductions are required in order to reach air quality standards that protect public health.

In recent years, there have been substantial reductions in $PM_{2.5}$ concentrations but increasing O_3 concentrations across China (Silver et al., 2018, 2020b). This highlights the need to identify policies that reduce both $PM_{2.5}$ and O_3 exposure. We found that the residential and regional industrial scenarios achieved reductions in both $PM_{2.5}$ and O_3 exposure, whereas the agricultural and regional land transport scenarios decreased $PM_{2.5}$ exposure at the expense of increased O_3 exposure. Reduced NO_x emissions can increase O_3 concentrations in urban regions due to less nitric oxide plus O_3 , and reduce $PM_{2.5}$ concentrations by reducing nitrate aerosol concentrations. K. Li et al. (2019c, 2020) suggested that recent increases in O_3 concentrations were attributable to decreased NO_x emissions in VOC-limited urban areas and to decreased $PM_{2.5}$ concentrations that have slowed aerosol radical sinks.

Emissions over China are uncertain, and future work reducing these uncertainties will improve the accuracy of comparative health impact assessments of different scenarios (M. Li et al., 2017; Saikawa et al., 2017). The residential scenario applied reductions across the whole sector, rather than by specific solid fuel type, due to limited input data availability. The impacts of this constraint will be small as the majority of residential emissions are from solid fuel use (Carter et al., 2019; S. Tao et al., 2018). However, the accuracy of this scenario would be improved by the availability of fuel specific emission inventories with time-activity data. The future availability of higher resolution emissions and baseline health data may improve the accuracy of this health impact assessment. Models with higher spatial resolution may better capture the spatial and temporal variation of $PM_{2.5}$ concentrations (Y. Li et al., 2016; Pungler & West, 2013; Thompson et al., 2014). Simulated O_3 concentrations may be less sensitive to increases in spatial resolution, relative to $PM_{2.5}$ concentrations (Pungler & West, 2013; Thompson & Selin, 2012; Wild & Prather, 2006).

Currently, there are three main types of RR model for health impact assessments of $PM_{2.5}$ exposure: log-linear (Pope III et al., 2002), integrated-exposure response (IER, R. Burnett et al., 2014), and the GEMM (Burnett et al., 2018). The IER and GEMM are both supra-linear models with steeper increases in RR at lower exposures which flatten off at higher exposures, while the log-linear models the logarithm of the RR as linear to the exposure change. The GEMM model has a larger increase in the RR at lower exposures ($<30 \mu\text{g m}^{-3}$) compared to the log-linear model (Burnett & Cohen, 2020). However, the log-linear model has substantially larger increases in the RR at higher exposures ($>50 \mu\text{g m}^{-3}$), which are considered biologically implausible and unsuitable for health impact assessments over areas with high $PM_{2.5}$ exposures (Burnett & Cohen, 2020). The IER risks are smaller than the GEMM risks due to the inclusion of epidemiological data from second-hand smoking and household air pollution in the IER, and because the GEMM model includes more causes of deaths than the six causes included in the IER (Burnett & Cohen, 2020). Hence, for single national assessments with high $PM_{2.5}$ exposures the GEMM model is a suitable choice of exposure-outcome association, though the disease burden estimates will be larger than those obtained from global comparative assessments using the IER. The differences between these RR models will be reduced when further data is included from epidemiological studies over areas with high $PM_{2.5}$ exposure (Pope III, Coleman, Pond, & Burnett, 2019), such as the China cohort (Yin et al., 2017).

Across the whole O_3 exposure range (35–200 ppb), the relationship between O_3 exposure and PAF is non-linear (Equation 5, Conibear et al., 2020; GBD 2017 Risk Factor Collaborators, 2018). However, for O_3 concentrations of 50–80 ppb, the association between O_3 exposure and PAF is approximately linear. Most of the

population in China is exposed to 6mDM8h O₃ exposures in this range, and hence, the relative impacts of O₃ exposure on disease burden are approximately proportionally linear (Figure 5).

In this study, we focused on the impacts of individual realistic air quality policy scenarios on ambient air pollution and human health. The impacts of these scenarios are policy-specific and do not represent general attributions to air quality exposure, where residential and industrial emissions have been found to dominate across China (Reddington et al., 2019). In reality, multiple policies will overlap and interact, and future work is needed to study these combined impacts.

5. Conclusion

In this study, we used a regional chemical transport model to explore the impacts of different emission scenarios on ambient air pollution exposure and public health across China. Our study focused on the GBA in South China, which suffers from poor air quality.

We find that in winter and autumn, emission sources outside the GBA contribute more to PM_{2.5} exposure in the GBA than emission sources inside the GBA. PM_{2.5} exposure in the GBA reduced by 8% in winter and 6% in autumn when emissions were reduced by 15% outside the GBA, larger than when emissions were reduced by 15% inside the GBA (4% in winter and 5% in autumn).

We quantified the potential impacts of six different air quality policy scenarios on ambient air pollution exposure and public health across China. Our selected policies addressed the land transport and industrial sectors where there are existing policies, as well as the residential and agricultural sectors where policy controls are less developed.

The scenario of a 50% transition from residential solid fuel use to LPG outside the GBA had the greatest benefit on public health across China and inside the GBA. PM_{2.5} exposure reduced by 10.6 μg m⁻³ (15%) and O₃ exposure reduced by 0.8 ppb (1%) across China, with large reductions in North, South West, South Central, East, and North East China. The residential scenario avoided 191,400 (95UI: 185,200–198,400) premature deaths each year across China, with 2,400 (95UI: 2,200–2,500) fewer premature deaths inside the GBA.

The next largest public health benefit came from the agricultural scenario of a 30% reduction in NH₃ emissions from fertilizer use outside the GBA. The agricultural scenario reduced PM_{2.5} exposure by 3.2 μg m⁻³ (4%) across China, with the largest reductions in North, South Central, and South West China. However, O₃ exposure increased by 0.5 ppb (1%) across China, mainly across North and South West China. As the O₃ disease burden is approximately 5% of the PM_{2.5} disease burden, the agricultural scenario still had a large net benefit to public health with 56,500 (95UI: 55,500–57,600) avoided premature deaths across China, of which 1,900 (95UI: 1,800–1,900) were inside the GBA.

The 2018–2020 3-year plan requires the PRD to enhance land transport standards and transition away from diesel to decrease NO_x emissions. We simulated an approximation to this transition by reducing land transport NO_x emissions by 80%. If this policy is applied only within the GBA, our simulations suggest PM_{2.5} and O₃ exposure remain relatively unchanged. If this policy is applied nationwide, PM_{2.5} exposure decreased by 1.9 μg m⁻³ (3%) across China, primarily from reductions in South Central and South West China. However, O₃ exposure increased by 1.4 ppb (2%) across China, where exposure increased in East, North, South Central, and North East China, and decreased in South West and North West China. This scenario resulted in 27,200 (95UI: 24,200–29,900) fewer premature deaths each year, of which 200 (95UI: 100–300) avoided premature deaths were inside the GBA.

The 2018–2020 3-year plan aims to tackle the recent rise in VOC emissions by requiring a 10% emission reduction in industrial VOC emissions. Our simulations suggest that if this policy is applied solely within the GBA, then there are only minor changes in PM_{2.5} and O₃ exposure. If this policy is applied nationwide, then PM_{2.5} exposure reduced by 0.9 μg m⁻³ (1%) and O₃ exposure reduced by 0.7 ppb (1%), both primarily due to reductions in East, North, and South Central China. The regional industrial scenario avoided 17,100 (95UI: 15,900–19,200) premature deaths across China, of which 900 (95UI: 900–1,000) avoided premature deaths were inside the GBA.

Overall, we find that controls on residential solid fuel and agricultural emissions would provide the largest public health benefits both across China and inside the GBA. There are currently no specific policies for reducing residential solid fuel use or agricultural emissions in South China, highlighting a major opportunity for targeted emission controls. Improving air quality inside the GBA will require coordinated regional emission reduction policies both inside and outside the GBA from key contributing sectors.

Conflict of Interest

The authors declare no conflicts of interest relevant to this study.

Data Availability Statement

The air pollution and health impact assessment data per Chinese province and GBA prefecture that support the findings of this study are available at doi.org/10.5518/919. Code to setup and run WRFChem (using WRFotron version 2.0) is available through Conibear and Knoté (2020).

Acknowledgments

The authors gratefully acknowledge support from the AIA Group Limited, a European Research Council Consolidator Grant (771492), and the Natural Environment Research Council (NE/S006680/1). This work was undertaken on Advanced Research Computing, part of the High Performance Computing facilities at the University of Leeds, UK. This work used WRFotron version 2.0, a tool to automatize WRFChem runs with re-initialized meteorology (Conibear & Knoté, 2020). The authors acknowledge the use of WRFChem preprocessor tools `mozbc`, `fire_emiss`, `anthro_emiss`, `bio_emiss` provided by the Atmospheric Chemistry Observations and Modeling Laboratory of the National Center for Atmospheric Research. The authors acknowledge the use of Model for Ozone and Related Chemical Tracers global model output available at <https://www.acom.ucar.edu/wrf%-chem/mo-zart.shtml>. We acknowledge the use of the emissions pre-processor available at github.com/douglowe/WRF_UoM_EMIT. The authors thank Qiang Zhang and Meng Li for providing MEIC data. The authors acknowledge the Python Software Foundation, Python Language Reference, available at python.org. We are particularly grateful to the Python libraries NumPy (Harris et al., 2020), Pandas (McKinney, 2010), Matplotlib (Hunter, 2007), SciPy (Virtanen et al., 2020), xarray (Hoyer & Hamman, 2017), Cartopy (Met Office, 2015), GeoPandas (Jordahl et al., 2020), Salem (Maussion, TimoRoth, Tbridel, Dusch, & Landmann, 2019), Jupyter (Kluyver et al., 2016), Rasterio (Gillies, 2013), Affine, and xESMF (Zhuang, Dussin, Jüling, & Rasp, 2020). The boundaries shown on any maps in this work do not imply any judgment concerning the legal status of any territory or the endorsement or acceptance of such boundaries.

References

- Anenberg, S. C., Miller, J., Minjares, R., Du, L., Henze, D. K., Lacey, F., et al. (2017). Impacts and mitigation of excess diesel-related NO_x emissions in 11 major vehicle markets. *Nature*, *545*, 467. <https://doi.org/10.1038/nature22086>
- Aunan, K., Ma, Q., Lund, M. T., & Wang, S. (2018). Population-weighted exposure to PM_{2.5} pollution in China: An integrated approach. *Environment International*, *120*, 111–120. <https://doi.org/10.1016/j.envint.2018.07.042>
- Aunan, K., & Wang, S. (2014). Internal migration and urbanization in China: Impacts on population exposure to household air pollution (2000–2010). *The Science of the Total Environment*, *481*(1), 186–195. [tagrole, struvcteam. https://doi.org/10.1016/j.scitotenv.2014.02.073](https://doi.org/10.1016/j.scitotenv.2014.02.073)
- Barrington-Leigh, C., Baumgartner, J., Carter, E., Robinson, B. E., Tao, S., & Zhang, Y. (2019). An evaluation of air quality, home heating and well-being under Beijing's programme to eliminate household coal use. *Nature Energy*, *4*(5), 416–423. <https://doi.org/10.1038/s41560-019-0386-2>
- Baumgartner, J., Zhang, Y., Schauer, J. J., Huang, W., Wang, Y., & Ezzati, M. (2014). Highway proximity and black carbon from cookstoves as a risk factor for higher blood pressure in rural China. *Proceedings of the National Academy of Sciences*, *111*(36), 13229–13234. <https://doi.org/10.1073/pnas.1317176111>
- Bian, Y., Huang, Z., Ou, J., Zhong, Z., Xu, Y., Zhang, Z., et al. (2019). Evolution of anthropogenic air pollutant emissions in Guangdong Province, China, from 2006 to 2015. *Atmospheric Chemistry and Physics*, *19*, 11701–11719. <https://doi.org/10.5194/acp-19-11701-2019>
- Burnett, R., Arden Pope, C., Ezzati, M., Olives, C., Lim, S. S., Mehta, S., et al. (2014). An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure. *Environmental Health Perspectives*, *122*(4), 397–403. <https://doi.org/10.1289/ehp.1307049>
- Burnett, R., Chen, H., Szyszkwicz, M., Fann, N., Hubbell, B., Pope, C. A., et al. (2018). Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter. *Proceedings of the National Academy of Sciences*, *115*(38), 9592–9597. <https://doi.org/10.1073/pnas.1803222115>
- Burnett, R., & Cohen, A. (2020). Relative risk functions for estimating excess mortality attributable to outdoor PM_{2.5} air pollution: Evolution and state-of-the-art. *Atmosphere*, *11*(589). <https://doi.org/10.3390/atmos11060589>
- Cai, S., Ma, Q., Wang, S., Zhao, B., Brauer, M., Cohen, A., et al. (2018). Impact of air pollution control policies on future PM_{2.5} concentrations and their source contributions in China. *Journal of Environmental Management*, *227*(August), 124–133. <https://doi.org/10.1016/j.jenvman.2018.08.052>
- Carter, E., Yan, L., Fu, Y., Robinson, B., Kelly, F., Elliott, P., et al. (2019). Household transitions to clean energy in a multiprovincial cohort study in China. *Nature Sustainability*, *3*, 42–50. <https://doi.org/10.1038/s41893-019-0432-x>
- Center for International Earth Science Information Network, & NASA Socioeconomic Data and Applications Center (2016). *Gridded population of the World, version 4 (GPWv4): Population count*. Palisades, NY: Columbia University. <https://doi.org/10.7927/H4NP22DQ>
- Chan, K. H., Kurmi, O. P., Bennett, D. A., Yang, L., Chen, Y., Tan, Y., et al. (2019). Solid fuel use and risks of respiratory diseases: A cohort study of 280,000 Chinese never-smokers. *American Journal of Respiratory and Critical Care Medicine*, *199*(3), 352–361. <https://doi.org/10.1164/rccm.201803-0432OC>
- Chen, R., Yin, P., Meng, X., Wang, L., Liu, C., Niu, Y., et al. (2019). Associations between coarse particulate matter air pollution and cause-specific mortality: A nationwide analysis in 272 Chinese cities. *Environmental Health Perspectives*, *127*(January), 5–10. <https://doi.org/10.1289/EHP2711>
- Chen, X., Cui, Z., Fan, M., Vitousek, P., Zhao, M., Ma, W., et al. (2014). Producing more grain with lower environmental costs. *Nature*, *514*(7253), 486–489. <https://doi.org/10.1038/nature13609>
- Chen, Y., Ebenstein, A., Greenstone, M., & Li, H. (2013). Evidence on the impact of sustained exposure to air pollution on life expectancy from China's Huai River policy. *Proceedings of the National Academy of Sciences*, *13*(15), 1–53. <https://doi.org/10.1073/pnas.1300018110>
- Chen, Y., Shen, H., Zhong, Q., Chen, H., Huang, T., Liu, J., et al. (2016). Transition of household cookfuels in China from 2010 to 2012. *Applied Energy*, *184*, 800–809. <https://doi.org/10.1016/j.apenergy.2016.07.136>
- Cheng, J., Su, J., Cui, T., Li, X., Dong, X., Sun, F., et al. (2019). Dominant role of emission reduction in PM_{2.5} air quality improvement in Beijing during 2013–2017: A model-based decomposition analysis. *Atmospheric Chemistry and Physics*, *19*(9), 6125–6146. <https://doi.org/10.5194/acp-19-6125-2019>
- China Ministry of Ecological Environment (2013). *China ecological environment status bulletin*. Retrieved from <http://www.mee.gov.cn/hjzl/zghjzkgb/lnzghjzkgb/>
- China Ministry of Ecological Environment (2017). *China ecological environment status bulletin*. Retrieved from <http://www.mee.gov.cn/hjzl/zghjzkgb/lnzghjzkgb/>

- Clark, S. N., Schmidt, A. M., Carter, E. M., Schauer, J. J., Yang, X., Ezzati, M., et al. (2019). Longitudinal evaluation of a household energy package on blood pressure, central hemodynamics, and arterial stiffness in China. *Environmental Research*, *177*, 108592. <https://doi.org/10.1016/j.envres.2019.108592>
- Conibear, L., Butt, E. W., Knote, C., Lam, N. L., Arnold, S. R., Tibrewal, K., et al. (2020). A complete transition to clean household energy can save one-quarter of the healthy life lost to particulate matter pollution exposure in India. *Environmental Research Letters*, *15*, 094096. <https://doi.org/10.1088/1748-9326/ab8e8a>
- Conibear, L., & Knote, C. (2020). *wrfchem-leeds/WRFotron: WRFotron 2.0*. Zenodo. <https://doi.org/10.5281/zenodo.3624087>
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., et al. (2011). The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, *137*(656), 553–597. <https://doi.org/10.1002/qj.828>
- Ding, D., Xing, J., Wang, S., Liu, K., & Hao, J. (2019). Estimated contributions of emissions controls, meteorological factors, population growth, and changes in baseline mortality to reductions in ambient PM_{2.5} and PM_{2.5}-related mortality in China, 2013–2017. *Environmental Health Perspectives*, *127*(6), 1–12. <https://doi.org/10.1289/EHP4157>
- Du, W., Cohen, A., Shen, G., Ru, M., Shen, H., & Tao, S. (2018). Fuel use trends for boiling water in rural China (1992–2012) and environmental health implications: A national cross-sectional study. *Environmental Science & Technology* acs.est.8b02389 <https://doi.org/10.1021/acs.est.8b02389>
- Ebenstein, A., Fan, M., Greenstone, M., He, G., & Zhou, M. (2017). New evidence on the impact of sustained exposure to air pollution on life expectancy from China's Huai River Policy. *Proceedings of the National Academy of Sciences*, *114*(39), 201616784. <https://doi.org/10.1073/pnas.1616784114>
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., et al. (2010). Description and evaluation of the model for ozone and related chemical Tracers, version 4 (MOZART-4). *Geoscientific Model Development*, *3*, 43–67. <https://doi.org/10.5194/gmd-3-43-2010>
- Fang, X., Fan, Q., Liao, Z., Xie, J., Xu, X., & Fan, S. (2019). Spatial-temporal characteristics of the air quality in the Guangdong–Hong Kong–Macau greater Bay area of China during 2015–2017. *Atmospheric Environment*, *210*, 14–34. <https://doi.org/10.1016/j.atmosenv.2019.04.037>
- GBD 2017 Risk Factor Collaborators (2018). Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: A systematic analysis for the global burden of disease stu. *The Lancet*, *392*, 1923–1994. [https://doi.org/10.1016/S0140-6736\(18\)32225-6](https://doi.org/10.1016/S0140-6736(18)32225-6)
- Gillies, S. (2013). *Rasterio: Geospatial raster I/O for Python programmers*. <https://doi.org/10.1016/j.github.com/mapbox/rasterio>
- Global Burden of Disease Study 2017. (2018). *Global burden of disease study 2017 (GBD 2017) population estimates 1950-2017*. Retrieved from <http://ghdx.healthdata.org/record/ihme-data/gbd-2017-population-estimates-1950-2017>
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., et al. (2005). Fully coupled “online” chemistry within the WRF model. *Atmospheric Environment*, *39*(37), 6957–6975. <https://doi.org/10.1016/j.atmosenv.2005.04.027>
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., & Geron, C. (2006). Estimates of global terrestrial isoprene emissions using MEGAN (model of emissions of gases and aerosols from nature). *Atmospheric Chemistry and Physics*, *6*(11), 3181–3210. <https://doi.org/10.5194/acpd-6-107-2006>
- Guo, X., Zhao, L., Chen, D., Jia, Y., Zhao, N., Liu, W., et al. (2018). Air quality improvement and health benefit of PM_{2.5} reduction from the coal cap policy in the Beijing–Tianjin–Hebei (BTH) region, China. *Environmental Science and Pollution Research*, *25*(32), 32709–32720. <https://doi.org/10.1007/s11356-018-3014-y>
- Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., et al. (2020). Array programming with NumPy. *Nature*, *585*, 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- Hedley, A. J., Wong, C. M., Thach, T. Q., Ma, S., Lam, T. H., & Anderson, H. R. (2002). Cardiorespiratory and all-cause mortality after restrictions on sulphur content of fuel in Hong Kong: An intervention study. *Lancet*, *360*(9346), 1646–1652. [https://doi.org/10.1016/S0140-6736\(02\)11612-6](https://doi.org/10.1016/S0140-6736(02)11612-6)
- Hijmans, R., Kapoor, J., Wieczorek, J., International Rice Research Institute, et al. (2016). *University of California, Berkeley Museum of Vertebrate Zoology. Global Administrative Areas (GADM) Boundaries without limits. Version 2.8*. Retrieved from <http://gadm.org/>
- Hodzic, A., & Jimenez, J. L. (2011). Modeling anthropogenically controlled secondary organic aerosols in a megacity: A simplified framework for global and climate models. *Geoscientific Model Development*, *4*(4), 901–917. <https://doi.org/10.5194/gmd-4-901-2011>
- Hodzic, A., & Knote, C. (2014). WRF-chem 3.6.1: MOZART gas-phase chemistry with MOSAIC aerosols (vol. 7). *National Center for Atmospheric Research (NCAR). Atmospheric Chemistry Division (ACD)*.
- Hong Kong Environment Bureau, & Ministry of Environmental Protection of China. (2019). *2025 air quality objectives review - public consultation*. Retrieved from https://www.epd.gov.hk/epd/sites/default/files/epd/english/environmentinhk/air/air_quality_objectives/files/AQOR_consultation_booklet_en.pdf
- Hou, X., Chan, C. K., Dong, G. H., & Yim, S. H. L. (2019). Impacts of transboundary air pollution and local emissions on PM_{2.5} pollution in the Pearl River Delta region of China and the public health, and the policy implications. *Environmental Research Letters*, *14*(3), 034005. <https://doi.org/10.1088/1748-9326/aaf493>
- Hoyer, S., & Hamman, J. J. (2017). xarray: N-D labeled Arrays and Datasets in Python. *Journal of Open Research Software*, *5*, 1–6. <https://doi.org/10.5334/jors.148>
- Huang, D., Xu, J., & Zhang, S. (2012). Valuing the health risks of particulate air pollution in the Pearl River Delta, China. *Environmental Science & Policy*, *15*(1), 38–47. <https://doi.org/10.1016/j.envsci.2011.09.007>
- Huang, J., Pan, X., Guo, X., & Li, G. (2018). Health impact of China's air pollution prevention and control action plan: An analysis of national air quality monitoring and mortality data. *The Lancet Planetary Health*, *2*(7), e313–e323. [https://doi.org/10.1016/S2542-5196\(18\)30141-4](https://doi.org/10.1016/S2542-5196(18)30141-4)
- Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science & Engineering*, *9*, 90–95. <https://doi.org/10.1109/MCSE.2007.55>
- Institute for Health Metrics and Evaluation (2020). *GBD compare data visualization*. Retrieved from vizhub.healthdata.org/gbd-compare
- Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., et al. (2015). HTAP-v2.2: A mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution. *Atmospheric Chemistry and Physics*, *15*(19), 11411–11432. <https://doi.org/10.5194/acp-15-11411-2015>
- Jiang, L., & O'Neill, B. C. (2017). Global urbanization projections for the shared socioeconomic pathways. *Global Environmental Change*, *42*, 193–199. <https://doi.org/10.1016/j.gloenvcha.2015.03.008>

- Jiang, X., Hong, C., Zheng, Y., Zheng, B., Guan, D., Gouldson, A., et al. (2015). To what extent can China's near-term air pollution control policy protect air quality and human health? A case study of the Pearl River Delta region. *Environmental Research Letters*, *10*(10), 104006. <https://doi.org/10.1088/1748-9326/10/10/104006>
- Jin, X., & Holloway, T. (2015). Spatial and temporal variability of ozone sensitivity over China observed from the Ozone Monitoring Instrument. *Journal of Geophysical Research: Atmospheres*, *120*, 7229–7246. <https://doi.org/10.1002/2015JD023250>
- Jordahl, K., Bossche, J. V. den, Wasserman, J., McBride, J., Fleischmann, M., & Gerard, J. (2020). *geopandas/geopandas: v0.7.0 (Version v0.7.0)*. Zenodo. <https://doi.org/10.5281/zenodo.3669853>
- Kim, C., Seow, W. J., Shu, X.-O., Bassig, B. A., Rothman, N., Chen, B. E., et al. (2016). Cooking coal use and all-cause and cause-specific mortality in a prospective cohort study of women in Shanghai, China. *Environmental Health Perspectives*, *124*(9), 1384–1389. <https://doi.org/10.1289/EHP236>
- Kluyver, T., Ragan-Kelley, B., Pérez, F., Granger, B., Bussonnier, M., Frederic, J., et al. (2016). *Jupyter Notebooks – a publishing format for reproducible computational workflows*. <https://doi.org/10.3233/978-1-61499-649-1-87>
- Knote, C., Hodzic, A., & Jimenez, J. L. (2015). The effect of dry and wet deposition of condensable vapors on secondary organic aerosol concentrations over the continental US. *Atmospheric Chemistry and Physics*, *15*(1), 1–18. <https://doi.org/10.5194/acp-15-1-2015>
- Knote, C., Hodzic, A., Jimenez, J. L., Volkamer, R., Orlando, J. J., Baidar, S., et al. (2014). Simulation of semi-explicit mechanisms of SOA formation from glyoxal in aerosol in a 3-D model. *Atmospheric Chemistry and Physics*, *14*(12), 6213–6239. <https://doi.org/10.5194/acp-14-6213-2014>
- Legrand, S. L., Polashenski, C., Letcher, T. W., Creighton, G. A., Peckham, E., & Cetola, J. D. (2019). The AFWA dust emissions scheme for the GOCART aerosol model in WRF-chem. *Geoscientific Model Development*, *12*, 131–166. <https://doi.org/10.5194/gmd-12-131-2019>
- Liang, F., Yang, X., Liu, F., Li, J., Xiao, Q., Chen, J., et al. (2019a). Long-term exposure to ambient fine particulate matter and incidence of diabetes in China: A cohort study. *Environment International*, *126*, 568–575. <https://doi.org/10.1016/j.envint.2019.02.069>
- Liang, L., Cai, Y., Barratt, B., Lyu, B., Chan, Q., Hansell, A. L., et al. (2019b). Associations between daily air quality and hospitalisations for acute exacerbation of chronic obstructive pulmonary disease in Beijing, 2013–17: An ecological analysis. *The Lancet Planetary Health*, *3*(6), e270–e279. [https://doi.org/10.1016/S2542-5196\(19\)30085-3](https://doi.org/10.1016/S2542-5196(19)30085-3)
- Liang, X., Zhang, S., Wu, Y., Xing, J., He, X., Zhang, K. M., et al. (2019c). Air quality and health benefits from fleet electrification in China. *Nature Sustainability*, *2*, 962–971. <https://doi.org/10.1038/s41893-019-0398-8>
- Li, J., Qin, C., Lv, J., Guo, Y., Bian, Z., Zhou, W., et al. (2019b). Solid fuel use and incident COPD in Chinese adults: Findings from the China kadoorie biobank. *Environmental Health Perspectives*, *127*(5), 1–10. <https://doi.org/10.1289/EHP2856>
- Li, J., Zhu, Y., Kelly, J. T., Jang, C. J., Wang, S., Hanna, A., et al. (2019a). Health benefit assessment of PM_{2.5} reduction in Pearl River Delta region of China using a model-monitor data fusion approach. *Journal of Environmental Management*, *233*, 489–498. <https://doi.org/10.1016/j.jenvman.2018.12.060>
- Li, K., Jacob, D. J., Liao, H., Shen, L., Zhang, Q., & Bates, K. H. (2019c). Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China. *Proceedings of the National Academy of Sciences*, *116*(2), 422–427. <https://doi.org/10.1073/pnas.1812168116>
- Li, K., Jacob, D. J., Liao, H., Zhu, J., Shah, V., Shen, L., et al. (2020). A two-pollutant strategy for improving ozone and particulate air quality in China. *Nature Geoscience*, *12*, 906–910. <https://doi.org/10.1038/s41561-019-0464-x>
- Li, M., Liu, H., Geng, G., Hong, C., Liu, F., Song, Y., et al. (2017). Anthropogenic emission inventories in China: A review. *National Science Review*, *4*(6), 834–866. <https://doi.org/10.1093/nsr/nwx150>
- Li, S., Yang, M., Carter, E., Schauer, J. J., Yang, X., Ezzati, M., et al. (2019d). Exposure–response associations of household air pollution and Buccal cell telomere length in women using biomass stoves. *Environmental Health Perspectives*, *127*, 1–11. <https://doi.org/10.1289/EHP4041>
- Li, T., Zhang, Y., Wang, J., Xu, D., Yin, Z., Chen, H., et al. (2018). All-cause mortality risk associated with long-term exposure to ambient PM_{2.5} in China: A cohort study. *The Lancet Public Health*, *3*(10), e470–e477. [https://doi.org/10.1016/S2468-2667\(18\)30144-0](https://doi.org/10.1016/S2468-2667(18)30144-0)
- Li, Y., Henze, D. K., Jack, D., & Kinney, P. L. (2016). The influence of air quality model resolution on health impact assessment for fine particulate matter and its components. *Air Quality, Atmosphere and Health*, *9*(1), 51–68. <https://doi.org/10.1007/s11869-015-0321-z>
- Liu, H., Wang, X. M., Pang, J. M., & He, K. B. (2013). Feasibility and difficulties of China's new air quality standard compliance: PRD case of PM_{2.5} and ozone from 2010 to 2025. *Atmospheric Chemistry and Physics*, *13*(23), 12013–12027. <https://doi.org/10.5194/acp-13-12013-2013>
- Liu, J., Kiesewetter, G., Klimont, Z., Cofala, J., Heyes, C., Schöpp, W., et al. (2019a). Mitigation pathways of air pollution from residential emissions in the Beijing-Tianjin-Hebei region in China. *Environment International*, *125*, 236–244. <https://doi.org/10.1016/j.envint.2018.09.059>
- Liu, M., Huang, X., Song, Y., Tang, J., Cao, J., Zhang, X., et al. (2019b). Ammonia emission control in China would mitigate haze pollution and nitrogen deposition, but worsen acid rain. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(16), 7760–7765. <https://doi.org/10.1073/pnas.1814880116>
- Liu, X., Vitousek, P., Chang, Y., Zhang, W., Matson, P., & Zhang, F. (2016). Evidence for a historic change occurring in China. *Environmental Science and Technology*, *50*(2), 505–506. <https://doi.org/10.1021/acs.est.5b05972>
- Lu, X., Yao, T., Fung, J. C. H., & Lin, C. (2016). Estimation of health and economic costs of air pollution over the Pearl River Delta region in China. *The Science of the Total Environment*, *566–567*, 134–143. <https://doi.org/10.1016/j.scitotenv.2016.05.060>
- Lyu, X. P., Zeng, L. W., Guo, H., Simpson, I. J., Ling, Z. H., Wang, Y., et al. (2017). Evaluation of the effectiveness of air pollution control measures in Hong Kong. *Environmental Pollution*, *220*, 87–94. <https://doi.org/10.1016/j.envpol.2016.09.025>
- Maussion, F., TimoRoth, T., Dusch, M., & Landmann, J. (2019). *fmaussion/salem: v0.2.4 (Version v0.2.4)*. Zenodo. <https://doi.org/10.5281/zenodo.2605265>
- McKinney, W. (2010). Data structures for statistical computing in Python. *Proceedings of the 9th Python in science conference* (pp. 51–56).
- MEIC Research Group, & Tsinghua University. (2019). *Multi-resolution emission inventory for China (MEIC) version 1.3*. Retrieved from <http://www.meicmodel.org/>
- Meng, W., Zhong, Q., Chen, Y., Shen, H., Yun, X., Smith, K. R., et al. (2019). Energy and air pollution benefits of household fuel policies in northern China. *Proceedings of the national academy of sciences*. 201904182. <https://doi.org/10.1073/pnas.1904182116>
- Met Office (2015). *Cartopy: A cartographic python library with a matplotlib interface*. <https://doi.org/scitools.org.uk/cartopy>
- Ministry of Environmental Protection of China (2013). *Air pollution prevention and control action plan*. Government of China.
- Ministry of Environmental Protection of China. (2017). *Beijing-Tianjin-Hebei and surrounding areas 2017 air pollution prevention and control work plan*. Retrieved from http://dqhj.mee.gov.cn/dtxx/201703/t20170323_408663.shtml
- Ministry of Environmental Protection of China. (2018). *Three-year action plan for winning the blue sky defence war*. Retrieved from http://www.gov.cn/zhengce/content/2018-07/03/content_5303158.htm

- Morrison, H., Thompson, G., & Tatarskii, V. (2009). Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: Comparison of one- and two-moment schemes. *Monthly Weather Review*, *137*(3), 991–1007. <https://doi.org/10.1175/2008MWR2556.1>
- National Center for Atmospheric Research. (2016). *ACOM MOZART-4/GEOS-5 global model output*. UCAR. Retrieved from <http://www.acom.ucar.edu/wrf-chem/mozart.shtml>
- National Development and Reform Commission of China. (2017). *Work plan for clean heating in winter in northern China (2017–2021)*. Retrieved from http://www.gov.cn/xinwen/2017-12/20/content_5248855.htm
- Nethery, R. C., & Dominici, F. (2019). Estimating pollution-attributable mortality at the regional and global scales: Challenges in uncertainty estimation and causal inference. *European Heart Journal*, *0*, 1–3. <https://doi.org/10.1093/eurheartj/ehz200>
- Ou, J., Yuan, Z., Zheng, J., Huang, Z., Shao, M., Li, Z., et al. (2016). Ambient ozone control in a photochemically active region: Short-term despike or long-term attainment?. *Environmental Science and Technology*, *50*(11), 5720–5728. <https://doi.org/10.1021/acs.est.6b00345>
- Peters, J., Hedley, A. J., Wong, C. M., Lam, T. H., Ong, S. G., Liu, J., et al. (1996). Effects of an ambient air pollution intervention and environmental tobacco smoke on children's respiratory health in Hong Kong. *International Journal of Epidemiology*, *25*(4), 821–828. <https://doi.org/10.1093/ije/25.4.821>
- Pope, C. A., III, Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Thurston, G. D., et al. (2002). Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *The Journal of the American Medical Association*, *287*(9), 1132–1141. <https://doi.org/10.1001/jama.287.9.1132>
- Pope, C. A., III, Coleman, N., Pond, Z. A., & Burnett, R. T. (2019). Fine particulate air pollution and human mortality: 25+ years of cohort studies. *Environmental Research*, 108924. <https://doi.org/10.1016/j.envres.2019.108924>
- Punger, E. M., & West, J. J. (2013). The effect of grid resolution on estimates of the burden of ozone and fine particulate matter on premature mortality in the United States. *Air Quality, Atmosphere and Health*, *6*(3), 1–22. <https://doi.org/10.1007/s11869-013-0197-8>
- Qi, J., Zheng, B., Li, M., Yu, F., Chen, C., Liu, F., et al. (2017). A high-resolution air pollutants emission inventory in 2013 for the Beijing-Tianjin-Hebei region, China. *Atmospheric Environment*, *170*, 156–168. <https://doi.org/10.1016/j.atmosenv.2017.09.039>
- Qin, Y., Wagner, F., Scovronick, N., Peng, W., Yang, J., Zhu, T., et al. (2017). Air quality, health, and climate implications of China's synthetic natural gas development. *Proceedings of the National Academy of Sciences*, *114*(19), 4887–4892. <https://doi.org/10.1073/pnas.1703167114>
- Reddington, C., Conibear, L., Silver, B., Knote, C., Arnold, S. R., & Spracklen, D. V. (2019). Exploring the impacts of anthropogenic emission sectors on PM_{2.5} and human health in South and East Asia. *Atmospheric Chemistry and Physics*, *19*, 11887–11910. <https://doi.org/10.5194/acp-19-11887-2019>
- Rich, D. Q., Liu, K., Zhang, J., Thurston, S. W., Stevens, T. P., Pan, Y., et al. (2015). Differences in birth weight associated with the 2008 Beijing olympics air pollution reduction: Results from a natural experiment. *Environmental Health Perspectives*, *123*(9), 880–887. <https://doi.org/10.1289/ehp.1408795>
- Rive, N., & Aunan, K. (2010). Quantifying the air quality cobenefits of the clean development mechanism in China. *Environmental Science and Technology*, *44*(11), 4368–4375. <https://doi.org/10.1021/es903546x>
- Saikawa, E., Kim, H., Zhong, M., Avramov, A., Zhao, Y., & Janssens-maenhout, G. (2017). Comparison of emissions inventories of anthropogenic air pollutants and greenhouse gases in China. *Atmospheric Chemistry and Physics*, *17*, 6393–6421. <https://doi.org/10.5194/acp-17-6393-2017>
- Silver, B., Conibear, L., Reddington, C. L., Knote, C., Arnold, S. R., & Spracklen, D. V. (2020a). Pollutant emission reductions deliver decreased PM_{2.5}-caused mortality across China during 2015–2017. *Atmospheric Chemistry and Physics*, *20*, 11683–11695. <https://doi.org/10.5194/acp-20-11683-2020>
- Silver, B., He, X., Arnold, S. R., & Spracklen, D. V. (2020b). The impact of COVID-19 control measures on air quality in China. *Environmental Research Letters*, *15*(8). <https://doi.org/10.1088/1748-9326/aba3a2>
- Silver, B., Reddington, C. L., Arnold, S. R., & Spracklen, D. V. (2018). Substantial changes in air pollution across China during 2015 to 2017. *Environmental Research Letters*, *13*, 114012.
- Skamarock, W. C., Klemp, J. B., Dudhi, J., Gill, D. O., Barker, D. M., Duda, M. G., et al. (2008). A description of the advanced research WRF version 3. *NCAR Tech*, *468+STR*, 113.
- Smith, K. R., & Haigler, E. (2008). Co-benefits of climate mitigation and health protection in energy systems: Scoping methods. *Annual Review of Public Health*, *29*, 11–25. <https://doi.org/10.1146/annurev.publhealth.29.020907.090759>
- Tao, S., Ru, M. Y., Du, W., Zhu, X., Zhong, Q. R., Li, B. G., et al. (2018). Quantifying the rural residential energy transition in China from 1992 to 2012 through a representative national survey. *Nature Energy*, *3*(7), 567–573. <https://doi.org/10.1038/s41560-018-0158-4>
- Tao, Y., Huang, W., Huang, X., Zhong, L., Lu, S. E., Li, Y., et al. (2012). Estimated acute effects of ambient ozone and nitrogen dioxide on mortality in the Pearl River Delta of southern China. *Environmental Health Perspectives*, *120*(3), 393–398. <https://doi.org/10.1289/ehp.1103715>
- Thompson, T. M., Saari, R. K., & Selin, N. E. (2014). Air quality resolution for health impact assessment: Influence of regional characteristics. *Atmospheric Chemistry and Physics*, *14*(2), 969–978. <https://doi.org/10.5194/acp-14-969-2014>
- Thompson, T. M., & Selin, N. E. (2012). Influence of air quality model resolution on uncertainty associated with health impacts. *Atmospheric Chemistry and Physics*, *12*(20), 9753–9762. <https://doi.org/10.5194/acp-12-9753-2012>
- Tian, Y., Liu, H., Liang, T., Xiang, X., Li, M., Juan, J., et al. (2019). Fine particulate air pollution and adult hospital admissions in 200 Chinese cities: A time-series analysis. *International Journal of Epidemiology*, 1142–1151. <https://doi.org/10.1093/ije/dyz106>
- Tsinghua University (2017). *How to Reach Air Quality Standard of PM_{2.5} in China by 2030 – A numerical simulation discussion based on energy and end-of-pipe control scenarios*. Energy Foundation (China Office).
- Turner, M. C., Jerrett, M., Pope, C. A., III, Krewski, D., Gapstur, S. M., Diver, R. W., et al. (2016). Long-term ozone exposure and mortality in a large prospective study. *American Journal of Respiratory and Critical Care Medicine*, *193*(10), 1134–1142. <https://doi.org/10.1164/rccm.201508-1633OC>
- United Nations Environment Programme (2019). *Synergizing action on the environment and climate. Good practise in China and around the globe*.
- Vermeulen, R., Downward, G. S., Zhang, J., Hu, W., Portengen, L., Bassig, B. A., et al. (2019). Constituents of household air pollution and risk of lung cancer among never-smoking women in Xuanwei and Fuyuan, China. *Environmental Health Perspectives*, *127*(9), 097001. <https://doi.org/10.1289/ehp4913>
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., et al. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*, *17*, 261–272.

- Wang, T., Xue, L., Brimblecombe, P., Lam, Y. F., Li, L., & Zhang, L. (2017). Ozone pollution in China: A review of concentrations, meteorological influences, chemical precursors, and effects. *The Science of the Total Environment*, 575, 1582–1596. <https://doi.org/10.1016/j.scitotenv.2016.10.081>
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., et al. (2011). The Fire INventory from NCAR (FINN) – A high resolution global model to estimate the emissions from open burning. *Geoscientific Model Development*, 4, 624–641. <https://doi.org/10.5194/gmd-4-625-2011>
- Wild, O., & Prather, M. J. (2006). Global tropospheric ozone modeling: Quantifying errors due to grid resolution. *Journal of Geophysical Research*, 111(11), 1–14. <https://doi.org/10.1029/2005JD006605>
- Wu, H., Jiang, B., Zhu, P., Geng, X., Liu, Z., Cui, L., et al. (2018). Associations between maternal weekly air pollutant exposures and low birth weight: A distributed lag non-linear model. *Environmental Research Letters*, 13(2). <https://doi.org/10.1088/1748-9326/aaa346>
- Xie, Y., Dai, H., Dong, H., Hanaoka, T., & Masui, T. (2016). Economic impacts from PM2.5 pollution-related health effects in China: A provincial-level analysis. *Environmental Science and Technology*, 50(9), 4836–4843. <https://doi.org/10.1021/acs.est.5b05576>
- Yang, C. J., & Jackson, R. B. (2013). China's synthetic natural gas revolution. *Nature Climate Change*, 3(10), 852–854. <https://doi.org/10.1038/nclimate1988>
- Yan, M., Wilson, A., Bell, M. L., Peng, R. D., Sun, Q., Pu, W., et al. (2019). The shape of the concentration – Response association between fine particulate health impact assessment. *Environmental Health Perspectives*, 127, 1–13. <https://doi.org/10.1289/EHP4464>
- Yin, P., Brauer, M., Cohen, A., Burnett, R. T., Liu, J., Liu, Y., et al. (2017). Long-term fine particulate matter exposure and nonaccidental and cause-specific mortality in a large national cohort of Chinese men. *Environmental Health Perspectives*, 125(11) 117002-1-117002–117011. <https://doi.org/10.1289/EHP1673>
- Yin, P., Brauer, M., Cohen, A. J., Wang, H., Li, J., Burnett, R. T., et al. (2020). The effect of air pollution on deaths, disease burden, and life expectancy across China and its provinces, 1990–2017: An analysis for the global burden of disease study 2017. *The Lancet Planetary Health*, 5196(20), 1–13. [https://doi.org/10.1016/S2542-5196\(20\)30161-3](https://doi.org/10.1016/S2542-5196(20)30161-3)
- Yu, K., Qiu, G., Chan, K. H., Lam, K. B. H., Kurmi, O. P., Bennett, D. A., et al. (2018). Association of solid fuel use with risk of cardiovascular and all-cause mortality in rural China. *JAMA - Journal of the American Medical Association*, 319(13), 1351–1361. <https://doi.org/10.1001/jama.2018.2151>
- Yu, S., Eder, B., Dennis, R., Chu, S.-H., & Schwartz, S. E. (2006). New unbiased symmetric metrics for evaluation of air quality models. *Atmospheric Science Letters*, 7(1), 26–34. <https://doi.org/10.1002/asl.125>
- Yun, X., Shen, G., Shen, H., Meng, W., Chen, Y., Xu, H., et al. (2020). Residential solid fuel emissions contribute significantly to air pollution and associated health impacts in China. *Science Advances*, 6(44), 1–8. <https://doi.org/10.1126/sciadv.aba7621>
- Zaveri, R. A., Easter, R. C., Fast, J. D., & Peters, L. K. (2008). Model for simulating aerosol Interactions and chemistry (MOSAIC). *Journal of Geophysical Research*, 113(D13204), 1–29. <https://doi.org/10.1029/2007JD008782>
- Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., et al. (2019). Drivers of improved PM2.5 air quality in China from 2013 to 2017. *Proceedings of the National Academy of Sciences of the United States of America*, 1–7. <https://doi.org/10.1073/pnas.1907956116>
- Zhang, Z., Wang, J., Chen, L., Chen, X., Sun, G., Zhong, N., et al. (2014). Impact of haze and air pollution-related hazards on hospital admissions in Guangzhou, China. *Environmental Science and Pollution Research*, 21(6), 4236–4244. <https://doi.org/10.1007/s11356-013-2374-6>
- Zhao, B., Zheng, H., Wang, S., Smith, K. R., Lu, X., Aunan, K., et al. (2018). Change in household fuels dominates the decrease in PM2.5 exposure and premature mortality in China in 2005–2015. *Proceedings of the National Academy of Sciences*, 115(49), 12401–12406. <https://doi.org/10.1073/pnas.1812955115>
- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., et al. (2018). Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions. *Atmospheric Chemistry and Physics*, 18(19), 14095–14111. <https://doi.org/10.5194/acp-18-14095-2018>
- Zheng, B., Zhang, Q., Tong, D., Chen, C., Hong, C., Li, M., et al. (2017). Resolution dependence of uncertainties in gridded emission inventories: A case study in Hebei, China. *Atmospheric Chemistry and Physics*, 17(2), 921–933. <https://doi.org/10.5194/acp-17-921-2017>
- Zheng, H., Zhao, B., Wang, S., Wang, T., Ding, D., Chang, X., et al. (2019). Transition in source contributions of PM2.5 exposure and associated premature mortality in China during 2005–2015. *Environment International*, 132, 105111. <https://doi.org/10.1016/j.envint.2019.105111>
- Zheng, Y., Xue, T., Zhang, Q., Geng, G., Tong, D., Li, X., et al. (2017b). Air quality improvements and health benefits from China's clean air action since 2013. *Environmental Research Letters*, 12(11), 114020. <https://doi.org/10.1088/1748-9326/aa8a32>
- Zhuang, J., Dussin, R., Jüling, A., & Rasp, S. (2020). JiaweiZhuang/xESMF: v0.3.0 adding ESMF.LocStream capabilities (version v0.3.0). Zenodo. <https://doi.org/10.5281/zenodo.3700105>
- Zhu, X., Yun, X., Meng, W., Xu, H., Du, W., Shen, G., et al. (2018). Stacked use and transition trends of rural household energy in mainland China. *Environmental Science & Technology*, 53. acs.est.8b04280. research-article <https://doi.org/10.1021/acs.est.8b04280>