



Large-scale controls of soil organic carbon in (sub)tropical soils

Sophie F. von Fromm^{1,2}, Alison M. Hoyt¹, Asmeret Asefaw Berhe³, Keith D. Shepherd⁴, Tor-Gunnar Vågen⁴, Leigh A. Winowicki⁴, Lulseged T. Desta⁵, Jerome E. Tondoh⁶, Andrew M. Sila⁴, Erick K. Towett⁴, Elvis Weullow⁴, Ermias Aynekulu⁴, Johan Six², Susan E. Trumbore¹, and Sebastian Doetterl²
¹Department of Biogeochemical Processes, Max-Planck-Institute for Biogeochemistry, Jena, Germany (sfromm@bgc-jena.mpg.de)

²Department of Environmental System Science, ETH Zurich, Switzerland

³University of California Merced, USA

⁴International Centre for Research in Agroforestry, Nairobi, Kenya

⁵International Centre for Tropical Agriculture, Addis Ababa, Ethiopia

⁶Université Nangui Abrougoua, Abidjan, Côte d'Ivoire

Soil organic carbon (SOC) is a key component of terrestrial ecosystems. Experimental studies have shown that soil texture and geochemistry have a strong effect on carbon stocks. However, those findings primarily rely on data from temperate regions or use model approaches that are often based on limited data from tropical and sub-tropical regions.

Here, we evaluate the controls on soil carbon stocks in Africa, using a dataset of 1,580 samples. These were collected across Sub-Saharan Africa (SSA) within the framework of the Africa Soil Information Service (AfSIS) project, which was built on the well-established Land Degradation Surveillance Framework (LDSF). Samples were taken from two depths (0–20 cm and 20–50 cm) at 46 LDSF sites that were stratified according to Koeppen-Geiger climate zones. The different pH-values, clay content, exchangeable cations and extractable elements across various soils of the different climatic zones (i.e. from arid to humid (sub)tropical) allow us to identify different soil and climate parameters that best explain SOC variance across SSA.

We tested if these SOC predictors differed across climatological conditions, using the ratio of potential evapotranspiration (PET) to mean annual precipitation (MAP) as indicator. For water-limited regions (PET/MAP > 1), the best predictors were climatic variables, likely because of their effect on the quantity of carbon inputs. Geochemistry dominated SOC storage in energy-limited systems (PET/MAP < 1), reflecting its effect on carbon protection. On a continental scale, climate (e.g. PET) is key to predicting SOC content in topsoil, whereas geochemistry, particularly iron-oxhydroxides and aluminum-oxides, is more important in subsoil. Clay content had little influence on SOC at both depths. These findings contribute to an improved understanding of the controls on SOC stocks in tropical and sub-tropical regions.