

Tocilizumab in COVID-19 therapy: who benefits, and how?

Avidan U Neumann, Mehmet Goekkaya, Karim Dorgham, Claudia Traidl-Hoffmann, Guy Gorochov

Angaben zur Veröffentlichung / Publication details:

Neumann, Avidan U, Mehmet Goekkaya, Karim Dorgham, Claudia Traidl-Hoffmann, and Guy Gorochov. 2021. "Tocilizumab in COVID-19 therapy: who benefits, and how?" *The Lancet* 398 (10297): 299-300.
[https://doi.org/10.1016/s0140-6736\(21\)01427-6](https://doi.org/10.1016/s0140-6736(21)01427-6).

Tocilizumab in COVID-19 therapy: who benefits, and how?

The RECOVERY Collaborative Group reported statistically significant improvement in survival of patients with COVID-19 who were receiving tocilizumab interleukin (IL)-6 inhibitor, albeit with very modest reduction of mortality (31% vs 35% with usual care, $p=0.0028$).¹ This result adds to a number of studies with tocilizumab and other IL-6 antagonists, such as sarilumab, which showed only minor, or no, reduction in mortality.² Given that IL-6 is associated with COVID-19 severity and mortality,³ the question arises as to why IL-6 antagonist therapy does not substantially improve survival.

In April, 2021, we showed that IL-6 serum concentrations are indeed associated with COVID-19 severity (appendix); however, a better classification of severity is obtained when IL-6 is combined with other cytokine concentrations.⁴ Moreover, within each respiratory severity group, IL-6 is not significantly associated with mortality (appendix). It is rather distinct combinations of interferon α , interferon β , IL-10, and tumour necrosis

factor α that are better predictors of mortality in different severity groups.⁴

Nevertheless, mortality in the low IL-6 group of patients is significantly lower than in the high IL-6 group of patients (appendix), suggesting that IL-6 inhibitors should be given only to patients with high IL-6. Indeed, a retrospective analysis of tocilizumab therapy as a function of baseline IL-6 concentrations showed a large reduction in mortality (from 36% to 16%) in patients with high-baseline IL-6, but no reduction in mortality in low-baseline IL-6 patients.⁵

In conclusion, clinical trials of IL-6 antagonist therapy, such as RECOVERY¹ and sarilumab COVID-19 global studies,² should consider reanalysis of their results as a function of IL-6 baseline concentrations. More generally, clinical trials of personalised precision medicine, based on cytokine profiling, are needed for optimisation of COVID-19 therapy.

We declare no competing interests.

*Avidan U Neumann,
Mehmet Goekkaya, Karim Dorgham,
Claudia Traidl-Hoffmann, Guy Gorochov
avidan.neumann@uni-a.de

Department of Environmental Medicine, Faculty of Medicine, Universitätsklinikum Augsburg, Augsburg, Germany (AUN, MG, CT-H); Institute of Environmental Medicine, Helmholtz Center Munich, Augsburg, Germany (AUN, CT-H); Environmental Medicine, Technical University Munich, Munich, Germany (MG, CT-H); Sorbonne Université, INSERM, Centre d'Immunologie et des Maladies Infectieuses, Paris, France. (KD, GG); Département d'Immunologie, Assistance Publique Hôpitaux de Paris, Hôpital Pitié-Salpêtrière, Paris, France (GG)

- 1 RECOVERY Collaborative Group. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet* 2021; **397**: 1637–45.
- 2 Lescure FX, Honda H, Fowler RA, et al. Sarilumab in patients admitted to hospital with severe or critical COVID-19: a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Respir Med* 2021; **9**: 522–32.
- 3 DelValle DM, Kim-Schulze S, Huang HH, et al. An inflammatory cytokine signature predicts COVID-19 severity and survival. *Nat Med* 2020; **26**: 1636–43.
- 4 Dorgham K, Quentric P, Goekkaya M, et al. Distinct cytokine profiles associated with COVID-19 severity and mortality. *J Allergy Clin Immunol* 2021; **147**: 2098–107.
- 5 Galvan-Roman JM, Rodriguez-Garcia SC, Roy-Vallejo E, et al. IL-6 serum levels predict severity and response to tocilizumab in COVID-19: an observational study. *J Allergy Clin Immunol* 2020; **147**: 72–80.