

An equilibrated a posteriori error estimator for an Interior Penalty Discontinuous Galerkin approximation of the p-Laplace problem

Ronald H. W. Hoppe, Yuri Iliash

Angaben zur Veröffentlichung / Publication details:

Hoppe, Ronald H. W., and Yuri Iliash. 2021. "An equilibrated a posteriori error estimator for an Interior Penalty Discontinuous Galerkin approximation of the p-Laplace problem." Russian Journal of Numerical Analysis and Mathematical Modelling 36 (6): 313-36. <https://doi.org/10.1515/rnam-2021-0026>.

Nutzungsbedingungen / Terms of use:

licgercopyright

Dieses Dokument wird unter folgenden Bedingungen zur Verfügung gestellt: / This document is made available under the following conditions:

Deutsches Urheberrecht

Weitere Informationen finden Sie unter: / For more information see:

<https://www.uni-augsburg.de/de/organisation/bibliothek/publizieren-zitieren-archivieren/publizieren>



Ronald H. W. Hoppe* and Youri Iliash

An equilibrated a posteriori error estimator for an Interior Penalty Discontinuous Galerkin approximation of the p -Laplace problem

<https://doi.org/10.1515/rnam-2021-0026>

Received June 2, 2021; accepted September 28, 2021

Abstract: We are concerned with an Interior Penalty Discontinuous Galerkin (IPDG) approximation of the p -Laplace equation and an equilibrated a posteriori error estimator. The IPDG method can be derived from a discretization of the associated minimization problem involving appropriately defined reconstruction operators. The equilibrated a posteriori error estimator provides an upper bound for the discretization error in the broken $W^{1,p}$ norm and relies on the construction of an equilibrated flux in terms of a numerical flux function associated with the mixed formulation of the IPDG approximation. The relationship with a residual-type a posteriori error estimator is established as well. Numerical results illustrate the performance of both estimators.

Keywords: Interior Penalty Discontinuous Galerkin method, p -Laplace problem, a posteriori error estimation, equilibration.

MSC 2010: 65N30, 65N15, 65N50

Adaptive finite element methods for the p -Laplace problem and generalizations thereof such as problems with power growth functionals based on either residual-type a posteriori error estimators or on error estimators derived by duality theory have been developed, analyzed, and implemented in [5, 7, 15, 18, 24, 34, 45, 50] (see also the monograph [47]), whereas Discontinuous Galerkin (DG) methods for such problems have been considered in [6, 13, 14, 21, 23, 33, 37, 40, 43, 44, 48] (see also [22] for the $p(x)$ -Laplacian).

On the other hand, equilibrated a posteriori error estimators for adaptive finite element approximations of linear and nonlinear second and fourth order elliptic boundary value problems have been suggested in [8–11, 16, 17, 20, 31, 32]; see also Chapter 12 in [52].

In the present paper, we consider an Interior Penalty Discontinuous Galerkin (IPDG) method for the p -Laplace problem similar to the ones in [5, 13, 14]. The method is the optimality condition for the minimization of an IPDG approximation of the primal energy functional associated with the p -Laplacian. We also consider a two-field formulation of the IPDG approximation which allows to specify a numerical flux function such that the IPDG method falls within the approach taken in [3]. The equilibrated a posteriori error estimator is based on a general approach from [46] which enables to estimate the global discretization error in terms of primal and dual energy functionals. It requires the construction of an equilibrated flux in Brezzi–Douglas–Marini finite element spaces involving the two-field formulation and the numerical flux function.

The paper is organized as follows: In Section 1, we provide some basic notations and auxiliary results. Then, in Section 2 we consider the p -Laplace problem and its associated primal and dual energy functionals. Section 3 is devoted to the IPDG approximation and its related two-field formulation. The equilibrated a posteriori error estimator based on the result from [46] is dealt with in Section 4, whereas Section 5 addresses the construction of an equilibrated flux using Brezzi–Douglas–Marini finite elements. Section 6 provides a comparison with a residual-based a posteriori error estimator for the IPDG approximation. Finally, in Section 7

*Corresponding author: Ronald H. W. Hoppe, Department of Mathematics, University of Augsburg, Germany; Department of Mathematics, University of Houston, USA. E-mail: rohop@math.uh.edu
Youri Iliash, Department of Mathematics, University of Augsburg, Germany

we present a documentation of numerical results for two examples that illustrates the performance of the suggested approach.

1 Basic notations and auxiliary results

We use standard notation from Lebesgue and Sobolev space theory (see, e.g., [49]). In particular, for a bounded domain $\Omega \subset \mathbb{R}^d, d \in \mathbb{N}$, we refer to $L^p(\Omega; \mathbb{R}^d), 1 < p < \infty$, as the Banach space of p th power Lebesgue integrable functions on Ω with norm $\|\cdot\|_{L^p(\Omega; \mathbb{R}^d)}$. In case $d = 1$ we will write $L^p(\Omega)$ instead of $L^p(\Omega; \mathbb{R})$. We denote by $W^{s,p}(\Omega), s \in \mathbb{R}_+, 1 < p < \infty$, the Sobolev spaces with norms $\|\cdot\|_{W^{s,p}(\Omega)}$ and by $W_0^{s,p}(\Omega)$ the closure of $C_0^\infty(\Omega)$ with respect to the norm $\|\cdot\|_{W^{s,p}(\Omega)}$. Functions $u \in W^{1,p}(\Omega)$ have a trace $u|_\Gamma$ on the boundary $\Gamma = \partial\Omega$ with $u|_\Gamma \in W^{1-1/p,p}(\Gamma)$. If $\Gamma = \bar{\Gamma}_D \cup \bar{\Gamma}_N, \Gamma_D \cap \Gamma_N = \emptyset$, the space $W_0^{1-1/p,p}(\Gamma_D)$ denotes the space of functions on Γ_D whose extension by zero to Γ belongs to $W^{1-1/p,p}(\Gamma)$ (for examples of functions that do belong to $W_0^{1-1/p,p}(\Gamma_D)$ and those that do not as well as for further discussion we refer to [35]). For $u_D \in W_0^{1-1/p,p}(\Gamma_D)$ we set

$$W_{u_D, \Gamma_D}^{1,p}(\Omega) := \{v \in W^{1,p}(\Omega) \mid v|_{\Gamma_D} = u_D\}.$$

Further, we define $\underline{\mathbf{H}}^{(p)}(\text{div}, \Omega), 1 < p < \infty$, as the Banach space

$$\underline{\mathbf{H}}^{(p)}(\text{div}, \Omega) = \{\underline{\boldsymbol{\tau}} \in L^p(\Omega)^2 \mid \nabla \cdot \underline{\boldsymbol{\tau}} \in L^p(\Omega)\}$$

with the graph norm

$$\|\underline{\boldsymbol{\tau}}\|_{\underline{\mathbf{H}}^{(p)}(\text{div}, \Omega)} := \left(\|\underline{\boldsymbol{\tau}}\|_{L^p(\Omega; \mathbb{R}^d)}^p + \|\nabla \cdot \underline{\boldsymbol{\tau}}\|_{L^p(\Omega)}^p \right)^{1/p}.$$

We refer to $\underline{\mathbf{H}}_{0, \Gamma_D}^{(p)}(\text{div}, \Omega)$ as the subspace

$$\underline{\mathbf{H}}_{0, \Gamma_D}^{(p)}(\text{div}, \Omega) = \{\underline{\boldsymbol{\tau}} \in \underline{\mathbf{H}}^{(p)}(\text{div}, \Omega) \mid \mathbf{n}_{\Gamma_D} \cdot \underline{\boldsymbol{\tau}} = 0 \text{ on } \Gamma_D\}.$$

For further properties of $\underline{\mathbf{H}}^{(p)}(\text{div}, \Omega)$ we refer to [1].

For later use we recall Young's inequality

$$\prod_{i=1}^2 a_i \leq \frac{\varepsilon}{p} a_1^p + \frac{\varepsilon^{-q/p}}{q} a_2^q \tag{1.1}$$

for $a_i > 0, 1 \leq i \leq 2$, and $1 < p, q < \infty, 1/p + 1/q = 1$, and any $\varepsilon > 0$, as well as the following inequality: Let $w_i \in \mathbb{R}, 1 \leq i \leq 2$, and $0 \leq r < \infty$. Then there exists a constant $C_r > 0$ such that it holds

$$(|w_1| + |w_2|)^r \leq C_r (|w_1|^r + |w_2|^r), \quad C_r^{1/r} C_{1/r} = 1. \tag{1.2}$$

2 The p -Laplace problem and the associated primal and dual energy functionals

Let $\Omega \subset \mathbb{R}^2$ be a bounded polygonal domain with boundary $\Gamma = \partial\Omega, \Gamma = \bar{\Gamma}_D \cup \bar{\Gamma}_N, \Gamma_D \cap \Gamma_N = \emptyset, \Gamma_D \neq \emptyset$, and exterior unit normal vectors $\mathbf{n}_r, \mathbf{n}_{r_D}, \mathbf{n}_{r_N}$. Further, let $1 < p, q < \infty, 1/p + 1/q = 1$, and $f \in L^q(\Omega), u_D \in W_0^{1-1/p,p}(\Gamma_D), u_N \in L^q(\Gamma_N)$. The p -Laplace problem with inhomogeneous Dirichlet and Neumann boundary conditions reads as follows:

$$-\nabla \cdot (|\nabla u|^{p-2} \nabla u) = f \quad \text{in } \Omega \tag{2.1a}$$

$$u = u_D \quad \text{on } \Gamma_D \tag{2.1b}$$

$$\mathbf{n}_{r_N} \cdot (|\nabla u|^{p-2} \nabla u) = u_N \quad \text{on } \Gamma_N. \tag{2.1c}$$

The variational formulation of (2.1) requires the computation of $u \in W_{u_D, \Gamma_D}^{1,p}(\Omega)$ such that for all $v \in W_{0, \Gamma_D}^{1,p}(\Omega)$ it holds

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v \, dx = \ell(v) \tag{2.2a}$$

where the functional $\ell : V \rightarrow \mathbb{R}$ is given by

$$\ell(v) := \int_{\Omega} f v \, dx + \int_{\Gamma_N} u_N v \, ds. \tag{2.2b}$$

It is well known that (2.2) admits a unique solution (see, e.g., [19]). Moreover, (2.2) represents the necessary and sufficient optimality condition for the minimization problem

$$J_P(u) = \inf_{v \in W_{u_D, \Gamma_D}^{1,p}(\Omega)} J_P(v) \tag{2.3a}$$

where the objective functional J_P is given by

$$J_P(v) := \frac{1}{p} \int_{\Omega} |\nabla v|^p \, dx - \int_{\Omega} f v \, dx - \int_{\Gamma_N} u_N v \, ds. \tag{2.3b}$$

The dual problem of (2.3) is given by (see Chapter 4, Section 2.2 in [29]):

$$J_D(\underline{\mathbf{p}}) = \inf_{\underline{\mathbf{q}} \in \underline{\mathbf{H}}^q(\text{div}; \Omega)} J_D(\underline{\mathbf{q}}) \tag{2.4a}$$

subject to the equilibrium conditions

$$-\nabla \cdot \underline{\mathbf{p}} = f \quad \text{in } L^q(\Omega), \quad \mathbf{n}_{\Gamma_N} \cdot \underline{\mathbf{p}} = u_N \quad \text{in } L^q(\Gamma_N) \tag{2.4b}$$

$$\tag{2.4c}$$

where the objective functional J_D is given by

$$J_D(\underline{\mathbf{q}}) := \frac{1}{q} \int_{\Omega} |\underline{\mathbf{q}}|^q \, dx - \int_{\Gamma_D} u_D \mathbf{n}_{\Gamma_D} \cdot \underline{\mathbf{q}} \, ds. \tag{2.4d}$$

3 IPDG approximation of the p -Laplace problem

Let \mathcal{T}_h be a geometrically conforming, locally quasi-uniform, simplicial triangulation of the computational domain Ω which on Γ aligns with Γ_D and Γ_N . Given $D \subset \bar{\Omega}$, we denote by $\mathcal{N}_h(D)$ and $\mathcal{E}_h(D)$ the set of vertices and edges of \mathcal{T}_h in D , and we refer to $P_k(D)$, $k \in \mathbb{N}$, as the set of polynomials of degree $\leq k$ on D . Moreover, h_K , $K \in \mathcal{T}_h$, and h_E , $E \in \mathcal{E}_h$, stand for the diameter of K and the length of E , respectively. We define $h := \min \{h_K \mid K \in \mathcal{T}_h\}$. Due to the local quasi-uniformity of the triangulation there exist constants $0 < c_R \leq C_R$ such that for all $K \in \mathcal{T}_h$ it holds

$$c_R h_K \leq h_E \leq C_R h_K, \quad E \in \mathcal{E}_h(\partial K). \tag{3.1}$$

For two quantities $a, b \in \mathbb{R}$ we will write $a \lesssim b$, if there exists a constant $C > 0$, independent of h , such that $a \leq Cb$.

We will further use the following trace inequality (see, e.g., [25]): For $1 \leq p \leq \infty$ there exists a constant $C_T > 0$, only depending on p , the polynomial degree k , and the local geometry of the triangulation, such that for $v_h \in P_k(K)$ and $K \in \mathcal{T}_h$ it holds

$$\|v_h\|_{L^p(\partial K)} \leq C_T h_K^{-1/p} \|v_h\|_{L^p(K)}. \tag{3.2}$$

For $E \in \mathcal{E}_h(\Omega)$, $E = K_+ \cap K_-$, $K_\pm \in \mathcal{T}_h(\Omega)$, and $v_h \in V_h$, we denote the average and jump of v_h across E by $\{v_h\}_E$ and $[v_h]_E$, i.e.,

$$\{v_h\}_E := \frac{1}{2}(v_h|_{E \cap K_+} + v_h|_{E \cap K_-}), \quad [v_h]_E := v_h|_{E \cap K_+} - v_h|_{E \cap K_-}$$

whereas for $E \in \mathcal{E}_h(\Gamma)$ we set

$$\{v_h\}_E := v_h|_E, \quad [v_h]_E := v_h|_E.$$

The averages $\{\nabla v_h\}_E$, $\{\underline{\boldsymbol{\tau}}_h\}_E$ and jumps $[\nabla v_h]_E$, $[\underline{\boldsymbol{\tau}}_h]_E$ of vector-valued functions ∇v_h and $\underline{\boldsymbol{\tau}}_h$ are defined analogously. For $E \in \mathcal{E}_h(\Omega)$ it holds

$$\int_E u_h v_h \, ds = \int_E (\{u_h\}_E [v_h]_E + [u_h]_E \{v_h\}_E) \, ds. \quad (3.3)$$

We further denote by \mathbf{n}_E , $E \in \mathcal{E}_h(\Omega)$, with $E = K_+ \cap K_-$ the unit normal on E pointing from K_+ to K_- and by \mathbf{n}_E , $E \in \mathcal{E}_h(\Gamma)$, the exterior unit normal on E .

We define the broken $W^{1,p}$ -space $W^{1,p}(\Omega; \mathcal{T}_h)$, $1 < p < \infty$, by

$$W^{1,p}(\Omega; \mathcal{T}_h) := \{v_h \in L^p(\Omega) \mid v_h|_K \in W^{1,p}(K), K \in \mathcal{T}_h\} \quad (3.4)$$

equipped with the norm

$$\|v_h\|_{W^{1,p}(\Omega; \mathcal{T}_h)} := \left(\sum_{K \in \mathcal{T}_h} \|v_h\|_{W^{1,p}(K)}^p \right)^{1/p} \quad (3.5)$$

and the broken \mathbf{H} -space $\mathbf{H}^{(p)}(\text{div}, \Omega; \mathcal{T}_h)$ by

$$\mathbf{H}^{(p)}(\text{div}, \Omega; \mathcal{T}_h) := \{\underline{\mathbf{q}}_h \in L^p(\Omega; \mathbb{R}^2) \mid \underline{\mathbf{q}}_h|_K \in \mathbf{H}^{(p)}(\text{div}, K), K \in \mathcal{T}_h\} \quad (3.6)$$

equipped with the norm

$$\|\underline{\mathbf{q}}_h\|_{\mathbf{H}^{(p)}(\text{div}, \Omega; \mathcal{T}_h)} := \left(\sum_{K \in \mathcal{T}_h} \|\underline{\mathbf{q}}_h\|_{\mathbf{H}^{(p)}(\text{div}, K)}^p \right)^{1/p}. \quad (3.7)$$

We redefine the primal energy functional (2.3b) according to

$$J_P(v) := \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla v|^p \, dx - \int_{\Omega} f v \, dx - \int_{\Gamma_N} u_N v \, ds, \quad v \in W^{1,p}(\Omega; \mathcal{T}_h) \quad (3.8)$$

and note that it reduces to (2.3b) for $v \in W^{1,p}(\Omega)$.

We consider the finite element approximation with the DG spaces

$$V_h := \{v_h : \bar{\Omega} \rightarrow \mathbb{R} \mid v_h|_K \in P_k(K), K \in \mathcal{T}_h\} \quad (3.9a)$$

$$\underline{\mathbf{V}}_h := \{\underline{\mathbf{q}}_h : \bar{\Omega} \rightarrow \mathbb{R}^2 \mid \underline{\mathbf{q}}_h|_K \in P_k(K)^2, K \in \mathcal{T}_h\}. \quad (3.9b)$$

We note that $V_h \subset W^{1,p}(\Omega; \mathcal{T}_h)$. Moreover, for $\underline{\mathbf{q}}_h \in \underline{\mathbf{V}}_h$, we have $(\nabla \cdot \underline{\mathbf{q}}_h)|_K \in P_{k-1}(K)$, $K \in \mathcal{T}_h$, and $\mathbf{n}_E \cdot \underline{\mathbf{q}}_h|_E \in P_k(E)$, $E \in \mathcal{E}_h(\Gamma)$.

For $u_h \in V_h$ we define the broken gradient $\nabla_h u_h$ by means of

$$\nabla_h u_h|_K := \nabla u_h|_K, \quad K \in \mathcal{T}_h. \quad (3.10)$$

Further, let u_D^* be chosen according to

$$u_D^* \in W^{1,p}(\Omega) \text{ such that } u_D^*|_{\Gamma_D} = u_D \text{ and } u_D^*|_{\Gamma_N} = 0. \quad (3.11)$$

Following [13, 23], we define recovery operators $\mathbf{R}_{h,i} : V_h \otimes W^{1,p}(\Omega) \rightarrow \underline{\mathbf{V}}_h$, $1 \leq i \leq 2$, according to

$$\int_{\Omega} \underline{\mathbf{R}}_{h,1}(u) \cdot \underline{\mathbf{q}}_h \, dx = \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E [u]_E \mathbf{n}_E \cdot \{\underline{\mathbf{q}}_h\}_E \, ds, \quad \underline{\mathbf{q}}_h \in \underline{\mathbf{V}}_h \quad (3.12a)$$

$$\int_{\Omega} \underline{\mathbf{R}}_{h,2}(u) \cdot \underline{\mathbf{q}}_h \, dx = \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u \mathbf{n}_E \cdot \underline{\mathbf{q}}_h \, ds, \quad \underline{\mathbf{q}}_h \in \underline{\mathbf{V}}_h. \quad (3.12b)$$

We define the broken DG gradients $\nabla_{DG,i} u_h$, $1 \leq i \leq 2$, as follows:

$$\nabla_{DG,1} u_h := \nabla_h u_h - \underline{\mathbf{R}}_{h,1}(u_h) \quad (3.13a)$$

$$\nabla_{DG,2} u_h := \nabla_{DG,1} u_h + \underline{\mathbf{R}}_{h,2}(u_h^*). \quad (3.13b)$$

The following auxiliary result from [13] will enable us to estimate the L^p norm of $\underline{\mathbf{R}}_{h,1}(u_h) - \underline{\mathbf{R}}_{h,2}(u_h^*)$ for $u_h \in V_h$ (see Lemma A2 in [13]).

Lemma 3.1. *For each $p, q \in (1, \infty)$ such that $1/p + 1/q = 1$ there exists a constant $C_{IS} > 0$, independent of h , such that it holds*

$$\inf_{u_h \in V_h} \sup_{\underline{\mathbf{q}}_h \in \underline{\mathbf{V}}_h} \frac{\int_{\Omega} (\underline{\mathbf{R}}_{h,1}(u_h) - \underline{\mathbf{R}}_{h,2}(u_h^*)) \cdot \underline{\mathbf{q}}_h \, dx}{\|u_h\|_{L^p(\Omega)} \|\underline{\mathbf{q}}_h\|_{L^q(\Omega; \mathbb{R}^2)}} \geq C_{IS}. \quad (3.14)$$

Theorem 3.1. *Under the assumptions of Lemma 3.1 there exists a constant $C_{rec} > 0$, independent of h , such that for $u_h \in V_h$ it holds*

$$\|\underline{\mathbf{R}}_{h,1}(u_h) - \underline{\mathbf{R}}_{h,2}(u_h^*)\|_{L^p(\Omega; \mathbb{R}^2)} \leq C_{rec} \left(\left(\sum_{E \in \mathcal{E}_h(\Omega)} h_E^{-p/q} \int_E |[u_h]_E|^p \, ds \right)^{1/p} + \left(\sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_D|^p \, ds \right)^{1/p} \right) \quad (3.15a)$$

$$\|\underline{\mathbf{R}}_{h,2}(u_h^*)\|_{L^p(\Omega; \mathbb{R}^2)} \leq C_{rec} \left(\sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_D|^p \, ds \right)^{1/p}. \quad (3.15b)$$

Proof. We have

$$\|\underline{\mathbf{R}}_{h,1}(u_h)\|_{L^p(\Omega; \mathbb{R}^2)} = \sup_{\underline{\mathbf{q}} \in L^q(\Omega; \mathbb{R}^2)} \frac{\int_{\Omega} \underline{\mathbf{R}}_{h,1}(u_h) \cdot \underline{\mathbf{q}} \, dx}{\|\underline{\mathbf{q}}\|_{L^q(\Omega; \mathbb{R}^2)}} \geq \sup_{\underline{\mathbf{q}}_h \in \underline{\mathbf{V}}_h} \frac{\int_{\Omega} \underline{\mathbf{R}}_{h,1}(u_h) \cdot \underline{\mathbf{q}}_h \, dx}{\|\underline{\mathbf{q}}_h\|_{L^q(\Omega; \mathbb{R}^2)}}. \quad (3.16)$$

The inf-sup property (3.14) implies

$$\|\underline{\mathbf{R}}_{h,1}(u_h)\|_{L^p(\Omega; \mathbb{R}^2)} \leq C_{IS}^{-1} \sup_{\underline{\mathbf{q}}_h \in \underline{\mathbf{V}}_h} \frac{\int_{\Omega} \underline{\mathbf{R}}_{h,1}(u_h) \cdot \underline{\mathbf{q}}_h \, dx}{\|\underline{\mathbf{q}}_h\|_{L^q(\Omega; \mathbb{R}^2)}}. \quad (3.17)$$

Now, observing (3.12), setting $E_1 := E_+$, $E_2 := E_-$ for $E \in \mathcal{E}_h(\Omega)$ and using (1.2), (3.1), the trace inequality (3.2) as well as Hölder's inequality and the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} \int_{\Omega} (\underline{\mathbf{R}}_{h,1}(u_h) - \underline{\mathbf{R}}_{h,2}(u_h^*)) \cdot \underline{\mathbf{q}}_h \, dx &\leq \sum_{E \in \mathcal{E}_h(\Omega)} \int_E h_E^{-1/q} |[u_h]_E| h_E^{1/q} |\{\underline{\mathbf{q}}_h\}_E| \, ds \\ &\quad + \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E h_E^{-1/q} |u_h - u_D| h_E^{1/q} |\{\underline{\mathbf{q}}_h\}_E| \, ds \\ &\leq \frac{1}{2} \sum_{E \in \mathcal{E}_h(\Omega)} \left(\int_E h_E^{-p/q} |[u_h]_E|^p \, ds \right)^{1/p} \left(\int_E h_E |\underline{\mathbf{q}}_h|_{E_+} + \underline{\mathbf{q}}_h|_{E_-}|^q \, ds \right)^{1/q} \\ &\quad + \sum_{E \in \mathcal{E}_h(\Gamma_D)} \left(\int_E h_E^{-p/q} |u_h - u_D|^p \, ds \right)^{1/p} \left(\int_E h_E |\underline{\mathbf{q}}_h|^q \, ds \right)^{1/q} \end{aligned}$$

$$\begin{aligned}
&\leq \frac{C_q^{1/q}}{2} \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \left(\int_E h_E^{-p/q} |[u_h]_E|^p ds \right)^{1/p} \left(\int_E h_E (|\underline{\mathbf{q}}_h|_{E^+}^q + |\underline{\mathbf{q}}_h|_{E^-}^q) ds \right)^{1/q} \\
&\quad + \sum_{E \in \mathcal{E}_h(\Gamma_D)} \left(\int_E h_E^{-p/q} |u_h - u_D|^p ds \right)^{1/p} \left(\int_E h_E |\underline{\mathbf{q}}_h|^q ds \right)^{1/q} \\
&\leq \frac{C_q^{1/q}}{2} \left(\sum_{E \in \mathcal{E}_h(\Omega)} \int_E h_E^{-p/q} |[u_h]_E|^p ds \right)^{1/p} \left(\sum_{E \in \mathcal{E}_h(\Omega)} \int_E h_E \sum_{i=1}^2 |\underline{\mathbf{q}}_h|_{E_i}|^q ds \right)^{1/q} \\
&\quad + \left(\sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E h_E^{-p/q} |u_h - u_D|^p ds \right)^{1/p} \left(\sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E h_E |\underline{\mathbf{q}}_h|^q ds \right)^{1/q} \\
&\leq C_R^{1/q} C_q^{1/q} \left(\sum_{E \in \mathcal{E}_h(\Omega)} \int_E h_E^{-p/q} |[u_h]_E|^p ds \right)^{1/p} \left(\sum_{K \in \mathcal{T}_h} \int_K h_K |\underline{\mathbf{q}}_h|_{\partial K}|^q ds \right)^{1/q} \\
&\quad + C_R^{1/q} \left(\sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E h_E^{-p/q} |u_h - u_D|^p ds \right)^{1/p} \left(\sum_{K \in \mathcal{T}_h} \int_K h_K |\underline{\mathbf{q}}_h|_{\partial K}|^q ds \right)^{1/q} \\
&\leq C_R^{1/q} C_q^{1/q} C_T \left(\sum_{E \in \mathcal{E}_h(\Omega)} \int_E h_E^{-p/q} |[u_h]_E|^p ds \right)^{1/p} \|\underline{\mathbf{q}}_h\|_{L^q(\Omega; \mathbb{R}^2)} \\
&\quad + C_R^{1/q} C_T \left(\sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E h_E^{-p/q} |u_h - u_D|^p ds \right)^{1/p} \|\underline{\mathbf{q}}_h\|_{L^q(\Omega; \mathbb{R}^2)}. \tag{3.18}
\end{aligned}$$

Using (3.18) in (3.17) gives (3.15a). The proof of (3.15b) follows along the same lines. \square

We refer to $\Pi_k^{(p)}$ as a contractive L^p projection of $L^p(\Omega)$ onto $\{v_h \in L^p(\Omega) \mid v_h|_K \in P_k(K), K \in \mathcal{T}_h\}$, which can be defined elementwise by

$$\begin{aligned}
\int_{\Omega} \Pi_k^{(p)}(v) v_h dx &= \sum_{K \in \mathcal{T}_h} \int_K \Pi_k^{(p)}(v) v_h dx, \quad v \in L^p(\Omega) \\
\int_K \Pi_k^{(p)}(v) p_k dx &= \int_K v p_k dx, \quad p_k \in P_k(K), K \in \mathcal{T}_h.
\end{aligned} \tag{3.19}$$

We note that $\Pi_k^{(q)} := (\Pi_k^{(p)})^*$ is a contractive L^q projection of $L^q(\Omega)$ onto $\{v_h \in L^q(\Omega) \mid v_h|_K \in P_k(K), K \in \mathcal{T}_h\}$ (see, e.g., [2]). We further refer to $\underline{\Pi}_k^{(p)}$ as a contractive L^p projection of $L^p(\Omega; \mathbb{R}^2)$ onto \underline{V}_h . We also denote by $\Pi_k^{(p)}$ a contractive L^p projection of $L^p(\Gamma')$, $\Gamma' = \Gamma_D$ or $\Gamma' = \Gamma_N$, onto $\{v_h \in L^p(\Gamma') \mid v_h|_E \in P_k(E), E \in \mathcal{E}_h(\Gamma')\}$.

We define approximations f_h , $u_{h,D}$, and $u_{h,N}$ of f , u_D , and u_N such that

$$\begin{aligned}
f_h|_K &\in P_{k-1}(K), \quad \int_K (f - f_h) p_0 dx = 0, \quad p_0 \in P_0(K), K \in \mathcal{T}_h \\
u_{h,D}|_E &\in P_k(E), \quad \int_E (u_D - u_{h,D}) p_0 ds = 0, \quad p_0 \in P_0(E), E \in \mathcal{E}_h(\Gamma_D) \\
u_{h,N}|_E &\in P_k(E), \quad \int_E (u_N - u_{h,N}) p_0 ds = 0, \quad p_0 \in P_0(E), E \in \mathcal{E}_h(\Gamma_N).
\end{aligned} \tag{3.20}$$

Remark 3.1. We may choose $f_h|_K$ and $u_{h,N}|_E$ as contractive L^q projections onto $P_{k-1}(K)$ and $P_k(E)$ and $u_{h,D}|_E$ as a contractive L^p projection onto $P_k(E)$.

We consider the discrete minimization problem

$$J_{h,p}(u_h) = \inf_{v_h \in \underline{V}_h} J_{h,p}(v_h) \tag{3.21a}$$

where the objective functional $J_{h,p}$ is given by

$$J_{h,p}(v_h) := \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} v_h|^p dx - \int_{\Omega} f_h v_h dx - \int_{\Gamma_N} u_{h,N} v_h ds + \frac{\alpha}{p} \left(\sum_{E \in \mathcal{E}_h(\Omega)} h_E^{-p/q} \int_E |[v_h]_E|^p ds + \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |v_h - u_D|^p ds \right) \quad (3.21b)$$

and $\alpha > 0$ is a penalization parameter. The existence and uniqueness of a solution of (3.21) follows by standard arguments from the calculus of variations. The necessary and sufficient optimality condition gives rise to a discrete variational equation which represents the IPDG approximation of the p -Laplace problem (2.1a)–(2.1c): Find $u_h \in V_h$ such that for all $v_h \in V_h$ it holds

$$a_h^{DG}(u_h, v_h) = \ell_h(v_h) \quad (3.22)$$

where, observing $\underline{\Pi}_k^{(p)}(\nabla_{DG,1} v_h) = \nabla_{DG,1} v_h$, the semilinear IPDG form $a_h^{DG}(\cdot, \cdot) : V_h \times V_h \rightarrow \mathbb{R}$ is given by

$$\begin{aligned} a_h^{DG}(u_h, v_h) &:= \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \underline{\Pi}_k^{(p)}(\nabla_{DG,1} v_h) dx \\ &\quad + \alpha \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} h_E^{-p/q} \int_E |[u_h - u_D^*]_E|^{p-2} [u_h - u_D^*]_E [v_h]_E ds \\ &= \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)}(|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h) \cdot \nabla_{DG,1} v_h dx \\ &\quad + \alpha \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} h_E^{-p/q} \int_E |[u_h - u_D^*]_E|^{p-2} [u_h - u_D^*]_E [v_h]_E ds \end{aligned} \quad (3.23)$$

and $\ell_h(\cdot) : V_h \rightarrow \mathbb{R}$ stands for the linear functional

$$\ell_h(v_h) := \sum_{K \in \mathcal{T}_h} \int_K f_h v_h dx + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E u_{h,N} v_h ds. \quad (3.24)$$

Lemma 3.2. *The IPDG approximation (3.22) is consistent with the p -Laplace problem (2.1a)–(2.1c) in the sense that if $f = f_h$ and $u_N = u_{h,N}$ in (2.2b) and u satisfies (2.1a)–(2.1c) pointwise almost everywhere, then for all $v_h \in V_h$ it holds*

$$a_h^{DG}(u, v_h) = \ell_h(v_h). \quad (3.25)$$

Proof. Since $[u]_E = [u_D^*]_E = 0$, $E \in \mathcal{E}_h(\Omega)$, and $(u - u_D^*)|_E = (u - u_D)|_E = 0$, $E \in \mathcal{E}_h(\Gamma_D)$, we have $\underline{\mathbf{R}}_{h,1}(u) - \underline{\mathbf{R}}_{h,2}(u_D^*) = 0$. It follows that

$$\begin{aligned} a_h^{DG}(u, v_h) &= \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)}(|\nabla u|^{p-2} \nabla u) \cdot (\nabla v_h - \underline{\mathbf{R}}_{h,1}(v_h)) dx \\ &= \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)}(|\nabla u|^{p-2} \nabla u) \cdot \nabla v_h dx - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E \mathbf{n}_E \cdot \{\underline{\Pi}_k^{(q)}(|\nabla u|^{p-2} \nabla u)\}_E [v_h]_E ds \\ &= \sum_{K \in \mathcal{T}_h} \int_K |\nabla u|^{p-2} \nabla u \cdot \underline{\Pi}_k^{(p)}(\nabla v_h) dx - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E \underline{\Pi}_k^{(q)}(\mathbf{n}_E \cdot \{|\nabla u|^{p-2} \nabla u\}_E) [v_h]_E ds \\ &= \sum_{K \in \mathcal{T}_h} \int_K |\nabla u|^{p-2} \nabla u \cdot \underline{\Pi}_k^{(p)}(\nabla v_h) dx - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E \mathbf{n}_E \cdot \{|\nabla u|^{p-2} \nabla u\}_E \underline{\Pi}_k^{(p)}([v_h]_E) ds \\ &= \sum_{K \in \mathcal{T}_h} \int_K |\nabla u|^{p-2} \nabla u \cdot \nabla v_h dx - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E \mathbf{n}_E \cdot \{|\nabla u|^{p-2} \nabla u\}_E [v_h]_E ds. \end{aligned} \quad (3.26)$$

An application of Green's formula gives

$$\int_K |\nabla u|^{p-2} \nabla u \cdot \nabla v_h dx = - \int_K (\nabla \cdot (|\nabla u|^{p-2} \nabla u)) v_h dx + \int_{\partial K} \mathbf{n}_{\partial K} \cdot (|\nabla u|^{p-2} \nabla u) v_h ds.$$

Summing over all $K \in \mathcal{T}_h$ and observing (3.3) yields

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} \int_K |\nabla u|^{p-2} \nabla u \cdot \nabla v_h \, dx &= \sum_{K \in \mathcal{T}_h} \int_K (-\nabla \cdot (|\nabla u|^{p-2} \nabla u)) v_h \, dx + \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E \mathbf{n}_E \cdot \{|\nabla u|^{p-2} \nabla u\}_E [v_h]_E \, ds \\ &\quad + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E \mathbf{n}_E \cdot (|\nabla u|^{p-2} \nabla u) v_h \, ds. \end{aligned} \quad (3.27)$$

Using (3.27) in (3.26) and observing (2.1a)–(2.1c) results in

$$a_h^{DG}(u, v_h) = \sum_{K \in \mathcal{T}_h} \int_K (-\nabla \cdot (|\nabla u|^{p-2} \nabla u)) v_h \, dx + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E \mathbf{n}_E \cdot |\nabla u|^{p-2} \nabla u v_h \, ds = \int_{\Omega} f_h v_h \, dx + \int_{\Gamma_N} u_{h,N} v_h \, ds$$

which is the assertion. \square

Observing (3.13a) and (3.12a), for the first term on the right-hand side of (3.23) we find

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)} (|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h) \cdot \nabla_{DG,1} v_h \, dx &= \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)} (|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h) \cdot \nabla v_h \, dx \\ &\quad - \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)} (|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h) \cdot \underline{\mathbf{R}}_{h,1}(v_h) \, dx \\ &= \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \underline{\Pi}_k^{(p)}(\nabla v_h) \, dx \\ &\quad - \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)} (|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h) \cdot \underline{\mathbf{R}}_{h,1}(v_h) \, dx \\ &= \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \nabla v_h \, dx \\ &\quad - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E [v_h]_E \mathbf{n}_E \cdot \{\underline{\Pi}_k^{(q)} (|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h)\}_E \, ds \\ &= \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \nabla v_h \, dx \\ &\quad - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E [v_h]_E \Pi_k(\mathbf{n}_E \cdot \{|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h\}_E) \, ds \\ &= \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \nabla v_h \, dx - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E [v_h]_E \mathbf{n}_E \cdot \{|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h\}_E \, ds \end{aligned}$$

and hence, we obtain

$$\begin{aligned} a_h^{DG}(u_h, v_h) &= \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \nabla v_h \, dx - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E [v_h]_E \mathbf{n}_E \cdot \{|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h\}_E \, ds \\ &\quad + \alpha \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} h_E^{-p/q} \int_E |[u_h - u_D^*]_E|^{p-2} [u_h - u_D^*]_E [v_h]_E \, ds. \end{aligned} \quad (3.28)$$

Remark 3.2. In case $p = 2$, i.e., for the Poisson problem, we have

$$\begin{aligned} a_h^{DG}(u_h, v_h) &= \sum_{K \in \mathcal{T}_h} \int_K \nabla u_h \cdot \nabla v_h \, dx - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E [v_h]_E \mathbf{n}_E \cdot \{\nabla u_h\}_E \, ds - \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} \int_E [u_h - u_D^*]_E \mathbf{n}_E \cdot \{\nabla v_h\}_E \, ds \\ &\quad + \alpha \sum_{E \in \mathcal{E}_h(\Omega \cup \Gamma_D)} h_E^{-p/q} \int_E |[u_h - u_D^*]_E|^{p-2} [u_h - u_D^*]_E [v_h]_E \, ds \\ &\quad + \sum_{K \in \mathcal{T}_h} \int_K (\underline{\mathbf{R}}_{h,1}(u_h) - \underline{\mathbf{R}}_{h,2}(u_D^*)) \cdot \underline{\mathbf{R}}_{h,1}(v_h) \, dx \end{aligned}$$

which, except for the last term, coincides with the IPDG approximation for the Poisson problem from [3].

Remark 3.3. We note that $u_h \notin W^{1,p}(\Omega)$, but a conforming finite element function $u_h^c \in V_h^c := V_h \cap W^{1,p}(\Omega)$ can be obtained from $u_h \in V_h$ by postprocessing in the following way (see [38]): Let \mathcal{N}^L be the set of Lagrangian nodal points for the elements in V_h^c and let κ_i be the number of triangles that share the nodal point $x_i \in \mathcal{N}^L$. We have $\kappa_i = 1$, if x_i is contained in the interior of an element, while $\kappa_i > 1$, if $x_i \in \mathcal{N}^L \cap \mathcal{E}_h(\Omega)$. The multiplicity κ_i is bounded, since the triangulation is locally quasi-uniform. Denoting by $\mathcal{N}(x_i)$ the set $\mathcal{N}(x_i) := \bigcup\{K \in \mathcal{T}_h(\Omega) \mid x_i \in K\}$, the associated conforming element is defined by its nodal values

$$u_h^c(x_i) := \begin{cases} \frac{1}{\kappa_i} \sum_{K \in \mathcal{N}(x_i)} u_h|_K(x_i), & x_i \in \Omega \cup \Gamma_N \\ u_{h,D}, & x_i \in \Gamma. \end{cases} \tag{3.29}$$

By a generalization of Theorem 2.2 in [38] to the case $p \neq 2$ there exists a constant $C_c > 0$, only depending on the local geometry of the triangulation, such that

$$\|u_h - u_h^c\|_{W^{1,p}(\Omega; \mathcal{T}_h)}^p \leq C_c \left(\sum_{E \in \mathcal{E}_h(\Omega)} h_E^{-p/q} \int_E |[u_h]_E|^p ds + \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_{h,D}|^p ds \right). \tag{3.30}$$

Next, we consider a two-field formulation of the IPDG approximation (3.22). We set

$$\mathbf{p}_h := \Pi_k^{(q)}(|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h) \tag{3.31a}$$

$$-\nabla_h \cdot \mathbf{p}_h = f_h. \tag{3.31b}$$

We consider (3.31a) elementwise for each $K \in \mathcal{T}_h$, multiply by $\mathbf{q}_h|_K, \mathbf{q}_h \in \mathbf{V}_h$, integrate over K , and finally sum over all $K \in \mathcal{T}_h$. Observing $\Pi_k^{(q)}(\mathbf{q}_h) = \mathbf{q}_h$, we thus obtain

$$\sum_{K \in \mathcal{T}_h} \int_K \mathbf{p}_h \cdot \mathbf{q}_h dx = \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \mathbf{q}_h dx. \tag{3.32a}$$

Likewise, we consider (3.31b) elementwise for each $K \in \mathcal{T}_h$, multiply by $v_h|_K, v_h \in V_h$, integrate over K , and finally sum over all $K \in \mathcal{T}_h$. An elementwise application of Green's formula gives

$$-\sum_{K \in \mathcal{T}_h} \int_K \nabla \cdot \mathbf{p}_h v_h dx = \sum_{K \in \mathcal{T}_h} \int_K \mathbf{p}_h \cdot \nabla v_h dx - \sum_{K \in \mathcal{T}_h} \int_{\partial K} \mathbf{n}_{\partial K} \cdot \mathbf{p}_h|_{\partial K} v_h ds = \sum_{K \in \mathcal{T}_h} \int_K f_h v_h dx. \tag{3.32b}$$

We replace $\mathbf{p}_h|_{\partial K}$ in (3.32b) by a numerical flux function $\hat{\mathbf{p}}_{\partial K}$. We thus obtain the following system of discrete variational equations: Find $(u_h, \mathbf{p}_h) \in V_h \times \mathbf{V}_h$ such that for all $(v_h, \mathbf{q}_h) \in V_h \times \mathbf{V}_h$ it holds

$$\sum_{K \in \mathcal{T}_h} \int_K \mathbf{p}_h \cdot \mathbf{q}_h dx = \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \mathbf{q}_h dx \tag{3.33a}$$

$$\sum_{K \in \mathcal{T}_h} \int_K \mathbf{p}_h \cdot \nabla v_h dx - \sum_{K \in \mathcal{T}_h} \int_{\partial K} \mathbf{n}_{\partial K} \cdot \hat{\mathbf{p}}_{\partial K} v_h ds = \sum_{K \in \mathcal{T}_h} \int_K f_h v_h dx. \tag{3.33b}$$

In particular, for the two-field formulation of the IPDG approximation (3.22) the numerical flux function $\hat{\mathbf{p}}_{\partial K}$ is chosen as follows:

$$\hat{\mathbf{p}}_{\partial K}|_E := \begin{cases} \{\mathbf{z}_h\}_E - \alpha h_E^{-p/q} |[u_h]_E|^{p-2} [u_h]_E \mathbf{n}_E, & E \in \mathcal{E}_h(\Omega) \\ \mathbf{z}_h - \alpha h_E^{-p/q} |u_h - u_D|^{p-2} (u_h - u_D) \mathbf{n}_E, & E \in \mathcal{E}_h(\Gamma_D) \\ u_N \mathbf{n}_E, & E \in \mathcal{E}_h(\Gamma_N) \end{cases} \tag{3.34}$$

where $\mathbf{z}_h := |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h$.

Theorem 3.2. *The two-field formulation (3.33) is equivalent with (3.22). In particular, if $u_h \in V_h$ is the solution of (3.22), there exists $\mathbf{p}_h \in \mathbf{V}_h$ such that the pair $(\mathbf{p}_h, u_h) \in \mathbf{V}_h \times V_h$ satisfies (3.33). Conversely, if the pair $(\mathbf{p}_h, u_h) \in \mathbf{V}_h \times V_h$ satisfies (3.33), then $u_h \in V_h$ solves (3.22).*

Proof. Let $u_h \in V_h$ be the solution of (3.22). We define $\mathbf{p}_h \in \mathbf{V}_h$ by means of (3.33a). Then, choosing $\mathbf{q}_h = \nabla v_h$ in (3.33a) and observing (3.28) and (3.34) yields (3.33b). Conversely, if the pair $(\mathbf{p}_h, u_h) \in \mathbf{V}_h \times V_h$ satisfies (3.33), we choose $\mathbf{q}_h = \nabla v_h$ in (3.33a) and insert (3.33a) into (3.33b). Taking (3.34) into account this shows that $u_h \in V_h$ satisfies (3.22). \square

4 An a posteriori error estimator for the global discretization error

Given reflexive Banach spaces V, Q with norms $\|\cdot\|_V, \|\cdot\|_Q$, convex and coercive objective functionals $C : V \rightarrow \mathbb{R}, D : Q \rightarrow \mathbb{R}$, and a bounded linear operator $\Lambda : V \rightarrow Q$, we consider the minimization problem

$$\inf_{u \in V} J(u) \tag{4.1}$$

for the objective functional

$$J(u) := C(u) + D(\Lambda u). \tag{4.2}$$

An abstract approach to the a posteriori error control for (4.1) has been provided in [46]. The a posteriori error control relies on the dual formulation of (4.1):

$$\sup_{q \in Q} J^*(q) \quad \text{or} \quad \inf_{q \in Q} (-J^*(q)) \tag{4.3}$$

in terms of the Fenchel conjugate J^* of J as given by

$$J^*(q) = -C^*(-\Lambda^* q) - D^*(q) \tag{4.4}$$

where C^* and D^* are the Fenchel conjugates of C and D and Λ^* stands for the adjoint of Λ .

Given some approximation $u_h \in V$ of the minimizer u of (4.1), the a posteriori error estimate from [46] states that for any admissible function $q \in Q$ it holds

$$\|u - u_h\|_V^2 \leq C(u_h) + C^*(-\Lambda^* q) + D(\Lambda u_h) + D^*(q). \tag{4.5}$$

Now, let $u_h^c \in V_h^c \subset W^{1,p}(\Omega)$ be the conforming finite element function obtained from the solution $u_h \in V_h$ of (3.22) by postprocessing according to Remark 3.3. Then it holds

$$\|u - u_h\|_{W^{1,p}(\Omega; \mathcal{T}_h)}^2 \leq 2 \left(\|u - u_h^c\|_{W^{1,p}(\Omega)}^2 + \|u_h - u_h^c\|_{W^{1,p}(\Omega; \mathcal{T}_h)}^2 \right). \tag{4.6}$$

In order to estimate the first term on the right-hand side of (4.6) we apply (4.5) with $V = W^{1,p}(\Omega), Q := L^p(\Omega; \mathbb{R}^2), \Lambda = \nabla$, and

$$C(u_h^c) := - \int_{\Omega} f u_h^c \, dx - \int_{\Gamma_N} u_N u_h^c \, ds \tag{4.7a}$$

$$D(\nabla u_h^c) := \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h^c|^p \, dx + I_{K_1}(u_h^c) \tag{4.7b}$$

where I_{K_1} is the indicator function of the closed convex set

$$K_1 := \{v \in W^{1,p}(\Omega) \mid v = u_D \text{ on } \Gamma_D\}. \tag{4.7c}$$

We obtain

$$C^*(-\Lambda^* \underline{q}) := I_{K_2}(\underline{q}), \quad \underline{q} \in \underline{H}^{(q)}(\text{div}, \Omega) \tag{4.8a}$$

$$D^*(\underline{q}) := \frac{1}{q} \int_{\Omega} |\underline{q}|^q \, dx - \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_D \mathbf{n}_E \cdot \underline{q} \, ds, \quad \underline{q} \in \underline{H}^{(q)}(\text{div}, \Omega) \tag{4.8b}$$

where I_{K_2} is the indicator function of the closed convex set

$$K_2 := \{\underline{q} \in \underline{H}^{(q)}(\text{div}; \Omega) \mid -\nabla \cdot \underline{q} = f \text{ in } \Omega, \mathbf{n}_{\Gamma_N} \cdot \underline{q} = u_N \text{ on } \Gamma_N\}. \tag{4.8c}$$

We call $\underline{p}_h^{\text{eq}} \in \underline{V}_h$ an equilibrated flux, if

$$\underline{p}_h^{\text{eq}} \in \underline{H}^{(q)}(\text{div}; \Omega) \tag{4.9a}$$

and $\underline{\mathbf{p}}_h^{\text{eq}}$ satisfies the equilibrium conditions

$$-\nabla \cdot \underline{\mathbf{p}}_h^{\text{eq}} = f_h \quad \text{in } \Omega \tag{4.9b}$$

$$\mathbf{n}_{\Gamma_N} \cdot \underline{\mathbf{p}}_h^{\text{eq}} = u_{h,N} \quad \text{on } \Gamma_N. \tag{4.9c}$$

Moreover, we choose $\underline{\mathbf{p}}_c \in \{\underline{\mathbf{q}} \in \mathbf{H}_{0,\Gamma_D}^{(q)}(\text{div}, \Omega) \mid \mathbf{n}_{\Gamma_N} \cdot \underline{\mathbf{q}} \in L^q(\Gamma_N)\}$ such that

$$-\nabla \cdot \underline{\mathbf{p}}_c = f - f_h \text{ in } \Omega, \quad \mathbf{n}_{\Gamma_N} \cdot \underline{\mathbf{p}}_c = u_N - u_{h,N} \text{ on } \Gamma_N. \tag{4.10}$$

It follows that $\underline{\mathbf{p}}_h^{\text{eq}} + \underline{\mathbf{p}}_c \in K_2$, i.e., $I_{K_2}(\underline{\mathbf{p}}_h^{\text{eq}} + \underline{\mathbf{p}}_c) = 0$, and hence, (4.5) reads as follows

$$\|u - u_h^c\|_{W^{1,p}(\Omega)}^2 \leq J_P(u_h^c) + I_{K_1}(u_h^c) + J_D(\underline{\mathbf{p}}_h^{\text{eq}} + \underline{\mathbf{p}}_c). \tag{4.11}$$

In view of (2.4d) and $\mathbf{n}_{\Gamma_D} \cdot \underline{\mathbf{p}}_c = 0$ on Γ_D we have

$$J_D(\underline{\mathbf{p}}_h^{\text{eq}} + \underline{\mathbf{p}}_c) = \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}} + \underline{\mathbf{p}}_c|^q dx - \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_D \mathbf{n}_{\Gamma_D} \cdot \underline{\mathbf{p}}_h^{\text{eq}} ds. \tag{4.12}$$

Using (1.2), we find

$$\frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}} + \underline{\mathbf{p}}_c|^q dx \leq \frac{1}{q} C_q \left(\sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}}|^q dx + \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_c|^q dx \right). \tag{4.13}$$

In order to estimate the second term on the right-hand side of (4.13) we use the Poincaré–Friedrichs inequalities

$$\|v - |K|^{-1} \int_K v dx\|_{L^p(K)} \leq C_{PF}^{(1)}(p) h_K \|\nabla v\|_{L^p(K)}, \quad v \in W^{1,p}(K), K \in \mathcal{T}_h \tag{4.14a}$$

$$\|v - |E|^{-1} \int_E v ds\|_{L^p(E)} \leq C_{PF}^{(2)}(p) h_E \|\nabla v\|_{L^p(E)}, \quad v \in W^{1,p}(E), E \in \mathcal{E}_h(\Gamma_N) \tag{4.14b}$$

where $C_{PF}^{(i)}(p)$, $1 \leq i \leq 2$, are positive constants depending only on p (see, e.g., [28]).

Lemma 4.1. *Suppose that the following regularity assumption is satisfied: For $\{\underline{\boldsymbol{\tau}} \in \mathbf{H}_{0,\Gamma_D}^{(p)}(\text{div}, \Omega) \mid \mathbf{n}_{\Gamma_N} \cdot \underline{\boldsymbol{\tau}} \in L^p(\Gamma_N)\}$ and the weak solution z of the elliptic boundary value problem*

$$-\Delta z = -\nabla \cdot \underline{\boldsymbol{\tau}} \quad \text{in } \Omega \tag{4.15a}$$

$$\mathbf{n}_{\Gamma_N} \cdot \nabla z = 0 \quad \text{on } \Gamma_D \tag{4.15b}$$

$$\mathbf{n}_{\Gamma_N} \cdot \nabla z = \mathbf{n}_{\Gamma_N} \cdot \underline{\boldsymbol{\tau}} \quad \text{on } \Gamma_N \tag{4.15c}$$

there exists a constant $C_z > 0$ such that

$$\|\nabla z|_{\Gamma_N} \in L^p(\Gamma_N), \quad \|\nabla z\|_{L^p(\Gamma_N; \mathbb{R}^2)} \leq C_z. \tag{4.16}$$

Then for $\underline{\mathbf{p}}_c \in \{\underline{\mathbf{q}} \in \mathbf{H}_{0,\Gamma_D}^{(q)}(\text{div}, \Omega) \mid \mathbf{n}_{\Gamma_N} \cdot \underline{\mathbf{q}} \in L^q(\Gamma_N)\}$ it holds

$$\|\underline{\mathbf{p}}_c\|_{L^q(\Omega; \mathbb{R}^2)}^q \leq C_q \left(C_{PF}^{(1)}(p)^q \text{osc}_{h,1} + C_z^q C_{PF}^{(2)}(p)^q \text{osc}_{h,2} \right) \tag{4.17}$$

where $\text{osc}_{h,1}$ and $\text{osc}_{h,2}$ refer to the data oscillations

$$\text{osc}_{h,1} := \sum_{K \in \mathcal{T}_h} \text{osc}_{K,1}, \quad \text{osc}_{K,1} := h_K^q \int_K |f - f_h|^q dx \tag{4.18a}$$

$$\text{osc}_{h,2} := \sum_{K \in \mathcal{T}_h} \text{osc}_{K,2}, \quad \text{osc}_{K,2} := \sum_{E \in \mathcal{E}_h(\partial K \cap \Gamma_N)} h_E^q \int_E |u_N - u_{h,N}|^q ds. \tag{4.18b}$$

Proof. We have

$$\|\underline{\mathbf{p}}_c\|_{L^q(\Omega; \mathbb{R}^2)} = \sup \left\{ \int_{\Omega} \underline{\mathbf{p}}_c \cdot \underline{\boldsymbol{\tau}} \, dx \mid \underline{\boldsymbol{\tau}} \in \underline{\mathbf{H}}_{0, \Gamma_D}^{(p)}(\operatorname{div}, \Omega), \|\underline{\boldsymbol{\tau}}\|_{L^p(\Omega; \mathbb{R}^2)} \leq 1 \right\}.$$

For $\underline{\boldsymbol{\tau}} \in \{\underline{\mathbf{H}}_{0, \Gamma_D}^{(p)} \mid \mathbf{n}_{\Gamma_N} \cdot \underline{\boldsymbol{\tau}} \in L^p(\Gamma_N)\}$ there exists $z \in W^{1,p}(\Omega)$ such that $\underline{\boldsymbol{\tau}} = \nabla z$. In fact, z can be chosen as the weak solution of the boundary value problem (4.15). Hence, we have

$$\|\underline{\mathbf{p}}_c\|_{L^q(\Omega; \mathbb{R}^2)} \leq \sup_{\|\nabla z\|_{L^p(\Omega; \mathbb{R}^2)} \leq 1} \int_{\Omega} \underline{\mathbf{p}}_c \cdot \nabla z \, dx. \tag{4.19}$$

Applying Green's formula locally on each $K \in \mathcal{T}_h$ and taking (3.20) into account, we get

$$\begin{aligned} \int_{\Omega} \underline{\mathbf{p}}_c \cdot \nabla z \, dx &= \sum_{K \in \mathcal{T}_h} \int_K \underline{\mathbf{p}}_c \cdot \nabla z \, dx = - \sum_{K \in \mathcal{T}_h} \int_K \nabla \cdot \underline{\mathbf{p}}_c \, z \, dx + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E \mathbf{n}_E \cdot \underline{\mathbf{p}}_c \, z \, ds \\ &= \sum_{K \in \mathcal{T}_h} \int_K (f - f_h)(z - p_{0,1}) \, dx + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E (u_N - u_{h,N})(z - p_{0,2}) \, ds \end{aligned} \tag{4.20}$$

where $p_{0,1} := |K|^{-1} \int_K z \, dx$ and $p_{0,2} := |E|^{-1} \int_E z \, ds$. Using Hölder's inequality, the Cauchy–Schwarz inequality, and the Poincaré–Friedrichs inequalities (4.14), we obtain

$$\begin{aligned} &\left| \sum_{K \in \mathcal{T}_h} \int_K (f - f_h)(z - p_{0,1}) \, dx + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E (u_N - u_{h,N})(z - p_{0,2}) \, ds \right| \\ &\leq \sum_{K \in \mathcal{T}_h} \left(\int_K |f - f_h|^q \, dx \right)^{1/q} \left(\int_K |z - p_{0,1}|^p \, dx \right)^{1/p} \\ &\quad + \left(\sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E |u_N - u_{h,N}|^q \, ds \right)^{1/q} \left(\sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E |z - p_{0,2}|^p \, ds \right)^{1/p} \\ &\leq C_{PF}^{(1)}(p) \left(\sum_{K \in \mathcal{T}_h} h_K^q \int_K |f - f_h|^q \, dx \right)^{1/q} \left(\sum_{K \in \mathcal{T}_h} \int_K |\nabla z|^p \, dx \right)^{1/p} \\ &\quad + C_{PF}^{(2)}(p) \left(\sum_{E \in \mathcal{E}_h(\Gamma_N)} h_E^q \int_E |u_N - u_{h,N}|^q \, ds \right)^{1/q} \left(\sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E |\nabla z|^p \, ds \right)^{1/p}. \end{aligned} \tag{4.21}$$

Using (4.20), (4.21), and (4.16) in (4.19), it follows that

$$\|\underline{\mathbf{p}}_c\|_{L^q(\Omega; \mathbb{R}^2)} \leq C_{PF}^{(1)}(p) \operatorname{osc}_{h,1}^{1/q} + C_2 C_{PF}^{(2)}(p) \operatorname{osc}_{h,2}^{1/q}. \tag{4.22}$$

Hence, using (1.2), we find

$$\|\underline{\mathbf{p}}_c\|_{L^q(\Omega; \mathbb{R}^2)}^q \leq C_q \left(C_{PF}^{(1)}(p)^q \operatorname{osc}_{h,1} + C_2^q C_{PF}^{(2)}(p)^q \operatorname{osc}_{h,2} \right) \tag{4.23}$$

which is the assertion. \square

Moreover, as far as $J_P(u_h^c)$ is concerned, we have

$$J_P(u_h^c) = J_P(u_h) + \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K (|\nabla u_h^c|^p - |\nabla u_h|^p) \, dx + \sum_{K \in \mathcal{T}_h} \int_K f(u_h - u_h^c) \, dx + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E u_N(u_h - u_h^c) \, ds. \tag{4.24}$$

Lemma 4.2. *Let $u_h \in V_h$ be the solution of (3.22) and let $u_h^c \in V_h^c$ be its postprocessed finite element function. Then it holds*

$$|J_P(u_h^c) - J_P(u_h)| \leq \sum_{K \in \mathcal{T}_h} \kappa_K^{\text{eq}} \tag{4.25}$$

where

$$\kappa_K^{\text{eq}} := \|u_h - u_h^c\|_{W^{1,p}(K)}^p + \left(\sum_{K \in \mathcal{T}_h} |\nabla u_h|_{DG,K}^{p/q} + \|f\|_{L^q(K)} + \sum_{E \in \mathcal{E}_h(\partial K \cap \Gamma_N)} \|u_N\|_{L^q(E)} \right) \|u_h - u_h^c\|_{W^{1,p}(K)} \tag{4.26}$$

and $|\nabla u_h|_{DG,K}$ is given by

$$|\nabla u_h|_{DG,K} := \left(\int_K |\nabla u_h|^p \, dx \right)^{1/p}. \tag{4.27}$$

Proof. By Taylor expansion and using (1.2) as well as Hölder's inequality we find

$$\begin{aligned}
& \left| \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K \left(|\nabla u_h^c|^p - |\nabla u_h|^p \right) \right| \\
&= \left| \sum_{K \in \mathcal{T}_h} \int_K \int_0^1 |\nabla u_h + \lambda \nabla(u_h^c - u_h)|^{p-2} (\nabla u_h + \lambda \nabla(u_h^c - u_h)) \, d\lambda \cdot \nabla(u_h^c - u_h) \, dx \right| \\
&\leq \sum_{K \in \mathcal{T}_h} \int_K \int_0^1 |\nabla u_h + \lambda \nabla(u_h^c - u_h)|^{p-1} |\nabla(u_h^c - u_h)| \, d\lambda \, dx \\
&\leq C_{p-1} \sum_{K \in \mathcal{T}_h} \int_K \int_0^1 \left(|\nabla u_h|^{p-1} + \lambda^{p-1} |\nabla(u_h - u_h^c)|^{p-1} |\nabla(u_h - u_h^c)| \right) \, d\lambda \, dx \\
&\leq C_{p-1} \sum_{K \in \mathcal{T}_h} \left(\int_K |\nabla u_h|^p \, dx \right)^{1/q} \left(\int_K |\nabla(u_h - u_h^c)|^p \, dx \right)^{1/p} + \frac{1}{p} C_{p-1} \sum_{K \in \mathcal{T}_h} \int_K |\nabla(u_h - u_h^c)|^p \, dx. \quad (4.28)
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
& \left| \sum_{K \in \mathcal{T}_h} \int_K f(u_h - u_h^c) \, dx + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E u_N(u_h - u_h^c) \, ds \right| \\
&\leq \sum_{K \in \mathcal{T}_h} \left(\int_K |f|^q \, dx \right)^{1/q} \left(\int_K |u_h - u_h^c|^p \, dx \right)^{1/p} + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \left(\int_E |u_N|^q \, ds \right)^{1/q} \left(\int_E |u_h - u_h^c|^p \, ds \right)^{1/p}. \quad (4.29)
\end{aligned}$$

The assertion now follows from (4.24) and (4.28), (4.29). \square

For practical purposes, we further replace $I_{K_1}(u_h^c)$ by the penalty term

$$\alpha \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h^c - u_D|^p \, ds. \quad (4.30)$$

In view of (3.29) we have $u_h^c|_E = u_{h,D}$ on $E \in \mathcal{E}_h(\Gamma_D)$ and hence, (4.30) gives rise to the data oscillation

$$\alpha \, \text{osc}_{h,3} := \alpha \sum_{K \in \mathcal{T}_h} \text{osc}_{K,3} \quad (4.31a)$$

$$\text{osc}_{K,3} := \sum_{E \in \mathcal{E}_h(\partial K \cap \Gamma_D)} h_E^{-p/q} \int_E |u_D - u_{h,D}|^p \, ds. \quad (4.31b)$$

Using Lemma 4.1 and Lemma 4.2 in (4.11) yields

$$\|u - u_h\|_{W^{1,p}(\Omega; \mathcal{T}_h)}^2 \lesssim \eta_{h,1}^{\text{eq}} + \eta_{h,2}^{\text{eq}}. \quad (4.32a)$$

Here, $\eta_{h,1}^{\text{eq}}$ and $\eta_{h,2}^{\text{eq}}$ are given by

$$\eta_{h,1}^{\text{eq}} := \sum_{K \in \mathcal{T}_h} \eta_{K,1}^{\text{res}}, \quad \eta_{h,2}^{\text{eq}} := \sum_{K \in \mathcal{T}_h} \eta_{K,2}^{\text{res}} \quad (4.32b)$$

where $\eta_{K,i}^{\text{res}}$, $1 \leq i \leq 2$, read as follows:

$$\eta_{K,1}^{\text{eq}} := \frac{1}{p} \int_K |\nabla u_h|^p \, dx - \int_K f u_h \, dx - \sum_{E \in \mathcal{E}_h(\partial K \cap \Gamma_N)} \int_E u_N u_h \, ds + \frac{1}{q} \int_K |\underline{\mathbf{p}}_h^{\text{eq}}|^q \, dx - \sum_{E \in \mathcal{E}_h(\partial K \cap \Gamma_D)} \int_E u_D \mathbf{n}_E \cdot \underline{\mathbf{p}}_h^{\text{eq}} \, ds \quad (4.32c)$$

$$\eta_{K,2}^{\text{eq}} := \|u_h - u_h^c\|_{W^{1,p}(K)}^2 + \chi_K^{\text{eq}} + \sum_{i=1}^3 \text{osc}_{K,i}. \quad (4.32d)$$

The right-hand side in (4.32) is then a computable and localizable quantity for the a posteriori estimation of

the global discretization error. It gives rise to the following equilibrated a posteriori error estimator

$$\eta_h^{\text{eq}} := \sum_{K \in \mathcal{T}_h(\Omega)} \eta_K^{\text{eq}} \quad (4.33)$$

where the local contributions η_K^{eq} , $K \in \mathcal{T}_h(\Omega)$, read as follows

$$\eta_K^{\text{eq}} := \eta_{K,1}^{\text{eq}} + \eta_{K,2}^{\text{eq}}.$$

The construction of an equilibrated flux will be dealt with in the subsequent section.

5 Construction of an equilibrated flux

We construct an equilibrated flux $\underline{\mathbf{p}}_h^{\text{eq}} \in \underline{\mathbf{H}}^{(q)}(\text{div}, \Omega)$ which allows us to apply the equilibrated a posteriori error estimator (4.33). The construction will be done locally by an interpolation on each element. In particular, we denote by $\text{BDM}_k(K)$, $k \in \mathbb{N}$, the Brezzi–Douglas–Marini element $\text{BDM}_k(K) := P_k(K)^2$ (see, e.g., [12]), and recall the following result.

Lemma 5.1. *Let $k \geq 1$. Any vector field $\underline{\mathbf{q}} \in \text{BDM}_k(K)$ is uniquely defined by the following degrees of freedom*

$$\int_E \mathbf{n}_E \cdot \underline{\mathbf{q}} p_k \, ds, \quad p_k \in P_k(E), \quad E \in \mathcal{E}_h(\partial K) \quad (5.1a)$$

$$\int_K \underline{\mathbf{q}} \cdot \nabla p_{k-1} \, dx, \quad p_{k-1} \in P_{k-1}(K) \quad (5.1b)$$

$$\int_K \underline{\mathbf{q}} \cdot \text{curl}(b_K p_{k-2}) \, dx, \quad p_{k-2} \in P_{k-2}(K) \quad (5.1c)$$

where b_K in (5.1c) is the element bubble function on K given by $b_K = \prod_{i=1}^3 \lambda_i^K$ and λ_i^K , $1 \leq i \leq 3$, are the barycentric coordinates of K .

Proof. We refer to [12]. □

Lemma 5.2. *Let $K \in \mathcal{T}_h$. There exists a constant $C_E > 0$, depending only on k and the local geometry of the triangulation, such that for any $\underline{\mathbf{q}} \in P_k(K)^2$ it holds*

$$\begin{aligned} \int_K |\underline{\mathbf{q}}|^q \, dx \leq C_E \left(\sum_{E \in \mathcal{E}_h(\partial K)} h_E \int_E |\mathbf{n}_E \cdot \underline{\mathbf{q}}|^q \, ds + h_K^q \int_K |\nabla \cdot \underline{\mathbf{q}}|^q \, dx \right. \\ \left. + h_K^q \max \left\{ \int_K |\underline{\mathbf{q}} \cdot \text{curl}(b_K p_{k-2})|^q \, dx \mid p_{k-2} \in P_{k-2}(K), \max_{x \in K} |p_{k-2}(x)| \leq 1 \right\} \right). \end{aligned} \quad (5.2)$$

Proof. The assertion can be proved by standard scaling arguments. □

We construct an equilibrated flux $\underline{\mathbf{p}}_h^{\text{eq}}|_K \in \text{BDM}_k(K)$ by the specifications

$$\int_E \mathbf{n}_E \cdot \underline{\mathbf{p}}_h^{\text{eq}} p_k \, ds = \int_E \mathbf{n}_E \cdot \hat{\underline{\mathbf{p}}}_{\partial K}|_E p_k \, ds, \quad p_k \in P_k(E), \quad E \in \mathcal{E}_h(\partial K) \quad (5.3a)$$

$$\int_K \underline{\mathbf{p}}_h^{\text{eq}} \cdot \nabla p_{k-1} \, dx = \int_K \underline{\mathbf{p}}_h \cdot \nabla p_{k-1} \, dx, \quad p_{k-1} \in P_{k-1}(K) \quad (5.3b)$$

$$\int_K \underline{\mathbf{p}}_h^{\text{eq}} \cdot \text{curl}(b_K p_{k-2}) \, dx = \int_K (\underline{\mathbf{H}}_k^{(q)}(|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h) \cdot \text{curl}(b_K p_{k-2})) \, dx, \quad p_{k-2} \in P_{k-2}(K) \quad (5.3c)$$

where $\underline{\mathbf{p}}_h \in \underline{\mathbf{V}}_h$ satisfies the two-field formulation (3.33).

Theorem 5.1. *The flux $\underline{\mathbf{p}}_h^{\text{eq}}$ as given by (5.3) is an equilibrated flux, i.e., $\underline{\mathbf{p}}_h^{\text{eq}} \in \underline{\mathbf{H}}^{(q)}(\text{div}, \Omega)$ and it satisfies (4.9b), (4.9c).*

Proof. Due to (5.3a) the normal components of $\underline{\mathbf{p}}_h^{\text{eq}}$ on $E \in \mathcal{E}_h(\Omega)$ are continuous across E , and hence, $\underline{\mathbf{p}}_h^{\text{eq}} \in \underline{\mathbf{H}}^{(q)}(\text{div}; \Omega)$, i.e., (4.9a) is satisfied.

Further, it follows from Gauss's theorem and (5.3a) that for $p_{k1} \in P_{k-1}(K)$, $K \in \mathcal{T}_h$, it holds

$$\int_K \nabla \cdot (\underline{\mathbf{p}}_h^{\text{eq}} p_{k-1}) \, dx = \int_{\partial K} \mathbf{n}_{\partial K} \cdot \underline{\mathbf{p}}_h^{\text{eq}} p_{k-1} \, ds = \int_{\partial K} \mathbf{n}_{\partial K} \cdot \hat{\underline{\mathbf{p}}}_{\partial K} p_{k-1} \, ds. \quad (5.4)$$

On the other hand, from (5.3b) we deduce

$$\int_K \nabla \cdot (\underline{\mathbf{p}}_h^{\text{eq}} p_{k-1}) \, dx = \int_K \nabla \cdot \underline{\mathbf{p}}_h^{\text{eq}} p_{k-1} \, dx + \int_K \underline{\mathbf{p}}_h^{\text{eq}} \cdot \nabla p_{k-1} \, dx = \int_K \nabla \cdot \underline{\mathbf{p}}_h^{\text{eq}} p_{k-1} \, dx + \int_K \underline{\mathbf{p}}_h \cdot \nabla p_{k-1} \, dx$$

Hence, observing (5.4) and using (3.33b) with $v_h|_K = p_{k-1}$ and $v_h|_{K'} = 0$, $K' \in \mathcal{T}_h$, $K' \neq K$, we obtain

$$0 = \int_K \nabla \cdot \underline{\mathbf{p}}_h^{\text{eq}} p_{k-1} \, dx + \int_K \underline{\mathbf{p}}_h \cdot \nabla p_{k-1} \, dx - \int_{\partial K} \mathbf{n}_{\partial K} \cdot \hat{\underline{\mathbf{p}}}_{\partial K} p_{k-1} \, ds = \int_K \nabla \cdot \underline{\mathbf{p}}_h^{\text{eq}} p_{k-1} \, dx + \int_K f_h p_{k-1} \, dx.$$

Since $\nabla \cdot \underline{\mathbf{p}}_h^{\text{eq}}$ and $f_h|_K$ are contained in $P_{k-1}(K)$, we readily deduce that (4.9b) holds true. Moreover, for $E \in \mathcal{E}_h(\Gamma_N)$ it follows from (5.3a) and (3.34) that

$$\int_E \mathbf{n}_E \cdot \underline{\mathbf{p}}_h^{\text{eq}} p_k \, ds = \int_E \mathbf{n}_E \cdot \hat{\underline{\mathbf{p}}}_{\partial K}|_E p_k \, ds = \int_E u_{h,N} p_k \, ds. \quad (5.5)$$

Since both $\mathbf{n}_E \cdot \underline{\mathbf{p}}_h^{\text{eq}}$ and $u_{h,N}|_E$ are polynomials of degree k on E , it follows from (5.5) that (4.9c) is satisfied. \square

6 Relationship with a residual type a posteriori error estimator

A residual-type a posteriori error estimator for the IPDG approximation of the p -Laplace equation with homogeneous Dirichlet boundary conditions in case $p = 2$ has been derived and analyzed in [36, 38, 39]. Its generalization to arbitrary $1 < p < \infty$ reads as follows:

$$\eta_h^{\text{res}} = \sum_{i=1}^5 (\eta_{h,i}^{\text{res}} + \hat{\eta}_{h,i}^{\text{res}}) + \sum_{i=3}^4 \hat{\eta}_{h,i}^{\text{res}}. \quad (6.1a)$$

Here, the element residuals $\eta_{h,1}^{\text{res}}$ and the edge residuals $\eta_{h,i}^{\text{res}}$, $2 \leq i \leq 5$, are given by

$$\eta_{h,1}^{\text{res}} := \sum_{K \in \mathcal{T}_h} h_K^q \int_K |f + \nabla \cdot \underline{\mathbf{H}}_k^{(q)} (|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h)|^q \, dx \quad (6.1b)$$

$$\eta_{h,2}^{\text{res}} := \sum_{E \in \mathcal{E}_h(\Omega)} h_E \int_E |\mathbf{n}_E \cdot [|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h]|_E|^q \, ds \quad (6.1c)$$

$$\eta_{h,3}^{\text{res}} := \sum_{E \in \mathcal{E}_h(\Omega)} h_E^{-p/q} \int_E |[u_h]_E|^p \, ds \quad (6.1d)$$

$$\eta_{h,4}^{\text{res}} := \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_D|^p \, ds \quad (6.1e)$$

$$\eta_{h,5}^{\text{res}} := \sum_{E \in \mathcal{E}_h(\Gamma_N)} h_E \int_E |u_N - \mathbf{n}_E \cdot |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h|^q \, ds. \quad (6.1f)$$

The residuals $\hat{\eta}_{h,i}^{\text{res}}$, $1 \leq i \leq 5$, and $\hat{\eta}_{h,i}^{\text{res}}$, $3 \leq i \leq 4$, read as follows:

$$\hat{\eta}_{h,i}^{\text{res}} := (\eta_{h,i}^{\text{res}})^{1/q} |\nabla_{DG,2} u_h|_{DG,\Omega}, \quad 1 \leq i \leq 5 \quad (6.1g)$$

$$\hat{\eta}_{h,i}^{\text{res}} := (\eta_{h,i}^{\text{res}})^{1/p} |\nabla u_h|_{DG,\Omega}^{1/q}, \quad 3 \leq i \leq 4 \quad (6.1h)$$

where

$$|\nabla_{DG,2} u_h|_{DG,\Omega} := \left(\sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx \right)^{1/p} \quad (6.2)$$

$$|\nabla u_h|_{DG,\Omega} := \sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^p dx. \quad (6.3)$$

In addition to (4.18) and (4.31) we define data oscillations $\overline{osc}_{h,i}$, $1 \leq i \leq 2$, according to

$$\overline{osc}_{h,1} := (osc_{h,1})^{1/q} |\nabla u_h|_{DG,\Omega}^{1/p} \quad (6.4a)$$

$$\overline{osc}_{h,2} := (osc_{h,2})^{1/q} \|\nabla u_h\|_{DG,\Gamma_D}^{1/p} \quad (6.4b)$$

where

$$|\nabla v_h|_{DG,\Gamma'} := \sum_{E \in \mathcal{E}_h(\Gamma')} \int_E |\nabla v_h|^p ds, \quad \Gamma' \in \{\Gamma_D, \Gamma_N\}. \quad (6.5)$$

Remark 6.1. Residual-type a posteriori error estimates for P1 conforming finite element approximations of the p -Laplace problem with homogeneous Dirichlet boundary conditions have been considered in [5] (see also [41, 42]). We note that in this case the residuals $\tilde{\eta}_{h,i}^{\text{res}}$, $3 \leq i \leq 5$, and $\eta_{h,i}^{\text{res}}$, $3 \leq i \leq 4$, as well as the data oscillations $osc_{h,i}$, $2 \leq i \leq 3$, and $\overline{osc}_{h,2}$ vanish. The residuals $\eta_{h,i}^{\text{res}}$, $1 \leq i \leq 2$, reduce to

$$\begin{aligned} \eta_{h,1}^{\text{res}} &= \sum_{K \in \mathcal{T}_h} h_K^q \int_K |f|^q dx \\ \eta_{h,2}^{\text{res}} &= \sum_{E \in \mathcal{E}_h(\Omega)} h_E \int_E |\mathbf{n}_E \cdot [|\nabla u_h|^{p-2} \nabla u_h]_E|^q ds. \end{aligned} \quad (6.6)$$

Using the relationships between the $\|\cdot\|_{W^{1,p}(\Omega)}$ norm and quasi-norms provided in [4], it can be shown that (6.6) and the data oscillation $\overline{osc}_{h,1}$ are closely related to those in [5].

The goal of this section is to establish the relationship between the equilibrated a posteriori error estimator η_h^{eq} and the residual-based a posteriori error estimator η_h^{res} . For notational convenience, throughout this section we set $\mathbf{z}_h := |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h$. In view of the definitions (2.3a) and (2.4a) of the primal and dual energies we have

$$\eta_{h,1}^{\text{eq}} = \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^p dx + \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\mathbf{p}_h^{\text{eq}}|^q dx - \sum_{K \in \mathcal{T}_h} \int_K f u_h dx - \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E u_N u_h ds - \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_D \mathbf{n}_E \cdot \mathbf{p}_h^{\text{eq}} ds. \quad (6.7)$$

For the first two terms on the right-hand side of (6.7) we obtain

$$\begin{aligned} \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^p dx + \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\mathbf{p}_h^{\text{eq}}|^q dx &= \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^p dx - \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx \\ &\quad + \left(1 - \frac{1}{q}\right) \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx + \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\mathbf{p}_h^{\text{eq}}|^q dx. \end{aligned} \quad (6.8)$$

Using (6.8) in (6.7) it follows that

$$\begin{aligned} \eta_{h,1}^{\text{eq}} &= \sum_{K \in \mathcal{T}_h} \int_K \mathbf{z}_h \cdot \nabla_{DG,2} u_h dx - \sum_{K \in \mathcal{T}_h} \int_K f u_h dx - \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_D \mathbf{n}_E \cdot \mathbf{p}_h^{\text{eq}} ds - \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E u_N u_h ds \\ &\quad + \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^p dx - \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx + \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\mathbf{p}_h^{\text{eq}}|^q dx - \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx. \end{aligned} \quad (6.9)$$

We will estimate the terms on the right-hand side in (6.9) by a series of Lemmas.

Lemma 6.1. *There exists a constant $\hat{C}_{\text{rel}}^{(1)} > 0$, depending only on $\alpha, C_p, C_{PF}^{(i)}(p), 1 \leq i \leq 2$, and on p, q , such that it holds*

$$\left| \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h \cdot \nabla_{DG,2} u_h \, dx - \sum_{K \in \mathcal{T}_h} \int_K f u_h \, dx - \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_D \mathbf{n}_E \cdot \underline{\mathbf{p}}_h^{\text{eq}} \, ds - \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E u_N u_h \, ds \right| \leq \hat{C}_{\text{rel}}^{(1)} \left(\eta_{h,4}^{\text{res}} + \text{osc}_{h,3} + \overline{\text{osc}}_{h,1} + \overline{\text{osc}}_{h,2} \right). \tag{6.10}$$

Proof. In view of (3.12b), (3.13b), and (3.22) we get

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} \int_K \mathbf{z}_h \cdot \nabla_{DG,2} u_h \, dx &= \sum_{K \in \mathcal{T}_h} \int_K \mathbf{z}_h \cdot \underline{\Pi}_k^{(p)}(\nabla_{DG,2} u_h) \, dx = \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)}(\mathbf{z}_h) \cdot \nabla_{DG,2} u_h \, dx \\ &= \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)}(\mathbf{z}_h) \cdot \nabla_{DG,1} u_h \, dx + \sum_{K \in \mathcal{T}_h} \int_K \underline{\Pi}_k^{(q)}(\mathbf{z}_h) \cdot \underline{\mathbf{R}}_{h,2}(u_D^*) \, dx \\ &= \sum_{K \in \mathcal{T}_h} \int_K f_h u_h \, dx + \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E u_{h,N} u_h \, ds - \alpha \sum_{E \in \mathcal{E}_h(\Omega)} h_E^{-p/q} \int_E |[u_h]_E|^{p-2} [u_h]_E [u_h]_E \, ds \\ &\quad - \alpha \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_D|^{p-2} (u_h - u_D) u_h \, ds + \sum_{E \in \mathcal{E}_h(\Gamma_D)} \mathbf{n}_E \cdot \mathbf{z}_h u_{h,D} \, ds \end{aligned} \tag{6.11}$$

where we have used that

$$\sum_{E \in \mathcal{E}_h(\Gamma_D)} \mathbf{n}_E \cdot \underline{\Pi}_k^{(q)}(\mathbf{z}_h) u_D \, ds = \sum_{E \in \mathcal{E}_h(\Gamma_D)} \Pi_k^{(q)}(\mathbf{n}_E \cdot \mathbf{z}_h) u_D \, ds = \sum_{E \in \mathcal{E}_h(\Gamma_D)} \mathbf{n}_E \cdot \mathbf{z}_h \Pi_k^{(p)}(u_D) \, ds = \sum_{E \in \mathcal{E}_h(\Gamma_D)} \mathbf{n}_E \cdot \mathbf{z}_h u_{h,D} \, ds.$$

On the other hand, observing (5.3a), (3.34), and $\mathbf{n}_E \cdot \underline{\mathbf{p}}_h^{\text{eq}} = \Pi_k(\mathbf{n}_E \cdot \hat{\underline{\mathbf{p}}}_{\partial K}|_E)$, it follows that

$$\begin{aligned} \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_D \mathbf{n}_E \cdot \underline{\mathbf{p}}_h^{\text{eq}} \, ds &= \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_D \Pi_k(\mathbf{n}_E \cdot \hat{\underline{\mathbf{p}}}_{\partial K}|_E) \, ds \\ &= \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E \Pi_k(u_D) \mathbf{n}_E \cdot \hat{\underline{\mathbf{p}}}_{\partial K}|_E \, ds = \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_{h,D} \mathbf{n}_E \cdot \hat{\underline{\mathbf{p}}}_{\partial K}|_E \, ds \\ &= \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_{h,D} \mathbf{n}_E \cdot \mathbf{z}_h \, ds - \alpha \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_D|^{p-2} (u_h - u_D) u_{h,D} \, ds. \end{aligned} \tag{6.12}$$

From (6.11) and (6.12) we deduce

$$\begin{aligned} \left| \sum_{K \in \mathcal{T}_h} \int_K \mathbf{z}_h \cdot \nabla_{DG,2} u_h \, dx - \sum_{K \in \mathcal{T}_h} \int_K f u_h \, dx - \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E u_D \mathbf{n}_E \cdot \underline{\mathbf{p}}_h^{\text{eq}} \, ds - \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E u_N u_h \, ds \right| \\ \leq \left| \sum_{K \in \mathcal{T}_h} \int_K (f_h - f) u_h \, dx \right| + \left| \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E (u_{h,N} - u_N) u_h \, ds \right| \\ + \alpha \left| \sum_{E \in \mathcal{E}_h(\Gamma_D)} \int_E h_E^{-p/q} |u_h - u_D|^{p-2} (u_h - u_D) (u_{h,D} - u_h) \, ds \right|. \end{aligned} \tag{6.13}$$

Applying Hölder’s inequality, the Cauchy–Schwarz inequality, observing (3.20) with $p_0 = \int_K u_h \, dx$, and the Poincaré–Friedrichs inequality (4.14), the first term on the right-hand side of (6.13) can be estimated from above as follows:

$$\begin{aligned} \left| \sum_{K \in \mathcal{T}_h} \int_K (f_h - f) u_h \, dx \right| &= \left| \sum_{K \in \mathcal{T}_h} \int_K (f_h - f)(u_h - p_0) \, dx \right| \leq \sum_{K \in \mathcal{T}_h} \left(\int_K |f - f_h|^q \, dx \right)^{1/q} \left(\int_K |u_h - p_0|^p \, dx \right)^{1/p} \\ &\leq C_{PF}^{(1)}(q) \sum_{K \in \mathcal{T}_h} \left(\int_K |f - f_h|^q \, dx \right)^{1/q} h_K \left(\int_K |\nabla u_h|^p \, dx \right)^{1/p} \\ &\leq C_{PF}^{(1)}(p) \left(\sum_{K \in \mathcal{T}_h} h_K^q \int_K |f - f_h|^q \, dx \right)^{1/q} |\nabla u_h|_{DG,\Omega} = C_{PF}^{(1)}(p) \overline{\text{osc}}_{h,1}. \end{aligned} \tag{6.14}$$

Likewise, with $p_0 = \int_E u_h ds$, $E \in \mathcal{E}_h(\Gamma_N)$, we obtain

$$\begin{aligned} \left| \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E (u_{h,N} - u_N) u_h ds \right| &= \left| \sum_{E \in \mathcal{E}_h(\Gamma_N)} \int_E (u_{h,N} - u_N) (u_h - p_0) ds \right| \\ &\leq C_{PF}^{(2)}(p) \left(\sum_{E \in \mathcal{E}_h(\Gamma_N)} h_E^q \int_E |u_N - u_{h,N}|^q dx \right)^{1/q} |\nabla u_h|_{DG, \Gamma_N} = C_{PF}^{(2)}(p) \overline{osc}_{h,2}. \end{aligned} \quad (6.15)$$

Likewise, using the same arguments and (1.2), for the last term on the right-hand side of (6.13) we obtain

$$\begin{aligned} \alpha \left| \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_D|^{p-2} (u_h - u_D) (u_{h,D} - u_h) ds \right| \\ \leq \alpha \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \left(\int_E |u_h - u_D|^{p-2} (u_h - u_D)^q ds \right)^{1/q} \left(\int_E |u_h - u_{h,D}|^p ds \right)^{1/p} \\ \leq \alpha \left(\sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_D|^p ds \right)^{1/q} \left(\sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_{h,D}|^p ds \right)^{1/p} \\ \leq \alpha \left(\frac{1}{q} \left(\frac{1}{2} \right)^{-q/p} \eta_{h,4}^{\text{res}} + \frac{1}{2p} C_p(\eta_{h,4}^{\text{res}} + osc_{h,3}) \right). \end{aligned} \quad (6.16)$$

The assertion now follows from (6.11)–(6.16). \square

Lemma 6.2. *There exists a constant $\hat{C}_{\text{rel}}^{(2)} > 0$, depending only on C_{p-1} , C_p , C_{rec} , and on p, q , such that it holds*

$$\left| \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^p dx - \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx \right| \leq \hat{C}_{\text{rel}}^{(2)} (\eta_{h,3}^{\text{res}} + \hat{\eta}_{h,3}^{\text{res}} + \eta_{h,4}^{\text{res}} + \hat{\eta}_{h,4}^{\text{res}}). \quad (6.17)$$

Proof. By Taylor expansion, observing (3.13), applying Hölder's inequality, the Cauchy–Schwarz inequality, and using (1.2) as well as (3.15), we find

$$\begin{aligned} \left| \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx - \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^p dx \right| \\ = \left| \sum_{K \in \mathcal{T}_h} \int_K \int_0^1 (|\nabla u_h + \lambda(\nabla_{DG,2} u_h - \nabla u_h)|^{p-2} (\nabla u_h + \lambda(\nabla_{DG,2} u_h - \nabla u_h)) d\lambda \cdot (\nabla_{DG,2} u_h - \nabla u_h) dx \right| \\ \leq \sum_{K \in \mathcal{T}_h} \int_K \int_0^1 |\nabla u_h + \lambda(\nabla_{DG,2} u_h - \nabla u_h)|^{p-1} |\nabla_{DG,2} u_h - \nabla u_h| d\lambda dx \\ \leq C_{p-1} \left(\sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^{p-1} \|\nabla_{DG,2} u_h - \nabla u_h\| dx + \frac{1}{p} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h - \nabla u_h|^p dx \right) \\ \leq C_{p-1} \left(\sum_{K \in \mathcal{T}_h} \int_K |\nabla u_h|^p dx \right)^{1/q} \left(\sum_{K \in \mathcal{T}_h} \int_K |\mathbf{R}_{h,1}(u_h) - \mathbf{R}_{h,2}(u_D^*)|^p dx \right)^{1/p} \\ + \frac{1}{p} C_{p-1} \sum_{K \in \mathcal{T}_h} \int_K |\mathbf{R}_{h,1}(u_h) - \mathbf{R}_{h,2}(u_D^*)|^p dx \\ \leq C_{p-1} C_{\text{rec}} (\hat{\eta}_{h,3}^{\text{res}} + \hat{\eta}_{h,4}^{\text{res}}) + \frac{1}{p} C_{p-1} C_p C_{\text{rec}} (\eta_{h,3}^{\text{res}} + \eta_{h,4}^{\text{res}}). \end{aligned} \quad (6.18)$$

This completes the proof. \square

Lemma 6.3. *There exists a constant $\hat{C}_{\text{rel}}^{(3)} > 0$, depending only on α , c_R , C_q , C_E , and on p, q , such that it holds*

$$\frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{p}_h^{\text{eq}}|^q dx - \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx \leq \hat{C}_{\text{rel}}^{(3)} \left(\sum_{i=1}^5 (\eta_{h,i}^{\text{res}} + \hat{\eta}_{h,i}^{\text{res}}) + osc_{h,1} + \overline{osc}_{h,1} \right). \quad (6.19)$$

Proof. We have

$$\begin{aligned} & \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}}|^q dx - \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx = \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}}|^q dx - \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx \\ & + \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx - \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx. \end{aligned} \quad (6.20)$$

For the first two terms on the right-hand side of (6.20), applying Taylor expansion and using (1.2) we obtain

$$\begin{aligned} & \left| \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}}|^q dx - \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx \right| \\ & = \left| \sum_{K \in \mathcal{T}_h} \int_K \int_0^1 |\underline{\Pi}_k^{(q)}(\mathbf{z}_h) + \lambda(\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h))|^{q-2} (\underline{\Pi}_k^{(q)}(\mathbf{z}_h) + \lambda(\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h))) d\lambda \cdot (\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h)) dx \right| \\ & \leq \sum_{K \in \mathcal{T}_h} \int_K \int_0^1 |\underline{\Pi}_k^{(q)}(\mathbf{z}_h) + \lambda(\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h))|^{q-1} |\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h)| d\lambda dx \\ & \leq C_{q-1} \left(\sum_{K \in \mathcal{T}_h} \int_K |\underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^{q-1} |\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h)| dx + \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx \right) \\ & \leq C_{q-1} \left(\left(\sum_{K \in \mathcal{T}_h} \int_K |\underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx \right)^{1/p} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx \right)^{1/q} \\ & + \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx. \end{aligned} \quad (6.21)$$

In view of (5.2) and (5.3c) we deduce

$$\sum_{K \in \mathcal{T}_h} \int_K |\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx \leq C_E \left(\sum_{K \in \mathcal{T}_h} h_K \int_{\partial K} |\mathbf{n}_{\partial K} \cdot (\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k(\mathbf{z}_h))|^q ds + \sum_{K \in \mathcal{T}_h} h_K^q \int_K |\nabla \cdot (\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k(\mathbf{z}_h))|^q dx \right). \quad (6.22)$$

Since (5.3a) implies $\mathbf{n}_{\partial K} \cdot \underline{\mathbf{p}}_h^{\text{eq}}|_E = \Pi_k^{(q)}(\mathbf{n}_{\partial K} \cdot \hat{\underline{\mathbf{p}}}_{\partial K})$, we get

$$\begin{aligned} & \sum_{K \in \mathcal{T}_h} h_K \int_{\partial K} |\mathbf{n}_{\partial K} \cdot (\underline{\mathbf{p}}_h^{\text{eq}} - \underline{\Pi}_k(|\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h))|^q ds \\ & = \sum_{K \in \mathcal{T}_h} h_K \int_{\partial K \cap \Omega} |\Pi_k^{(q)}(\mathbf{n}_{\partial K} \cdot (\hat{\underline{\mathbf{p}}}_{\partial K} - |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h))|^q ds \\ & + \sum_{K \in \mathcal{T}_h} h_K \int_{\partial K \cap \Gamma} |\Pi_k^{(q)}(\mathbf{n}_{\partial K} \cdot (\hat{\underline{\mathbf{p}}}_{\partial K} - |\nabla_{DG,2} u_h|^{p-2} \nabla_{DG,2} u_h))|^q ds. \end{aligned} \quad (6.23)$$

Now, using (3.1), (3.34) and observing $1 - p = -p/q$ it follows that

$$\begin{aligned} & \left| \sum_{K \in \mathcal{T}_h} h_K \int_{\partial K \cap \Omega} |\Pi_k^{(q)}(\mathbf{n}_{\partial K} \cdot (\hat{\underline{\mathbf{p}}}_{\partial K} - \mathbf{z}_h)|^q ds \right| \\ & \leq c_R^{-1} \sum_{E \in \mathcal{E}_h(\Omega)} h_E \int_E (|\{\mathbf{z}_h\}_E - \mathbf{z}_h|_{E,+}|^q + |\{\mathbf{z}_h\}_E - \mathbf{z}_h|_{E,-}|^q) ds \\ & + c_R^{-1} \alpha^q \sum_{E \in \mathcal{E}_h(\Omega)} h_E \int_E |h_E^{-p/q} |[u_h]_E|^{p-2} [u_h]_E|^q ds \\ & \leq c_R^{-1} \left(\sum_{E \in \mathcal{E}_h(\Omega)} h_E \int_E |[\mathbf{z}_h]_E|^q ds + \alpha^q \sum_{E \in \mathcal{E}_h(\Omega)} h_E^{-p/q} \int_E |[u_h]_E|^p ds \right) = c_R^{-1} (\eta_{h,2}^{\text{res}} + \alpha^q \eta_{h,3}^{\text{res}}). \end{aligned} \quad (6.24)$$

Using (3.1) and (3.34) again, we obtain

$$\begin{aligned}
 & \left| \sum_{K \in \mathcal{T}_h} h_K \int_{\partial K \cap \Gamma} |\Pi_k^{(q)}(\mathbf{n}_{\partial K} \cdot (\hat{\mathbf{p}}_{\partial K} - \mathbf{z}_h))^q ds \right| \\
 & \leq c_R^{-1} \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E \int_E |\mathbf{n}_E \cdot (\hat{\mathbf{p}}_{\partial K}|_E - \mathbf{z}_h)|^q ds + c_R^{-1} \sum_{E \in \mathcal{E}_h(\Gamma_N)} h_E \int_E |\mathbf{n}_E \cdot (\hat{\mathbf{p}}_{\partial K}|_E - \mathbf{z}_h)|^q ds \\
 & \leq \alpha^q c_R^{-1} \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E \int_E |h_E^{-p/q} |u_h - u_D|^{p-2} (u_h - u_D)|^q ds + c_R^{-1} \sum_{E \in \mathcal{E}_h(\Gamma_N)} h_E \int_E |(u_N - \mathbf{n}_E \cdot \mathbf{z}_h)|^q ds \\
 & \leq \alpha^q c_R^{-1} \sum_{E \in \mathcal{E}_h(\Gamma_D)} h_E^{-p/q} \int_E |u_h - u_D|^p ds + c_R^{-1} \sum_{E \in \mathcal{E}_h(\Gamma_N)} h_E \int_E |(u_N - \mathbf{n}_E \cdot \mathbf{z}_h)|^q ds = c_R^{-1} (\alpha^q \eta_{h,4}^{\text{res}} + \eta_{h,5}^{\text{res}}).
 \end{aligned} \tag{6.25}$$

Moreover, due to (1.2) and (4.9a) we have

$$\begin{aligned}
 \sum_{K \in \mathcal{T}_h} h_K^q \int_K |\nabla \cdot (\mathbf{p}_h^{\text{eq}} - \underline{\Pi}_k^{(q)}(\mathbf{z}_h))|^q dx &= \sum_{K \in \mathcal{T}_h} h_K^q \int_K |f_h + \nabla \cdot \underline{\Pi}_k(\mathbf{z}_h)|^q dx \\
 &\leq C_q \sum_{K \in \mathcal{T}_h} h_K^q \int_K |f - f_h|^q dx + C_q \sum_{K \in \mathcal{T}_h} h_K^q \int_K |f + \nabla \cdot \underline{\Pi}_k(\mathbf{z}_h)|^q dx = C_q (\eta_{h,1}^{\text{res}} + \text{osc}_{h,1}).
 \end{aligned} \tag{6.26}$$

Finally, we have

$$\sum_{K \in \mathcal{T}_h} \int_K |\underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx \leq \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx$$

and hence,

$$\frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\underline{\Pi}_k^{(q)}(\mathbf{z}_h)|^q dx - \frac{1}{q} \sum_{K \in \mathcal{T}_h} \int_K |\nabla_{DG,2} u_h|^p dx \leq 0. \tag{6.27}$$

The assertion now follows from (6.20)–(6.27). □

The following result establishes the relationship between the equilibrated and the residual a posteriori error estimator.

Theorem 6.1. *Let $u_h \in V_h$ be the IPDG approximation as given by (3.22) and let $\eta_{h,1}^{\text{eq}}, \eta_{h,i}^{\text{res}}, \tilde{\eta}_{h,i}^{\text{res}}, 1 \leq i \leq 5, \hat{\eta}_{h,i}^{\text{res}}, 3 \leq i \leq 4$, and $\text{osc}_{h,i}, 1 \leq i \leq 3, \overline{\text{osc}}_{h,i}, 1 \leq i \leq 2$, be the equilibrated and the residual a posteriori error estimators as well as the data oscillations as given by (4.32b), (6.1), and (4.18), (4.31), (6.4). Then there exists a constant $C_{\text{res}} > 0$, depending on $\alpha, c_R, C_{p-1}, C_p, C_q, C_{\text{rec}}, C_E, C_{\text{PF}}^{(i)}(p), 1 \leq i \leq 2$, and on p, q , such that*

$$\eta_{h,1}^{\text{eq}} \leq C_{\text{res}} \left(\eta_h^{\text{res}} + \sum_{i=1}^3 \text{osc}_{h,i} + \sum_{i=1}^2 \overline{\text{osc}}_{h,i} \right). \tag{6.28}$$

Moreover, if we use (3.30) in (4.32b), then $\eta_{h,2}^{\text{eq}}$ can be estimated from above in terms of the residuals $\eta_{h,3}^{\text{res}}, \eta_{h,4}^{\text{res}}$, and the data oscillations $\text{osc}_{h,i}, 1 \leq i \leq 3$.

Proof. The estimate (6.28) follows from (6.9) and Lemmas 6.1, 6.2, and 6.3, whereas the second assertion can be established by means of the definition of $\eta_{h,2}^{\text{eq}}$ in (4.32b) and (3.30). □

Remark 6.2. Theorem 6.1 implies the reliability of the residual-type a posteriori error estimator via the reliability of the equilibrated error estimator. The constants in the estimate are specified in the theorem and are computable. In general, the reliability estimate for residual-type error estimators involves interpolation constants which can be computed as well. The number of constants increases with the polynomial degree k . The computations involve eigenvalue-type problems whose solutions may require a substantial amount of computational time (see, for example, [47] for details).

The efficiency of the residual-type error estimator can be established using techniques as in [51] involving suitably chosen bubble functions. The constants in the efficiency estimate depend on the local geometry of the triangulation and can be computed as well.

7 Numerical results

We have implemented the IPDG approximation (3.22) with the penalty parameter α chosen as $\alpha = 12 k^2$. Further, we have implemented the adaptive algorithm based on the equilibrated error estimator η_h^{eq} by Dörfler marking [27], i.e., given a bulk parameter $\theta \in (0, 1)$, we have selected a set $\mathcal{M}_h \subset \mathcal{T}_h$ according to

$$\theta \sum_{K \in \mathcal{T}_h} \eta_K^{\text{eq}} \leq \sum_{K \in \mathcal{M}_h} \eta_K^{\text{eq}}$$

and we have refined elements $K \in \mathcal{M}_h$ by newest vertex bisection. We note that for the evaluation of $\eta_{h,2}^{\text{eq}}$ we have not computed u_h^c by postprocessing according to (3.29) but used the estimate (3.30) instead. In case of the residual-based error estimator η_h^{res} we have implemented the adaptive refinement likewise.

As numerical examples, we have chosen Ω as the L-shaped domain $\Omega := (-1, +1)^2 \setminus ([0, 1] \times (-1, 0])$ with Dirichlet boundary $\Gamma_D = (\{0\} \times (-1, 0)) \cup ((0, 1) \times \{0\})$ and Neumann boundary $\Gamma_N = \partial\Omega \setminus \bar{\Gamma}_D$ with interior angle $\varphi = 3\pi/2$ at the origin. In [26] (see also [5]) it has been shown that in polar coordinates (r, φ) the solution u of the p -Laplace problem (2.1) behaves as $u \sim r^\gamma$ with

$$\gamma = \sigma(p) - \sqrt{\sigma^2(p) - 4/3}, \quad \sigma(p) = \frac{7p - 6}{6(p - 1)}.$$

We have considered the cases $p = 1.5$ and $p = 3.0$ with the solution given by

$$u(r, \varphi) = r^\gamma \sin\left(\frac{2}{3}\varphi\right)$$

and the right-hand side f in (2.1a), the Dirichlet data u_D in (2.1b), and the Neumann data u_N in (2.1c) given accordingly. This results in homogeneous Dirichlet data $u_D = 0$. We note that $u \in W^{1+\gamma-\varepsilon,p}(\Omega)$ for any $\varepsilon > 0$ and u has a singularity at the origin. The same applies to the solution z of the boundary value problem (4.15). However, since Γ_N is located off the singularity at the origin, the trace of ∇z on Γ_N is more regular and the regularity assumption (4.16) is satisfied.

We have performed computations for $p = 1.5$ and $p = 3.0$ and the polynomial degrees $k = 1$ and $k = 3$. The discrete data $f_h|_K$, $K \in \mathcal{T}_h$, and $u_{h,N}|_E$, $E \in \mathcal{E}_h(\Gamma_N)$, have been obtained according to Remark 3.1. Moreover, the numerical solution of the nonlinear IPDG approximation (3.22) has been done by Newton's method with a relative tolerance of $\text{tol} = 10^{-3}$ as termination criterion for the Newton iterates. As outlined in [5], the expected convergence rate for the discretization error in the broken $W^{1,p}$ norm is 0.5.

Figure 1 shows the adaptively generated mesh in case $p = 1.5$ and bulk parameter $\theta = 0.5$ for polynomial degree $k = 1$ (left) and polynomial degree $k = 3$ (right) where the adaptive mesh refinements were based on the equilibrated error estimator. As expected, we observe a pronounced refinement around the reentrant corner and substantially less refinement off the singularity for the higher polynomial degree $k = 3$. The meshes in case $p = 3.0$ and bulk parameter $\theta = 0.5$ for $k = 1$ and $k = 3$ as well as the meshes obtained by the residual-based error estimator look similarly and are therefore omitted.

Figure 2 displays the discretization error in the broken $W^{1,p}$ norm, the equilibrated error estimator η_h^{eq} , and the residual-based error estimator η_h^{res} as a function of the total number of degrees of freedom (DOFs) on a logarithmic scale. As expected by the theory, the convergence rate appears to be 0.5. The equilibrated error estimator is smaller than the residual-based error estimator by approximately 1/2 of an order of magnitude. Figure 3 shows the corresponding results for the polynomial degree $k = 3$. Here, we observe that in a pre-asymptotic phase the error decays faster, but asymptotically approaches the predicted convergence rate of 0.5.

Figures 4 and 5 contain the corresponding results for $p = 3.0$, $\theta = 0.5$, and the polynomial degrees $k = 1$ and $k = 3$. We observe a similar behavior as for $p = 1.5$.

Remark 7.1. As far as the robustness of the a priori error estimators is concerned, in our numerical examples the equilibrated error estimator turned out to be robust with respect to the termination criterion for the approximate solution by Newton's method and the choice of the penalty parameter α . In contrast, the




Fig. 1: $p = 1.5, \theta = 0.5$: Adaptively generated mesh (equilibrated error estimator) for polynomial degree $k = 1$ (left) and $k = 3$ (right).




Fig. 2: $p = 1.5, \theta = 0.5$, polynomial degree $k = 1$: The error in the broken $W^{1,p}$ norm (black), the equilibrated error estimator η_h^{eq} (red), and the residual-based error estimator η_h^{res} (blue).




Fig. 3: $p = 1.5, \theta = 0.5$, polynomial degree $k = 3$: The error in the broken $W^{1,p}$ norm (black), the equilibrated error estimator η_h^{eq} (red), and the residual-based error estimator η_h^{res} (blue).




Fig. 4: $p = 3.0, \theta = 0.5$, polynomial degree $k = 1$: The error in the broken $W^{1,p}$ norm (black), the equilibrated error estimator η_h^{eq} (red), and the residual-based error estimator η_h^{res} (blue).




Fig. 5: $p = 3.0, \theta = 0.5$, polynomial degree $k = 3$: The error in the broken $W^{1,p}$ norm (black), the equilibrated error estimator η_h^{eq} (red), and the residual-based error estimator η_h^{res} (blue).

residual-type estimator has shown less robustness with the efficiency index (estimated error versus true error) increasing slightly with a reduction of the tolerance in the termination criterion and for choosing larger penalty parameters.

Funding: The work has been supported by the NSF grant DMS-1520886.

References

- [1] C. Amrouche and N. E. H. Seloula, L^p -theory for vector potentials and Sobolev's inequalities for vector fields: Application to the Stokes equations with pressure boundary conditions. *Math. Models Meth. Appl. Sci.* **23** (2013), 37–92.
- [2] T. Ando, Contractive projections in L^p spaces. *Pacific J. Math.* **17** (1966), 391–405.
- [3] D. Arnold, F. Brezzi, B. Cockburn, and D. Marini, Unified analysis of discontinuous Galerkin methods for elliptic problems. *SIAM J. Numer. Anal.* **39** (2002), 1749–1779.
- [4] J. W. Barrett and W. B. Liu, Finite element approximation of the p -Laplacian. *Math. Comput.* **61** (1993), 523–537.
- [5] L. Belenki, L. Diening, and C. Kreuzer, Optimality of an adaptive finite element method for the p -Laplacian equation. *IMA J. Numer. Anal.* **32** (2012), 484–510.
- [6] C. Bi and Y. Lin, Discontinuous Galerkin method for monotone nonlinear elliptic problems. *Int. J. Numer. Anal. Mod.* **4** (2012), 999–1024.
- [7] M. Bildhauer and S. Repin, Estimates for the deviation from exact solutions of variational problems with power growth functionals. *Zapiski Nauchnykh Seminar POMI* **336** (2006), 5–24.
- [8] D. Braess, *Finite Elements, Theory, Fast Solvers and Applications in Solid Mechanics*. 3rd edition. Cambridge University Press, Cambridge, 2007.
- [9] D. Braess, T. Fraunholz, and R. H. W. Hoppe, An equilibrated a posteriori error estimator for the interior penalty discontinuous Galerkin method. *SIAM J. Numer. Anal.* **52** (2014), 2121–2136.
- [10] D. Braess, R. H. W. Hoppe, and C. Linsenmann, A two-energies principle for the biharmonic equation and an a posteriori error estimator for an interior penalty discontinuous Galerkin approximation. *ESAIM: M2AN* **52** (2019), 2479–2504.
- [11] D. Braess, V. Pillwein, and J. Schöberl, Equilibrated residual error estimates are p -robust. *Comp. Meth. Applied Mech. Engrg.* **198** (2009), 1189–1197.
- [12] F. Brezzi and M. Fortin, *Mixed and Hybrid Finite Element Methods*. Springer, Berlin–Heidelberg–New York, 1991.
- [13] A. Buffa and C. Ortner, Compact embeddings of broken Sobolev spaces and applications. *IMA J. Numer. Anal.* **29** (2009), 827–855.
- [14] E. Burman and A. Ern, Discontinuous Galerkin approximations with discrete variational principle for the nonlinear Laplacian. *C. R. Acad. Sci. Paris Ser. I* **346** (2008), 1013–1016.
- [15] R. Bustinza, G. N. Gatica, and B. Cockburn, An a posteriori error estimate for the local discontinuous Galerkin method applied to linear and nonlinear diffusion problems. *J. Sci. Comput.* **22/23** (2005), 147–185.
- [16] Z. Cai and S. Zhang, Flux recovery and a posteriori error estimators: Conforming elements for scalar elliptic equations. *SIAM J. Numer. Anal.* **48** (2010), 578–602.
- [17] Z. Cai and S. Zhang, Robust equilibrated residual error estimator for diffusion problems: conforming elements. *SIAM J. Numer. Anal.* **50** (2012), 151–170.
- [18] C. Carstensen and R. Klose, A posteriori finite element control for the p -Laplace problem. *SIAM J. Sci. Comput.* **25** (2003), 792–814.
- [19] P. G. Ciarlet, *The Finite Element Method for Elliptic Problems*. SIAM, Philadelphia, 2002.
- [20] S. Cochez-Dhondt and S. Nicaise, Equilibrated error estimators for discontinuous Galerkin methods. *Numer. Methods Partial Differ. Equations* **24** (2008), 1236–1252.
- [21] B. Cockburn and J. Shen, A hybridizable discontinuous Galerkin method for the p -Laplacian. *SIAM J. Sci. Comput.* **38** (2016), 545–566.
- [22] L. M. Del Pezzo, A. L. Lombardi, and S. Martinez, Interior Penalty Discontinuous Galerkin FEM for the $p(x)$ -Laplacian. *SIAM J. Numer. Anal.* **50** (2012), 2497–2521.
- [23] L. Diening, D. Kröner, M. Ruzicka, and I. Touloupoulos, A local discontinuous Galerkin approximation for systems with p -structure. *IMA J. Numer. Anal.* **34** (2013), 1447–1488.
- [24] L. Diening and C. Kreuzer, Linear convergence of an adaptive finite element method for the p -Laplacian equation. *SIAM J. Numer. Anal.* **46** (2008), 614–638.
- [25] D. A. DiPietro and A. Ern, *Mathematical Aspects of Discontinuous Galerkin Methods*. Springer, Berlin–Heidelberg–New York, 2012.
- [26] M. Dobrowolski, On finite element methods for nonlinear elliptic problems with corners. In: *Springer Lecture Notes in Mathematics, Vol. 1121*. Springer, Berlin–Heidelberg–New York, 1986, pp. 85–103.
- [27] W. Dörfler, A convergent adaptive algorithm for Poisson's equation. *SIAM J. Numer. Anal.* **33** (1996), 1106–1124.

- [28] M. R. Dostanić, Inequality of Poincaré–Friedrichs on L^p spaces. *Matematički Vesnik* **57** (2005), 11–14.
- [29] I. Ekeland and R. Temam, *Convex Analysis and Variational Problems*. SIAM, Philadelphia, 1999.
- [30] A. Ern and J. L. Guermond, Finite element quasi-interpolation and best approximation. *ESAIM: M2AN* **51** (2017), 1367–1385.
- [31] A. Ern, S. Nicaise, and M. Vohralík, An accurate $H(\text{div})$ flux reconstruction for discontinuous Galerkin approximations of elliptic problems. *C. R. Acad. Sci. Paris Ser. I* **345** (2007), 709–712.
- [32] A. Ern and M. Vohralík, Flux reconstruction and a posteriori error estimation for discontinuous Galerkin methods on general nonmatching grids. *C. R. Acad. Sci. Paris Ser. I* **347** (2009), 4441–4444.
- [33] R. Eymard and C. Guichard, Discontinuous Galerkin gradient discretisations for the approximation of second-order differential operators in divergence form. *Comput. Appl. Math.* **37** (2018), 4023–4054.
- [34] M. Fuchs and S. Repin, Estimates for the deviation from the exact solutions of variational problems modeling certain classes of generalized Newtonian fluids. *Math. Methods Appl. Sci.* **29** (2010), 2225–2244.
- [35] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*. Pitman, Boston, 1985.
- [36] R. H. W. Hoppe, G. Kanschat, and T. Warburton, Convergence analysis of an adaptive interior penalty discontinuous Galerkin method. *SIAM J. Numer. Anal.* **47** (2009), 534–550.
- [37] P. Huston, J. Robson, and E. Süli, Discontinuous Galerkin finite element approximation of quasilinear elliptic boundary value problems I: the scalar case. *IMA J. Numer. Anal.* **25** (2005), 726–749.
- [38] O. A. Karakashian and F. Pascal, A posteriori error estimates for a discontinuous Galerkin approximation of second-order elliptic problems. *SIAM J. Numer. Anal.* **41** (2003), 2374–2399.
- [39] O. A. Karakashian and F. Pascal, Convergence of adaptive discontinuous Galerkin approximations of second-order elliptic problems. *SIAM J. Numer. Anal.* **45** (2007), 641–665.
- [40] R. Lazarov, S. Repin, and S. Tomar, Functional a posteriori error estimates for discontinuous Galerkin approximations of elliptic problems. *Numer. Methods Partial Differ. Equ.* **25** (2009), 952–971.
- [41] W. Liu and N. Yan, Quasi-norm local error estimators for p -Laplacian. *SIAM J. Numer. Anal.* **39** (2001), 100–127.
- [42] W. Liu and N. Yan, On quasi-norm interpolation error estimation and a posteriori error estimates for p -Laplacian. *SIAM J. Numer. Anal.* **40** (2002), 1870–1895.
- [43] C. Ortner and E. Süli, Discontinuous Galerkin finite element approximation of nonlinear second-order elliptic and hyperbolic systems. *SIAM J. Numer. Anal.* **45** (2007), 1370–1395.
- [44] W. Qiu and K. Shi, Analysis on an HDG method for the p -Laplacian equations. *J. Sci. Comp.* **80** (2019), 1019–1032.
- [45] S. I. Repin, A posteriori error estimation for variational problems with power growth functionals based on duality theory. *Zapiski Nauchnykh Seminarov POMI* **249** (1997), 244–255.
- [46] S. I. Repin, A posteriori error estimation for variational problems with uniformly convex functionals. *Math. Comp.* **69** (2000), 481–500.
- [47] S. I. Repin, *A Posteriori Estimates for Partial Differential Equations*. De Gruyter, Berlin, 2008.
- [48] S. I. Repin and S. Tomar, Guaranteed and robust error bounds for nonconforming approximations of elliptic problems. *IMA J. Numer. Anal.* **31** (2011), 597–615.
- [49] L. Tartar, *Introduction to Sobolev Spaces and Interpolation Theory*. Springer, Berlin–Heidelberg–New York, 2007.
- [50] A. Veiser, Convergent adaptive finite elements for the nonlinear Laplacian. *Numer. Math.* **92** (2002), 743–770.
- [51] R. Verfürth, *A Posteriori Error Estimation Techniques for Finite Element Methods*. Oxford University Press, Oxford, 2013.
- [52] M. Vohralík, A posteriori error estimates for efficiency and error control in numerical simulations. *Lecture Notes NM 497*. Université Pierre et Marie Curie, Paris, 2015.